

genome consensus sequence data provides information that represents the dominant virus subtype. It does not provide sufficient information to resolve transmission events particularly for rapidly spreading viruses. However, changes in the composition of minor variants between hosts and the pattern of minor variants fixation during outbreaks, could provide additional high-resolution data on who is infecting whom. The same data could also potentially inform the extent of within-host virus diversity as well as the proportion of diversity that is transmitted between individuals. We have developed a reproducible semi-automated whole genome variant calling pipeline to explore the role of minority variants in resolving transmission patterns and within host viral evolution. The pipeline is available as modular Bash scripts that run on a Linux cluster environment.

#### **A28 Frequent co-infection among human group A rotaviruses in Thailand**

M.A. Rabaa,<sup>1,2</sup> T.T.N. Dung,<sup>1</sup> P.T. Duy,<sup>1</sup> L. Bodhidatta,<sup>3</sup> B. Swierczewski,<sup>4</sup> O.M. Sessions,<sup>4</sup> C.J. Mason,<sup>3</sup> S. Baker,<sup>1,2,5</sup>

<sup>1</sup>The Hospital for Tropical Diseases, Wellcome, Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Vietnam, <sup>2</sup>Centre for Tropical Medicine, Oxford University, Oxford, UK, <sup>3</sup>U.S. Armed Forces Research Institute of Medical Science, Bangkok, Thailand, <sup>4</sup>Program in Emerging Infectious Diseases, Duke-National University of Singapore Graduate Medical School, Singapore, Singapore and <sup>5</sup>The London School of Hygiene and Tropical Medicine, London, UK

Rotavirus (RoV) is a non-enveloped dsRNA virus in the Reoviridae family, with a 18.5-kb genome of 11 segments encoding six structural (VP1–4, VP6 and VP7) and five or six non-structural proteins (NSP1–NSP5/6). Reassortment between human and/or animal RoVs plays an important role in the generation of genetic diversity in these viruses, and is presumed to result from co-infection in human or animal reservoirs. However, co-infection with heterologous RoV has rarely been documented, in part due to inadequate detection methods and a lack of large-scale genomic investigations. Despite the availability of an efficacious vaccine, the burden of rotaviral diarrhea remains high in many developing countries, with rotavirus infection detected in 40–50% of all pediatric patients hospitalized with diarrhea. In addition to its cost, reduced vaccine effectiveness in developing country settings has contributed to its low uptake and the lack of government support for vaccination programs across Southeast Asia. The genetics and dynamics of rotavirus (RoV) have rarely been systematically investigated in these settings. The government of Thailand is preparing to add the rotavirus vaccine to its immunization program but has expressed concern regarding its effectiveness in the population and its long-term impact on rotavirus diversity and disease burden. To investigate the diversity of Group A RoV prior to the initiation of immunization programs, we performed full genome sequencing of 200 RoV from infected children across the country from 2004 to 2010 using novel in-house rotavirus capture and sequencing methods. Whilst the majority of samples (76%) showed infection with RoV of the common G1-P[8]-I1-C1-M1-A1-N1-T1-E1-H1 (Wa-like) genotype constellation (representing VP7-VP4-VP6-VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5, respectively), 41% of all samples additionally showed heterologous rotavirus co-infection, containing one or more segments representing an additional, less common genotype. Co-infection of G1-P[8] Wa-like viruses together with segments representing G9-P[19]-I5 lineages was particularly common and phylogenetic analysis suggests that these lineages may have spread through one region of Thailand for three years via serial co-transmission. G1-P[8]

Wa-like infection with additional G3 or G9 VP7 segments was also common across the country. Despite the high frequency of RoV co-infection in this setting, reassortant viruses were rarely observed (in ~8% of cases), and none of these appeared to represent reassortment among the common co-infecting viral segments detected within this study. The impacts of co-infection and reassortment on transmission fitness, the epidemiology of rotaviral diarrhea, and vaccine efficacy and long-term viral diversity warrant further investigation via full genome sequencing following the large-scale introduction of immunization programs in Southeast Asia.

#### **A29 Wolbachia for dengue control; will dengue viruses evolve resistance?**

T.T. Vi,<sup>1</sup> D.T. Kien,<sup>1</sup> B. Wills,<sup>1,2</sup> C.P. Simmons,<sup>1,2,3</sup>

<sup>1</sup>Oxford University Clinical Research Unit, Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam, <sup>2</sup>Centre for Tropical Medicine & Global Health, Oxford University, Oxford, UK and <sup>3</sup>Dept. of Microbiology and Immunology, University of Melbourne, Parkville, Victoria, Australia

Dengue viruses (DENV) are major human pathogens, transmitted between people mainly by *Aedes aegypti* mosquitoes. Recently, the endosymbiont bacteria *Wolbachia* was horizontally introduced into *Ae. Aegypti*; *Wolbachia* infection non-specifically impairs DENV replication in these mosquitoes opening up a potential new strategy for dengue control. This project will test the hypothesis that DENV cannot easily develop resistance to the mechanisms by which *Wolbachia* blocks virus replication. Using contemporary viruses currently circulating in Vietnam, we will perform 30 cycles of repeated passage of DENV in wild-type and *Wolbachia*-infected *Ae. Aegypti* (wMel strain). This experimental system maximizes the possibility of observing selective pressure by wMel on DENV populations in mosquito tissues. We will use intermittent quantitative virology and virus genome sequencing to detect changes over time in the genotype and phenotype of DENV in *Ae. aegypti*. The results will provide valuable insights into how easily DENV adapt to *Wolbachia* in a controlled system. If *Wolbachia* specific evolution is observed, then we will test the infectiousness and replication profile of such viruses in cultured human cells. The results will deliver the first evidence of the likelihood of DENV “escape” from wMel, with implications for future and ongoing field releases of wMel in dengue-endemic countries.

#### **A30 Recombination & evolution in two viral families: effective steps or a random walk?**

R. Bingham,<sup>1,\*</sup> G. Leonov,<sup>1</sup> N. Patel,<sup>2</sup> E. Weiss,<sup>1</sup> S. White,<sup>2</sup> E. Dykeman,<sup>1</sup> P. Stockley,<sup>2</sup> R. Twarock,<sup>1</sup>

<sup>1</sup>York Centre for Complex Systems Analysis, University of York, York YO10 5GE, UK and <sup>2</sup>Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds LS2 9JT, UK

Recombination or reassortment occurs in many different viral families and is understood as one of the predominant mechanisms of evolution. We perform a broad analysis of the mechanisms and recombinants of two viral families, Picornaviridae and Hepadnaviridae, which exhibit very different viral behaviour. Combining information about recombinants with knowledge of the evolutionary selection pressures on the genome and insights from the conservation of functional aspects of the genome, we find that viruses can dynamically exploit recombination in different ways, either to evade antigens and prolong infection, or to enhance viral diversity and allow the virus to infect new hosts. Better understanding of the mechanisms of

recombination may lead to better ways to inhibit infections and develop new antiviral treatments.

#### A31 A large-scale screening for hepaciviruses in African rodents

M. Bletsa,<sup>1\*</sup> B. Vrancken,<sup>1</sup> S. Gryseels,<sup>2</sup> H. Leirs,<sup>2</sup> J. Göyü de Bellocq,<sup>3</sup> A. Laudisoit,<sup>4</sup> E. Wollants,<sup>1</sup> M. Van Ranst,<sup>1</sup> J.F. Drexler,<sup>5</sup> P. Lemey,<sup>1</sup>

<sup>1</sup>Clinical and Epidemiological Virology, Department of Microbiology and Immunology, KU Leuven, Rega Institute for Medical Research, Leuven, Belgium, <sup>2</sup>Department of Biology, Evolutionary Ecology Group, University of Antwerp, Antwerp, Belgium, <sup>3</sup>Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, Studenec, Czech Republic, <sup>4</sup>Ecology, Evolution and Genomics of Infectious Disease Research Group, Institute of Integrative Biology, University of Liverpool, Liverpool, UK and <sup>5</sup>Institute of Virology, University of Bonn Medical Centre, Bonn, Germany

Hepatitis C virus (HCV) is considered to be a major public health problem, infecting >3% of the human population and causing acute and chronic liver disease. To date, the zoonotic origins of HCV remain elusive, since no animal population has been identified with closely related hepaciviruses. Inspired by recent findings of divergent hepaciviruses in rodents, we have screened a comprehensive set of African rodent samples that have been collected through various collaborations. Screening was performed by employing a highly sensitive nested PCR assay directed against the NS3 protease-helicase gene. From the 2,361 samples that were screened, 74 were found positive for the presence of hepaciviruses. At this stage, we focus on generating longer stretches of genomic information based on previously described protocols that make use of host rRNA depletion and a simple viral enrichment methodology followed by NGS approaches. By generating and analyzing sequence data from these samples we aim to perform in-depth phylogenetic and phylogeographic analyses gaining valuable insight into the evolutionary origins and epidemic emergence of HCV. Emphasis will be put on the identification of novel hepacivirus lineages more closely related to HCV, as well as the examination of host-specific adaptation and geographic structuring of these viruses.

#### A32 Search for viral integration insertion sites into the human genome—strategy matters

T.M. Zorec,<sup>1</sup> L. Hošnjak,<sup>1</sup> B. Kocjan,<sup>1</sup> N. Gale,<sup>2</sup> N. Zidar,<sup>2</sup> P. Strojjan,<sup>3</sup> M. Poljak,<sup>1</sup>

<sup>1</sup>Institute of Microbiology and Immunology, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia, <sup>2</sup>Institute of Pathology, Faculty of Medicine, University of Ljubljana, Slovenia and <sup>3</sup>Department of Radiation Oncology, Institute of Oncology, Ljubljana, Slovenia

During one of our previous works we investigated the integration of a low-risk human papillomavirus' (HPV) genetic material into the human genome, we aimed to elucidate whether the integration event(s) had helped facilitate the manifestation of cancer in an individual with a HPV11-positive sinonasal squamous cell carcinoma. To elucidate viral integration sites, whole-genome sequencing (WGS) was applied to the tissue sample of sinonasal carcinoma. We employed three different analysis strategies – data pipelines with different topologies that should intuitively all yield qualitatively the same result with minor variations. One of the pipeline topologies was adopted after Li et al. (2013) with slight modifications, whereas the other two were simple meta-algorithms that should generally answer the same question. Although the bulk of the breakpoints yielded through the three different topologies did overlap to some degree, a great deal of breakpoints varied greatly between the approaches.

#### A33 The cervico-vaginale microbiota in chlamydia trachomatis notified women: a case-control study at the sexually transmitted infection outpatient clinic in Amsterdam

Charlotte van der Veer,<sup>1</sup> S.M. Bruisten,<sup>1</sup> J.J. van der Helm,<sup>2</sup> H.J.C. de Vries,<sup>2,3</sup> R. van Houdt,<sup>4</sup>

<sup>1</sup>Public Health Laboratory, Public Health Service Amsterdam, Amsterdam, The Netherlands, <sup>2</sup>Public Health Service Amsterdam, Sexually transmitted infections outpatient clinic, Amsterdam, The Netherlands, <sup>3</sup>Department of Dermatology, Amsterdam Medical Centre, University of Amsterdam, Amsterdam, The Netherlands and <sup>4</sup>Department of Medical Microbiology and Infection Prevention, VU University Medical Centre, Amsterdam, The Netherlands

Increasing evidence suggests that the cervico-vaginal microbiome (CVM) plays an important role in acquiring sexually transmitted infections (STIs). Here we studied the CVM in women exposed to *Chlamydia trachomatis* (Ct). We included 98 women who were notified by Ct-positive sex partners via contact-tracing at the STI outpatient clinic in Amsterdam, the Netherlands. Cervico-vaginal samples and clinical data were collected for all women. CVM compositions were characterized by sequencing of the V3/V4 region of the 16srRNA gene using the Illumina MiSeq platform. High quality reads were assigned to operational taxonomic units and classified using a vaginal reference package. Hierarchical clustering delineated CVM clusters based on microbial relative abundances. Possible determinants for acquiring Ct were analyzed using multivariable logistic regression. The CVM was characterized for 93 women, of whom 52 were Ct positive and 41 Ct negative. We identified three major CVM clusters. Clustered CVM predominantly comprised either diverse anaerobic bacteria ( $n=39$ ; 42%), *Lactobacillus iners* ( $n=32$ ; 34%) or *Lactobacillus crispatus* ( $n=22$ ; 24%). In multivariable analysis, we found that the CVM was significantly associated with *C. trachomatis* infection (OR = 4.2 (95% confidence interval, CI: 1.2–15.4) for women with diverse anaerobic CVM and OR = 4.4 (CI: 1.3–15.6), for women with *L. iners*-dominated CVM, compared to women with *L. crispatus*-dominated CVM), as was younger age (OR = 3.1, CI: 1.1–8.7, for those  $\leq 21$  years old) and reporting a steady sex partner (OR = 3.6, CI: 1.4–9.4). Women who tested positive for *Chlamydia trachomatis* infection after having been contact-traced by a chlamydia positive partner were more likely to have CVM dominated by *L. iners* or by diverse anaerobic bacteria, than by *L. crispatus*.

#### A34 Automated profiling of the human virome from raw metagenomic data

Yumna Moosa,<sup>1,2</sup> Michael Vilsker,<sup>3</sup> Ewout Vanden Eyden,<sup>4</sup> Vagner Fonseca,<sup>5</sup> Sam Nooij,<sup>6</sup> Koen Deforche,<sup>3</sup> Tulio de Oliveira,<sup>1,2</sup>

<sup>1</sup>Nelson R Mandela School of Medicine College of Health Sciences, University of KwaZulu-Natal (UKZN), South Africa, <sup>2</sup>Centre for the AIDS Programme of Research in South Africa (CAPRISA), UKZN, South Africa, <sup>3</sup>Emweb, Herent, Belgium, <sup>4</sup>Rega Instituut, KU Leuven, Belgium, <sup>5</sup>Universidade do Estado da Bahia (UNEB), Brazil and <sup>6</sup>Rijksinstituut voor Volksgezondheid en Milieu (RIVM), The Netherlands

Viruses influence human health as conventional pathogens, as modulators of gene expression and through their involvement in complex host-microbiome interactions. Next generation sequencing (NGS) has enabled us to explore the role of the microbiome in human health and disease. Metagenomic sequencing should allow us to profile all biological elements in a clinical sample in an unbiased, hypothesis-free way. However viruses display much greater variation than all other elements, and the existing tools and methods for virus identification and