

# Subtrochanteric fractures after long-term treatment with bisphosphonates: a European Society on Clinical and Economic Aspects of Osteoporosis and Osteoarthritis, and International Osteoporosis Foundation Working Group Report

R. Rizzoli • K. Åkesson • M. Bouxsein • J. A. Kanis •  
N. Napoli • S. Papapoulos • J.-Y. Reginster • C. Cooper

Received: 10 August 2010 / Accepted: 30 September 2010 / Published online: 18 November 2010  
© The Author(s) 2010. This article is published with open access at Springerlink.com

## Abstract

**Summary** This paper reviews the evidence for an association between atypical subtrochanteric fractures and long-term bisphosphonate use. Clinical case reports/reviews and case-control studies report this association, but retrospective phase III trial analyses show no increased risk. Bisphosphonate use may be associated with atypical subtrochanteric fractures, but the case is yet unproven.

**Introduction** A Working Group of the European Society on Clinical and Economic Aspects of Osteoporosis and Osteoarthritis and the International Osteoporosis Foundation has reviewed the evidence for a causal association between subtrochanteric fractures and long-term treatment with bisphosphonates, with the aim of identifying areas for further research and providing recommendations for physicians.

R. Rizzoli (✉)  
Division of Bone Diseases,  
Department of Rehabilitation and Geriatrics,  
University Hospitals and Faculty of Medicine of Geneva,  
1211 Geneva 14, Switzerland  
e-mail: Rene.Rizzoli@unige.ch

K. Åkesson  
Department of Orthopedics Malmö, Skåne University Hospital,  
Lund University,  
Lund, Sweden

M. Bouxsein  
Department of Orthopaedic Surgery,  
Harvard Medical School Center for Advanced Orthopaedic Studies,  
Beth Israel Deaconess Medical Center,  
Boston, MA, USA

J. A. Kanis  
WHO Collaborating Centre for Metabolic Bone Diseases,  
University of Sheffield Medical School,  
Sheffield, UK

N. Napoli  
Division of Endocrinology,  
Universita' Campus Bio-Medico di Roma,  
Rome, Italy

S. Papapoulos  
Department of Endocrinology and Metabolic Diseases,  
Leiden University Medical Center,  
Leiden, The Netherlands

J.-Y. Reginster  
Department of Public Health Sciences  
and Department of Physical Medicine  
and Rehabilitation, University of Liège,  
CHU Sart-Tilman and BRULL,  
Liège, Belgium

C. Cooper  
MRC Epidemiology Resource Centre,  
University of Southampton,  
Southampton, UK

C. Cooper  
NIHR Musculoskeletal Biomedical Research Unit,  
University of Oxford,  
Oxford, UK

**Methods** A PubMed search of literature from 1994 to May 2010 was performed using key search terms, and articles pertinent to subtrochanteric fractures following bisphosphonate use were analysed.

**Results** Several clinical case reports and case reviews report a possible association between atypical fractures at the subtrochanteric region of the femur in bisphosphonate-treated patients. Common features of these ‘atypical’ fractures include prodromal pain, occurrence with minimal/no trauma, a thickened diaphyseal cortex and transverse fracture pattern. Some small case–control studies report the same association, but a large register-based study and retrospective analyses of phase III trials of bisphosphonates do not show an increased risk of subtrochanteric fractures with bisphosphonate use. The number of atypical subtrochanteric fractures in association with bisphosphonates is an estimated one per 1,000 per year. It is recommended that physicians remain vigilant in assessing their patients treated with bisphosphonates for the treatment or prevention of osteoporosis and advise patients of the potential risks.

**Conclusions** Bisphosphonate use may be associated with atypical subtrochanteric fractures, but the case is unproven and requires further research. Were the case to be proven, the risk–benefit ratio still remains favourable for use of bisphosphonates to prevent fractures.

**Keywords** Atypical · Bisphosphonate · Femur · Low trauma · Osteoporosis · Subtrochanteric

## Introduction

Treatment with bisphosphonates significantly reduces the risk of fractures in men and women with osteoporosis. The evidence is based on high-quality phase III randomized controlled trials (RCTs) with fracture as an endpoint [1–10]. The benefits of bisphosphonates also extend to other disorders of bone metabolism such as glucocorticoid-induced osteoporosis [11], Paget’s disease [12] and bone metastases [13, 14].

Treatment with bisphosphonates is not without adverse effects, but they are generally minor and occur in a minority of patients. The most common adverse effect is gastrointestinal upset with the oral formulations, the frequency of which decreases with intermittent treatment such as once-weekly or monthly regimens. Intravenous (IV) administration of nitrogen-containing bisphosphonates may induce an acute phase reaction which manifests as fever, myalgia and arthralgia, although these side effects usually resolve within a few days of onset [3, 7, 15]. High doses of bisphosphonates given intravenously may impair renal function, and the kidney is a major route of elimination of the

bisphosphonates. For this reason, bisphosphonates are not recommended for use in patients with severe renal impairment [16–18]. The use of bisphosphonates has been associated with osteonecrosis of the jaw, but most cases have occurred in patients receiving high-dose IV bisphosphonates for neoplastic bone disease, and osteonecrosis of the jaw has rarely been reported in patients with benign bone diseases [19, 20]. An increased risk of atrial fibrillation has been reported for zoledronic acid [3], but the association may be coincidental [7]. Other uncommon or rare side effects of bisphosphonates include anaemia [21], urticaria [22, 23] and symptomatic hypocalcaemia [22].

In recent years, several clinical case reports and case reviews have reported an association between atypical fractures in patients receiving treatment with bisphosphonates. The majority of these cases have described fractures at the subtrochanteric region of the femur [24–31].

Against this background, the aim of this report was to critically review the evidence for an increased incidence of subtrochanteric fractures after long-term treatment with bisphosphonates, to identify gaps in our knowledge that warrant further research and to provide guidance for healthcare professionals. A PubMed search of literature from 1994 to May 2010 was performed using the search terms ‘bisphosphonate(s)’ AND/OR ‘alendronate’ AND/OR ‘risedronate’ AND/OR ‘ibandronate/ibandronic acid’ AND/OR ‘zoledronate/zoledronic acid’ AND/OR ‘subtrochanter(ic)’ AND ‘fracture’ AND/OR ‘femur/femoral’ AND/OR ‘atypical’ AND/OR ‘low-trauma’ AND/OR ‘low-energy’. Scientific papers pertinent to subtrochanteric fractures following bisphosphonate use were analysed and included in the evidence base.

## Characteristics of subtrochanteric fractures

Subtrochanteric fractures have been defined as occurring in a zone extending from the lesser trochanter to 5 cm distal to the lesser trochanter [32]. However, this anatomical classification of subtrochanteric fracture has several variations [33, 34], resulting in variable definitions in published studies [26, 30, 35].

Regardless of the definition used, many case reports and case reviews have suggested that there are several common features of subtrochanteric fractures associated with bisphosphonate use. Major features were that the fractures arose with minimal or no trauma and, on radiography, the fracture line was transverse. Minor features were that fractures were commonly preceded by prodromal pain and, on radiographs, there appeared beaking of the cortex on one side and bilateral thickened diaphyseal cortices [26, 28, 36–39]. This fracture pattern has often been referred to as an ‘atypical subtrochanteric fracture’ [40–42] although,

as reviewed below, the distinction between typical and atypical subtrochanteric fractures has not yet been firmly established.

It is worth noting that, on radiography, the appearance of atypical subtrochanteric fractures is similar to that of stress fractures, including a periosteal reaction, linear areas of bone sclerosis and a transverse fracture line. Prodromal pain prior to diagnosis is also common [43]. However, stress fractures are more commonly associated with repeated episodes of increased activity (e.g. participation in sports). Nevertheless, stress fractures due to non-traumatic loading in patients with low bone density do occur (insufficiency fractures) [43]; thus, the terms ‘atypical’ and ‘stress’ are often used interchangeably or in conjunction to describe subtrochanteric fractures in this context. However, this terminology also requires clarification, as not all stress fractures are atypical.

### Epidemiology of subtrochanteric fractures

Subtrochanteric fractures are a relatively rare type of hip fracture [44–46], usually resulting from high-energy trauma, pathologic fracture or, in the elderly, low-energy injury involving osteoporotic bone. Several series report the incidence of this fracture [25–28, 30, 36, 37, 47], although the definition of the subtrochanteric site has varied. Nieves et al. reported a large, 11-year epidemiological study of fractures of the hip, subtrochanter, femoral shaft and distal femur in the US population aged  $\geq 50$  years using National Hospital Discharge Survey data from the National Center for Health Statistics and MarketScan® (medical claims experience) data [46]. Of all femoral fractures, 3% were at the subtrochanteric region, 5% at the femoral shaft, 5% at the distal femur and 87% were at the proximal femur (i.e. hip). Importantly, this study classified fractures solely according to their location in the femur and did not evaluate the fracture patterns radiographically. Thus, they were not

able to determine the incidence of ‘typical’ vs ‘atypical’ subtrochanteric fractures. In men and in women, the incidence rate of each type of fracture remained stable over 5 years but increased exponentially with age (Fig. 1). Each fracture type was more prevalent in women than in men. Seventy-five percent of all femur fracture cases were in women. The mean age at fracture was 80 years old, and those with a subtrochanteric fracture were of a similar age to those with a hip fracture.

Leung et al. published a retrospective analysis that aimed to document the incidence of low-trauma subtrochanteric or femoral diaphyseal fractures in a Hong Kong hospital over a 5-year period [42]. In all, 88 cases of subtrochanteric fractures and 66 of diaphyseal fractures were identified, accounting for 3.9% and 2.9% of all recorded osteoporotic fractures, respectively.

Thus, although the incidence of subtrochanteric fractures is much lower than other femoral fractures, they are not rare and account for about 3% of all femoral fractures in the elderly. If these estimates were applied to the UK, then more than 2,000 subtrochanteric fractures are expected to occur each year [48], and approximately 48,000 are expected annually worldwide [49].


### Subtrochanteric fractures and bisphosphonate exposure

#### Case reports and case reviews

Twenty-six published case reports and case reviews were identified that were considered relevant, a similar number to that identified in a recent review by Giusti et al. [50].

Concern about bisphosphonate use in relation to atypical subtrochanteric fractures arose from case reports that described patients with subtrochanteric fractures who had been exposed to bisphosphonates, particularly long-term treatment with alendronate (Fosamax®/Fosavance®, alendronate sodium, Merck Sharp & Dohme Limited). The

**Fig. 1** Age-specific incidence of femoral fractures according to fracture site in men (X) and women (O) aged  $\geq 50$  years (adapted from Nieves et al. [46])



association between long-term bisphosphonate use and unusual diaphyseal fractures was first described by Odvina et al. in 2005 [31] who reported nine patients with osteoporosis or osteopenia who had been treated with alendronate for 3–8 years and sustained atraumatic fractures in the course of their normal daily activities. Three patients had fractures of the femoral shaft and two had fractures of the proximal femur. Of these five patients, fracture healing was radiographically assessed in four. All four patients had delayed or absent fracture healing ranging from 4 months to 2 years while on alendronate treatment.

This and subsequent case reports are summarized in Table 1. The mean and median age of patients was 65 years (range 35–85). All cases involved treatment with alendronate, except for five patients who took risedronate (Actonel®, risedronate sodium, Procter and Gamble Pharmaceuticals) and three who took pamidronate (Aredia®, pamidronate disodium, Novartis Pharmaceuticals Limited). One patient had been taking ibandronate (Bonviva®/Boniva®, ibandronic acid, Roche) for 1 year following long-term alendronate use, and one had been taking risedronate for 5 years following 7 years of pamidronate use. There were no published case reports of subtrochanteric fractures following the use of once-yearly zoledronic acid 5 mg (Aclasta®/Reclast®, zoledronic acid anhydrous, Novartis Pharmaceuticals Limited), although cases following treatment with the monthly 4-mg dose have been reported [36, 38]. The mean and median duration of bisphosphonate use was 7.3 and 7.5 years, respectively (range 1–16), and the majority of patients had unilateral fractures (29 out of 43; 67.4%).

In addition to case reports, several case reviews have been published, which are summarized in Table 2. For example, the characteristics of low-trauma subtrochanteric and diaphyseal fractures were studied retrospectively by Neviaser et al. in all patients admitted to a US trauma centre over a 5-year interval (Table 2) [30]. Radiographs were examined by independent experts to identify fractures with a simple, transverse or short oblique pattern in areas of cortical hypertrophy with a cortical beak. The observers were blinded to patient characteristics, including alendronate use. Seventy patients were identified, of whom 25 were treated with alendronate. Nineteen out of 25 (76%) alendronate-treated patients had the radiographic pattern compared with one out of 45 (2%) non-alendronate-treated patients. Thus, the risk of having an ‘atypical’ subtrochanteric fracture pattern was significantly associated with alendronate use (odds ratio=139; 95% confidence interval (CI) 19–939;  $p<0.0001$ ). The mean duration of treatment with alendronate was 6.2 years (6.9 years in those who had the fracture pattern vs 2.5 years in those who did not) [30]. The authors concluded that there are unique features to bisphosphonate-associated fractures.

## Controlled studies

Six studies that utilized control groups were identified that have investigated the association of subtrochanteric fractures with the use of bisphosphonates. In the study of Nieves et al. described above, the rate of subtrochanteric and femoral shaft fractures appeared to be higher than that of other fractures in women taking oral bisphosphonates (Fig. 2) [46], although there is no statistical information provided. It is not known whether excess fractures were due to trauma or not. The study concluded, however, that there was no evidence of an increase in the incidence of subtrochanteric or femoral shaft fracture between 1996 (around the time that bisphosphonates were first introduced) and 2006. Limitations of these data include the lack of radiological and clinical verification and no information on the type of bisphosphonate used or the duration of treatment.

In a study by Leung et al., ten patients with subtrochanteric fractures who had received alendronate were identified over a 5-year period. This included one patient who had taken alendronate for 1 year followed by ibandronate for 2 years [42]. The crude incidence of subtrochanteric/femoral diaphyseal fractures associated with prior bisphosphonate use increased over 5 years from 0% in 2003/2004 to 6% in 2004/2005, 8.6% in 2006/2007 and 25% in 2007/2008. This trend was despite a steady annual incidence of subtrochanteric/femoral diaphyseal fractures. It is difficult to draw meaningful conclusions from these data because of the very small sample size (ten subtrochanteric fractures in patients exposed to a bisphosphonate) and the lack of information on bisphosphonate use at other fracture sites. At best, the study documents the increasing use of bisphosphonates over the time of study.

In a small retrospective case–control study, Lenart et al. aimed to identify an association between low-energy subtrochanteric/femoral shaft fractures (according to the Müller AO classification) and long-term bisphosphonate use [29]. Forty-one low-energy subtrochanteric or femoral shaft fracture cases were identified and matched by age, body mass index and race to one low-energy intertrochanteric and femoral neck fracture each.

Fifteen out of the 41 (37%) cases of subtrochanteric or femoral shaft fracture cases were taking bisphosphonates, compared with nine out of 82 (11%) controls (OR=4.4; 95% CI 1.8–11.4;  $p=0.002$ ). Alendronate was the bisphosphonate taken in all cases. Eight out of nine cases in the control group were taking alendronate (one had previously taken etidronate). A radiographic pattern of a simple transverse or oblique fracture, beaking of the cortex on one side and cortical thickening at the fracture site, was observed in ten of the 15 (67%) subtrochanteric/femoral shaft fracture cases taking bisphosphonate and three of the

**Table 1** Case reports of incidents of subtrochanteric fracture following bisphosphonate use (all cases in women unless otherwise indicated)

| Reference                        | Total patients (patients ST/FS/PF fracture) | Age (years)     | Location                                                   | Radiographic features                                                          | Bilateral? | Prodromal symptoms (duration)                            | Osteoporosis diagnosis?             | Prior bisphosphonate | Duration of use (years) | Concomitant therapy                           | Healing (months of follow-up)              |
|----------------------------------|---------------------------------------------|-----------------|------------------------------------------------------------|--------------------------------------------------------------------------------|------------|----------------------------------------------------------|-------------------------------------|----------------------|-------------------------|-----------------------------------------------|--------------------------------------------|
| Odvina et al. [31]               | 9 (5)                                       | 52              | Femoral shaft                                              |                                                                                | No         |                                                          | No (osteopenia)                     | ALN                  | 8                       | Ca, D                                         | No (9)                                     |
|                                  |                                             | 68 <sup>a</sup> | Femoral shaft                                              |                                                                                | Yes        |                                                          | Yes                                 | ALN                  | 8                       | Ca, D                                         | No (8)                                     |
|                                  |                                             | 67              | Femoral shaft                                              |                                                                                | Yes        |                                                          | No (osteopenia)                     | ALN                  | 5                       | Oestrogen, Ca, D                              | Yes (5)                                    |
|                                  |                                             | 49              | Proximal femur                                             |                                                                                | No         |                                                          | Yes (GIO)                           | ALN                  | 3                       | Pred, Ca, D                                   | No (8)                                     |
|                                  |                                             | 64              | Proximal femur                                             |                                                                                | No         |                                                          | Yes (GIO)                           | ALN                  | 4                       | Pred, Ca, D                                   | Yes (3)                                    |
| Husada et al. [51]               | 1                                           | 72              | Femoral shaft                                              | Cortical thickening in lateral mid-shaft of contralateral femur                | Yes        | Severe pain in back and hip (1 month)                    | Yes                                 | ALN                  | Not specified           | Ca, amlodipine, metoprolol, aspirin           |                                            |
| Schneider [52]                   | 1                                           | 59              | Upper femur                                                | Cortical thickening                                                            | No         | Moderate pain in thigh (3 months)                        | No (family history of osteoporosis) | ALN                  | 7                       | Ca, hormone replacement therapy               | Yes (>9)                                   |
| Arriamanto-Villareal et al. [53] | 1                                           | 35 <sup>a</sup> | Subtrochanteric femur                                      |                                                                                | No         |                                                          |                                     | ALN                  | 6                       | Ca                                            | No (36) <sup>c</sup>                       |
| Cheung et al. [54]               | 1                                           | 82              | Femoral shaft                                              |                                                                                | No         |                                                          | Yes                                 | ALN                  | 10                      | Ca, glucosamine, chondroitin                  |                                            |
| Demiralp et al. [55]             | 1                                           | 65              | Femoral shaft                                              | Fracture line, callus, cortical thickening, bowing deformity                   | Yes        | Incapacitating bilateral femoral shaft pain (1.5 months) | Yes                                 | ALN                  | 7                       | Ca, D, steroid, thyroxine replacement therapy |                                            |
| Lee et al. [56]                  | 1                                           | 73              | Femoral diaphysis                                          |                                                                                | No         | Bilateral groin pain, difficulty walking (10 months)     | Yes                                 | ALN                  | 1.5                     |                                               | Yes                                        |
| Sayed-Noor and Sjoden [57]       | 1                                           | 72              | Subtrochanteric femur                                      | Cortical thickening of lateral femoral cortex, medial beaking at fracture site | No         | Diffuse pain in hips and thighs (18 months)              | Yes                                 | ALN                  | 7                       | Ca                                            | No (3)yes (6)                              |
| Visserkrana et al. [39]          | 3                                           | 51              | Femoral metadiaphysis                                      |                                                                                | Yes        | Bilateral, lateral hip pain                              |                                     | ALN                  | 5                       | Pred                                          | No (3 while on ALN; 12 after stopping ALN) |
| Odvina et al. [58]               | 13 (11)                                     | 62              | Femoral metadiaphysis                                      |                                                                                | Yes        | Bilateral thigh pain                                     |                                     | ALN                  | 10                      | Raloxifene, pred                              | Yes (12) <sup>d</sup>                      |
|                                  |                                             | 75              | Femoral metadiaphysis                                      |                                                                                | No         |                                                          |                                     | ALN                  | 10                      | Pred                                          | No (22)                                    |
|                                  |                                             | 57              | Subtrochanteric, contralateral femur shaft (3 years later) | Cortical thickening                                                            | Yes        | Pain at fracture site (1–6 months)                       | No (osteopenia)                     | ALN                  | 6                       | Ca, D                                         | Yes (36)                                   |
|                                  |                                             | 74              | Femoral shaft                                              | Cortical thickening                                                            | No         |                                                          | Yes                                 | ALN                  | 10                      | Ca, D                                         | No                                         |

**Table 1** (continued)

| Reference                  | Total patients (patients ST/FS/PF fracture) | Age (years) | Location                                   | Radiographic features                                                                  | Bilateral? | Prodromal symptoms (duration)                                 | Osteoporosis diagnosis? | Prior bisphosphonate | Duration of use (years) | Concomitant therapy | Healing (months of follow-up) |
|----------------------------|---------------------------------------------|-------------|--------------------------------------------|----------------------------------------------------------------------------------------|------------|---------------------------------------------------------------|-------------------------|----------------------|-------------------------|---------------------|-------------------------------|
|                            |                                             | 67          | Femoral shaft                              | Cortical thickening                                                                    | No         | Pain at fracture site (1–6 months)                            | Yes                     | RIS                  | >5                      | Ca, D               | Yes (6)                       |
|                            |                                             | 58          | Femoral shaft (fractured twice in 3 years) | Cortical thickening                                                                    | No         | Pain at fracture site (1–6 months)                            | No                      | ALN                  | 7                       | Ca, D, tamoxifen    | Yes (6)                       |
|                            |                                             | 62          | Femoral shaft                              | Cortical thickening                                                                    | No         |                                                               | No (osteopenia)         | RIS                  | 2                       | Ca, D, tamoxifen    |                               |
|                            |                                             | 63          | Femoral shaft                              | Cortical thickening                                                                    | No         |                                                               | Yes                     | ALN                  | 10                      | Ca, D, oestrogen    | Yes (6)                       |
|                            |                                             | 72          | Femoral shaft                              | Cortical thickening                                                                    | No         | Pain at fracture site (1–6 months)                            | Yes                     | ALN                  | 9                       | Ca, D, oestrogen    | Yes                           |
|                            |                                             | 76          | Femoral shaft                              | Cortical thickening                                                                    | No         |                                                               | Yes (GIO)               | ALN                  | 11                      | Ca, D, pred         | Yes (12)                      |
|                            |                                             | 72          | Left and right femoral shaft               | Cortical thickening                                                                    | Yes        | Pain at fracture site (1–6 months)                            | Yes (GIO)               | ALN                  | 10                      | Ca, D, pred         | Yes                           |
|                            |                                             | 77          | Femoral shaft                              | Cortical thickening                                                                    | No         |                                                               | Yes (GIO)               | ALN                  | 9                       | Ca, D, pred         | Yes                           |
|                            |                                             | 38          | Left and right femoral shaft               | Cortical thickening                                                                    | Yes        |                                                               | Yes (GIO)               | ALN                  | 3                       | Ca, D, pred         | Yes                           |
| Ali and Jay [59]           | 1                                           | 82          | Femoral shaft                              | Cortical thickening                                                                    | No         |                                                               |                         | ALN                  | 8                       |                     | Yes (3)                       |
| Goddard et al. [60]        | 1                                           | 67          | Femoral diaphysis                          | Cortical thickening, uncortical beaking                                                | No         |                                                               |                         | ALN                  | 16                      |                     | Yes (12)                      |
| Sayed-Noor and Sjoden [61] | 2                                           | 78          | Tip of femoral stem                        | Cortical thickening                                                                    | No         |                                                               | Yes                     | ALN                  | 9                       |                     | No (6)                        |
|                            |                                             | 55          | Subtrochanteric femur                      | Cortical thickening, medial beaking, cortical thickening on contralateral femur        | No         | Diffuse pain in thighs, walking difficulties (several months) | Yes                     | ALN                  | 9                       | D                   | Yes (9)                       |
| Cermak et al. [62]         | 4                                           | 64          | Subtrochanteric femur                      | Cortical thickening                                                                    | No         | Pain in left thigh (3 months)                                 | No                      | ALN                  | 5.5                     |                     | Yes (6)                       |
|                            |                                             | 70          | Femur                                      | Medial cortical beaking                                                                | Yes        | Pain in thighs                                                | Yes                     | ALN                  | 6                       |                     | Yes (4)                       |
|                            |                                             |             | Other femur (3 months later)               |                                                                                        |            |                                                               |                         |                      |                         |                     | Yes (7)                       |
|                            |                                             | 77          | Femoral shaft                              | Cortical thickening                                                                    | No         | Pain in right thigh                                           | Yes                     | ALN                  | 12                      |                     | Yes (12)                      |
|                            |                                             | 59          | Subtrochanteric femur                      | Cortical thickening, medial cortical beaking                                           | No         | None                                                          | Yes                     | ALN                  | 10                      |                     | Yes (5)                       |
| Bush and Chew [63]         | 1                                           | 85          | Subtrochanteric femur                      | Focal beak of cortical thickening of lateral cortex                                    | No         | Limp, persistent pain in anterior thigh (2–3 months)          | Yes (GIO)               | RIS                  | >6                      | Ca, D, pred         | Yes (2)                       |
| Lee [64]                   | 1                                           | 82          | Left femoral diaphysis                     | Horizontal fracture lines at thickest part of femoral cortex extending lateral–medial, | Yes        |                                                               | Yes                     | ALN                  | 8                       | Ca, D               | Yes (5)                       |

|                     |    |    |                                                                                                 |                                                                                                 |                                                                           |           |                                  |     |                                                                                           |          |
|---------------------|----|----|-------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|-----------|----------------------------------|-----|-------------------------------------------------------------------------------------------|----------|
| Edwards et al. [65] | 1  | 60 | Right femoral diaphysis<br>Left femoral diaphysis (2 years later)                               | Yes<br>(Taken after initial, right fracture)<br>Minor lateral cortical thickening on left femur | Mild pain in right thigh before right fracture, none before left fracture | Yes (GIO) | ALN                              | 8   | Pred                                                                                      | No (6)   |
| Giusti et al. [50]  | 8  | 60 | Right subtrochanteric femur<br>Left subtrochanteric femur (9 months later)                      | Yes                                                                                             | Pain in right hip                                                         | No        | ALN                              | 4   | Ca, D, pred, inhaled GCs, esomeprazole, repaglinide, metformin, azathioprine, rosvastatin | Yes      |
|                     | 36 |    | Femoral shaft                                                                                   | No                                                                                              |                                                                           | Yes       | ALN                              | 8   | D, pred, simvastatin, cyclosporine, amlodipine, atenolol, lisinopril                      | Yes      |
|                     | 64 |    | Left and right subtrochanteric femur (1 complete, 1 insufficiency fracture)                     | Yes                                                                                             | Pain in right thigh                                                       | No        | ALN                              | 2.5 | Ca, D, pred, omeprazole, azathioprine, losartan, triamteren, HCT                          | No (18)  |
|                     | 62 |    | Right and left femoral shaft <sup>b</sup>                                                       | Yes                                                                                             | Pain in right thigh and hip                                               | Yes       | Oral pamidronate                 | 4   | Ca, D,                                                                                    | Yes      |
|                     | 58 |    | Femoral shaft                                                                                   | No                                                                                              | Pain in left thigh                                                        | Yes       | Intravenous pamidronate          | 3   | Ca, D                                                                                     | No (12)  |
|                     | 58 |    | Subtrochanteric femur                                                                           | No                                                                                              | Pain in left hip                                                          | No        | RIS                              | 5.5 | Ca, D, pred, inhaled GCs, omeprazole, pravastatin, ibuprofen                              | No (12)  |
|                     | 72 |    | Left subtrochanteric femur<br>Right subtrochanteric femur (insufficiency fracture 1 year later) | Yes                                                                                             | Pain in left thigh and hip                                                | Yes (GIO) | Oral pamidronate followed by ALN | 7+5 | Ca, D, inhaled GCs, esomeprazole, simvastatin, captopril, irbesartan, clopidogrel         | Yes (12) |
|                     | 75 |    | Femoral shaft (insufficiency fracture)<br>Femoral shaft (insufficiency fracture 1 year later)   |                                                                                                 | Severe pain in left thigh and hip<br>Pain in hip                          | Yes       | RIS                              | 6   | Ca, D, esomeprazole, etoricoxib                                                           |          |

ALN alendronate, BP bisphosphonate, Ca calcium, D vitamin D, FS femoral shaft, GCs glucocorticoids, GIO glucocorticoid-induced osteoporosis, HCT hydrochlorothiazide, PF proximal femur, RIS risedronate, ST subtrochanteric, Pred prednisone

<sup>a</sup> Male patient

<sup>b</sup> First fracture prior to BP treatment; contralateral fracture following 4 years' BP treatment; refracture of contralateral femoral shaft 4 years after second fracture

<sup>c</sup> Patient was prescribed alendronate in 1996 and took it for 6 years. Fracture occurred 1 year after discontinuation and had not completely healed when reported in 2006

<sup>d</sup> Patient began teriparatide immediately after fracture

26 (11%) subtrochanteric/femoral shaft fracture cases not taking bisphosphonate (OR=15.3; 95% CI=3.1–76.9;  $p<0.001$ ). The duration of bisphosphonate exposure was significantly longer in patients with this X-ray pattern [29].

Koh et al. carried out a retrospective clinical and radiological review of geriatric hip fracture patients at a Singapore tertiary centre over 4 years to assess features that predispose to complete stress fractures [38]. Thirty-two patients with spontaneous or low-energy fractures with metaphyseal–diaphyseal involvement and on bisphosphonate therapy were identified. All were on alendronate therapy except for one who was on monthly zoledronic acid 4 mg and one who had been on risedronate for 6 years following 4 years of alendronate. Of these, 16 patients (median duration of therapy 4.5 years) had radiographic evidence of lateral cortical thickening. Four had cortical stress lesions on the prefracture radiograph (group F) and 12 had cortical stress lesions on the contralateral femur (group C). The type of bisphosphonate taken by patients according to group was not detailed. All patients in group F experienced prodromal thigh discomfort, compared with 25% of patients in group C ( $p=0.019$ ), and radiographic evidence of a stress line across the cortical thickening occurred in 100% and 8% of patients, respectively ( $p=0.003$ ). At a median follow-up of 23 months, none of the patients in group C had developed a complete fracture. All of these patients except for one had discontinued bisphosphonate therapy; five had not taken any alternative therapy since discontinuation. Nevertheless, eight out of the 11 were asymptomatic, and no new cortical thickening was detected in any of the patients. The authors concluded that, in people taking long-term bisphosphonate therapy, symptomatic cortical stress reactions accompanied by evidence of a stress line across the cortical thickening suggest an increased risk of a complete stress fracture [38].

In the only population-based study that included radiological review of all cases, Schilcher and Aspenberg studied the incidence of stress fractures at the femoral shaft in bisphosphonate-treated patients in four hospitals in Sweden. Women aged over 55 years with fractures of the femoral diaphysis or subtrochanteric region were identified from the operation registry. Preoperative radiographs were examined to identify stress fractures, defined as a transverse fracture of the femoral shaft with cortical thickening. Of 91,956 women identified, 3,087 bisphosphonate users were identified, of whom five had femoral stress fractures. All of these five patients were aged >75 years, and their mean duration of treatment was 5.8 years [66]. Three patients that were not treated with bisphosphonates had stress fractures. All were aged <75 years. The annual incidence of femoral shaft stress fractures in bisphosphonate users was 1/1,000 per year (95% CI 0.3–2) vs 0.02/1,000 (0.004–0.1) per year in control patients. Thus, the risk of such fractures was

estimated to be 46 times greater with bisphosphonate use (95% CI 11–200) [65]. An obvious weakness of the study is that, although the confidence intervals were corrected for sample size, the findings were based on just eight femoral shaft stress fractures. The results thus raise a hypothesis to be tested on larger samples.

A larger study is provided by Abrahamsen et al. who studied the epidemiology of subtrochanteric and diaphyseal femur fractures in patients in Denmark treated with alendronate [67]. However, in contrast to the Schilcher and Aspenberg report, in this study, radiographic fracture patterns were not reviewed, and thus, fractures were identified purely based on their location. In patients aged  $\geq 60$  years that had subtrochanteric, diaphyseal femur and hip fractures in 2005, the incidence of subtrochanteric ( $n=898$ ) and diaphyseal fractures ( $n=720$ ) were similar, and the ratio of high-to-low-energy trauma fractures was the same for each of these fracture types (approximately 2.5:1 for each). Exposure to alendronate was also similar between fracture types (approximately 7% each). Patients with subtrochanteric fractures and diaphyseal fractures were more likely to have taken glucocorticoids in the year before fracture than patients with hip fracture (10.9%, 8.4% and 6.5% of patients, respectively).

In a register-based matched cohort analysis, Abrahamsen et al. investigated whether the increase in risk of ‘atypical’ femur fracture in alendronate-treated patients was greater than the increase in risk of ‘typical’ osteoporotic femur fractures (‘typical’ and ‘atypical’ were not defined). In total, 15,187 patients who took alendronate for  $\geq 6$  months after the fracture event (the treatment cohort) were compared with two randomly assigned sex-, age- and fracture-matched controls ( $n=10,374$ ). The use of alendronate was associated with an increase in the hazard ratio (HR; adjusted for baseline comorbidities) for both subtrochanteric/diaphyseal fractures (HR=1.46; 95% CI 0.91–2.35;  $p=0.12$ ) and hip fracture (HR=1.45; 95% CI 1.21–1.74;  $p<0.001$ ). Subtrochanteric/diaphyseal fractures were equally common in the alendronate-treated (14% of hip fractures) and control patients (13%;  $p=0.70$ ). Both hip fractures and subtrochanteric/diaphyseal fractures were significantly lower in patients with higher adherence (HR=0.47 [0.34–0.65;  $p<0.001$ ] and 0.28 [0.12–0.63;  $p<0.01$ ], respectively). In a sub-analysis of 178 compliant (medication possession ratio >80%) patients who took alendronate for >6 years, long-term alendronate use was associated with no change in both hip (HR=1.24 [0.66–2.34];  $p=0.52$ ) and subtrochanteric/diaphyseal fractures (HR=1.37 [0.22–8.62];  $p=0.74$ ). The incidence of subtrochanteric/diaphyseal fractures was similar in the long-term alendronate (10%) and control (12.5%) groups (10% vs 12.5%, respectively) [67].

This study, in a large number of patients, does not support the hypothesis that exposure to alendronate is

**Table 2** Case reviews of incidents of subtrochanteric fracture following bisphosphonate use (all cases in women unless otherwise indicated)

| Reference                              | Review location/<br>period                              | Inclusion criteria                                                                                         | Patients<br>eligible (n)       | Mean age<br>(years<br>[range]) | Fracture location                                                                                                                                                              | Radiographic<br>features (n)                                                | Bilateral? (n)                                                                                              | Prodromal<br>symptoms<br>(duration)                                  | OP<br>diagnosis?<br>(n)          | Prior BP (duration<br>of use, years)                                                                                                                                                        | Concomitant<br>therapy (n)                                                                                                                                |
|----------------------------------------|---------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------|--------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------|----------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Goh et al. [26]                        | 2 Singapore<br>hospitals/May<br>2005–February<br>2006   | ST fracture <sup>a</sup> due<br>to low-energy<br>trauma                                                    | 13<br>ALN (9)                  | 66.9 (55–82)                   | NA                                                                                                                                                                             | Cortical thickening<br>(6=lateral, 3=<br>contralateral)                     | NR                                                                                                          | 5 pts<br>(2–6<br>months)                                             | Yes (3)<br>No (4)<br>Unknown (2) | ALN (4.2 [2.5–5])                                                                                                                                                                           | Ca (all); long-<br>term oral<br>steroids (1)                                                                                                              |
| Kwek et al. [28]                       | Singapore<br>hospital/May<br>2005–January<br>2007       | ST fracture <sup>b</sup> due<br>to low-energy<br>trauma in patients<br>taking ALN                          | No ALN (4)<br>17               | 80.3 (64–92)<br>66 (53–82)     | NA<br>NA                                                                                                                                                                       | Lateral cortical<br>thickening, medial<br>cortical<br>beaking (all)         | NR<br>ST stress<br>fracture (2)<br>Femoral shaft<br>stress<br>fracture (1)<br>Femoral shaft<br>fracture (1) | None<br>Yes, 13 pts<br>(1 week–<br>24 years)                         | Yes (all)<br>Yes (10)<br>No (6)  | NA<br>ALN (4.4 [2–8])<br>RIS after 4 years<br>on ALN                                                                                                                                        | Ca (2)<br>Ca (all); long-<br>term<br>prednisolone (1)                                                                                                     |
| Neviaser<br>et al. [30]                | US trauma<br>centre/January<br>2002–March<br>2007       | Low-energy ST<br>and mid-shaft<br>femur fractures <sup>c</sup>                                             | 70 (11 male)                   | 74.7                           | ST femur (50)<br>Femoral shaft<br>(20)                                                                                                                                         | Lateral cortical<br>thickening,<br>uncortical<br>beaking (20) <sup>d</sup>  | NR                                                                                                          | NR                                                                   | Yes (31) <sup>e</sup>            | ALN (6.2 [1–10])<br>[25 pts] <sup>f</sup>                                                                                                                                                   | NR                                                                                                                                                        |
| Glennon [47]                           | Australian<br>tertiary<br>hospital,<br>12 months        | ST stress fracture<br>with characteristic<br>radiological/<br>clinical features                            | 6                              | 60–87                          | NA                                                                                                                                                                             | Transverse fracture,<br>uncortical<br>beaking, cortical<br>thickening (all) | 1 patient                                                                                                   | Pain in 5 pts<br>(1 week to<br>6 months)                             | NR                               | ALN (1.5–16)<br>[5 pts]<br>RIS (>3) [1 pt]                                                                                                                                                  | NR                                                                                                                                                        |
| Ing-Lorenzini<br>et al. [27]           | Swiss university<br>hospital/<br>2 years                | Low-energy ST<br>fracture, history<br>of BP use                                                            | 8 (7 females)                  | 67.5                           | ST femur (7)<br>Femoral shaft<br>(1)                                                                                                                                           | Cortical thickening,<br>also in<br>contralateral<br>femur in 4 patients     | Yes, in 4 patients<br>0.5–5 years<br>after<br>first fracture                                                | Pain in 2 pts,<br>1 lateral<br>side, 1 both<br>sides                 | Yes (all; 1 pt GIO)              | ALN alone<br>(1.5–8) [3 pts]<br>ALN (3–10)<br>switched<br>to ibandronate<br>(1 NK) <sup>g</sup> [3 pts]<br>RIS (NK) switched<br>to ALN (2) [1 pt]<br>Pamidronate (5) <sup>h</sup><br>[1 pt] | Ca (all),<br>glucocorticoids<br>(4), proton-<br>pump<br>inhibitors (7)                                                                                    |
| Armamento-<br>Villareal<br>et al. [25] | US medical<br>school/<br>November<br>2004–March<br>2007 | Low-energy<br>fracture,<br>mainly at cortical<br>sites, 2 years <sup>i</sup><br>BP therapy,<br>bone biopsy | 15 (12<br>females,<br>3 males) | 43–75                          | Femoral shaft<br>(7) [1 male]                                                                                                                                                  | Cortical thickening,<br>medial beaking<br>(all)                             | Yes (2)                                                                                                     | NR                                                                   | NR                               | ALN (4–10) [6 pts]                                                                                                                                                                          | Ca (6); vitamin D (6);<br>infliximab (1);<br>triamcinolone (1);<br>tamoxifen (1);<br>levothyroxine (1);<br>fluticasone (1);<br>HCT (1);<br>mometazone (1) |
| Capecci and<br>Tejwani [37]            | US university<br>hospital/4 years                       | Bilateral low-energy<br>femoral<br>diaphyseal<br>or ST fracture,<br>long-term ALN                          | 7                              | 61 (53–75)                     | Other (9)<br>Simultaneous<br>femoral<br>diaphysis (1)<br>Sequential ST<br>femur (2)<br>ST and impending<br>contralateral ST<br>femur (3)<br>Femoral diaphysis<br>and impending | Cortical thickening,<br>medial beaking<br>(all)                             | Yes (all)                                                                                                   | Thigh pain<br>(4 pts<br>with<br>impending<br>ST stress<br>fractures) | NR                               | RIS (2) [1 pt]<br>ALN (8.6 [5–13])                                                                                                                                                          | None affecting<br>bone<br>metabolism                                                                                                                      |

**Table 2** (continued)

| Reference              | Review location/<br>period            | Inclusion criteria                                | Patients<br>eligible (n) | Mean age<br>(years<br>[range]) | Fracture location                                                                                         | Radiographic<br>features (n)      | Bilateral? (n) | Prodromal<br>symptoms<br>(duration)                          | OP<br>diagnosis?<br>(n) | Prior BP (duration<br>of use, years)                                                                                  | Concomitant<br>therapy (n) |
|------------------------|---------------------------------------|---------------------------------------------------|--------------------------|--------------------------------|-----------------------------------------------------------------------------------------------------------|-----------------------------------|----------------|--------------------------------------------------------------|-------------------------|-----------------------------------------------------------------------------------------------------------------------|----------------------------|
| Bunning<br>et al. [36] | US rehabilitation<br>hospital/7 years | Atypical low- or<br>no-impact femoral<br>fracture | 4 (1 male)               | 49–59                          | contralateral ST<br>femur (1)<br>Diaphyseal<br>femoral (3);<br>left ST/right<br>diaphyseal<br>femoral (1) | Medial cortical<br>thickening (1) | 1 pt           | Pain in hip<br>(1–3 months)<br>[all], pain in<br>knee [1 pt] | Yes (all)               | None [1 patient]<br>Pamidronate (0.5)<br>zoledronic acid<br>4 mg (>4.5)<br>[1 pt]<br>ALN (5) [1 pt]<br>ALN (6) [1 pt] | NR                         |

ALN alendronate, BP bisphosphonate, Ca calcium, GIO glucocorticoid-induced osteoporosis, HCT hydrochlorothiazide, NA not applicable (described in inclusion criteria), NK not known, NR not reported, OP osteoporosis, Pt patient, RIS risedronate, ST subtrochanteric

<sup>a</sup> In the region of the femur which extended from the lesser trochanter to the junction of the proximal and middle third of the femoral shaft

<sup>b</sup> Within the region of the femur 5 cm distal to the lesser trochanter

<sup>c</sup> Muller AO classification type 32 and type 31 A3 fractures involving or extending distally to the lesser trochanter

<sup>d</sup> Nineteen had been treated with alendronate


<sup>e</sup> Twenty-one had been treated with alendronate

<sup>f</sup> All females. Eighteen cases confirmed through physician/patient contact. Duration of use established in 16 cases

<sup>g</sup> One patient had been on ibandronate for 1 year. One switched to ibandronate 4 months before first fracture in February 2006; one switched 1 year before second fracture in Jan 2008

<sup>h</sup> Stopped 1 year before fracture

**Fig. 2** Medical and prescription drug history in US female fracture patients (2002–2006) during the 1 year before index date (adapted from Nieves et al. [46])



associated with an increased frequency of subtrochanteric fractures compared with controls. However, the same study reported that treatment with alendronate was associated with an increased risk of hip fracture. This should not be interpreted as ‘alendronate causes hip fracture’, but only that high-risk patients are exposed to alendronate. The finding also illustrates the difficulties in the interpretation of retrospective observational studies, particularly accounting for selection bias that likely confounds the other much smaller observational studies.

#### Randomized controlled trials

Black et al. recently reported an analysis of subtrochanteric and diaphyseal fractures in the Fracture Intervention Trial (FIT) of alendronate and its extension [1, 2, 5, 68] and the HORIZON Pivotal Fracture Trial (PFT) of zoledronic acid 5 mg [3]. Twelve fractures in ten patients were documented in the subtrochanteric or diaphyseal region (Table 3) a combined rate of 2.3 per 10,000 patient-years [69]. However, radiographs were not available to confirm typical vs atypical radiographic features. There was no significant increase over placebo in the risk of subtrochanteric/diaphyseal fractures during the FIT, FIT Long-Term Extension (FLEX) or HORIZON-PFT trials. Compared with placebo, the relative hazard was 1.03 (95% CI 0.1–16.5) for alendronate use in the FIT trial, 1.5 (95% CI 0.3–9.0) for zoledronic acid in the HORIZON-PFT and 1.3 (95% CI 0.1–14.7) for continued alendronate use in the FLEX trial. The interpretation of this analysis is limited by the small number of events and the large confidence intervals.

Bilezikian et al. reported the incidence of subtrochanteric fractures in the randomized, placebo-controlled phase III studies of risendronate in post-menopausal osteoporosis, which enrolled more than 15,000 patients. In trials of up to 3 years duration, the mean incidence of subtrochanteric fractures was 0.14% in risendronate 2.5-mg treated patients ( $n=4,998$ ), 0.13% in risendronate 5-mg treated patients ( $n=5,395$ ) and 0.17% in placebo-treated patients ( $n=5,363$ ) [70]. In active control studies of risendronate involving various doses (35 mg once weekly, 75 mg on two consecutive days per month, 150 mg once monthly), no subtrochanteric fractures were reported, and the incidence of hip/femoral fractures was similar to that in the placebo-controlled studies [70].

The manufacturers of ibandronate have assessed their clinical trials database to determine the incidence of subtrochanteric and diaphyseal femoral fractures in women taking ibandronate for post-menopausal osteoporosis. Atypical fractures were defined as ‘mostly non-spine fractures including hip or femur fractures in the subtrochanteric region or shaft and occurring without trauma or in association with low-energy trauma’. For femur fractures, subtrochanteric fracture location was considered as atypical for osteoporosis-related fractures, defined as a region below the lesser trochanter and a junction between the proximal and middle third of the femoral shaft. In the pivotal trials (MF 4380, BONE, MOBILE and DIVA) [4, 71–73], there were nine fracture cases corresponding to these defined locations and characteristics (subtrochanteric, femoral shaft, stress or multiple fractures): six occurred in placebo-treated patients ( $n=1,924$ ) and three in ibandronate-treated patients ( $n=6,830$ ). In addition, there was one identified case of a

femoral shaft fracture in an ibandronate-treated patient in the extension and major phase IIIb trials (MOBILE LTE, DIVA LTE, MOTION and PREVENTION;  $n=2,451$ ) [74–77]. Some fractures were reported without identifying the precise location. However, all of these fractures were associated with trauma and thus did not meet the definition for atypical fractures. An additional 5-year analysis of the marketed regimens of ibandronate (150 mg once monthly and 3 mg IV quarterly) was also carried out from the active comparator-controlled trials and their extensions (MOBILE, DIVA, MOTION, MOBILE LTE and DIVA LTE) [71, 72, 74, 75, 77]. No atypical subtrochanteric/diaphyseal femoral fractures were found for either of the marketed regimens (150 mg,  $n=1,279$ ; 3 mg,  $n=469$ ).

#### Pharmacovigilance data

Since fractures are the clinical outcome of osteoporosis and no treatments are fully effective, fractures are expected in treated patients. It is likely, however, that the number of reports through pharmacovigilance will be small. The number of postmarketing reports of atypical stress fractures in association with alendronate to circa July 2008 was 115 (of which 84 were femur fractures) and included a large number of the cases reported in the literature [78].

Bilezikian et al. have reported that in more than 10 years of risedronate post-approval surveillance to September

2008 (18 million patient-years of exposure), the reporting rate for subtrochanteric fractures was  $<0.1$  per 100,000 patient treatment years of exposure [70].

Postmarketing data from the manufacturers of zoledronic acid have revealed a similarly low rate of subtrochanteric fractures with zoledronic acid 5 mg. Using the last cutoff date for worldwide Health Authority Reporting prior to January 2010 (Periodic Safety Update Report v6) and assessing all adverse event reports for zoledronic acid 5 mg (579,501 patient-years of exposure), the rate of femoral subtrochanteric fracture reporting was three per 1,000,000 patient treatment years of exposure.

Postmarketing data from the manufacturers of ibandronate have also revealed a low rate of possible atypical fractures occurring in patients receiving ibandronate for the management of postmenopausal osteoporosis. According to their global safety database as of June 2009, cumulative postmarketing exposure of ibandronate yielded a crude reporting rate of possible atypical fractures of approximately one per 1,000,000 patients. Three of the cases involved alendronate treatment followed by ibandronate treatment and were reported in the case series of Ing-Lorenzini et al. [27].

#### Regulatory perspective

In July 2008, the Pharmacovigilance Working Party (PhVWP) of the Committee for Medicinal Products for

**Table 3** Characteristics of ten patients with 12 low-trauma subtrochanteric or femoral diaphyseal fractures in the FIT, FLEX and HORIZON-PFT trials (adapted from Black et al. [69])

| Study   | Age (years) | Study medication              | Time from randomization to fracture (days [years]) | Bilateral? | Prodromal symptoms | Compliance                            | Concomitant therapy                                               |
|---------|-------------|-------------------------------|----------------------------------------------------|------------|--------------------|---------------------------------------|-------------------------------------------------------------------|
| FIT     | 75          | Placebo                       | 962 (2.6)                                          |            |                    | >75%                                  | None                                                              |
| FIT     | 69          | Alendronate                   | 1,682 (4.6)                                        |            |                    | >75%                                  | None                                                              |
| FLEX    | 79          | Alendronate (first fracture)  | 1,250 (3.4)                                        |            |                    | Stopped 3 years before first fracture | Alendronate, 6 years (in FIT before FLEX)                         |
|         |             | Alendronate (second fracture) | 1,369 (3.8)                                        |            |                    |                                       |                                                                   |
| FLEX    | 80          | Alendronate/ placebo          | 1,257 (3.4)                                        |            |                    | Stopped 3 years before fracture       | Alendronate, 6 years (in FIT before FLEX)                         |
| FLEX    | 83          | Alendronate/ alendronate      | 1,006 (2.8)                                        |            |                    | >75%                                  | Alendronate, 5 years (in FIT before FLEX)                         |
| HORIZON | 65          | Zoledronic acid               | 454 (1.2)                                          |            | Hip pain           | 100%                                  | Raloxifene                                                        |
| HORIZON | 78          | Placebo                       | 1,051 (2.9)                                        |            | Hip pain           | 100%                                  | None                                                              |
| HORIZON | 65          | Zoledronic acid               | 732 (2.0)                                          |            |                    | 100%                                  | None                                                              |
| HORIZON | 72          | Placebo                       | 321 (0.9)                                          |            |                    | 100%                                  | Calcitonin                                                        |
| HORIZON | 71          | Zoledronic acid (2 fractures) | 934 (2.6)                                          | Yes        | Bone pain          | 100%                                  | Bisphosphonate and hormone replacement therapy, both before study |

Human Use (CHMP) initiated a class review on bisphosphonates and atypical stress fractures. Marketing Authorization Holders supplied information about all preclinical, clinical and future studies, published case reports, post-marketing data, possible mechanisms and proposed risk-minimization activities. Following a PhVWP review of these data in December 2008, the CHMP concluded that there was an association between atypical stress fractures and long-term use of alendronate, due to the distinct fracture pattern, prodromal pain and poor fracture healing. However, the benefit–risk balance of alendronate use was considered favourable. The CHMP highlighted that there was uncertainty concerning a class effect for other bisphosphonates and that switching of bisphosphonates should be avoided at this time. Ultimately, the CHMP recommended that information about atypical stress fractures should be added to the product information for medicinal products containing alendronate [78]. Consequently, the labelling for alendronate (Fosamax®/Fosavance®, Merck Sharp & Dohme Limited) now includes a special warning/precaution for alendronate use, advising discontinuation of bisphosphonate therapy in patients with stress fracture pending evaluation, based on an individual benefit–risk assessment [22, 79]. Alendronate is the only bisphosphonate for osteoporosis treatment that currently carries this warning.

In addition to the 2008 class review, the EMEA released a statement in August 2009 highlighting their 2010 priorities for drug safety research with regards to the long-term adverse skeletal effects of bisphosphonates: (1) generate methodologies to study the link between bisphosphonate use and long-term adverse skeletal events in human populations and (2) measure the incidence of stress/insufficiency fractures in association with high-dose/long-term use of bisphosphonates by class, compound, mode of administration, dose etc. Methods could include meta-analysis or nested case–control studies [80].

In June 2008, the US Food and Drug Administration (FDA) initiated a review of bisphosphonates for a possible association with increased risk of atypical subtrochanteric femur fractures. All available case reports and clinical trial data were requested from all bisphosphonate drug manufacturers and were reviewed alongside the registry data from the large observational study of Abrahamsen et al. [67]. In March 2010, the FDA announced that the data reviewed had not shown a clear connection between bisphosphonate use and the risk of atypical subtrochanteric fractures. Physicians were urged to continue to follow the labelling when prescribing bisphosphonates and patients were instructed not to discontinue their medication unless instructed to do so by their physician [81].

Pathophysiology of subtrochanteric fractures associated with bisphosphonate use

The pathophysiology of atypical low-trauma subtrochanteric fractures following bisphosphonate use is not known. However, preclinical and clinical studies of the effects of bisphosphonates on bone suggest that there are several possible mechanisms that work either alone or in tandem. The organic matrix of the bone determines its toughness, and this matrix is partly made up of bone collagen, which impacts on the bone's mechanical properties. Bisphosphonate use may negatively affect collagen by preventing or reducing its maturation [82], although this finding has not been consistently replicated [83]. Bisphosphonates may also affect bone mineralization density distribution (BMDD). The more heterogeneous the BMDD, the slower that cracks in the bone will develop and the lower the risk of new cracks and fractures forming [84]. As bisphosphonate treatment reduces bone turnover, the increase in overall mineralization leads to more homogeneous bone—as evidenced by a narrow BMDD [85, 86]—and thus an increased risk of cracks and fractures. Reduced bone turnover also increases the accumulation of microdamage, as cracks are not repaired [87], and reduces bone toughness, which contributes to the increased susceptibility of bone to new cracks [88–90]. Finally, bisphosphonates have differing impacts on different types of fracture. Acute fractures of long bone are not affected by bisphosphonates in the initial healing stages [91–93], as they heal via endochondral ossification. However, stress fractures heal by normal bone remodelling, and thus, bisphosphonates may prevent or delay healing, increasing the likelihood of a complete fracture with little or no trauma. Several reports have reported on bone quality in people with low-trauma fractures taking bisphosphonate therapy.

For example, Odvina et al. reported that cancellous bone histomorphometry in alendronate-treated patients (3–8 years) who sustained spontaneous non-vertebral fractures showed markedly suppressed bone formation, with reduced or absent osteoblastic surface in most patients. Osteoclastic surface was also low in most patients, and eroded surface decreased in half [31]. Odvina et al. reported similar findings in a later report in a comparable patient population [58]. In a case report by Armamento-Villareal et al. of a man who had a low-trauma subtrochanteric fracture after discontinuing 6 years of alendronate treatment, a bone biopsy showed severely decreased trabecular connectivity, a lack of osteoid on trabecular surfaces and an absence of tetracycline labelling [53]. Armamento-Villareal et al. later reported that of 15 bisphosphonate-treated patients (2–10 years; Table 2) who underwent bone biopsies following a low-energy cortical (femoral shaft, pelvis, rib, metatarsal, ankle) fracture, ten had an absence of double-tetracycline

label, reduced osteoid surface and thickness suggestive of suppressed trabecular bone remodelling. However, there was no difference in cortical thickness between patients with suppressed ( $n=10$ ) and normal ( $n=5$ ) turnover [25]. Recent findings by Somford et al., however, suggest an alternative pathophysiology for subtrochanteric fractures associated with bisphosphonate treatment. In a patient who was treated with alendronate for 8 years and subsequently developed spontaneous bilateral subtrochanteric/diaphyseal fractures, biopsies showed a marked decrease in bone formation as expected; however, this was not coupled with the expected decrease in bone resorption. In fact, bone resorption parameters such as osteoclast number were markedly increased in the femur sample. In addition, there was no evidence of hypermineralized bone. This suggests that an imbalance between bone resorption and bone formation at the affected femur—the cause of which is currently unknown—rather than excessive suppression of bone turnover may be the underlying mechanism for subtrochanteric/diaphyseal femoral fractures in bisphosphonate-treated patients [94].

### Summary of evidence

The view that bisphosphonates increase the risk of subtrochanteric femoral fractures arises from the case reports and retrospective case reviews that have reported ‘atypical’ subtrochanteric and diaphyseal fractures in patients exposed to bisphosphonates. In all, these data highlight the scope of the problem, i.e. a trend that warrants further investigation. However, the data in their entirety are insufficient proof that long-term bisphosphonate use is the only cause of atypical low-trauma subtrochanteric fractures.

There are several limitations to the existing evidence base: lack of a consensus definition of an atypical subtrochanteric fracture, small numbers of patients involved, lack of radiographs which precludes characterization of the radiographic features of the fractures and incomplete reporting of subject characteristics. In addition, subtrochanteric fractures in general are not atypical fractures; rather, they are part of the natural history of fragility fractures in osteoporosis. They increase in frequency with age in much the same way as does the incidence of other osteoporotic fractures [95]. Although their incidence is much lower than other femoral fractures, they are not rare and account for approximately 3% of all femoral fractures [46]. Thus, the term ‘atypical’ is not synonymous with ‘unexpected’ which is the common interpretation. Rather, the term should be reserved for subtrochanteric fractures that have atypical features, of which some are similar to with those associated with stress.

Therein lies an additional problem in that it has been difficult to provide characteristics of the fracture that are associated with the use of bisphosphonates. Candidate features, which include the prodromal manifestation of incomplete (fissure) fractures, a thickened cortex and a transverse fracture pattern with cortical beaking may be associated with the use of bisphosphonates but, in the absence of blinded evaluation in all cases, may be subject to large observer biases. In addition, in many instances, cases have been complicated, for example, by concomitant exposure to glucocorticoids [25–28, 31, 39, 50, 55, 58, 63, 65], which appears to be a risk factor for subtrochanteric fractures [46].

In terms of evidence-based medicine, the ultimate arbiter for a causal relationship between subtrochanteric fractures and exposure to bisphosphonates might be expected to derive from information from RCTs. All the information available fail to show an association of this fracture with exposure to bisphosphonates, although all RCTs were completed before attention was drawn to the problem, so the documentation of the sites of fracture and any associated features is inevitably incomplete. Furthermore, the frequency of the event is sufficiently low that even large RCTs are insufficiently powered to identify meaningful associations with drug exposure. Finally, the duration of exposure to bisphosphonates may be too short in the setting of RCTs if, as has been suggested, the complication were to increase in frequency with exposure time. Against this background, data from observational studies might be expected to contribute to our understanding, but such studies are fraught with biases and limitations for which it may be difficult to adjust.

### Research agenda

The ultimate question for physicians is what type of patient is at the highest risk of an atypical low-trauma subtrochanteric fracture. Thus far, apart from long-term alendronate therapy, suggested risk factors include glucocorticoid, proton-pump inhibitor or calcitonin use and female gender [26, 46, 67]. Thus, a number of urgent issues and areas for research have been identified as follows:

1. Standardized definition of ‘subtrochanteric fracture’, including a definition of ‘atypical’ and ‘typical’ fractures
2. Provision of descriptive epidemiology based on large-scale studies with characterization of radiographic features
3. Definition of fracture incidence by femoral location, mechanism of injury and underlying pathology
4. Identification of risk factors, with greater clarity as to the precise risk factors in patients taking bisphosphonates

5. Pathophysiological studies in relation to risk factors
6. Pathophysiological studies at the tissue level, i.e. is the mechanism of atraumatic (insufficiency) fractures different to that of low-trauma fractures?
7. Long-term, large, prospective, observational studies to assess incidence of subtrochanteric fractures in bisphosphonate-treated vs bisphosphonate-naïve patients. Methods should include (1) futility analysis and (2) radiographic measurements. Outcomes should include (1) adherence, (2) number needed to harm and (3) assessment of temporal relationship between bisphosphonate treatment and fracture type
8. Long-term, large, prospective, observational studies allowing for systematic follow-up of patients with subtrochanteric fractures treated long-term with bisphosphonates, in order to assess fracture healing characteristics (e.g. time to healing, choice of fracture treatment device, adjuvant bone anabolic intervention etc.)
9. Large, prospective, randomized, controlled clinical trials of the efficacy and safety of pharmacological treatment (e.g. strontium ranelate, teriparatide) for patients with subtrochanteric fractures

## Conclusions and recommendations

A sense of proportion may be helpful in alleviating the concerns of the medical community. A plausible scenario is that long-term exposure to bisphosphonates (more than 5 years) increases the risk of subtrochanteric femoral fractures twofold. In the UK, using the guidance of the National Osteoporosis Guideline Group, the relative risk of hip fracture is expected to be approximately threefold increased in postmenopausal women identified for treatment [96]. Assuming that the average population risk of hip fracture is 1% per year in postmenopausal women, then 300 hip fractures are expected for every 10,000 patients identified to be at high risk. If these patients were treated and assuming an effectiveness of bisphosphonates of 36% (RR=0.64) [97], then 108 hip fractures are averted by treatment (and approximately 750 fractures at other sites). On the debit side, three subtrochanteric fractures (both typical and atypical) are to be expected, which might increase to six if bisphosphonates doubled the risk of all subtrochanteric fractures. Under the assumptions of this scenario, the risk–benefit ratio remains very favourable.

Evidence, including that from an EMEA class review, suggests that alendronate use may potentially increase the risk for atypical, low-trauma subtrochanteric fractures, although it is unclear whether this applies to other

bisphosphonates. Irrespective of exposure to bisphosphonates, the occurrence of subtrochanteric fractures is an expected finding in patients with osteoporosis. If atypical fractures do occur, then their characteristics are poorly defined, their causality with bisphosphonate exposure insecure and their frequency rare. Bisphosphonates as the cause of atypical fractures at the subtrochanteric site is therefore still merely a hypothesis, though no less important for that. Clearly more research is required from well-designed prospective observational studies, meta-analyses and nested case–control studies.

Thus, the available evidence does not suggest that the well-known benefits of bisphosphonate treatment are outweighed by the risk of these rare, atypical, low-trauma subtrochanteric fractures. Nevertheless, it is recommended that physicians remain vigilant in assessing their patients treated with bisphosphonates for osteoporosis or associated conditions. They should continue to follow the recommendations on the drug label when prescribing bisphosphonates and advise patients of the potential risks. Patients with pain in the hips, thighs or femur should be radiologically assessed and, where a stress fracture is evident, the physician should decide whether bisphosphonate therapy should be discontinued pending a full evaluation, based on an individual benefit–risk assessment. The radiographic changes should be evaluated for orthopaedic intervention—since surgery prior to fracture completion might be advantageous—or be closely monitored.

**Acknowledgements** The Working Group meeting was supported by an unrestricted educational grant from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis. Editorial assistance for the manuscript was provided by Sola Neunie of BioScience Communications, supported by a financial grant from Novartis Pharmaceuticals.

**Conflicts of interest** Rene Rizzoli has attended paid advisory boards and received consultancy and lecturing fees from Servier, Novartis, Eli Lilly, Amgen, Roche, Nycomed, Merck Sharp and Dohme and Danone. Kristina Åkesson has received lecturing fees from Medtronic, Novartis, Amgen, Merck and Nycomed. Mary Bouxsein has undertaken consultancy and lecturing commitments for Amgen and Merck & Co. John A. Kanis consults or has received research support from a large number of pharmaceutical companies involved in marketing products for treatment of osteoporosis. He is president of the International Osteoporosis Foundation and serves on its Committee of Scientific Advisors. Nicola Napoli has received grant support from Merck Sharpe and Dohme. Socrates Papapoulos has received consultancy and lecturing fees from Alliance for Better Bone Health, Amgen, Eli Lilly, GSK, Merck & Co, Novartis, Pfizer and Roche. Jean-Yves Reginster has received consulting fees and attended paid advisory boards for Servier, Novartis, Negma, Lilly, Wyeth, Amgen, GlaxoSmithKline, Roche, Merckle, Nycomed, NPS, Theramex and UCB. He has received invited lecture fees from Merck Sharp and Dohme, Lilly, Rottapharm, IBSA, Genevrier, Novartis, Servier, Roche, GlaxoSmithKline, Teijin, Teva, Ebewee Pharma, Zodiac, Analis, Theramex, Nycomed and Novo Nordisk. He has received grant support from Bristol Myers Squibb, Merck Sharp & Dohme, Rottapharm, Teva, Lilly, Novartis, Roche, GlaxoSmithKline,

Amgen and Servier. Cyrus Cooper has undertaken consultancy and lecturing commitments for Alliance for Better Bone Health, Eli Lilly, Novartis, GSK Roche, Servier, MSD and Amgen.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- Black DM, Cummings SR, Karpf DB, Cauley JA, Thompson DE, Nevitt MC, Bauer DC, Genant HK, Haskell WL, Marcus R, Ott SM, Torner JC, Quandt SA, Reiss TF, Ensrud KE (1996) Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. Fracture Intervention Trial Research Group. *Lancet* 348:1535–1541
- Black DM, Schwartz AV, Ensrud KE, Cauley JA, Levis S, Quandt SA, Satterfield S, Wallace RB, Bauer DC, Palermo L, Wehren LE, Lombardi A, Santora AC, Cummings SR (2006) Effects of continuing or stopping alendronate after 5 years of treatment: the Fracture Intervention Trial Long-term Extension (FLEX): a randomized trial. *JAMA* 296:2927–2938
- Black DM, Delmas PD, Eastell R, Reid IR, Boonen S, Cauley JA, Cosman F, Lakatos P, Leung PC, Man Z, Mautalen C, Mesenbrink P, Hu H, Caminis J, Tong K, Rosario-Jansen T, Krasnow J, Hue TF, Sellmeyer D, Eriksen EF, Cummings SR (2007) Once-yearly zoledronic acid for treatment of postmenopausal osteoporosis. *N Engl J Med* 356:1809–1822
- Chesnut CH III, Skag A, Christiansen C, Recker R, Stakkestad JA, Hoiseth A, Felsenberg D, Huss H, Gilbride J, Schimmer RC, Delmas PD (2004) Effects of oral ibandronate administered daily or intermittently on fracture risk in postmenopausal osteoporosis. *J Bone Miner Res* 19:1241–1249
- Cummings SR, Black DM, Thompson DE, Applegate WB, Barrett-Connor E, Musliner TA, Palermo L, Prineas R, Rubin SM, Scott JC, Vogt T, Wallace R, Yates AJ, LaCroix AZ (1998) Effect of alendronate on risk of fracture in women with low bone density but without vertebral fractures: results from the Fracture Intervention Trial. *JAMA* 280:2077–2082
- Harris ST, Watts NB, Genant HK, McKeever CD, Hangartner T, Keller M, Chesnut CH III, Brown J, Eriksen EF, Hoesly MS, Axelrod DW, Miller PD (1999) Effects of risendronate treatment on vertebral and nonvertebral fractures in women with postmenopausal osteoporosis: a randomized controlled trial. Vertebral Efficacy With Risendronate Therapy (VERT) Study Group. *JAMA* 282:1344–1352
- Lyles KW, Colon-Emeric CS, Magaziner JS, Adachi JD, Pieper CF, Mautalen C, Hyldstrup L, Recknor C, Nordsletten L, Moore KA, Laveccchia C, Zhang J, Mesenbrink P, Hodgson PK, Abrams K, Orloff JJ, Horowitz Z, Eriksen EF, Boonen S (2007) Zoledronic acid and clinical fractures and mortality after hip fracture. *N Engl J Med* 357:1799–1809
- Reginster J, Minne HW, Sorensen OH, Hooper M, Roux C, Brandi ML, Lund B, Ethgen D, Pack S, Roumagnac I, Eastell R (2000) Randomized trial of the effects of risendronate on vertebral fractures in women with established postmenopausal osteoporosis. Vertebral Efficacy with Risendronate Therapy (VERT) Study Group. *Osteoporos Int* 11:83–91
- Sorensen OH, Crawford GM, Mulder H, Hosking DJ, Gennari C, Mellstrom D, Pack S, Wenderoth D, Cooper C, Reginster JY (2003) Long-term efficacy of risendronate: a 5-year placebo-controlled clinical experience. *Bone* 32:120–126
- McClung MR, Geusens P, Miller PD, Zippel H, Bensen WG, Roux C, Adami S, Fogelman I, Diamond T, Eastell R, Meunier PJ, Reginster JY (2001) Effect of risendronate on the risk of hip fracture in elderly women. Hip Intervention Program Study Group. *N Engl J Med* 344:333–340
- Reid DM, Devogelaer JP, Saag K, Roux C, Lau CS, Reginster JY, Papanastasiou P, Ferreira A, Hartl F, Fashola T, Mesenbrink P, Sambrook PN (2009) Zoledronic acid and risendronate in the prevention and treatment of glucocorticoid-induced osteoporosis (HORIZON): a multicentre, double-blind, double-dummy, randomised controlled trial. *Lancet* 373:1253–1263
- Devogelaer JP (2002) Modern therapy for Paget's disease of bone: focus on bisphosphonates. *Treat Endocrinol* 1:241–257
- Lipton A (2007) Treatment of bone metastases and bone pain with bisphosphonates. *Support Cancer Ther* 4:92–100
- Polascik TJ (2009) Bisphosphonates in oncology: evidence for the prevention of skeletal events in patients with bone metastases. *Drug Des Devel Ther* 3:27–40
- Roche Registration Limited (2009) Bonviva summary of product characteristics. Roche Registration, Hertfordshire
- Merck SaDL (2007) Fosamax summary of product characteristics. Merck SaDL, Hertfordshire
- Procter & Gamble Pharmaceuticals (2007) Actonel summary of product characteristics. Procter & Gamble Pharmaceuticals, Weybridge
- US Food and Drug Administration (FDA) (2009) Drug safety newsletter. Volume 2, Number 2. <http://www.fda.gov/Drugs/DrugSafety/DrugSafetyNewsletter/default.htm>. Accessed 23 Sep 2010
- Bilezikian JP (2006) Osteonecrosis of the jaw—do bisphosphonates pose a risk? *N Engl J Med* 355:2278–2281
- Rizzoli R, Burlet N, Cahall D, Delmas PD, Eriksen EF, Felsenberg D, Grbic J, Jontell M, Landesberg R, Laslop A, Wollenhaupt M, Papapoulos S, Sezer O, Sprafka M, Reginster JY (2008) Osteonecrosis of the jaw and bisphosphonate treatment for osteoporosis. *Bone* 42:841–847
- Novartis Europharm Limited (2009) Aclasta summary of product characteristics. Novartis Europharm, Horsham
- Merck Sharp & Dohme Limited (2009) Fosavance summary of product characteristics. Merck Sharp & Dohme, Hertfordshire
- Roche Pharmaceuticals (2009) Boniva (ibandronate sodium) injection prescribing information. Roche Pharmaceuticals, Nutley
- Guanabens N, Peris P, Monegal A, Pons F, Collado A, Munoz-Gomez J (1994) Lower extremity stress fractures during intermittent cyclical etidronate treatment for osteoporosis. *Calcif Tissue Int* 54:431–434
- Armamento-Villareal R, Napoli N, Diemer K, Watkins M, Civitelli R, Teitelbaum S, Novack D (2009) Bone turnover in bone biopsies of patients with low-energy cortical fractures receiving bisphosphonates: a case series. *Calcif Tissue Int* 85:37–44
- Goh SK, Yang KY, Koh JS, Wong MK, Chua SY, Chua DT, Howe TS (2007) Subtrochanteric insufficiency fractures in patients on alendronate therapy: a caution. *J Bone Joint Surg Br* 89:349–353
- Ing-Lorenzini K, Desmeules J, Plachta O, Suva D, Dayer P, Peter R (2009) Low-energy femoral fractures associated with the long-term use of bisphosphonates: a case series from a Swiss university hospital. *Drug Saf* 32:775–785
- Kwek EB, Goh SK, Koh JS, Png MA, Howe TS (2008) An emerging pattern of subtrochanteric stress fractures: a long-term complication of alendronate therapy? *Injury* 39:224–231
- Lenart BA, Neviaser AS, Lyman S, Chang CC, Edobor-Osula F, Steele B, van der Meulen MC, Lorich DG, Lane JM (2009) Association of low-energy femoral fractures with prolonged bisphosphonate use: a case control study. *Osteoporos Int* 20:1353–1362

30. Neviaser AS, Lane JM, Lenart BA, Edobor-Osula F, Lorch DG (2008) Low-energy femoral shaft fractures associated with alendronate use. *J Orthop Trauma* 22:346–350
31. Odvina CV, Zerwekh JE, Rao DS, Maalouf N, Gottschalk FA, Pak CY (2005) Severely suppressed bone turnover: a potential complication of alendronate therapy. *J Clin Endocrinol Metab* 90:1294–1301
32. Fielding JW, Magliato HJ (1966) Subtrochanteric fractures. *Surg Gynecol Obstet* 122:555–560
33. Muller ME, Nazarian S, Koch P, Schatzker J (1990) The AO classification of long bones. [http://aofoundation.com/AOFileServer/PortalFiles/FilePath=/Extranet/en/\\_att/wor/act/fracture\\_-classif/mueller\\_ao\\_class.pdf](http://aofoundation.com/AOFileServer/PortalFiles/FilePath=/Extranet/en/_att/wor/act/fracture_-classif/mueller_ao_class.pdf). Accessed 23 Sep 2010
34. Seinsheimer F (1978) Subtrochanteric fractures of the femur. *J Bone Joint Surg Am* 60:300–306
35. Black DM, Genant HK, Bucci-Rechtweg C, Bauer DC, Mesenbrink PG, Palermo L, Nussgartner L, Eastell R (2009) Does zoledronic acid increase risk of atypical subtrochanteric femoral shaft fractures? Results from the HORIZON-PFT. *J Bone Miner Res* 24 (Suppl 1). <http://www.asbmr.org/Meetings/AnnualMeeting/AbstractDetail.aspx?aid=918d35dd-6a3d-43f6-b35f-b484a15b81cf>. Accessed 23 Sep 2010
36. Bunning RD, Rentfro RJ, Jelinek JS (2010) Low-energy femoral fractures associated with long-term bisphosphonate use in a rehabilitation setting: a case series. *PM&R* 2:76–80
37. Capeci CM, Tejwani NC (2009) Bilateral low-energy simultaneous or sequential femoral fractures in patients on long-term alendronate therapy. *J Bone Joint Surg Am* 91:2556–2561
38. Koh JS, Goh SK, Png MA, Kwek EB, Howe TS (2010) Femoral cortical stress lesions in long-term bisphosphonate therapy: a herald of impending fracture? *J Orthop Trauma* 24:75–81
39. Visekruna M, Wilson D, McKiernan FE (2008) Severely suppressed bone turnover and atypical skeletal fragility. *J Clin Endocrinol Metab* 93:2948–2952
40. Kwek EB, Koh JS, Howe TS (2008) More on atypical fractures of the femoral diaphysis. *N Engl J Med* 359:316–317
41. Lenart BA, Lorch DG, Lane JM (2008) Atypical fractures of the femoral diaphysis in postmenopausal women taking alendronate. *N Engl J Med* 358:1304–1306
42. Leung F, Lau T-W, To M, Luk K-K, Kung A (2010) Atypical femoral diaphyseal and subtrochanteric fractures and their association with bisphosphonates. *BMJ Case Reports*. doi:10.1136/bcr.10.2008.1073
43. Brandser EA, Buckwalter JA (1996) Imaging studies for diagnosing stress and insufficiency fractures. *Iowa Orthop J* 16:70–78
44. Lunsjo K, Ceder L, Tidermark J, Hamberg P, Larsson BE, Ragnarsson B, Knebel RW, Allvin I, Hjalmarsson K, Norberg S, Fornander P, Hauggaard A, Stigsson L (1999) Extramedullary fixation of 107 subtrochanteric fractures: a randomized multicenter trial of the Medoff sliding plate versus 3 other screw-plate systems. *Acta Orthop Scand* 70:459–466
45. Whitelaw GP, Segal D, Sanzone CF, Ober NS, Hadley N (1990) Unstable intertrochanteric/subtrochanteric fractures of the femur. *Clin Orthop Relat Res* 252:238–245
46. Nieves JW, Bilezikian JP, Lane JM, Einhorn TA, Wang Y, Steinbuch M, Cosman F (2010) Fragility fractures of the hip and femur: incidence and patient characteristics. *Osteoporos Int* 21:299–408
47. Glennon DA (2009) Subtrochanteric stress fractures in six patients on long term bisphosphonate therapy: a case series. *Bone* 44:S68–S98
48. National Institute for Health and Clinical Excellence (2010) Final appraisal determination. Alendronate, etidronate, risedronate, raloxifene and strontium ranelate for the primary prevention of osteoporotic fragility fractures in postmenopausal women (amended). NICE technology appraisal guidance 160 (amended). <http://www.nice.org.uk/nicemedia/pdf/TA160guidance.pdf>. Accessed 23 Sep 2010
49. Johnell O, Kanis JA (2006) An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 17:1726–1733
50. Giusti A, Hamdy NA, Papapoulos SE (2010) Atypical fractures of the femur and bisphosphonate therapy. A systematic review of case/case series studies. *Bone* 47:169–180
51. Husada G, Libberecht K, Peeters T, Populaire J (2005) Bilateral mid-diaphyseal femoral stress fractures in the elderly. *Eur J Trauma* 31:68–71
52. Schneider JP (2006) Should bisphosphonates be continued indefinitely? An unusual fracture in a healthy woman on long-term alendronate. *Geriatrics* 61:31–33
53. Armamento-Villareal R, Napoli N, Panwar V, Novack D (2006) Suppressed bone turnover during alendronate therapy for high-turnover osteoporosis. *N Engl J Med* 355:2048–2050
54. Cheung RK, Leung KK, Lee KC, Chow TC (2007) Sequential non-traumatic femoral shaft fractures in a patient on long-term alendronate. *Hong Kong Med J* 13:485–489
55. Demiralp B, Ilgan S, Ozgur KA, Cicek EI, Yildirim D, Erler K (2007) Bilateral femoral insufficiency fractures treated with inflatable intramedullary nails: a case report. *Arch Orthop Trauma Surg* 127:597–601
56. Lee P, van der Wall H, Seibel MJ (2007) Looking beyond low bone mineral density: multiple insufficiency fractures in a woman with post-menopausal osteoporosis on alendronate therapy. *J Endocrinol Invest* 30:590–597
57. Sayed-Noor AS, Sjoden GO (2008) Subtrochanteric displaced insufficiency fracture after long-term alendronate therapy—a case report. *Acta Orthop* 79:565–567
58. Odvina CV, Levy S, Rao S, Zerwekh JE, Sudhaker RD (2009) Unusual mid-shaft fractures during long term bisphosphonate therapy. *Clin Endocrinol (Oxf)* 72:161–168
59. Ali T, Jay RH (2009) Spontaneous femoral shaft fracture after long-term alendronate. *Age Ageing* 38:625–626
60. Goddard MS, Reid KR, Johnston JC, Khanuja HS (2009) Atraumatic bilateral femur fracture in long-term bisphosphonate use. *Orthopedics* 32:607. doi:10.3928/01477447-20090624-27
61. Sayed-Noor AS, Sjoden GO (2009) Case reports: two femoral insufficiency fractures after long-term alendronate therapy. *Clin Orthop Relat Res* 467:1921–1926
62. Cermak K, Shumelinsky F, Alexiou J, Gebhart MJ (2009) Case reports: subtrochanteric femoral stress fractures after prolonged alendronate therapy. *Clin Orthop Relat Res* 468: 1991–1996
63. Bush LA, Chew FS (2009) Subtrochanteric femoral insufficiency fracture in woman on bisphosphonate therapy for glucocorticoid-induced osteoporosis. *Radiol Case Rep* 4. doi:1.2484/rcr.v4i1.261
64. Lee JK (2009) Bilateral atypical femoral diaphyseal fractures in a patient treated with alendronate sodium. *Int J Rheum Dis* 12:149–154
65. Edwards MH, McCrae FC, Young-Min SA (2010) Alendronate-related femoral diaphysis fracture—what should be done to predict and prevent subsequent fracture of the contralateral side? *Osteoporos Int* 21:701–703
66. Schilcher J, Aspenberg P (2009) Incidence of stress fractures of the femoral shaft in women treated with bisphosphonate. *Acta Orthop* 80:413–415
67. Abrahamsen B, Eiken P, Eastell R (2009) Subtrochanteric and diaphyseal femur fractures in patients treated with alendronate: a register-based national cohort study. *J Bone Miner Res* 24:1095–1102
68. Black DM, Thompson DE, Bauer DC, Ensrud K, Musliner T, Hochberg MC, Nevitt MC, Suryawanshi S, Cummings SR (2000) Fracture risk reduction with alendronate in women with osteoporosis: the Fracture Intervention Trial. FIT Research Group. *J Clin Endocrinol Metab* 85:4118–4124

69. Black DM, Kelly MP, Genant HK, Palermo L, Eastell R, Bucci-Rechtweg C, Cauley J, Leung PC, Boonen S, Santora A, de Papp A, Bauer DC (2010) Bisphosphonates and fractures of the subtrochanteric or diaphyseal femur. *N Engl J Med* 362:1761–1771
70. Bilezikian J, Klemes A, Silverman S, Cosman F (2009) Subtrochanteric fracture reports coincident with risenedronate use. *J Bone Miner Res* 24(Suppl 1). <http://www.asbmr.org/Meetings/AnnualMeeting/AbstractDetail.aspx?aid=0367cfaa-4d0d-47d8-a57a-ff76098839a2>. Accessed 23 Sep 2010
71. Eisman JA, Civitelli R, Adami S, Czerwinski E, Recknor C, Prince R, Reginster JY, Zaidi M, Felsenberg D, Hughes C, Mairon N, Masanaukaite D, Reid DM, Delmas PD, Recker RR (2008) Efficacy and tolerability of intravenous ibandronate injections in postmenopausal osteoporosis: 2-year results from the DIVA study. *J Rheumatol* 35:488–497
72. Miller PD, McClung MR, Macovei L, Stakkestad JA, Luckey M, Bonvoisin B, Reginster JY, Recker RR, Hughes C, Lewiecki EM, Felsenberg D, Delmas PD, Kendler DL, Bolognese MA, Mairon N, Cooper C (2005) Monthly oral ibandronate therapy in postmenopausal osteoporosis: 1-year results from the MOBILE study. *J Bone Miner Res* 20:1315–1322
73. Recker R, Stakkestad JA, Chesnut CH III, Christiansen C, Skag A, Hoiseth A, Ettinger M, Mahoney P, Schimmer RC, Delmas PD (2004) Insufficiently dosed intravenous ibandronate injections are associated with suboptimal antifracture efficacy in postmenopausal osteoporosis. *Bone* 34:890–899
74. Miller PD, Epstein S, Sedarati F, Reginster JY (2008) Once-monthly oral ibandronate compared with weekly oral alendronate in postmenopausal osteoporosis: results from the head-to-head MOTION study. *Curr Med Res Opin* 24:207–213
75. Stakkestad JA, Lakatos P, Lorenc R, Sedarati F, Neate C, Reginster JY (2008) Monthly oral ibandronate is effective and well tolerated after 3 years: the MOBILE long-term extension. *Clin Rheumatol* 27:955–960
76. McClung MR, Bolognese MA, Sedarati F, Recker RR, Miller PD (2009) Efficacy and safety of monthly oral ibandronate in the prevention of postmenopausal bone loss. *Bone* 44:418–422
77. Bianchi G, Felsenberg D, Czerwinski E, Reid D, Kenwright A, Burdeska A, Recker R (2009) Efficacy of IV ibandronate is maintained over 5 years: the DIVA LTE study. *Ann Rheum Dis* 68 (Suppl 3):494
78. European Medicines Agency (2009) Assessment report for Fosavance. EMEA/CHMP/188952/2009. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Assessment\\_Report\\_-\\_Variation/human/000619/WC500024252.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Assessment_Report_-_Variation/human/000619/WC500024252.pdf). Accessed 23 Sep 2010
79. Merck Sharp & Dohme Limited (2010) Fosamax summary of product characteristics. Merck Sharp & Dohme, Hertfordshire
80. European Medicines Agency (2009) EMEA 2010 priorities for drug safety research. Long-term adverse skeletal effects of bisphosphonates. Doc.Ref: EMEA/493711/2009 Rev.1. European Medicines Agency, London
81. US Food and Drug Administration (FDA) (2010) FDA drug safety communication: ongoing safety review of oral bisphosphonates and atypical subtrochanteric fractures. <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientandProviders/ucm203891.htm>. Accessed 23 Sep 2010
82. Durchschlag E, Paschalis EP, Zoehrer R, Roschger P, Fratzl P, Recker R, Phipps R, Klaushofer K (2006) Bone material properties in trabecular bone from human iliac crest biopsies after 3- and 5-year treatment with risenedronate. *J Bone Miner Res* 21:1581–1590
83. Boskey AL, Spevak L, Weinstein RS (2009) Spectroscopic markers of bone quality in alendronate treated postmenopausal women. *Osteoporos Int* 20:793–800
84. Turner CH, Burr DB (2006) Principles of bone biomechanics. In: Lane NE, Sambrook PN (eds) *Osteoporosis and the osteoporosis of rheumatic diseases*. Mosby Elsevier, Philadelphia, pp 41–53
85. Boivin GY, Chavassieux PM, Santora AC, Yates J, Meunier PJ (2000) Alendronate increases bone strength by increasing the mean degree of mineralization of bone tissue in osteoporotic women. *Bone* 27:687–694
86. Roschger P, Rinnerthaler S, Yates J, Rodan GA, Fratzl P, Klaushofer K (2001) Alendronate increases degree and uniformity of mineralization in cancellous bone and decreases the porosity in cortical bone of osteoporotic women. *Bone* 29:185–191
87. Allen MR, Burr DB (2007) Three years of alendronate treatment results in similar levels of vertebral microdamage as after one year of treatment. *J Bone Miner Res* 22:1759–1765
88. Allen MR, Iwata K, Phipps R, Burr DB (2006) Alterations in canine vertebral bone turnover, microdamage accumulation, and biomechanical properties following 1-year treatment with clinical treatment doses of risenedronate or alendronate. *Bone* 39:872–879
89. Allen MR, Reinwald S, Burr DB (2008) Alendronate reduces bone toughness of ribs without significantly increasing microdamage accumulation in dogs following 3 years of daily treatment. *Calcif Tissue Int* 82:354–360
90. Iwata K, Allen MR, Phipps R, Burr DB (2006) Microcrack initiation occurs more easily in vertebrae from beagles treated with alendronate than with risenedronate. *Bone* 38(Suppl):42
91. Cao Y, Mori S, Mashiba T, Westmore MS, Ma L, Sato M, Akiyama T, Shi L, Komatsubara S, Miyamoto K, Norimatsu H (2002) Raloxifene, estrogen, and alendronate affect the processes of fracture repair differently in ovariectomized rats. *J Bone Miner Res* 17:2237–2246
92. MacDonald MM, Schindeler A, Little DG (2007) Bisphosphonate treatment and fracture repair. *BoneKey* 4:236–251
93. Martinez MD, Schmid GJ, McKenzie JA, Ornitz DM, Silva MJ (2010) Healing of non-displaced fractures produced by fatigue loading of the mouse ulna. *Bone* 46:1604–1612
94. Somford MP, Draijer FW, Thomassen BJ, Chavassieux PM, Boivin G, Papapoulos SE (2009) Bilateral fractures of the femur diaphysis in a patient with rheumatoid arthritis on long-term treatment with alendronate: clues to the mechanism of increased bone fragility. *J Bone Miner Res* 24:1736–1740
95. O'Neill TW (2005) Looking back: developments in our understanding of the occurrence, aetiology and prognosis of osteoporosis over the last 50 years. *Rheumatology (Oxford)* 44:iv33–iv35
96. Kanis JA, McCloskey EV, Johansson H, Strom O, Borgstrom F, Oden A (2008) Case finding for the management of osteoporosis with FRAX—assessment and intervention thresholds for the UK. *Osteoporos Int* 19:1395–1408
97. Kanis JA, Borgstrom F, Zethraeus N, Johnell O, Oden A, Jonsson B (2005) Intervention thresholds for osteoporosis in the UK. *Bone* 36:22–32