

# Multi-factorial influences on sex ratio: a spatio-temporal investigation of endocrine disruptor pollution and neighborhood stress

Ewan McDonald<sup>1</sup>, Andrew Watterson<sup>1</sup>, Andrew N. Tyler<sup>2</sup>, John McArthur<sup>2</sup>, E. Marion Scott<sup>3</sup>

<sup>1</sup>Occupational and Environmental Health Research Group, Centre for Public Health and Population Health Research, University of Stirling, UK, <sup>2</sup>Biological and Environmental Sciences, School of Natural Sciences, University of Stirling, UK, <sup>3</sup>School of Mathematics and Statistics, University of Glasgow, UK

**Background:** It is suggested the declining male birth proportion in some industrialized countries is linked to ubiquitous endocrine disruptor exposure. Stress and advanced parental age are determinants which frequently present positive findings. Multi-factorial influences on population sex ratio are rarely explored or tested in research.

**Objectives:** To test the hypothesis that dual factors of pollution and population stress affects sex proportion at birth through geographical analysis of Central Scotland.

**Methods:** The study incorporates the use of Geographical Information Systems (GIS) tools to overlay modeled point source endocrine disruptor air emissions with “small-area” data on multiple deprivation (a proxy measurement of stress) and birth sex. Historical review of regional sex ratio trends presents additional data on sex ratio in Scotland to consider.

**Results:** There was no overall concentration in Central Scotland of low sex ratio neighborhoods with areas where endocrine disruptor air pollution and deprivation or economic stress were high. Historical regional trends in Scotland (from 1973), however, do show significantly lower sex ratio values for populations where industrial air pollution is highest (i.e. Eastern Central Scotland).

**Conclusions:** Use of small area data sets and pollution inventories is a potential new method of inquiry for reproductive environmental and health protection monitoring and has produced interesting findings.

**Keywords:** Sex ratio, Male births, Endocrine disruption, Reproductive health, Air pollution, Multiple deprivation, Neighborhood stress, Maternal age

## Introduction

It has been suggested that the declining proportion of male births in some Western industrialized countries post-World War II may be attributed to human exposure to endocrine disrupting compounds.<sup>1,2</sup> This hypothesis sits alongside other reproductive epidemiological trends that have been linked to endocrine disruption and pollution, including: the rising incidence of testicular cancer and declining sperm quality for men;<sup>3,4</sup> earlier sexual development in adolescents;<sup>5</sup> and the increasing occurrence of cryptorchidism in male newborns.<sup>6</sup> Some of these reported trends could also be a result of social determinants that reflect lifestyles in Western countries, such as poor nutrition and sedentary behavior.<sup>7</sup>

The secondary sex ratio is the ratio between male to female births per 100 or 1000 live births. The male

birth proportion marker is calculated by dividing male births into total live births and can be used to describe demographic trends in countries and regions.<sup>8,9</sup> For the purposes of this paper, the term “sex ratio” refers to both of these calculations and “sex proportion at birth” (male births/total births) is the unit of analysis in the spatio-temporal investigation. The primary sex ratio is the ratio of males to females at the time of conception and the secondary sex ratio is the ratio at birth. Various reports have demonstrated proportionally greater male fetal death rates, resulting in the adjustment of the primary sex ratio downwards to the secondary sex ratio.<sup>10–12</sup>

Motivation for measurement and analysis of the human sex ratio arises from multiple hypotheses across social biology;<sup>13–15</sup> environmental, medical and behavioral science;<sup>16–19</sup> demography;<sup>20,21</sup> and epidemiology.<sup>22,23</sup> It has been suggested that because the sex ratio can easily be calculated, there has been an over-proliferation of such research with highly speculative

Correspondence to: E. McDonald, Occupational and Environmental Health Research Group, RG Bomont Bldg, University of Stirling, Stirling, Scotland FK9 4LA, UK. Email: e.w.mcdonald@stir.ac.uk

hypotheses built from sex ratio alteration reports.<sup>24</sup> However, the sex ratio offers an alternative “response variable” for many environmental toxicology and epidemiological study designs. For example, there are clear ethical difficulties in recording adverse reproductive health events, such as pregnancy loss, in small geographical areas. Furthermore, a snapshot of sex ratio imbalance in a population may be instructive for new paradigms of biology, whereby teratogenic exposures are possibly linked to the later on-set of disease in the life course.<sup>25</sup>

Scientists have identified persistent organic pollutants, particularly chronic exposure to polychlorinated bi-phenyls (PCBs) and acute releases of dioxins, as likely to be linked to human sex ratio alteration. For PCBs, research suggests the effects are bi-phasic whereby the sex ratio increase occurs with acute exposure and declines with persistent and chronic exposure.<sup>26–28</sup> Bio-measurement of toxin serum levels add validity to findings, though this is qualified by the lack of any STROBE assessment of particular bio-markers.<sup>29</sup> The use of bio-measurement is particularly instructive in studies designed with other reproductive parameters such as sperm quality. Analysis of socio-economic and health-related confounders and obtaining serum samples almost immediately upon exposure can also add validity to investigations. The direct recruitment of cohorts with acute dioxin exposure from the Seveso industrial accident permitted robust results showing significant downward sex ratio alteration, particularly among young men in a comprehensive follow-up study.<sup>19,30</sup> The triple investigation of sperm production, 1,2-dibromo-3-chloropropane (DBCP) levels (previously used as a soil fumigant) and low secondary sex ratio, as well as a case-control study of testicular cancer incidence and reduced sex ratio, add further weight to the environmental hypothesis on reduced male births.<sup>31,32</sup> However, well-designed toxicological investigations of PCBs in breast milk and maternal lead pollution exposure found no effect on sex ratio.<sup>33,34</sup> These non-significant results may reflect different endocrine disrupting properties for particular compounds or congeners, or altering effects on sex ratio may be dependent on dose, timing, or male or female exposure. Studies adopting mapping techniques and geographical analysis, including estimates of pesticide exposure, air pollution, and multiple source pollution, have produced mixed results with skewing of sex ratios upwards, downwards, and within normal confidence limits.<sup>35–37</sup> Such studies present significant methodological challenges in modeling or measuring environmental exposure which usually include surrogate data, particularly given the geographic and temporal variation and composition of multiple and potentially interacting pollutants.

Substantial evidence from large demographic ecological studies suggests that advancing parental age contributes to a lowering of sex ratio.<sup>20,21,38–42</sup> Stress may also alter the secondary sex ratio, with declines in the proportion of male births during and after acutely hazardous events.<sup>43–48</sup> The greater vulnerability of the male fetus to spontaneous loss in pregnancy has been consistently demonstrated and may be a factor in the decline in sex ratios.<sup>10,12</sup> In a recent UK cohort study, advanced maternal age was independently associated with greater risk of miscarriage in the first trimester of pregnancy.<sup>49</sup> There is also evidence that women with older male partners are more susceptible to early pregnancy loss.<sup>50,51</sup> Maternal endocrine changes and stressor responses linked to adverse birth outcomes, including fetal mortality, have received much attention in peri-natal epidemiology. Furthermore, exposure to stress during pregnancy as a result of living in deprived urban areas with elevated crime and physical degradation has been examined in a number of studies focusing on birth outcomes.<sup>52–54</sup>

Many analyses have been conducted on socio-economic status and sex ratio that test the adaptive Trivers–Willard hypothesis that proposes, with natural selection, the “strongest” and wealthiest parents are more likely to sire sons owing to their reproductive advantage over daughters i.e. producing more grandchildren. Results from such types of studies, however, are contradictory with methodological weaknesses attached to using occupation and educational levels as singular de-facto measurements of socio-economic status.<sup>55</sup>

Determination of the secondary sex ratio is multifaceted with evidence suggesting environmental pollution, population stress, and parental age as likely influential factors. Yet there is a paucity of research on sex ratios that investigates the environmental, health-related, and socio-economic influences. Frequently, individual studies tend to be “locked” in a single discipline approach though exceptions are some well-designed environmental epidemiological studies that include adjustments for confounders. The aim of this study is to test the hypothesis that endocrine disruptor pollution combined with population stress influences the proportion of sex at birth in Scotland. The authors utilized Geographical Information Systems (GIS) to combine data on point source endocrine disruptor pollution emissions, area-based data-sets of socio-economic and public health indicators, and compatible area measurement of sex proportion at birth. Starting in 2003, pollution data from Scotland are available in the Scottish Pollutant Release Inventory. All registered emitters are mandated to report total emissions (in kg) of individual pollutants at a site address. Extensive socio-economic and health-related area-based data collected at the neighborhood level

are also available. Scottish birth registration records are available starting in the year 1855 with some regional areas retaining consistent boundaries from 1973 and neighborhood data also available from 2001. The extent and consistency of birth sex ratio data in Scotland thus permits broad analysis over historical time periods and regions complementing the neighborhood multi-factorial approach.

## Methods

The study was divided into three stages: (1) a review of temporal and regional trends in the secondary sex ratio in Scotland; (2) a stratified analysis of sex ratios and levels of multiple deprivation for small areas, i.e. datazones, in Scotland; and (3) a GIS spatial analysis of datazone sex ratios with modeled endocrine disruptor exposure for Central Scotland combined with the levels of multiple deprivation.

### Data sources

For stage (1), data were sourced from the General Register Office (GRO), which compiles birth statistics from local registrars in Scotland. Births by sex from 1855 to 2010 were obtained from the GRO and GRO statisticians provided birth totals by health board area from 1973 to 2010. The “Advanced Reporter” database at Scottish Neighborhood Statistics (SNS) (<http://www.sns.gov.uk>) was utilized for stages (2) and (3). Data were constructed from multiple sources, including the Scottish Index of Multiple Deprivation (SIMD) and geo-coding from Information Services Division (NHS Scotland). The smallest area abstracted in SNS is the datazone level, which was selected as the major geographical unit of reference for stages (2) and (3). Datazones comprise 6505 areas in Scotland formed by amalgamating census enumeration districts to establish distinctive neighborhood zones with populations ranging from 600 to 1000 persons. Homogeneity of birth data sets was achieved for all stages, as the GRO (Scotland) geo-codes and reports the sex at birth for Scottish Neighborhood Statistics data zones. The Scottish Pollutant Release Inventory (SPRI), produced by the Scottish Environmental Protection Authority (SEPA), was the primary data source for modeled endocrine disruptor pollution in stage (3). The interactive database contains annual declarations in kilograms for separate pollutants by registered emitters, principally industrial companies, or landfill/waste water operators. Data are tabulated for the years 2002 and 2004–2010 and searchable according to local authority area, pollution media type (air, land, water, wastewater), and specific pollutant.

With the exception of non-compliance with birth registration in some rural areas in the late nineteenth century, the GRO (Scotland) statistics are robust and accurate.<sup>56</sup> We used the SIMD for the year 2006 to

create the stratified and spatial analyses categorical values (“vigintiles” — divide the distribution of individuals into twenty groups of equal frequency). SIMD is an area-based composite index of deprivation with “equalizing” of varied forms of data on income, employment, education, crime, access, and housing. Deprivation indices have been widely used in Scotland as geographical measurements of poverty, undergoing numerous methodological and data refinements since the 1980s.<sup>57</sup> SIMD was produced according to the rank of datazones from 1 to 6505 and also *vigintiles* — group 1 with most deprivation to 20 for the lowest deprivation. Stratification was performed by cross-tabulating and aggregating the datazones by SIMD *vigintile* with total male births and total births. This allowed us to calculate secondary sex ratios for each level of deprivation (1–20). All statistical analyses were performed using SPSS version 19.

### Statistical methods

Significance testing was performed by hypothesis testing of two proportions (*z*-test). The two proportions tested were: (1) SIMD *vigintile*/regional (health board area) sex ratios; and (2) “control” samples of other regional sex ratios in the UK. Male births and total births in the Yorkshire & Humber region were selected as the control region for cumulative health board sex ratios (1973–2010) and for the 20 levels of multiple deprivation. This region was selected as the control due to the roughly equivalent birth totals per year (approx. 30 000), comparable post-war downward trend in sex ratio and similar industrial geography of heavy industry concentrated in urban areas contrasted by expansive rural catchments. Where sex ratios for five separate years of health board areas in Scotland were compared, cumulative birth figures from an earlier national period in Scotland (1974–1979) were used as the control sample. Independence between the two proportions is thus achieved, as time periods are not commensurate. The null hypothesis was that regional sex ratio, or sex ratio for an area-measured deprivation level, is no different than the sex ratio for a similar region or earlier national calculation.

### Spatial analysis

The spatial analysis was confined to a central region of Scotland consisting of six areas (Falkirk, South Lanarkshire, North Lanarkshire, Stirling, Fife, Clackmannanshire) covering 1640 datazones (Fig. 2). Central Scotland is the most industrial area in the country with historically high levels of pollution from old industries, but also with rural areas with low pollution. It contains major sites of industrial pollution in Scotland, such as the Grangemouth INEOS complex and Longannet coal-fired power station, as well as large tracts of contaminated and remediated

land, a legacy of Scotland's heavy manufacturing industry. There is also socio-economic diversity with both urban and rural localities and deprived and non-deprived areas.

Mapping of potential multi-factorial influences on local sex ratios involves creating a projection of modeled endocrine disruptor air pollution emissions from SPRI registered emitter sites over datazone areas in Central Scotland. All datazones were assigned multiple deprivation levels. Significantly low sex ratios ( $P < 0.1$ ) in these small-areas are also shown. A simplified modeling of endocrine disruptor air pollution emissions was achieved using the following methodology:

1. Identification of endocrine disruptors on the SPRI Pollutant Search list. Cross-referencing with the Endocrine Disruption Exchange — *List of Potential Endocrine Disruptors* with SPRI Pollutant List.
2. Systematic search of SPRI for air pollutant releases of identified endocrine disruptors for each year from 2002 to 2010 (no records 2003) by six local authorities. Recording of cumulative total emissions (kg) for each site listed.
3. Grid reference obtained for each SPRI emission site using address and postcode from search results and <http://www.gridreferencefinder.com>. The  $X$  and  $Y$  co-ordinates were plotted using ArcMap.
4. Simplified modeling of source air pollution by approximating from (1) Met Office wind-roses and (2) air quality scenarios from environmental impact reports on Grangemouth Renewable Plant (2012) and Dunbar Energy from Waste Facility (2008). Three polygons were created using ArcGIS of wind-rose directions (degrees) and predicted distance of pollutants (meters) for SPRI emission sites in (1) Falkirk and Clackmannanshire, (2) Stirling, North Lanarkshire and South Lanarkshire, and (3) Fife.
5. Projection of air pollution polygons layered onto the Central Region datazone layer.

The SPRI emissions data utilized are from declarations made by industrial operators. The SEPA sets wide, flexible criteria for determining the pollutant releases for emitters in SPRI, suggesting some results may be imprecise. Occasional over-estimation errors have also been traced in SPRI, though ostensibly emission values are likely to easily graduate endocrine disruptor pollution between sites.<sup>58</sup> Significances for low datazone sex ratios were calculated by hypothesis testing of two proportions with the second comparator proportion being the national Scotland sex ratio for the same period (2002–2010).

## Results

Figure 1 shows the historical secondary sex ratio linear trend in Scotland from 1865 to 2010. There is a statistically significant downward shift in national birth sex ratio observed starting in 1960 ( $R^2 = 0.243$ ,  $P < 0.01$ ). There is a peak birth sex ratio of 0.519 in 1973. The pattern of declining sex proportions between 1855 and 1905 was similar to the trend of

declining sex ratios observed for the period 1970–2010, though birth registration around this time was not reliable. The change in the sex ratio between 1905 and 1973 reflects a similar pattern as in other industrial countries. An explanation offered for this trend is that improvements in obstetrics and midwifery have led to lower rates of still births.<sup>2</sup> As fetal mortality disproportionately affects male babies, the sex ratio has climbed over this long period. The cumulative secondary sex ratios for health regions in Scotland from 1973 to 2010 (Tables 1 and 2) provide a regional breakdown, which is in temporal alignment with the declining national male birth proportion period. The figures show significant upward skewing of the sex ratio for the Highland region and lower sex ratios for the Borders and Forth Valley. Although the Forth Valley difference in male birth proportion is not significant ( $Z$  score: 1.152), it is potentially noteworthy given that the region's total births constitute over five percent of the national figure.

Table 2 provides “two proportions” testing on health region birth totals for five separate years where Scotland's sex ratio was below 0.510. The three statistically lowest birth sex ratio regions are shown in Table 2. The Forth Valley Health Board area is identified as one of these three lowest regions in four of the 5 years of low national sex ratio. For 2008, the sex ratio value for Forth Valley is statistically significant ( $P < 0.05$ ) when compared to the control sample of Scotland's national ratio from 1974 to 1979. From the years 1973–2010, the Forth Valley region contributes to the downward trajectory of the national sex ratio. The larger Lothian region had the lowest sex proportion at birth for all regions in 2001 and had a significantly lower sex ratio in 2010 compared with the earlier control sample ( $P < 0.01$ ).

In the stratified analysis of sex ratio and SIMD quintiles (Table 3), there is no step-by-step change in sex proportion from lower to higher deprivation categories. There is a significantly elevated male birth/total birth proportion of 0.5235 for the highest quintile, the least deprived and wealthiest sub-group of the Scottish population ( $P < 0.01$ ). This upwards skewing also occurs for the second most deprived group with a sex ratio of 0.5185 ( $P < 0.05$ ). Furthermore, significantly reduced male birth proportion was calculated for quintiles 13 ( $P < 0.05$ ) and 17 ( $P < 0.05$ ).

Figure 3 shows the spatial distribution of modeled endocrine disruptor air pollutants, low skewed sex ratio datazones and datazone SIMD levels for the mid area of the Central Region, which encompasses North Lanarkshire, Falkirk, southern Stirling, western Fife, and the northern area of South Lanarkshire. The air pollution polygons reflect the likely dispersal of pollutants based on local windroses




Figure 1 Sex proportion at birth in Scotland: 1855–2010. Data source: General Register Office for Scotland 2010 (accessed through <http://www.groscotland.gov.uk/statistics/births/births-time-series-data.html>).

and other modeling data. They are gradated according to the cumulative annual (2002, 2004–2010) releases in kilograms recorded on SPRI. Low sex ratio datazones are identified based on cumulative male and total births records from 2001 to 2010, with significance at the low tail of the 90% confidence interval. Multiple deprivation measurement of datazones is by five levels based on equal intervals of vigintile categories (1–4, 5–8, 9–12, 13–16, and 17–20).

The maps display a concentration of low skewed sex ratio datazones for some urban and central rural areas within the region. The map does not show a general spatial concentration of endocrine disruptor air pollution with small areas of low sex ratio or for high levels of multiple deprivation and pollution. However, there are some localities with low sex ratio markers that overlap or are proximate to the polygon models of endocrine disruptor air pollution. Of note are; (1) low sex ratio figures for neighborhoods in the Falkirk local authority area, in Grangemouth close to the industrial complex and Bonnybridge (West of Falkirk) replicating results from an earlier study with

1980s data and (2) the large datazone to the east of Rutherglen, which registers a large number of births with an excess of 230 females per 188 males (birth sex ratio=0.45,  $P=0.010$ ).<sup>59</sup>

## Discussion

A spatio-temporal analysis of sex proportion at birth and secondary data on pollution and deprivation reveal important regional differences, which may contribute to Scotland's declining male birth proportion over the last 40 years. Results also show an upwards skewing in the sex ratio among the least deprived and wealthiest populations. We found no combined effect for pollution and socio-economic status, as there was not a consistent concentration of low sex ratio neighborhoods with endocrine disruptor air pollution and deprivation.

For the low "male birth proportion" regions of Forth Valley and Lothian, it is noteworthy that SEPA stated in their 2008 National Air Quality Report that "the majority of Scotland's industrial oxides of nitrogen are generated in east central Scotland."<sup>60</sup> The current study also found that the

### Central Region Case Study: 6 Local Authority Areas




Figure 2 Central region case study: 6 local authority areas.

**Table 1 Large sample significance testing of sex proportion at birth by health region, 1973–2010**

Health board region	Male births	Total births	Sex proportion at birth	Z score	Two-tail significance test
Ayrshire + Arran	83 308	162 732	0.5119	0.4842	0.6282
Borders	21 015	41 372	0.5080	1.8561	0.0634
Dumfries + Galloway	30 505	59 640	0.5115	0.5159	0.6059
Fife	79 231	154 753	0.5120	0.4354	0.6632
Forth Valley	62 268	121 888	0.5109	1.1522	0.2492
Grampian	116 390	226 240	0.5144	1.7244	0.0846
Greater Glasgow + Clyde	300 875	586 978	0.5126	0.0383	0.9694
Highland (incl. A+B)	67 503	130 407	0.5177	3.5705	0.0004*
Lanarkshire	136 182	265 541	0.5128	0.2854	0.7754
Lothian	173 421	338 582	0.5122	0.3883	0.6978
Orkney, Shetland, and Western Isles	15 738	30 946	0.5086	1.3953	0.1629
Tayside					

Note: "Control" and second largest sample=Yorkshire & Humber 1974–2010 male births/total births (1 185 114/2 312 167).

\*Significant at the 0.01 level.

largest air pollution emitters in Scotland are located in the Forth Valley and Lothian regions. Scotland's only air quality management area for industrial pollution is in Grangemouth in the Forth Valley region, the location of a large manufacturing industrial complex and refinery. The identification of these regional birth sex ratio trends tentatively suggests a possible environmental link to the national male birth proportion adjustment. Localized incidences of a downward sex ratio skewing coupled with high exposure to endocrine disruptor pollution (as shown in Fig. 3) add to this evidence. The 90% significance levels for the neighborhood concentration of low sex ratios however, mean caution must be exercised in drawing such a link. Importantly though, by analyzing data across geographical areas, both

regional (Forth Valley) and local (datazones), the methodological limitations of the "modifiable areal unit" problem is partly averted.

A limitation of the stage (3) analyses is that estimating exposure to pollutants in populations with geographical area-based data is in effect a "moving target." The normal movement of persons and households in and out of neighborhoods and regional areas means that assuming transference of pollution and reproductive effects for a given locality is not straightforward. In Scotland, approximately 13% of all households were classified as 'moving households' in the 2001 Census. However, in North Lanarkshire, South Lanarkshire, Falkirk, Clackmannanshire, and Fife, council area populations are less mobile, with percentages below the national figure.<sup>61</sup> We thus

**Table 2 Large sample significance testing of sex proportion at birth by health region for five 'downward' years of national birth sex ratio**

National/health authority	Male births	Female births	Total births	Sex proportion at birth	Z score	Two-tail significance test
Scotland	33 144	31 962	65 106	0.5091	2.5752	0.01**
Ayrshire + Arran	2401	2401	4802	0.5000	2.0001	0.0455*
Tayside	2411	2390	4801	0.5022	1.6987	0.0894
Highland	1932	1900	3832	0.5042	1.2745	0.2025
Year 1994						
Scotland	31 399	30 257	61 656	0.5093	2.4306	0.0151**
Fife	2028	2082	4110	0.4934	2.6901	0.0071**
Forth Valley	1615	1594	3209	0.5033	1.2691	0.2044
Lanarkshire	3506	3409	6915	0.5070	1.2377	0.2158
Year 2001						
Scotland	26 786	25 741	52 527	0.5099	1.9705	0.0488*
Lothian	4208	4123	8331	0.5051	1.7016	0.0888
Forth Valley	1414	1403	2817	0.5020	1.3291	0.1838
Borders	529	537	1066	0.4962	1.1911	0.2336
Year 2008						
Scotland	30 570	29 471	60 041	0.5092	2.4526	0.0142**
Forth Valley	1711	1743	3454	0.4954	2.2413	0.025*
Borders	558	580	1138	0.4903	1.6291	0.1033
Grampian	3193	3130	6323	0.5050	1.5050	0.1323
Year 2010						
Scotland	29 872	28 919	58 791	0.5081	2.9042	0.0037**
Lothian	4921	4903	9824	0.5009	2.6639	0.0077**
Grampian	3158	3096	6254	0.5050	1.5008	0.1334
Forth Valley	1672	1658	3330	0.5021	1.4270	0.1536

Note: "Control" and second largest sample for Z score=male births/total births for Scotland from 1974 to 1979 (204 745/397 934).

\*Significant at the 0.05 level.

\*\*Significant at the 0.01 level.

hypothesize that mobilization of the population was minimal and that the study populations have likely been living in the study sites and exposed to endocrine disruptors for enough of their reproductive life to potentially impact the primary and secondary sex ratios. Further, the English control region, although similar in several respects in terms of population size, range of industrial activity, and urban/rural settings, is also not identical to Scotland. The available environmental data relate to specific point sources with possible under or over-estimates of releases, and not all endocrine disruptor releases were measured or recorded. Biological monitoring or modeling of exposures is not available.

Area-based measurement can also be imprecise as not all persons residing in an area display the characteristics shown by the indicator, i.e. individuals experiencing deprivation can live in non-deprived areas. There is also a difficulty in temporally aligning all the indicators from the data sets. For example, the SPRI begins in 2002 and no releases were published for 2003. Despite these shortcomings, area-based measurement is predominantly how public health and socio-economic data are constructed in Scotland. Meanwhile, in Sweden, individually tabulated medical and demographic data sets permit comprehensive ecological studies.<sup>47,62</sup> Given the global extent of reproductive health changes observed, countries with different approaches to public health data collection must also be documented. A crucial advantage of utilizing geographical area-based measurement is that it facilitates a *comprehensive* inquiry, both spatially and temporally, of secondary sex ratio change in national and regional populations, rather than by acute events or selected cohort groups.

It is also feasible that confounding, in the form of other pollutants that are not recorded as endocrine disruptors, may be present in the spatial analysis. Recent research undertaken in the Teplice area of the Czech Republic on potential male reproductive effects from exposure to heavy concentrations of sulfur-polluted air is instructive.<sup>63</sup> Damage to the chromatin of sperm initially mediated through environmental chemicals and then transferred genetically through the paternal line an additional hypothesis for consideration on the sex ratio question. Environmentally mediated genetic damage to sperm has been proffered as an alternative explanation to the endocrine disruptor hypothesis for rises in testicular cancers and declines in sperm quality.<sup>64</sup> Exposure to pollutants not identified as endocrine disruptors may therefore compromise some of the spatial concentrations of pollution and low sex ratio flagged in this study. In common with both hypotheses, however, the mediation of chronic long-term exposure of pollution, potentially over generations, and including mixing of low doses of chemicals, requires additional scientific research. Further, the inclusion of secondary sex ratio monitoring with new methods of environmental assessment, on cumulative impacts from pollution, is also worthy of consideration.<sup>65,66</sup> It is noteworthy that in the context of long standing proliferation of industrial pollution in the Central Region, the Lanarkshire and Forth Valley health regions display the highest crude rates (9.9 and 8.8) of cancer of the testis (per 100 000 person-years at risk) in Scotland for the 2004–2008 period.

Furthermore, studies also suggest that increasing male age is associated with greater sperm DNA fragmentation as well as greater probability of a

**Table 3 Large Sample Significance Testing of Sex Proportion at Birth 2001 to 2009 by Datazone SIMD Vigintile**

Vigintile	Deprivation	Male births	Total births	Sex proportion at birth	Z score	Two-tail significance test
1	Most deprived	17 529	34 272	0.5115	0.3297	0.7417
2		16 736	32 278	0.5185	2.1355	0.0327*
3		15 081	29 401	0.5129	0.1862	0.8523
4		13 636	26 695	0.5108	0.5035	0.6146
5		13 208	25 865	0.5107	0.5449	0.5858
6		13 087	25 535	0.5125	0.0399	0.9682
7		12 685	24 983	0.5077	1.4345	0.1514
8		12 245	23 787	0.5148	0.7228	0.4698
9		11 880	23 133	0.5136	0.3480	0.7278
10		12 130	23 593	0.5141	0.5269	0.5982
11		11 943	23 379	0.5108	0.4618	0.6442
12		11 245	21 856	0.5145	0.6148	0.5387
13		10 830	21 422	0.5056	1.9614	0.0498*
14		12 004	23 361	0.5138	0.4383	0.6612
15		12 490	24 180	0.5165	1.2663	0.2054
16		11 963	23 529	0.5084	1.1863	0.2355
17		11 138	22 041	0.5053	2.0537	0.04*
18		12 416	24 285	0.5113	0.3424	0.732
19		11 211	22 129	0.5066	1.6816	0.0926
20	Least deprived	11 206	21 404	0.5235	3.2078	0.0013**

Note: "Control" and second largest sample= male births/total births for Yorkshire & Humber 2001–2009 (281 708/549 798).

\*Significant at the 0.05 level.

\*\*Significant at the 0.01 level.



Central Scotland - Cumulative Endocrine Disruptor Pollution (2002, 2004-2010),  
SIMD (2006) Vigintile and Low Secondary Sex Ratios (2002 to 2010) by Datazone




Figure 3 Central Scotland: cumulative endocrine disruptor pollution (2002, 2004–2010). SIMD (2006) vigintile and low secondary sex ratios (2002–2010) by datazone.

partner's miscarriage.<sup>50,51</sup> With the social trend of delayed fatherhood for many men in industrialized countries, increasing paternal age as a contributing factor to the declining male birth proportion is another reasonable hypothesis. Future environmental studies of sex ratio skewing following post-hazardous events and/or heavily polluted areas should consider parental age and stress as confounding factors.

With regard to multiple deprivation, the "spike" in sex ratio for the wealthiest SIMD group would initially seem to reinforce the evolutionary psychological theory of Trivers–Willard described earlier. However, it is contradicted by the stratified results for deprivation and the sex ratio not displaying a graded trend across the other 19 groups. Furthermore, there was a significant increase in the sex ratio for highly deprived group (vignette 2). The pollution and SIMD mapping show no evidence of a combined "environmental deprivation" effect on secondary sex ratio. Genomic imprinting during embryogenesis before clinically recognized pregnancy may be a key reproductive process to examine.<sup>67</sup> Adoption and advancement of reproductive medicine and technologies in Western countries may also be linked. There is anecdotal evidence of richer American couples purchasing pre-implantation genetic diagnosis to balance females in families, though empirical research does not indicate any bias towards either daughters or sons among couples.<sup>68,69</sup> In the UK, non-medical sex selection is illegal but may occur. The ethical debate on pre-implantation genetic diagnosis and sperm sorting practices is ongoing.<sup>70</sup>

Relevant and available socio-economic and public health, pollution inventory emissions, and birth sex ratios data sets in Scotland were brought together using GIS. Their analysis provides further tentative evidence for the hypothesis that declines in male birth proportion in some industrialized countries are possibly linked to chronic exposure to pollution by endocrine disrupting compounds. The historical and regional trends and localized skewing of sex ratios found in this study are in line with other studies linking male birth proportion decline with air pollution. The findings have implications for the availability, frequency, and range of future environmental pollution monitoring. They further highlight the importance of continuity and consistency in such sets for meaningful time series analysis to inform health protection policy, especially of vulnerable populations exposed to a range of environmental and socio-economic conditions that may affect ratios. This study provides a foundation for further work to examine the possible relationships between social, economic and environmental factors in a diverse demographic setting with regard to sex proportions. Advanced geo-statistical analyses can be used to help

shed light both on public health issues linked to reproductive health and the significance of particular exposures to pollution and the related effectiveness of environmental regulations in protecting reproductive health.

### Disclaimer statements

**Contributors** EM completed the analysis and interpretation of the data and drafted the manuscript. AW conceived the study, and jointly edited the manuscript with EM. AW and AT supervised the study on which the paper is based. AT contributed to editing of the manuscript. JM assisted with mapping and contributed intellectually to spatial analysis in the study. MS assisted with statistical review of the analysis described in the manuscript. All authors reviewed the manuscript.

**Funding** The research was funded through a joint PhD studentship from the School of Nursing Midwifery and Health, and School of Natural Sciences, at the University of Stirling.

**Conflicts of interest** The authors declare that they have no competing interests.

**Ethics approval** No ethical approval required.

### Acknowledgements

The authors wish to acknowledge the Scottish Environmental Protection Authority for the use of Pollutant Release inventory information. The Scottish Environment Protection Agency reserves all copyright and database rights. The authors also acknowledge the helpful contribution of General Register Office (Scotland), in particular David Shelton, for guidance on appropriate birth statistics.

### References

- 1 van Larebeke NA, Sasco AJ, Brophy JT, Keith MM, Gilbertson M, Watterson A. Sex ratio changes as sentinel health events of endocrine disruption. *Int J Occup Environ Health*. 2008;14(2):138–43.
- 2 Davis DL, Gottlieb MB, Stampnitzky JR. Reduced ratio of male to female births in several industrial countries — a sentinel health indicator? *JAMA*. 1998;279(13):1018–23.
- 3 Skakkebaek NE, Rajpert-De Meyts E, Main KM. Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Hum Reprod*. 2001;16(5):972–8.
- 4 Swan SH, Elkin EP, Fenster L. The question of declining sperm density revisited: an analysis of 101 studies published 1934–1996. *Environ Health Perspect*. 2000;108(10):961–6.
- 5 Schoeters G, Den Hond E, Dhooze W, van Larebeke N, Leijts M. Endocrine disruptors and abnormalities of pubertal development. *Basic Clin Pharmacol Toxicol*. 2008;102(2):168–75.
- 6 Virtanen HE, Adamsson A. Cryptorchidism and endocrine disrupting chemicals. *Mol Cell Endocrinol*. 2012;355(2):208–20.
- 7 Sharpe RM. Environmental/lifestyle effects on spermatogenesis. *Philos Trans R Soc B Biol Sci*. 2010;365(1546):1697–712.
- 8 Smith D, von Behren J. Trends in the sex ratio of California births, 1960–1996. *J Epidemiol Community Health*. 2005;59(12):1047–53.
- 9 Martuzzi M, Di Tanno N, Bertollini R. Declining trends of male proportion at birth in Europe. *Arch Environ Health*. 2001;56(4):358–64.

- 10 Zeitlin J, Saurel-Cubizolles MJ, de Mouzon J, Rivera L, Ancel P, Blondel A. Fetal sex and preterm birth: are males at greater risk? *Hum Reprod.* 2002;17(10):2762–8.
- 11 Miliaras DE, K, Meditskou S. The sex ratio of fetal deaths and its potential relation with the decline of sex ratio at birth in Greece. *Arist Univ Med J.* 2011;38(3):8.
- 12 Bruckner TA, Catalano R, Ahern J. Male fetal loss in the US following the terrorist attacks of September 11, 2001. *BMC Public Health.* 2010;10:273.
- 13 Mealey L, Mackey W. Variation in offspring sex-ratio in women of differing social-status. *Ethol Sociobiol.* 1990;11(2):83–95.
- 14 Trivers RL, Willard DE. Natural selection of parental ability to vary the sex ratio of offspring. *Science.* 1973;179(4068):90–2.
- 15 Ellis L, Bonin S. Social status and the secondary sex ratio: new evidence on a lingering controversy. *Soc Biol.* 2002;49(1–2):35–43.
- 16 Klinger HP, Glasser M. Contraceptives and the conceptus: sex of the fetus and neonate after oral-contraceptive use. *Contraception.* 1981;23(4):367–74.
- 17 Grant V, Wise L, Palmer J, Hatch E, Troisi R, Titus-Ernstoff L, Herbst A. Maternal personality, evolution and the sex ratio. London: Routledge; 1998.
- 18 Wise LA, Palmer JR, Hatch EE, Troisi R, Titus-Ernstoff L, Herbst A. Secondary sex ratio among women exposed to diethylstilbestrol in utero. *Environ Health Perspect.* 2007;115(9):1314–9.
- 19 Mocarelli P, Brambilla P, Gerthoux PM, Patterson DG, Needham LL. Change in sex ratio with exposure to dioxin. *Lancet.* 1996;348(9024):409.
- 20 Matsuo K, Ushioda N, Udoff LC. Parental aging synergistically decreases offspring sex ratio (vol 35, pg 164, 2009). *J Obstet Gynaecol Res.* 2009;35(2):3911.
- 21 Ulizzi L, Zonta LA. Factors affecting the sex-ratio in humans — multivariate-analysis of the Italian population. *Hum Biol.* 1995;67(1):59–67.
- 22 Mills JL, England L, Granath F, Cnattingius S. Cigarette smoking and the male-female sex ratio. *Fertil Steril.* 2003;79(5):1243–5.
- 23 Mathews F, Johnson PJ, Neil A. You are what your mother eats: evidence for maternal preconception diet influencing foetal sex in humans. *Proc R Soc B Biol Sci.* 2008;275(1643):1661–8.
- 24 Bonde JP, Wilcox A. Ratio of boys to girls at birth — is not related to the time taken to conceive, or exposure to environmental agents. *Br Med J.* 2007;334(7592):486–7.
- 25 Gluckman PD, Hanson MA, Beedle AS. Early life events and their consequences for later disease: a life history and evolutionary perspective. *Am J Hum Biol.* 2007;19(1):1–19.
- 26 Weisskopf MG, Anderson HA, Hanrahan LP and the Great Lakes Consortium. Maternal exposure to Great Lakes sport-caught fish and dichlorodiphenyl dichloroethylene, but not polychlorinated biphenyls, is associated with reduced birth weight. *Environ Res.* 2005;97(2):149–62.
- 27 Karmaus W, Huang SY, Cameron L. Parental concentration of dichlorodiphenyl dichloroethene and polychlorinated biphenyls in Michigan fish eaters and sex ratio in offspring. *J Occup Environ Med.* 2002;44(1):8–13.
- 28 Hertz-Picciotto I, Jusko TA, Willman EJ, Baker R, Teplin S, Charles M. A cohort study of in utero polychlorinated biphenyl (PCB) exposures in relation to secondary sex ratio. *Environ Health.* 2008;7:37.
- 29 Gallo V, Egger M, McCormack V, Farmer P, Kirsch-Volders M, Matullo G. Strengthening the reporting of Observational studies in Epidemiology-Molecular Epidemiology (STROBE-ME): an extension of the STROBE statement. *Eur J Epidemiol.* 2011;26(10):797–810.
- 30 Mocarelli P, Gerthoux PM, Ferrari E, Patterson D, Kieszak S, Brambilla P. Paternal concentrations of dioxin and sex ratio of offspring. *Lancet.* 2000;355(9218):1858–63.
- 31 Potashnik G, Goldsmith J, Insler V. Dibromochloropropane-induced reduction of the sex-ratio in man. *Andrologia.* 1984;16(3):213–8.
- 32 Jacobsen R, Bostofte E, Engholm G, Hansen J, Skakkebaek NE, Moller H. Fertility and offspring sex ratio of men who develop testicular cancer: a record linkage study. *Hum Reprod.* 2000;15(9):1958–61.
- 33 Khanjani N, Sim MR. Maternal contamination with PCBs and reproductive outcomes in an Australian population. *J Expo Sci Environ Epidemiol.* 2007;17(2):191–5.
- 34 Jarrell JF, Weisskopf MG, Weuve J, Tellez-Rojo MM, Hu H, Hernandez-Avila M. Maternal lead exposure and the secondary sex ratio. *Hum Reprod.* 2006;21(7):1901–6.
- 35 Gibson G, Koifman S. Agricultural toxic use and temporal distribution of male birth rate in the state of Parana, Brazil. *Revista Panamericana De Salud Publica-Pan American Journal of Public Health.* 2008;24(4):240–7.
- 36 Viel JF, Floret N, Mauny F. Spatial and space-time scan statistics to detect low rate clusters of sex ratio. *Environ Ecol Stat.* 2005;12(3):289–99.
- 37 Lloyd OL, Smith G, Lloyd MM, Holland Y, Gailey F. Raised mortality from lung-cancer and high sex-ratios of births associated with industrial-pollution. *Br J Ind Med.* 1985;42(7):475–80.
- 38 Garfinkel J, Selvin S. Multivariate-analysis of relationship between parental age and birth-order and human secondary sex-ratio. *J Biosoc Sci.* 1976;8(2):113–21.
- 39 Jacobsen R, Moller H, Mouritsen A. Natural variation in the human sex ratio. *Hum Reprod.* 1999;14(12):3120–5.
- 40 Moran PAP, Novitski E, Novitski C. Paternal age and secondary sex ratio in humans. *Ann Hum Genet.* 1969;32:315.
- 41 Nicolich MJ, Huebner WW, Schnatter AR. Influence of parental and biological factors on the male birth fraction in the United States: an analysis of birth certificate data from 1964 through 1988. *Fertil Steril.* 2000;73(3):487–92.
- 42 Ruder A. Paternal-age and birth-order effect on the human secondary sex-ratio. *Am J Hum Genet.* 1985;37(2):362–72.
- 43 Fukuda M, Fukuda K, Shimizu T, Moller H. Decline in sex ratio at birth after Kobe earthquake. *Hum Reprod.* 1998;13(8):2321–22.
- 44 Catalano R, Bruckner T, Marks AR, Eskenazi B. Exogenous shocks to the human sex ratio: the case of September 11, 2001 in New York City. *Hum Reprod.* 2006;21(12):3127–31.
- 45 Obel C, Henriksen TB, Secher NJ, Eskenazi B, Hedegaard M. Psychological distress during early gestation and offspring sex ratio. *Hum Reprod.* 2007;22(11):3009–12.
- 46 Khashan AS, Mortensen PB, McNamee R, Baker PN, Abel KM. Sex ratio at birth following prenatal maternal exposure to severe life events: a population-based cohort study. *Hum Reprod.* 2009;24(7):1754–7.
- 47 Hansen D, Moller H, Olsen J. Severe periconceptional life events and the sex ratio in offspring: follow up study based on five national registers. *Br Med J.* 1999;319(7209):548–9.
- 48 Catalano R, Bruckner T, Gould J, Eskenazi B, Anderson E. Sex ratios in California following the terrorist attacks of September 11, 2001. *Hum Reprod.* 2005;20(5):1221–7.
- 49 Maconochie N, Doyle P, Prior S, Simmons R. Risk factors for first trimester miscarriage — results from a UK-population-based case-control study. *BJOG.* 2007;114(2):170–86.
- 50 Kleinhans K, Perrin M, Friedlander Y, Paltiel O, Malaspina D, Harlap S. Paternal age and spontaneous abortion. *Obstet Gynecol.* 2006;108(2):369–77.
- 51 Sartorius GA, Nieschlag E. Paternal age and reproduction. *Hum Reprod Update.* 2010;16(1):65–79.
- 52 Auger N, Daniel M, Platt RW, Wu YQ, Luo ZC, Choiniere R. Association between perceived security of the neighbourhood and small-for-gestational-age birth. *Paediatr Perinat Epidemiol.* 2008;22(5):467–77.
- 53 Elo IT, Culhane JF, Kohler IV, O'Campo P, Burke J, Messer L. Neighbourhood deprivation and small-for-gestational-age term births in the United States. *Paediatr Perinat Epidemiol.* 2009;23(1):87–96.
- 54 Farley TA, Mason K, Rice J, Habel JD, Scribner R, Cohen DA. The relationship between the neighbourhood environment and adverse birth outcomes. *Paediatr Perinat Epidemiol.* 2006;20(3):188–200.
- 55 Freese J, Powell B. Making love out of nothing at all? Null findings and the Trivers-Willard hypothesis. *Am J Sociol.* 2001;106(6):1776–88.
- 56 Centre for the History of Medicine. Scottish History of Medicine (ND) Scottish way of birth and death. Glasgow: University of Glasgow. Available from: <http://www.gla.ac.uk/schools/socialpolitical/research/economicsocialhistory/historymedicine/scottishwayofbirthanddeath/death/stillbirths/date> accessed 27.05.14.
- 57 Bailey NF, Goodlad R, Shucksmith M, Fitzpatrick S, Pryce G. Measuring deprivation in Scotland: developing a long-term strategy. Glasgow: Scottish Centre for Research; 2003.
- 58 Edwards R. SEPA: The Sleeping Watchdog. *Herald Scotland.* 24 July 2011.

- 59 Williams FL, Lawson AB, Lloyd OL. Low sex-ratios of births in areas at risk from air-pollution from incinerators, as shown by geographical analysis and 3-dimensional mapping. *Int J Epidemiol.* 1992;21(2):311–9.
- 60 Scottish Environmental Protection Authority. SEPA's national air quality report. Stirling: SEPA; 2008.
- 61 GRO Scotland. Scotland's Census 2001: statistics on moving households and moving groups. Edinburgh: GRO Scotland; 2007.
- 62 Magnuson A, Bodin L, Montgomery SM. Father's occupation and sex ratio of offspring. *Scand J Public Health.* 2007;35(5):454–9.
- 63 Rubes J, Selevan SG, Evenson DP, Zudova D, Vozdova M, Zudova Z. Episodic air pollution is associated with increased DNA fragmentation in human sperm without other changes in semen quality. *Hum Reprod.* 2005;20(10):2776–83.
- 64 Joffe M. What has happened to human fertility? *Hum Reprod.* 2010;25(2):295–307.
- 65 August LM, Faust JB, Cushing L, Zeise L, Alexeeff GV. Methodological considerations in screening for cumulative environmental health impacts: lessons learned from a pilot study in California. *Int J Environ Res Public Health.* 2012;9(9):3069–84.
- 66 Alexeeff GV, Faust JB, August LM, Milanec C, Randles K, Zeise L. A screening method for assessing cumulative impacts. *Int J Environ Res Public Health.* 2012;9(2):648–59.
- 67 Boklage CE. The epigenetic environment: secondary sex ratio depends on differential survival in embryogenesis. *Hum Reprod.* 2005;20(3):583–7.
- 68 Hvistendahl, M. *Unnatural selection.* New York: Public Affairs; 2011.
- 69 Sharp RR, McGowan ML, Verma JA, Landy D, McAdoo S, Carson S. Moral attitudes and beliefs among couples pursuing PGD for sex selection. *Reprod Biomed Online.* 2010;21(7):838–47.
- 70 Strange H, Cesagen EC. A. Non-medical sex selection: ethical issues. *Br Med Bull.* 2010;94(1):7–20.