



Published in final edited form as:

*J Mol Cell Cardiol.* 2011 December ; 51(6): 894–901. doi:10.1016/j.yjmcc.2011.08.026.

## Upregulation of Cardiomyocyte Ribonucleotide Reductase Increases Intracellular 2 deoxy-ATP, Contractility, and Relaxation

F. Steven Korte, Ph.D.<sup>1,5</sup>, Jin Dai, Ph.D.<sup>1</sup>, Kate Buckley, B.S.<sup>1</sup>, Erik R. Feest, B.S.<sup>1</sup>, Nancy Adamek, Ph.D.<sup>4</sup>, Michael A. Geeves, Ph.D.<sup>4</sup>, Charles E. Murry, M.D., Ph.D.<sup>1,2,3,5</sup>, and Michael Regnier, Ph.D.<sup>1,5</sup>

<sup>1</sup>Department of Bioengineering, University of Washington, Seattle WA 98195 USA

<sup>2</sup>Department of Pathology, University of Washington, Seattle WA 98195 USA

<sup>3</sup>Department of Medicine/Cardiology, University of Washington, Seattle WA 98195 USA

<sup>4</sup>Department of Biosciences, University of Kent, Canterbury, Kent CT2 7NJ UK

<sup>5</sup>Center for Cardiovascular Biology, Institute for Stem Cell and Regenerative Medicine, University of Washington, Seattle WA 98109 USA

### Abstract

We have previously demonstrated that substitution of ATP with 2 deoxy-ATP (dATP) increased the magnitude and rate of force production at all levels of Ca<sup>2+</sup>-mediated activation in demembrated cardiac muscle. In the current study we hypothesized that cellular [dATP] could be increased by viral-mediated over expression of the ribonucleotide reductase (Rrm1 and Rrm2) complex, which would increase contractility of adult rat cardiomyocytes. Cell length and ratiometric (fura2) Ca<sup>2+</sup> fluorescence were monitored by video microscopy. At 0.5 Hz stimulation, the extent of shortening was increased ~40% and maximal rate of shortening was increased ~80% in cardiomyocytes overexpressing Rrm1+Rrm2 as compared to non-transduced cardiomyocytes. The maximal rate of relaxation was also increased ~150% with Rrm1+Rrm2 over expression, resulting in decreased time to 50% relaxation over non-transduced cardiomyocytes. These differences were even more dramatic when compared to cardiomyocytes expressing GFP-only. Interestingly, Rrm1+Rrm2 over expression had no effect on minimal or maximal intracellular [Ca<sup>2+</sup>] (Fura2 fluorescence), indicating increased contractility is primarily due to increased myofilament activity without altering Ca<sup>2+</sup> release from the sarcoplasmic reticulum. Additionally, functional potentiation was maintained with Rrm1+Rrm2 over expression as stimulation frequency was increased (1 Hz and 2 Hz). HPLC analysis indicated cellular [dATP] was increased by approximately 10-fold following

Open Access under [CC BY 3.0](https://creativecommons.org/licenses/by/3.0/) license.

Corresponding Author: Michael Regnier, Ph.D., University of Washington Department of Bioengineering, Institute for Stem Cell and Regenerative Medicine, Box 355061, Seattle, WA 98195-7962 USA, Phone (206) 616-4325 Fax (206) 685-3300, mregnier@u.washington.edu.

### DISCLOSURES

Dr. Regnier holds a provisional patent application (UW ref. 45511.01US1) on RIR2 over expression to improve cardiac contractile function.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

transduction, becoming ~1.5% of the adenine nucleotide pool. Furthermore, 2% dATP was sufficient to significantly increase crossbridge binding and contractile force during sub-maximal  $\text{Ca}^{2+}$  activation in demembranated cardiac muscle. These experiments demonstrate the feasibility of directly targeting the actin-myosin chemomechanical crossbridge cycle to enhance cardiac contractility and relaxation without affecting minimal or maximal  $\text{Ca}^{2+}$ .

## Keywords

Cardiomyocyte; Contractility; Ribonucleotide Reductase

---

## 1. INTRODUCTION

Systolic and/or diastolic cardiac function is compromised in a number of cardiovascular diseases including myocardial infarction, ischemia/reperfusion injury, diabetes, high blood pressure and hypertrophic and dilated cardiomyopathy. These pathophysiological conditions often involve alterations in the  $\text{Ca}^{2+}$  cycle<sup>[1]</sup>,  $\beta$ -adrenergic responsiveness<sup>[2]</sup>, and/or the contractile apparatus of cardiomyocytes<sup>[3, 4]</sup>. To date, therapeutic efforts have focused primarily on approaches that increase  $[\text{Ca}^{2+}]_i$  that can be pro-arrhythmogenic and may impair ventricular filling by slowing diastolic relaxation<sup>[5]</sup>. Other approaches involving adrenergic agents can have undesirable long-term consequences, including significant side-effects due to drug actions in non-target areas, pro-arrhythmogenic triggered activity, and potential for accelerated progression into heart failure<sup>[2]</sup>. Thus, new approaches to combat cardiac dysfunction are desirable.

We have previously shown that replacing ATP with 2 deoxy-ATP (dATP) as the substrate for contraction of demembranated cardiac muscle increased isometric force and the rate of force development and shortening at all levels of  $\text{Ca}^{2+}$  activation, including saturating  $[\text{Ca}^{2+}]_i$  (pCa 4.0) <sup>[6–9]</sup>. The presence of dATP results in enhanced myosin binding to actin and an increase in the rates of Pi and dADP release and myosin detachments. As such, contractile properties can be improved by >50% over the range of  $[\text{Ca}^{2+}]_i$  seen *in vivo*. Thus, replacement of ATP with dATP offers the potential to improve contraction independent of changes in  $[\text{Ca}^{2+}]_i$  or adrenergic signaling.

To date, the effect of dATP has only been studied in demembranated cardiac tissue and with isolated contractile proteins. As such, its potential to improve intact cardiomyocyte contraction or cardiac function *in situ* is unknown. Cellular production of dATP occurs in the cytoplasm of mammalian cells by ribonucleotide reductase (Rrm), which removes a hydroxyl moiety from the 2-position on the ribose ring of ADP to produce dADP. dADP is then rapidly converted to dATP. Rrm consists of two subunit proteins, a catalytic activator (Rrm1) and free radical containing (Rrm2) subunit and is regulated by nucleoside triphosphate allosteric effectors<sup>[10]</sup>. While the details of regulating cellular RR content, enzymatic activity and cellular concentration  $[\text{dATP}]$  are unclear, it is known that both subunits are necessary for activity.<sup>[11]</sup>

In the current study, we produced adenoviral vectors expressing cytomegalovirus (CMV) promoter driven Rrm1 or Rrm2, each along with green fluorescent protein (GFP) as a

transduction reporter. Cultured adult rat cardiomyocytes were transduced with these vectors, and the rate and extent of myocyte contraction and relaxation and  $\text{Ca}^{2+}$  transient rise and decay (Fura2 fluorescence) were monitored by video microscopy following a 48 hour viral incubation period. Here we show that these treatments significantly increased cellular [dATP], rate and extent of shortening, and rate of relaxation, with minimal effects on  $\text{Ca}^{2+}$  transients, at 0.5 Hz, 1 Hz and 2 Hz stimulation. Additionally, the [dATP] found in transduced cells (1–2% of adenine nucleotide content) was sufficient to increase sub-maximal  $\text{Ca}^{2+}$  activated force in skinned cardiac trabeculae. These experiments suggest that increases in cardiac intracellular Rrm and/or the dATP pool can significantly alter the actin-myosin crossbridge cycle to enhance cardiac contractility without impairing diastolic function or cardiomyocyte  $\text{Ca}^{2+}$  handling.

## 2. METHODS

Greater details of plasmid design and vector production, cell culture, contractile assessment, nucleotide binding affinity, and western blot analysis are provided in online supporting information.

### 2.1 Animal and Tissue Handling

These studies were approved by the University of Washington (UW) Animal Care Committee and conducted in accordance with federal guidelines. Animals were cared for in accordance with US NIH Policy on Humane Care and Use of Laboratory Animals in the Department of Comparative Medicine at UW. Adult rat (Fischer 344) cardiomyocytes (ARCs) were isolated from heart using aortic retrograde perfusion for enzymatic (collagenase/protease) dispersion of cells[12]. Neonatal Rat Cardiomyocytes (NRCs) were isolated by enzymatic dispersion from 1–3-day old newborn Fischer 344 rats as previously described[13]. Rat cardiac trabeculae were dissected from the right ventricle of male Sprague-Dawley rats, chemically demembrated, and prepared for mechanical measurements as previously described [13]. Trabeculae averaged  $1.3 \pm 0.2$  mm in length by  $170 \pm 30$   $\mu\text{m}$  in width.

### 2.2 Plasmid design and virus production

HEK293 cells were used to generate adenoviral vectors<sup>[14]</sup> expressing Rrm1 or Rrm2 from the CMV promoter. Both vectors contained a second expression cassette for green fluorescent protein (GFP) as a transduction reporter protein, and we also expressed a vector for GFP-only. Virus was introduced to cardiomyocytes at ~250 particles per cell.

### 2.3 Nucleotide Binding Affinity

Rapid kinetic measurements of nucleotide binding and actin-myosin dissociation were taken at 10°C and 20°C (Hi-Tech Scientific SF-61 DX2 stopped-flow system) as previously described[15] using pyrene labeled actin and myosin S1. Myosin was purified from mouse hearts, rabbit soleus, and rabbit bulk fast muscle as previously described[16, 17]. Actin was purified from rabbit skeletal muscle<sup>[18]</sup>. The stopped-flow transients were fitted to one or two exponentials by non-linear least squares curve fitting using the Kinetic Studio software (TgK Scientific). All experiments were carried out in 20 mM Cacodylate buffer, pH 7.0

containing 100 mM KCl, and 5 mM MgCl<sub>2</sub>. The rate constant for ATP-induced actin-S1 dissociation ( $k_{\text{obs}}$ ) was determined from equation 1 based on SI Scheme 1:

$$k_{\text{obs}} = k_{+2} K_1 \frac{[\text{ATP}]}{1 + K_1 [\text{ATP}]} \quad \text{Equation 1}$$

## 2.4 Contractile Assessments

In modified Tyrodes buffer at ambient temperature (22–24°C) and at 37°C, cell shortening and relaxation of arbitrarily selected stimulated cardiomyocytes was recorded using IonOptix system video microscopy. (IonOptix, Milton, MA, USA). Calcium transients induced by electrical stimulation were measured in Fura2 loaded cells using IonOptix equipment as described [19]. Fura2 fluorescence was measured using an IonOptix spectrophotometer (Stepper Switch) attached to a fluorescence microscope. Emitted Fura2 fluorescence was collected by the 40X objective, passed through a 510nm filter and detected by a photomultiplier tube. For demembrated trabeculae, steady-state force and high frequency sinusoidal stiffness (to determine crossbridge binding) were measured in a custom built mechanical apparatus at 15° and 22° C during sub-maximal (pica 5.6) and maximal (pica 4.0) Ca<sup>2+</sup> activation as previously described [20]. Experimental physiological Ca<sup>2+</sup> solutions were calculated as previously described for trabeculae mechanics [21].

## 2.5 Data Processing and Statistical Analysis

Maximal cardiomyocyte shortening and relengthening and calcium transient rise and decay were calculated offline using IonOptix software to determine the maximum of the first derivative of these transients. Times to peak shortening and 50% and 90% return to baseline were also calculated offline. Statistical differences were determined by ANOVA, with Student-Newman-Keuls as a *post-hoc* pairwise test (SigmaPlot 11). Trabeculae were compared using paired t-tests. Differences at p-value < 0.05 were considered statistically significant. Data is displayed as mean ± s.e.m.

## 3. RESULTS

Transduction with recombinant adenovirus containing appropriate coda constructs driven by the CMV promoter was used to induce over expression of muscle rib nucleotide reeducates 1 (Rrm1) and 2 (Rrm2) in cultured adult and neonatal rat cardiomyocytes. Each adenovirus also contained a second expression cassette for green fluorescent protein (GFP), which was used as a reporter protein identifying successful transduction. Cardiomyocytes were infected with adenovirus containing genes for [Rrm1 + GFP and Rrm2 + GFP] or [GFP] for 2 days. Successful gene transfer, grossly indicated by green fluorescence with microscopy, indicated nearly 100% transduction efficiency (Supplemental Figure 1). This is consistent with previous studies using cardiomyocytes [22]. Cell survival over this period was similar for all groups, including non-transduced control cells, suggesting these viral vectors did not compromise cardiomyocyte viability. Cardiomyocyte numbers and sarcomere lengths are summarized in Table 1. There was no difference in resting sarcomere length between groups,

indicating that over expression of Rrm1+Rrm2 (or GFP) did not increase calcium independent activation.

### 3.1 Contractile Analysis of Cultured Cardiomyocytes

The effects of Rrm1+Rrm2 over expression on extent and rate of stimulated shortening-relengthening of adult rat cardiomyocytes were determined using video length-detection (IonOptix). Figure 1a shows representative shortening traces, and Figure 1b shows representative  $\text{Ca}^{2+}$  transients (Fura2 fluorescence), for non-transduced (black), GFP-only (green), and Rrm1+Rrm2 (red) transduced cardiomyocytes. The data for all measurements at 0.5 Hz is summarized in Table 2. Cardiomyocytes transduced with Rrm1+Rrm2 (+GFP) had a significantly greater magnitude and rate of shortening vs. non-transducer cardiomyocytes and GFP-only transducer controls. This is illustrated in Figure 1c, which shows % differences in rate and extent of shortening, relaxation rate, and time to 50% and 90% relaxation. While GFP has been reported to have a deleterious<sup>[23]</sup> or no effect<sup>[19, 24]</sup> on contractility, it did not appear to act as a contractile inhibitor in this study. However, GFP did slow the 90% relaxation time, which was accompanied by a slower time to 50% and 90% decay of the  $\text{Ca}^{2+}$  transient (Figure 1c, Table 2). Regardless, Rrm1+Rrm2 over expression increased the rate of relaxation and decreased the time to 50% relaxation, and this effect may be somewhat underestimated due to the presence of GFP. Figure 1d illustrates the % difference in  $\text{Ca}^{2+}$  transient properties, including minimal and maximal  $\text{Ca}^{2+}$ , and the time to 50% and 90%  $\text{Ca}^{2+}$  decay. There was no significant effect from either GFP or Rrm1+Rrm2 + GFP on minimal and maximal  $\text{Ca}^{2+}$ , indicating that enhanced contractility with Rrm1+Rrm2 was primarily due to increased myofilament responsiveness to activating  $\text{Ca}^{2+}$ . Interestingly, Rrm1+Rrm2 over expression did speed  $\text{Ca}^{2+}$  re-sequestration, as indicated by a reduction the time to 50% and 90% decay. This could be due to increased SERCA activity and, in part, explain the increased maximal rate of cardiomyocyte relaxation at 0.5 Hz (Table 2) stimulation. Faster relaxation in Rrm1+Rrm2 overexpressing cardiomyocytes could also be due to faster crossbridge cycling, that leads to shortening induced thin filament inactivation and  $\text{Ca}^{2+}$  release from troponin C.

It is important to determine whether Rrm1+Rrm2 over expression affects normal cellular response to increased stimulation frequency, as changes in heart rate are a normal physiological adaptation to systemic demand. Figure 2 summarizes the effect of increased stimulation frequency (0.5 to 1 to 2 Hz) on fractional shortening (2a), shortening velocity (2b), relaxation velocity (2c), and time to 90% relaxation (2c). The contractile response to stimulation frequency was similar between groups, and Rrm1+Rrm2 transducer cardiomyocytes maintained functional potentiation at all frequencies. Importantly, increased pacing frequency is associated with a positive lusitropic effect, shortening the time to 90% relaxation in all groups. There was little difference in non-transducer myocytes vs. GFP-only transducer myocytes, except that time to 90% relaxation is longer at 0.5 Hz in GFP-only myocytes. The effect of stimulation frequency on  $\text{Ca}^{2+}$  transients was also assessed, and is summarized in Figure 3 for minimal  $\text{Ca}^{2+}$  (3a), maximal  $\text{Ca}^{2+}$  (3b) and time to 50% (3c) and 90% (3d)  $\text{Ca}^{2+}$  decay ( $\text{DT}_{50}$ ,  $\text{DT}_{90}$ ). As with contraction, there was no difference in  $\text{Ca}^{2+}$  transient behavior with increased stimulation frequency between non-transducer and Rrm1+Rrm2 transducer myocytes. GFP-only transducer cardiomyocytes had a slight

increase in minimal  $\text{Ca}^{2+}$  at 2 Hz, and an increase in maximal  $\text{Ca}^{2+}$  at 1 Hz and 2 Hz, as compared to non-transducer myocytes, but the times to 50% and 90% decay were similar. As at 0.5 Hz, the times to 50% and 90% decay were decreased (faster decay) in Rrm1+Rrm2 transducer myocytes at both 1 and 2 Hz. Most importantly, although higher stimulating frequencies increased relaxation parameters in all groups, the relative increase in relaxation kinetics was maintained with Rrm1+Rrm2 over expression, such that relaxation was improved, not impaired. Results for 1 Hz and 2 Hz stimulation are summarized in Supplemental Table 1 and 2, respectively.

We chose to perform these experiments at room temperature (22–24° C) to compare with the predominant number of reports for cultured cardiomyocytes in the literature [25–29]. However, a subset of measurements was made at 37°C to determine if the effects persist at physiological temperature. At 37°C (Supplemental Table 3), shortening and  $\text{Ca}^{2+}$  transients were faster than at 22–24°C, but were similarly increased in cardiomyocytes transducer with Rrm1+Rrm2 vs. GFP-only transducer and non-transducer cells. Similarly, the rates of  $\text{Ca}^{2+}$  release and re-uptake were also increased at 37°C vs. room temperature, but with Rrm1+Rrm2 over expression resulting in faster  $\text{Ca}^{2+}$  transient decay as was observed at ambient temperature.

Since there was little difference between groups in minimal and maximal  $\text{Ca}^{2+}$ , changes in contractility can best be explained by a change in myofilament responsiveness to activating  $\text{Ca}^{2+}$ . This is illustrated in Figure 4 as contractile response, defined here as cardiomyocyte fractional shortening divided by maximal fura2 fluorescence (peak  $\text{Ca}^{2+}$ ). Cardiomyocytes expressing Rrm1+Rrm2 had significantly higher contractile response than non-transducer or GFP transducer cardiomyocytes at all stimulation frequencies. There was no difference in contractile response between GFP only or non-transducer myocytes except at 2 Hz, which can be primarily be attributed to increased maximal  $\text{Ca}^{2+}$  in GFP only myocytes with no increase in fractional shortening, reducing response.

### 3.2 Protein and Nucleotide Analysis

To verify increased Rrm mRNA, Rrm protein, and dATP production in Rrm1+Rrm2 transducer cells, neonatal rat cardiomyocytes were collected and processed for RT-PCR, western blotting and HPLC analysis of intracellular [ATP] and [dATP]. Neonatal cardiomyocytes were used to achieve high enough cell density for accurate nucleotide content analysis, as intracellular [dATP] is known to be in the pM range. Although there are structural differences between neonatal and adult cardiomyocytes, it is important to note that Rrm1+Rrm2 over-expression increased contractility to a similar extent in both cell types (Supplemental Figure 2). Interestingly, as neonatal cardiomyocytes have been used to study the effects of cellular engraftment following myocardial infarction[13], improved contractility in these cells may be another mechanism to improve cardiac function following an infarct. Rrm1 and Rrm2 mRNA was significantly increased following adenoviral transduction (Supplemental Figure 3). Concomitant with this, Figure 5a and 5b illustrate that Rrm1 and Rrm2 transducer cardiomyocytes had greater than 24-fold and 46-fold increased Rrm1 and Rrm2 protein content, respectively. GAPDH was used as a loading control. Figure 5c illustrates that Rrm1+Rrm2 transducer cardiomyocytes had ~10-fold increased cellular



[dATP] as compared to GFP transducer cardiomyocytes (an increase to 0.35 nmol/mg protein). While this is robust, since [dATP] normally comprises less than 0.2% of total adenine triphosphate nucleotide, this increase in [dATP] represents only ~1.5% of the total adenine nucleotide pool. This suggests that only a small amount of dATP is required to significantly increase cardiomyocyte contractility.

To determine how the relatively small increase in cellular [dATP] might influence crossbridge binding and contraction, we compared the rates of nucleotide binding + actomyosin S1 dissociation ( $k_{obs}$ ) for ATP vs. dATP. Figure 6 shows the effect of increasing [ATP] and [dATP] on  $k_{obs}$  at 10°C and 20°C for mouse cardiac (alpha) myosin. There was no difference in  $k_{obs}$  between ATP and dATP at any [NTP] at either temperature. This was also true for fast and slow skeletal S1 myosin (Supplemental Figure 4). This data indicates NTP binding to S1, and subsequent S1 dissociation from actin, is not different for dATP vs. ATP. Thus, it appears that enhanced contractility of R1R2 overexpressing cardiomyocytes is not likely due to a greater myosin affinity for dATP.

### 3.3 Crossbridge Binding and Force in Demembrated Trabeculae

To determine if small amounts of dATP increase force production, we activated contraction of demembrated rat cardiac trabeculae at pica 5.6 (sub-maximal) and pica 4.0 (maximal) in solutions containing 5 mM NTP, composed of either 100% ATP or 2% dATP, 98% ATP. The sub-maximal  $Ca^{2+}$  activation approximates the force levels attained during twitch activity, and was  $30 \pm 7\%$  of maximal force ( $88.4 \pm 1.9$  mN/mm<sup>2</sup>). The example force trace in Figure 7A demonstrates that moving a trabeculae from 100% ATP solution to the 2% dATP, 98% ATP solution resulted in a significant increase in force, which was reversible upon transfer back to the 100% ATP solution. Figure 7B summarizes this increase for all trabeculae activated and shows that 2% dATP, 98% ATP increased force  $17.1 \pm 0.02\%$  ( $p < 0.05$ ) at pica 5.6 but not during maximal  $Ca^{2+}$  (pica 4.0) activation. Similarly, crossbridge binding, assessed by high frequency sinusoidal stiffness measurements, increased  $16.0 \pm 0.03\%$  ( $p < 0.05$ ), which indicates that the increased force with 2% dATP, 98% ATP is due to increased crossbridge binding. Thus the data demonstrate that a relatively small increase in cellular [dATP] (1–2% of adenine nucleotide) is sufficient to significantly increase the contractile strength of intact cardiomyocytes by increasing the number of strong crossbridges.

## 4. DISCUSSION

The main objective of this study was to determine if over expression of rib nucleotide reeducates (Rrm1+Rrm2) increases cellular [dATP] and, in turn, increases contractility in intact cardiomyocytes without adversely affecting cardiomyocyte relaxation. Over expression of Rrm1+Rrm2 resulted in increased cellular [dATP] to ~1.5% of the total adenine nucleotide pool, and this dramatically increased the extent and rate of myocyte shortening and rate of myocyte relaxation, while having no apparent effect  $Ca^{2+}$  transient properties.

Previous experiments in our laboratory using skinned cardiac trabeculae showed dATP increased isometric force and the rate of force development and shortening at all levels of

Author Manuscript  
Author Manuscript  
Author Manuscript

$\text{Ca}^{2+}$  activation, including saturating  $[\text{Ca}^{2+}]$  (pica 4.0), but these studies were performed with 100% replacement of 5 mM ATP with 5 mM dATP in bathing solutions[6–8]. For our current study in intact cardiomyocytes we did not expect over expression of Rrm1+Rrm2 to result in high (mM) levels of dATP. Others have shown that as little as 10% replacement of ATP with dATP is sufficient to see a gain of force in demembrated porcine trabeculae (15°C)[30] and replacement of ~30% increased contractility in intact embryonic chick cardiomyocytes[31]. In our studies the observed large increases in contractility of adult rat cardiomyocytes occurred with a relatively small increase in cellular [dATP] resulting from over expression of Rrm1+Rrm2, and a similar amount (2% dATP) significantly increased sub-maximal force in demembrated trabeculae. It is possible there was a small population of contaminating cells (e.g., fibroblasts) that were either not as easily transducer or overexpressed less Rrm1+Rrm2, which would lead to underestimation of cardiomyocyte [dATP] from the HPLC analysis. However, considering the relative scarcity of non-cardiomyocyte cells in our culture, this confounding effect should be minimal. Our force measurements with demembrated trabeculae suggest this relatively small concentration of cellular [dATP] is sufficient to result in the increased contractility seen in intact cardiomyocytes. This may be advantageous in that large increases in [dATP] are not required to achieve contractile potentiation, thus reducing the potential for negative side effects[31, 32].

Author Manuscript  
Author Manuscript  
Author Manuscript

It is interesting to speculate on how a relatively small increase in cellular dATP can improve cardiomyocyte function. Contractile response estimations (Figure 4) indicate increased contractility in Rrm1+Rrm2 transducer myocytes is primarily myofilament based, thus dATP likely has its effect primarily by improving myosin binding (to actin) and crossbridge cycling. The increase in sub-maximal steady-state force and stiffness seen in demembrated trabeculae activated with 2% dATP, 98% ATP supports this idea. This is similar to experiments where faster (alpha) myosin has been expressed in cardiomyocytes that normally express slower (beta) myosin, resulting in functional potentiation with no effect on  $\text{Ca}^{2+}$  transient amplitude [24]. One possibility we examined was that dATP affinity for cardiac myosin is much greater than ATP affinity for myosin, such that the increased level in cells (with Rrm1+Rrm2 over expression) was utilized almost specifically by myosin. We have previously shown that ATP and dATP have similar binding affinity to skeletal myosin and actomyosin and a similar  $\gamma$ phosphate cleavage equilibrium by myosin [33]. Here we report that NTP binding and subsequent dissociation of cardiac  $\alpha$ -myosin from actin does not differ for dATP vs. ATP, as assessed by  $k_{\text{obs}}$  (Figure 6). This suggests dATP has a similar binding affinity for cardiac myosin as ATP (Supplemental Scheme 1). For skeletal myosin we have also shown post-hydrolysis crossbridge binding and the rate of crossbridge detachment is increased with dATP[33]. This can explain an increase in the  $\text{Ca}^{2+}$  sensitivity of tension development, and a faster rate of tension development and shortening velocity in skinned skeletal muscle[33–35]. While we have not performed a detailed chemo-mechanical analysis with dATP in cardiac muscle, we have shown that it increases maximal crossbridge binding (as indicated from stiffness measurements) and isometric force by >40%, in addition to increasing  $k_{\text{tr}}$  and unloaded shortening velocity[6]. We have also shown that dATP significantly increases isometric force and  $k_{\text{tr}}$  in cardiac muscle at all levels of  $\text{Ca}^{2+}$ , whether the demembrated cardiac muscle was expressing primarily  $\alpha$ - or  $\beta$ -myosin heavy chain[8].



This is important because, unlike skeletal muscle, the intracellular  $[Ca^{2+}]$  during a cardiac muscle twitch only reaches a level that is approximately within the half-maximally activating range. Additionally, cooperative thin filament activation in cardiac muscle is strongly influenced by strong binding crossbridges[7]. Based on our results that 2% dATP, 98% ATP significantly enhanced crossbridge binding and force in demembranated trabeculae, we propose that dATP results in the formation of a few additional strong binding crossbridges early on during the twitch, which provides a positive feedback amplification of thin filament activation. This results in greater total crossbridge binding, including crossbridges using ATP, during the cardiomyocyte twitch. Therefore, for our current experiments with cultured cardiomyocytes, it may be that a small increase in initial binding of myosin S1 heads was enough to cooperatively increase thin filament activation, resulting in the increased magnitude and rate of shortening. Future studies will be required to determine how R1R2 over expression and resulting increases in cellular [dATP] affect cardiac function *in situ*, but previous experiments have demonstrated that even demembranated cardiomyocyte function translates to cardiac organ function[36]. Interestingly, a recent study investigating a small molecule myosin activator (omecantiv mecarbil) demonstrated functional potentiation at the cardiomyocyte level, via increased crossbridge binding, that was very similar to results in the present study[37]. However, this molecule appeared to slow the times to peak shortening and 50% and 90% relaxation, which would likely increase time spent in systole while decreasing time spent in diastole. In contrast, Rrm1+Rrm2 over-expression increased fractional shortening without increasing the time to peak shortening, and shortened the times to 50% and 90% relaxation (Figure 1a, Table 1, Supplemental Tables 1–4), which would allow more time for ventricular filling.

There was no adverse effect on relaxation with over expression of Rrm1+Rrm2 (and the subsequent increase in [dATP]), in fact myocyte relaxation was enhanced. It is possible that this resulted, at least in part, from a faster decay of the  $Ca^{2+}$  transient. dATP could be used by other ATPases (besides myosin) such as the sarcoplasmic  $Ca^{2+}$  ATPase (SERCA), the plasma membrane  $Ca^{2+}$  ATPase (PMCA), and may also indirectly effect activity of the sodium/calcium exchanger (NCX)[38]. An increase in SERCA activity could explain the increased decay rate of the  $Ca^{2+}$  transient, especially at 0.5 and 1.0 Hz stimulation. However, increased SERCA activity is known to increase SR  $Ca^{2+}$  stores[39], making more  $Ca^{2+}$  available for release during activation, which was not observed in Rrm1+Rrm2 transducer cardiomyocytes (Figure 1d). Furthermore, increased PMCA and NCX activity should result in a  $Ca^{2+}$  transient decay over time by extruding  $Ca^{2+}$  out of the cell. Because ~95% of activating  $Ca^{2+}$  is released from the SR in rat cardiomyocytes[40, 41],  $Ca^{2+}$  extrusion from the cell would lead to progressively decreased  $Ca^{2+}$  transient amplitudes and contraction, which was not observed over the duration of these experiments. However, the specific mechanism behind increased  $Ca^{2+}$  transient decay rate warrants future investigation.

Since dATP increases the rate of crossbridge detachment[6, 34] this may also explain a faster rate of relaxation in the current experiments with cultured cardiomyocytes. Although specific mechanisms that govern relaxation in intact cardiac muscle are not known, early phase relaxation in cardiac and skeletal myofibrils has been shown to be governed by the rate of crossbridge dissociation[42–45]. This would also be consistent with our finding that cardiomyocyte contractility was increased with Rrm1+Rrm2, because shortening rate in

unloaded cells (as in culture) is primarily determined by crossbridge detachment rates[46, 47], although cultured cardiomyocytes still contract against a small internal load. It is also possible that increased crossbridge detachment rate with dATP accelerates cooperative thin filament inactivation, by more rapidly decreasing the bound crossbridge population as thin filament  $\text{Ca}^{2+}$  binding decreases during relaxation. A more rapid decrease in bound crossbridge population could also increase the rate of  $\text{Ca}^{2+}$  dissociation from troponin, as demonstrated by Tikunova *et al.* (2010) [48], further accelerating relaxation.

In summary, dATP provides a dual benefit of positive inotropy and lusitropy in cultured rat cardiomyocytes, with little alteration of  $\text{Ca}^{2+}$  transient properties, and also increases isometric force production at physiologically relevant  $[\text{Ca}^{2+}]$ . These results warrant progression to animal studies to determine its potential to improve global cardiac function in normal and diseased hearts.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank Dr. Hans Reinecke for assistance in adenoviral production, and Scott Lundy for assistance in preparing the manuscript. Funding support for this work was provided by from NIH HL07828 and AHA 2310117 (FSK), NIH HL65497 (MR), R21 HL091368 (MR and CEM), R01 HL084642 (CEM) and P01 HL094374 (CEM). MR is an Established Investigator of the American Heart Association.

## Non-standard Abbreviations and Acronyms

<b>Rrm1</b>	muscle ribonucleotide reductase 1
<b>Rrm2</b>	muscle ribonucleotide reductase 2
<b>ARC</b>	adult rat cardiomyocyte
<b>GFP</b>	green fluorescent protein
<b>RT<sub>50</sub>, RT<sub>90</sub></b>	time to 50% and 90% relaxation
<b>DT<sub>50</sub>, DT<sub>90</sub></b>	time to 50% and 90% $\text{Ca}^{2+}$ decay

## References


1. Benitah JP, Alvarez JL, Gomez AM. L-type  $\text{Ca}^{2+}$  current in ventricular cardiomyocytes. *J Mol Cell Cardiol.* 2010; 48(1):26–36. [PubMed: 19660468]
2. Sabbah HN. Biologic rationale for the use of beta-blockers in the treatment of heart failure. *Heart Fail Rev.* 2004; 9(2):91–7. [PubMed: 15516856]
3. LeWinter MM. Functional consequences of sarcomeric protein abnormalities in failing myocardium. *Heart Fail Rev.* 2005; 10(3):249–57. [PubMed: 16416047]
4. Palmer BM. Thick filament proteins and performance in human heart failure. *Heart Fail Rev.* 2005; 10(3):187–97. [PubMed: 16416042]
5. Rubart M, Zipes DP. Mechanisms of sudden cardiac death. *J Clin Invest.* 2005; 115(9):2305–15. [PubMed: 16138184]

6. Regnier M, et al. 2-deoxy-ATP enhances contractility of rat cardiac muscle. *Circ Res.* 2000; 86(12): 1211–7. [PubMed: 10864910]
7. Regnier M, et al. Cross-bridge versus thin filament contributions to the level and rate of force development in cardiac muscle. *Biophys J.* 2004; 87(3):1815–24. [PubMed: 15345560]
8. Adhikari BB, et al. Cardiac length dependence of force and force redevelopment kinetics with altered cross-bridge cycling. *Biophys J.* 2004; 87(3):1784–94. [PubMed: 15345557]
9. Clemmens EW, Regnier M. Skeletal regulatory proteins enhance thin filament sliding speed and force by skeletal HMM. *J Muscle Res Cell Motil.* 2004; 25(7):515–25. [PubMed: 15711882]
10. Kashlan OB, Cooperman BS. Comprehensive model for allosteric regulation of mammalian rib nucleotide reeducates: refinements and consequences. *Biochemistry.* 2003; 42(6):1696–706. [PubMed: 12578384]
11. Kashlan OB, et al. A comprehensive model for the allosteric regulation of mammalian rib nucleotide reeducates. Functional consequences of ATP-and dATP-induced oligomerization of the large subunit. *Biochemistry.* 2002; 41(2):462–74. [PubMed: 11781084]
12. Santana LF, Kranias EG, Lederer WJ. Calcium sparks and excitation-contraction coupling in phospholamban-deficient mouse ventricular myocytes. *J Physiol.* 1997; 503(1):21–29. [PubMed: 9288671]
13. Moreno-Gonzalez A, et al. Cell therapy enhances function of remote non-infarcted myocardium. *Journal of Molecular and Cellular Cardiology.* 2009; 47(5):603–13. [PubMed: 19683533]
14. He TC, et al. A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci U S A.* 1998; 95:2509–14. [PubMed: 9482916]
15. Iorga B, Adamek N, Geeves MA. The slow skeletal muscle isoform of myosin shows kinetic features common to smooth and non-muscle myosins. *J Biol Chem.* 2007; 282(6):3559–3570. [PubMed: 17130133]
16. Margossian SS, Lowey S. Preparation of myosin and its subfragments from rabbit skeletal muscle. *Methods Enzymol.* 1982; 85(B):55–71. [PubMed: 6214692]
17. Weeds AG, Taylor RS. Separation of subfragment-1 isoenzymes from rabbit skeletal muscle myosin. *Nature.* 1975; 257(5521):54–56. [PubMed: 125854]
18. Spudich JA, Watt S. The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J Biol Chem.* 1971; 246(15):4866–4871. [PubMed: 4254541]
19. Herron TJ, et al. Calcium-independent negative inotropy by beta-myosin heavy chain gene transfer in cardiac myocytes. *Circ Res.* 2007; 100(8):1182–90. [PubMed: 17363698]
20. Kreuziger KL, et al. Cooperative activation and tension kinetics in cardiac muscle are strongly modulated by calcium binding kinetics of troponin C. *Journal of Molecular and Cellular Cardiology.* 2011; 50(1):165–174. [PubMed: 21035455]
21. Martyn DA, et al. Unloaded shortening of skinned muscle fibers from rabbit activated with and without Ca<sup>2+</sup>. *Biophys J.* 1994; 67(5):1984–93. [PubMed: 7858135]
22. Badrian B, Bogoyevitch MA. Changes in the transcriptional profile of cardiac myocytes following green fluorescent protein expression. *DNA Cell Biology.* 2007; 26(10):727–736. [PubMed: 17723104]
23. Nishimura S, et al. Expression of green fluorescent protein impairs the force-generating ability of isolated rat ventricular cardiomyocytes. *Mol Cell Biochem.* 2006; 286(1–2):59–85. [PubMed: 16532258]
24. Herron TJ, et al. Ca<sup>2+</sup>-independent positive molecular inotropy for failing rabbit and human cardiac muscle by alpha-myosin motor gene transfer. *Faseb J.* 2010; 24(2):415–424. [PubMed: 19801488]
25. Cazorla O, et al. The cAMP binding protein Epac regulates cardiac myofilament function. *PNAS.* 2009; 106(33):14144–14149. [PubMed: 19666481]
26. Sheehan KA, et al. Functional effects of a tropomyosin mutation linked to FHC contribute to maladaptation during acidosis. *Journal of Molecular and Cellular Cardiology.* 2011; 50(3):442–450. [PubMed: 21047515]

27. Kawashima H, et al. Protein phosphatase inhibitor-1 augments a protein kinase A-dependent increase in the Ca<sup>2+</sup> loading of the sarcoplasmic reticulum without changing its Ca<sup>2+</sup> release. *Circ J.* 2009; 73:1133–1140. [PubMed: 19377265]
28. Shah AM, et al. The influence of endothelium-derived nitric oxide on myocardial contractile function. *Int J Cardiol.* 1995; 50(3):225–31. [PubMed: 8537145]
29. Vogelpohl I, et al. Transgenic over expression of heart-specific adenine nucleotide translocase 1 positively affects contractile function in cardiomyocytes. *Cell Physiol Biochem.* 2011; 27(2):121–128. [PubMed: 21325829]
30. Schoffstall B, Clark A, Chase P. Positive inotropic effects of low dATP/ATP ratios on mechanics and kinetics of porcine cardiac muscle. *Biophys J.* 2006; 91(6):2216–26. [PubMed: 16798797]
31. Schoffstall B, Chase P. Increased intracellular [dATP] enhances cardiac contraction in embryonic chick cardiomyocytes. *Journal of Cellular Biochemistry.* 2008; 104(6):2217–2227. [PubMed: 18452163]
32. Xu X, et al. Broad over expression of rib nucleotide reeducates genes in mice specifically induces lung neoplasms. *Cancer Res.* 2008; 68(8):2652–2660. [PubMed: 18413732]
33. Regnier M, Lee DM, Homsher E. ATP analogs and muscle contraction: mechanics and kinetics of nucleoside triphosphate binding and hydrolysis. *Biophys J.* 1998; 74(6):3044–58. [PubMed: 9635759]
34. Regnier M, Homsher E. The effect of ATP analogs on posthydrolytic and force development steps in skinned skeletal muscle fibers. *Biophys J.* 1998; 74(6):3059–71. [PubMed: 9635760]
35. Regnier M, Martyn DA, Chase PB. Calcium regulation of tension redevelopment kinetics with 2-deoxy-ATP or low [ATP] in rabbit skeletal muscle. *Biophys J.* 1998; 74(4):2005–15. [PubMed: 9545059]
36. Korte FS, et al. Power output is linearly related to MyHC content in rat skinned myocytes and isolated working hearts. *Am J Physiol Heart Circ Physiol.* 2005; 289(2):H801–812. [PubMed: 15792987]
37. Malik FI, et al. Cardiac myosin activation: a potential therapeutic approach for systolic heart failure. *Science.* 2011; 331:1439–1443. [PubMed: 21415352]
38. Hilgemann DW, Ball R. Regulation of Na<sup>+</sup>,Ca<sup>2+</sup> exchange and KATP potassium channels by PIP<sub>2</sub>. *Science.* 1996; 273(5277):956–959. [PubMed: 8688080]
39. Janssen PM, Periasamy M. Determinants of frequency-dependent contraction and relaxation of mammalian myocardium. *J Mol Cell Cardiol.* 2007; 43(5):523–531. [PubMed: 17919652]
40. Bassani JW, Bassani RA, Bers DM. Relaxation in rabbit and rat cardiac cells: species-dependent differences in cellular mechanisms. *J Physiol.* 1994; 476:279–293. [PubMed: 8046643]
41. Bers DM. Cardiac excitation-contraction coupling. *Nature.* 2002; 415:198–205. [PubMed: 11805843]
42. Luo Y, et al. Determinants of relaxation rate in rabbit skinned skeletal muscle fibres. *J Physiol.* 2002; 545(Pt 3):887–901. [PubMed: 12482894]
43. Luo Y, et al. Myofibrillar determinants of rate of relaxation in skinned skeletal muscle fibers. *Adv Exp Med Biol.* 2003; 538:573–581. [PubMed: 15098700]
44. Piroddi N, et al. Tension generation and relaxation in single myofibrils from human atrial and ventricular myocardium. *Pflugers Arch.* 2007; 454(1):63–73. [PubMed: 17123098]
45. Tesi C, et al. Relaxation kinetics following sudden Ca<sup>2+</sup> reduction in single myofibrils from skeletal muscle. *Biophys J.* 2002; 83(4):2142–51. [PubMed: 12324431]
46. Edman KAP. The velocity of unloaded shortening and its relation to sarcomere length and isometric force in vertebrate muscle fibers. *Journal of Physiology.* 1979; 291:143–159. [PubMed: 314510]
47. Hinken AC, McDonald KS. Inorganic phosphate speeds loaded shortening in rat skinned cardiac myocytes. *American Journal of Physiology.* 2004; 287:C500–C507. [PubMed: 15084471]
48. Tikunova SB, et al. Effect of calcium-sensitizing mutations on calcium binding and exchange with troponin C in increasingly complex biochemical systems. *Biochemistry.* 2010; 49(9):1975–1984. [PubMed: 20128626]

### Highlights


- dATP enhances myocardial contraction via increased crossbridge binding and kinetics
- [dATP] can be increased >10-fold via rib nucleotide reeducates (Rrm) overexpression
- Rrm over expression increased intact cardiomyocyte contractility and relaxation
- Rrm over expression had no effect on Ca<sup>2+</sup> transient magnitude
- Myofilament targeted gene manipulations can improve cardiac function



### Figure 1. Representative traces and data summary


Representative cell length traces (a) and Ca<sup>2+</sup> transients (b, Fura-2 fluorescence) of non-transducer (black), GFP-only (green), and Rrm1+Rrm2+GFP (red) transducer cardiomyocytes. Percentage change in contractile (c) and Ca<sup>2+</sup> transient (d) properties of GFP-only and Rrm1+Rrm2+GFP transducer myocytes, stimulated at 0.5 Hz, as compared to non-transducer myocytes.  $V_{short}$  = velocity of shortening; FS = fractional shortening;  $V_{rel}$  = maximal relaxation velocity; RT<sub>50,90</sub> = time to 50% and 90% relaxation, respectively; FL = fluorescence; DT<sub>50,90</sub> = time to 50% and 90% Ca<sup>2+</sup> decay, respectively \*p<0.05 as compared to Non-Transducer.






**Figure 2. Effect of stimulation frequency on contractile properties**

Rrm1+Rrm2 transducer myocytes (open triangles) respond similarly to stimulation frequency as GFP-only transduce open circles) and non-transducer myocytes (closed circles) but show elevated fractional shortening (a) and shortening velocity (b) at all frequencies. Relaxation velocity (c) and time to 90% relaxation (d) are also similar between groups, with time to relaxation shortening as stimulation frequency increases. \* =  $p < 0.05$  as compared to Non-Transducer, † =  $p < 0.05$  as compared to GFP, ‡ =  $p < 0.05$  as compared to 0.5 Hz for all groups.




### Figure 3. Effect of stimulation frequency on Ca<sup>2+</sup> handling properties

Rrm1+Rrm2 transducer myocytes (open triangles) respond similarly to stimulation frequency as non-transducer myocytes (closed circles) in minimal (a) and maximal (b) fluorescence, while GFP-only transducer myocytes (closed circles) showed a greater increase in both as frequency increased. As with cardiomyocyte relaxation, Ca<sup>2+</sup> transient decay time (DT) to 50% (c) and 90% (d) is shortened with increased stimulation frequency, but both are dramatically shortened in R1R2 transducer cardiomyocytes. \* = p<0.05 as compared to Non-Transducer, † = p<0.05 as compared to GFP, ‡ = p<0.05 as compared to 0.5 Hz for all groups.




**Figure 4. Contractile Responsiveness**

Contractile response as assessed as fractional shortening divided by maximal fura fluorescence (peak  $\text{Ca}^{2+}$ ) indicates Rrm1+Rrm2 transducer cardiomyocytes (open triangles) are significantly more responsive to  $\text{Ca}^{2+}$  at all stimulation frequencies, while GFP-only transducer cardiomyocytes (open circles) are less responsive to  $\text{Ca}^{2+}$  only at 2Hz stimulation frequency as compared to non-transducer cardiomyocytes (closed circles). \* =  $p < 0.05$  as compared to Non-transducer, † =  $p < 0.05$  as compared to GFP.




**Figure 5. Increased Rrm and dATP**

(a) Western blot of Rrm1 transducer neonatal rat cardiomyocytes probed with anti-Rrm1 antibody indicates a >24-fold increase in Rrm1. (b) Western blot of Rrm2 transducer neonatal rat cardiomyocytes probed with anti-Rrm2 antibody indicates a > 46-fold increase in Rrm2. (c) Rrm1+Rrm2 over expression significantly increased intracellular [dATP] by >10-fold in neonatal rat cardiomyocytes as assessed by HPLC analysis. \* =  $p < 0.05$  as compared to GFP transducer cardiomyocytes.



**Figure 6. Nucleotide binding and actin-myosin dissociation**

Rapid kinetic measurements of nucleotide binding and actin-myosin dissociation of mouse cardiac myosin taken at 10°C (top) and 20°C (bottom). There was no difference in  $k_{obs}$  between ATP and dATP at any [NTP] at either temperature.



**Figure 7. Isometric force increases with 2% dATP**

Isometric force development of demembrated cardiac trabeculae activated with 2% dATP, 98% ATP vs. 100% ATP (5 mM [NTP] total). (a) Force trace with pica 5.6 activation. (b) Summary of % increase in force with 2% dATP in activation solutions for sub-maximal (pica 5.6), but not maximal (pica 4.0) Ca<sup>2+</sup> activation.



Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1**

	N	n	SL ( $\mu\text{m}$ )	Cell length( $\mu\text{m}$ )
Non-transducer	5	51	1.88 $\pm$ 0.03	90.4 $\pm$ 1.8
Control(GFP)	5	50	1.84 $\pm$ 0.03	89.1 $\pm$ 1.5
R1R2(GFP)	5	52	1.82 $\pm$ 0.02	91.6 $\pm$ 1.7

**Table 2**

Contractile and Ca<sup>2+</sup> transient values at 0.5 Hz stimulation.

	Fractional Shortening(%)	Maximal Shortening Rate (µm/s)	Time to Peak(ms)	Maximal Relaxation Rate (µm/s)	RT <sub>50</sub> (ms)	RT <sub>90</sub> (ms)	Minimal Ca <sup>2+</sup> /Fura ratio units)	Maximal Ca <sup>2+</sup> /Fura ratio units)	DT <sub>50</sub> (ms)	DT <sub>90</sub> (ms)
Non-transducer	6.2 ± 0.4	61.1 ± 4.4	173 ± 12	46.8 ± 6.5	208 ± 28	330 ± 59	1.10 ± 0.02	1.22 ± 0.04	246 ± 26	666 ± 74
Control(GFP)	5.5 ± 0.5	56.5 ± 4.4	217 ± 15 <sup>*</sup>	37.5 ± 4.3	202 ± 25	518 ± 42 <sup>*</sup>	1.12 ± 0.03	1.25 ± 0.04	297 ± 24	893 ± 63 <sup>*</sup>
R1R2(GFP)	8.9 ± 0.5 <sup>*</sup>	109.5 ± 8.7 <sup>*</sup>	177 ± 7	117.9 ± 13.1 <sup>*</sup>	113 ± 7 <sup>*‡</sup>	265 ± 23 <sup>‡</sup>	1.14 ± 0.02	1.23 ± 0.03	153 ± 10 <sup>*‡</sup>	435 ± 34 <sup>*‡</sup>

\* p<0.05 as compared to No Treat.

<sup>‡</sup> p<0.05 as compared to GFP.

<sup>‡</sup> p<0.05 as compared to 0.5 Hz for all groups.