

Accuracy of self-referral in health anxiety: comparison of patients self-referring to internet-delivered treatment versus patients clinician-referred to face-to-face treatment

Ditte Hoffmann, Charlotte Ulrikka Rask, Erik Hedman-Lagerlöf, Trine Eilenberg and Lisbeth Frostholt

Background

Severe health anxiety is a chronic and costly disorder if untreated. Patient self-referral may lower barriers to treatment and decrease diagnostic delay.

Aims

This study evaluated the accuracy of self-referral for severe health anxiety and compared characteristics of patients self-referred to internet-delivered treatment with patients referred by a clinician to face-to-face treatment.

Method

Two trials in the same clinic employed different referral methods for health anxiety, namely self-referral and clinician-referral (trial registration: clinicaltrials.gov: NCT01158430 and NCT02735434). The trials were conducted at different time points but with largely comparable eligibility criteria. The accuracy of the recruitment methods was compared by looking at the number of eligible patients in the two trials. Patients completed a baseline questionnaire and subsequently underwent a diagnostic interview by experienced clinicians. Mean differences in self-report and clinical data explored between-group demographic and clinical characteristics.

Results

In total, 101/151 (67%) self-referred patients were eligible compared with 126/254 (50%) clinician-referred patients ($P = 0.001$). Self-referred patients were 3.4 years older ($P = 0.008$) and had a somewhat higher educational level ($P = 0.030$). Patients who self-referred reported significantly higher levels of health anxiety,

emotional distress and somatic symptoms compared with clinician-referred patients. Yet, they had less clinician-assessed comorbid anxiety disorders ($P < 0.001$) and better physical health-related quality of life ($P < 0.001$) suggesting a more distinct symptom profile.

Conclusions

Self-referral was found to be an accurate method to recruit highly relevant patients with treatment-demanding health anxiety. Thus, both self-referral and clinician-referral seem feasible and valid referral methods, but they may recruit patients with slightly different characteristics.

Declaration of interest

None.

Keywords

Health anxiety; hypochondriasis; self-referral; barriers to treatment; internet intervention.

Copyright and usage

© The Royal College of Psychiatrists 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Background

Severe health anxiety is a highly disabling and costly disorder^{1,2} affecting approximately 1–5% of the adult population.^{3,4} It is characterised by exaggerated rumination with fears of suffering from a serious illness.³ In DSM-5, health anxiety is now classified as illness anxiety disorder or somatic symptom disorder if severe somatic symptoms are present, respectively.⁵ Even though empirically supported psychological therapies for health anxiety exist,⁶ the limited availability and barriers to treatment challenge the dissemination and potential impact. Common barriers to psychological treatment are the geographical distance to treatment, the interference with work and daily life, and the fear of stigmatisation.^{7–9} In patients with self-identified health anxiety, a study found that only 20.7% currently received therapy or counselling, and 53.4% reported past experiences with either of these.¹⁰ This treatment gap is also widely recognised for common mental disorders.¹¹ Although patients with health anxiety are frequently seen in medical care settings, they are hesitant to mention their illness worries^{2,12} and thus risk a chronic, undetected course.¹ These barriers call for new approaches to improve access to psychological treatment.

The potential role of self-referral

In many countries, referrals are initiated by a general practitioner (GP) in primary care who serves as a gatekeeper to more specialised services. One of the explicit ideas is that this will result in reduced demand for specialised care, since health clinicians are better informed than patients to make the right referrals.¹³ However, the use of patient self-referral and internet-delivered treatment programmes are recent developments that may improve treatment access and reduce barriers for patients with severe health anxiety.¹⁴ These innovative programmes consist of highly structured online modules with homework assignments and communication with a clinician via email, telephone or an embedded online forum.^{15,16} The content mirrors the same techniques and material as in traditional face-to-face treatment. Additionally, self-referral where the patients themselves and not healthcare clinicians initiate the treatment contact may help reduce barriers to treatment.

In recent years, self-referral has become increasingly used partly because of the increased amount and quality of online health information making patients more knowledgeable about their conditions and treatment options.¹⁷ However, the accuracy is dependent on patients' ability to preliminarily self-assess their problems and their

need for treatment. At best, self-referral may provide access for patients who for years have remained untreated for their health anxiety.¹¹ Still, a common concern among healthcare providers is whether the already limited services will primarily be used by people with mild symptoms and limited impairment instead of by those with the highest treatment demand.¹⁸ Therefore, knowledge about the appropriateness of self-referral among patients with health anxiety as well as which patients use it is needed. This is important for several reasons. First, this question relates to the generalisability of findings from treatment studies. That is, if there are marked differences between patient populations self-referred and clinician-referred, comparisons of effect estimates may be done with caution. Second, knowledge about potential differences or similarities is of great relevance for healthcare providers and policymakers when making prioritisations within a limited healthcare budget. Third, this information is valuable for treatment developers as treatments can be tailored to fit the patients most likely to receive it.

Aims

The present study aims to (a) evaluate the accuracy of self-referral for health anxiety based on the number of patients being clinically assessed to have sufficient levels of health anxiety after a self-referral procedure in relation to a trial of internet-delivered treatment; and (b) explore self-referred patients' demography (for example age, gender, education) and clinical characteristics in comparison with clinician-referred patients recruited to a trial of face-to-face treatment in the same clinical setting.

Method

Setting and participants

This study took place at the Research Clinic for Functional Disorders and Psychosomatics at Aarhus University Hospital, Denmark, which is a specialised unit with a combined research objective and a routine clinical function. Participants were retrieved from two separate randomised controlled trials (RCTs, trial registration: clinicaltrials.gov: NCT01158430 and NCT02735434) conducted in the research clinic. In the primary trial, the patients were recruited through self-referral from April 2016 to May 2017 to an internet-delivered acceptance and commitment therapy programme (iACT),¹⁴ whereas in the comparison trial the patients were recruited from March 2010 to April 2012 by clinicians working in other hospital settings or in general practice referring patients to a group-based acceptance and commitment therapy programme (ACT-G).¹⁹ Study information was available on the clinic's webpage and sent to GPs in the catchment areas. In the iACT trial, information was also provided on a nationwide anxiety association webpage. In both trials, the recruitment, diagnostic interview and treatment were performed at the research clinic. The treatment content in iACT was based on ACT-G. Thus, the main differences between the two trials were the mode of referral and the face-to-face versus online treatment delivery.

The eligibility criteria were, in general, comparable in the two trials (Table 1). The main exception was the different age limit, which was 18 years or older in the iACT trial compared with 20–60 years in the ACT-G trial. The different catchment areas were both large thus minimising the risk of geographic interference. Furthermore, recent changes in anxiety medication were only an exclusion criterion in the iACT trial where patients, if on pharmacological treatment for their anxiety, had to be on a stable dose for 2 months prior to treatment initiation. In total, 227 patients with severe health anxiety were included; 101 self-referred patients from the iACT trial (the self-referred group) and 126 clinician-

referred patients from the ACT-G trial (the clinician-referred group).¹⁹

The two studies were approved by the Ethical Committee in Central Region Denmark (iACT ID: 1-10-72-321-15 and ACT-G ID: 20090201) and the Danish Data Protection Agency (iACT case: 1-16-02-51-16 and ACT-G case: 1-16-02-40-13). All participants provided written informed consent.

Referral and assessment procedures

In the iACT trial, patient self-referral was carried out through a web portal through the research clinic's webpage. The self-assessment and referral was enhanced by extensive online information encompassing two patient videos about health anxiety and internet-delivered treatment, a guide on how to self-refer, and core criteria on health anxiety. Interested patients logged in using their unique Danish personal identification number and completed a baseline questionnaire. All self-referrals were preliminarily screened by a psychologist from the research clinic (D.H.) in accordance with the eligibility criteria. Subsequently, patients were invited for a diagnostic interview.

In the ACT-G trial, the clinician-referrals were primarily carried out by the patient's GP or by a clinician in another hospital department. The referrals were subsequently screened by medical doctors from the research clinic in accordance with the eligibility criteria and eligible patients were invited for a diagnostic interview. Patients completed a baseline questionnaire prior to this interview. This procedure has been thoroughly described elsewhere.¹⁹

Patients in both groups underwent a thorough diagnostic interview with the semi-structured interview Schedules for Clinical Assessment in Neuropsychiatry (SCAN) assessing both health anxiety and comorbid disorders.²⁰ In both groups, experienced clinicians trained in using SCAN conducted the diagnostic interviews (one psychiatrist and four to five psychologists in each trial). However, patients in the clinician-referred group underwent the interview face-to-face, whereas patients in the self-referred group underwent the interview via video conferencing. The video-based SCAN interview was modified and shortened based on a procedure from a large Danish epidemiology study²¹ (1–2 h) compared with the longer face-to-face interview (3–4 h). This modification pertained primarily to the length of history taking and amount of psychoeducation provided at the time of the interview. Both versions covered clinical questions regarding health anxiety, depression and obsessive-compulsive disorder (OCD). The number of questions regarding other anxiety disorders and somatoform disorders were reduced in the video session. Video-recorded diagnostic interviews were discussed, and consensus rating among the assessors was frequently performed in order to secure diagnostic validity and reliability. Non-face-to-face assessment of diagnoses has proven to be a reliable method.^{22,23}

Study design

This study was a secondary and exploratory analysis based on data from two separate RCTs. We evaluated the accuracy of self-referral by comparing the number of excluded patients in the group of self-referred and clinician-referred patients, respectively. The accuracy of referrals was evaluated based on (a) the number of clinician-excluded patients prior to the diagnostic interview following the initial screening at the research clinic in accordance with the eligibility criteria, (b) the number of patients failing to attend the interview, (c) the number of patients excluded by clinicians following the interview because of ineligibility, (d) the number of patients declining participation, and (e) the total number of eligible patients in the two trials. To explore demographic and clinical characteristics of self-referred patients (for example do they represent less severe

Table 1 Recruitment and eligibility criteria

Study procedures	Clinician-referral, ACT-G (<i>n</i> = 126) ^a	Self-referral, iACT (<i>n</i> = 101)
Recruitment		
Enrolment	March 2010 to April 2012	April 2016 to May 2017
Study information	On the clinic's webpage, letters with information to the general practitioners in the western part of Denmark	On the clinic's webpage, electronic information to the general practitioners nationwide, notice on a webpage for a nationwide anxiety association
Catchment area	The Region of Southern Denmark and Central Region Denmark (2.5 million people, approximately 40% of the Danish population)	Nationwide (5.7 million persons)
Referral	Patients referred by a clinician from primary or secondary care	Patients self-referred through the clinic's webpage
Eligibility criteria		
Inclusion criteria	Severe health anxiety >6 months ^b Whiteley Index-7 score >21.4 (0–100) ^c Health anxiety is the primary diagnosis if comorbid disorders are present 20–60 years Patients of Scandinavian origin who speak, read, write and understand Danish	Severe health anxiety >6 months ^b Whiteley Index-7 score >21.4 (0–100) ^c Health anxiety is the primary diagnosis if comorbid disorders are present 18 years or older Patients who speak, read and write Danish Computer and internet access
Exclusion criteria	Acute suicidal risk Abuse of narcotics or alcohol and (non-prescribed) medicine Life-time diagnosis of psychoses, bipolar affective disorder or depression with psychotic symptoms (ICD-10: F20-29, F30-31, F32.3, F33.3) Pregnancy No informed consent	Acute suicidal risk Abuse of narcotics or alcohol and (non-prescribed) medicine Life-time diagnosis of psychoses, bipolar affective disorder or depression with psychotic symptoms (ICD-10: F20-29, F30-31, F32.3, F33.3) Pregnancy No informed consent Former treatment at the clinic i.e. ACT-G Changes in anxiety medication within past 2 months
Assessment	Full version SCAN ^d interview conducted face-to-face (3–4 h.)	Shortened version of the SCAN ^d interview conducted through video conferencing (1–2 h.)

ACT-G, group-based acceptance and commitment therapy programme; iACT, internet-delivered acceptance and commitment therapy programme; SCAN, Schedules for Clinical Assessment in Neuropsychiatry.
a. Trial reported by Eilenberg *et al.*¹⁹
b. Diagnostic, empirically-based criteria by Fink and colleagues.³
c. Established cut-off for non-clinical cases based on Whiteley Index-7 score >21.4 (scale range 0–100).¹
d. Schedules for Clinical Assessment in Neuropsychiatry: a semi-structured clinical diagnostic interview.

cases of health anxiety or do they differ in terms of age, gender, education), eligible self-referred patients were compared with the other sample of clinician-referred patients with severe health anxiety. Demographic variables were measured by self-report questionnaires. Clinical characteristics partly comprised self-report measures, as well as clinician-assessed illness onset, duration and comorbidity assessed at the diagnostic interview. In both groups, all questionnaires were administered before treatment start.

Measures

Accuracy of self-referral versus clinician-referral

The accuracy of self-referral was evaluated based on the number of excluded self-referred and clinician-referred patients prior to the diagnostic interview, following the diagnostic interview and in total.

Sociodemographic characteristics

This included the self-reported variables of age, gender, living status, highest education qualification, employment status and current sick leave.

Clinical characteristics

Health anxiety severity. We measured health anxiety severity using a seven-item version of the Whiteley Index (Whiteley-7),²⁴ assessing worrying and convictions about illness. The Whiteley-7 employs a five-point Likert scale from 1, 'not at all' to 5, 'a lot' with a scale range from 7 to 35. It includes items such as 'Do you worry a lot about your health'. Higher scores indicate more health anxiety. The Whiteley-7 has demonstrated satisfactory internal validity in a Danish primary care sample and good external validity for screening DSM-IV somatisation disorder and hypochondriasis.^{24,25}

In the present study, the internal consistency was $\alpha = 0.88$ and 0.77 for ACT-G and iACT, respectively.

Health anxiety onset and duration. This was calculated based on the patient-reported onset of health anxiety noted by the clinician at the diagnostic SCAN interview.

Clinically assessed comorbidity. Comorbidity encompassing depression, anxiety disorders and OCD was summarised based on the diagnostic SCAN interview using the diagnostic classification of the 10th edition of the International Classification of Diseases and Related Health Problems (ICD-10).²⁶

Emotional distress, somatic symptoms and obsessive-compulsive symptoms. These were all measured using questions from the Symptom Checklist scale (SCL-92).²⁷ All questions employ a five-point Likert scale from 1, 'not at all' to 5, 'a lot'. Emotional distress was measured on eight items assessing anxiety and depression (SCL-8).²⁸ Items include 'feeling fearful' and 'feelings of worthlessness' with a scale range from 8 to 40. Higher scores indicate more emotional distress. This brief screening tool has shown good validity in non-psychiatric medical settings in Denmark,²⁹ and in the present study, the internal consistency was $\alpha = 0.86$ in both samples. Somatic symptoms were measured on the 12-item somatisation subscale (SCL-SOM)³⁰ with a scale range from 12 to 60. This subscale checks for 12 common physical symptoms such as headaches or pains in the heart or chest. Higher scores indicate more symptoms. The internal consistency of the SCL-SOM in this study was $\alpha = 0.84$ for ACT-G and $\alpha = 0.83$ for iACT.

Obsessive-compulsive symptoms were measured on the ten-item obsessive-compulsive subscale (SCL-OCD)³¹ with a scale

range from 5 to 50. Higher scores indicate more symptoms, and the subscale includes items such as 'Having to check and double check what you do'. In the present study, the internal consistency of SCL-OCD was $\alpha = 0.85$ and 0.88 for ACT-G and iACT, respectively.

Health-related quality of life. We measured on two dimensions of the 12-item Short-Form Health Survey (SF-12),^{32,33} namely the physical component summary (PCS) and the mental component summary (MCS) with a scale range from 12 to 47. Higher scores indicate better physical and mental health. The internal consistency was high in a large Danish population-based sample, namely $\alpha = 0.90$ for PCS and $\alpha = 0.85$ for MCS.³⁴

Psychological flexibility. We measured psychological flexibility with the Acceptance and Action Questionnaire (AAQ-II),^{35,36} which is a seven-item measure encompassing questions like 'I'm afraid of my feelings'. Overall, psychological flexibility assesses patients' ability to behave flexibly when distressing thoughts and feelings are present. Items are rated on a seven-point Likert scale from 1, 'never true' to 7, 'always true' with a scale range from 7 to 49. The scale was reversed with higher scores indicating more flexibility and less experiential avoidance. The AAQ-II has shown a mean Cronbach's alpha coefficient of 0.84 (ranging from 0.78 to 0.88) and good test-retest reliability at 12 months ($\alpha = 0.79$) across six different samples.³⁵ In the present study, the internal consistency was $\alpha = 0.88$ for ACT-G and $\alpha = 0.86$ for iACT.

Non-reactivity. We measured non-reactivity with the seven-item subscale of the Five Facet Mindfulness Questionnaire (FFMQ non-react).³⁷ This facet of mindfulness covers the ability to be non-reactive towards inner experiences such as thoughts, feelings and bodily sensations. Questions such as 'I perceive my feelings and emotions without having to react to them' are rated on a five-point Likert scale from 1, 'never or very rarely true' to 5, 'very often or always true' with a scale range from 7 to 35. Higher scores of non-reactivity indicate less avoidance. The FFMQ has shown good psychometric properties with a Cronbach's alpha coefficient ranging from 0.72 to 0.92 .³⁷ Non-reactivity is the only subscale of the FFMQ found to mediate a positive treatment outcome in treatment for severe health anxiety across two studies.^{38,39} In the present study, the internal consistency of the FFMQ non-react was $\alpha = 0.76$ for ACT-G and $\alpha = 0.88$ for iACT.

Statistical analysis

The number of patients with clinically relevant health anxiety consenting to treatment was compared using equality of proportions z -test based on excluded patients following self-referral and clinician-referral. The distribution of all other data was judged by inspection of QQ plots. Sum scores that followed a normal distribution were summarised using the mean and standard deviation, whereas sum scores not following a normal distribution were summarised using the median and interquartile range. The mean differences in continuous data were compared between the two groups using independent two-sample t -tests, whereas categorical data were analysed using χ^2 -tests or Fisher's exact test for small sample sizes. All symptom scales were converted into a scale range 0–100 points ($((\text{score} - \text{min}) / (\text{max} - \text{min})) \times 100$). One variable – duration of health anxiety – was not normally distributed and was thus analysed using the Wilcoxon rank-sum (Mann-Whitney) test. To check the impact of the different age criteria in the two groups, a subanalysis comparing only participants from 20 to 60 years of age across the groups was performed. The multiple variables compared between these study populations are exploratory; hence we did not do any formal correction for P -values. Instead, we

provided exact P -values for each test. Analyses were performed using Stata version 13 for Windows. Statistical significance was set at $P < 0.05$ using two-tailed tests.

Results

Accuracy of self-referral versus clinician-referral

In total, 151 patients self-referred to the iACT trial and 254 patients were referred by a clinician to the ACT-G trial (Fig. 1). Out of these patients, 101/151 (67%) self-referred patients were finally included for the primary trial compared with 126/254 (50%) of the clinician-referred patients for the comparison trial ($z = -3.39$, $P = 0.001$). This significant difference pertained to pre-assessment where clinician-referred patients to a larger extent failed to attend the diagnostic interview ($z = -3.03$, $P = 0.003$) and were excluded based on the eligibility criteria by the screening clinicians in the research clinic ($z = -2.68$, $P = 0.007$). There were no differences found in eligibility for trial participation between patients undergoing the diagnostic interview in the two groups. The overall participation rate was high in both trials, i.e. 98/101 (97%) self-referred patients and 116/126 (92%) clinician-referred patients completed questionnaires at the end of treatment ($z = -1.08$, $P = 0.282$).

Sociodemographic characteristics

Sociodemographic characteristics of the included patients in the two trials are displayed in Table 2. The mean age of self-referred patients was significantly higher than those referred by a clinician with a mean difference of 3.4 years (95% CI 0.9–5.8, $t(225) = 2.69$, $P = 0.008$). A secondary analysis controlling for the different age criterion in the two trials by excluding patients in the iACT trial younger than 20 years and older than 60 years found a significant 3.3-year mean difference in age between groups (95% CI 0.9–5.7, $t(219) = 2.76$, $P = 0.006$). Furthermore, there was a significant difference in educational qualification between the groups showing that 25/101 (25%) of self-referred patients had more than 4 years of higher education (for example university-level education) compared with 15/123 (12%) of the clinician-referred patients ($\chi^2(4) = 10.69$, $P = 0.030$). There were no differences in gender distribution, living status or employment status. Even though not significant, a higher proportion of clinician-referred patients were on full-time sick leave (15%) than the self-referred patients (7%) ($\chi^2(3) = 3.91$, $P = 0.271$).

Clinical characteristics

Self-referred patients reported more symptoms of health anxiety compared with the clinician-referred patients (Table 3). The mean difference between groups was 5.6 points (95% CI 1.0–10.2, $t(224.80) = 2.38$, $P = 0.018$). Patients in both groups had an average onset of health anxiety in their early twenties, and no difference in health anxiety duration was found. In terms of clinical comorbidity, self-referred patients had fewer clinician-assessed comorbid anxiety disorders ($\chi^2(1) = 12.86$, $P < 0.001$).

As opposed to this, the self-referred patients reported higher levels of emotional distress and somatic symptoms compared with the clinician-referred patients. Also, they had a lowered mental health-related quality of life with a mean difference of 10.0 points (95% CI 7.1–12.9, $t(201) = -6.85$, $P < 0.001$) in comparison with the clinician-referred patients, whom in contrast reported 6.3 points lower physical health-related quality of life (95% CI 3.1 to 9.4, $t(201) = 3.92$, $P < 0.001$). There was no difference between the two groups' level of obsessive-compulsive symptoms.

Moreover, the self-referred patients were more reactive compared with the clinician-referred patients as shown by a 6.0 points




Fig. 1 Flow chart of recruitment.

a. Eilenberg *et al.*¹⁹

ACT-G, group-based acceptance and commitment therapy programme; iACT, internet-delivered acceptance and commitment therapy programme.

lower non-reactivity towards distressing thoughts, feelings and images (95% CI 1.8 to 10.2, $t(224) = -2.83, P = 0.005$). In addition, both groups had comparable and low psychological flexibility namely 40.0 (s.d. = 17.7) in the self-referred group and 41.6 (s.d. = 21.7) in the clinician-referred group. Both groups were below the suggested cut-off score for clinically relevant distress of 50.0–59.5 (scale range 0–100, i.e. >24–28 (scale range 7–49) on the original non-reversed and transformed scale).³⁵

Discussion

This study evaluated the accuracy of self-referral in patients with severe health anxiety. The main findings were the greater percentage of patients found to have sufficient levels of health anxiety after a self-referral procedure in relation to a trial of internet-delivered treatment compared with clinician-referred patients recruited to a trial of face-to-face treatment in the same clinical setting. The

broader question for this study was whether patients with severe health anxiety seeking treatment through self-referral differed on clinical characteristics from clinician-referred patients in these two trials. Overall, the results suggest that relevant patients self-referred. In fact, the self-referred patients had more symptoms of health anxiety, emotional distress and poorer mental health-related quality of life. In addition, they were more reactive towards distressing thoughts, feelings and images. This indicates that their need for treatment was at least comparable with the clinician-referred patients.

On the other hand, the self-referred patients had less clinician-assessed psychiatric comorbidity and better physical-health-related quality of life, which might suggest a more distinct symptom profile of self-referred patients. Furthermore, the self-referred group had a somewhat higher educational level and was slightly older, but there was no statistically significant difference in age at onset and duration of health anxiety between the groups. Even though the two patient groups differed, the differences were relatively minor. All

Table 2 Demographic characteristics					
	Clinician-referral, ACT-G ^a (n = 126)	Self-referral, iACT (n = 101)	t-test (d.f.)	Pearsons, χ^2 (d.f.)	P
Age, years: mean (s.d.)	36.2 (8.9)	39.6 (9.9)	2.69 (225)		0.008
Gender, n (%)			–	0.72 (1)	0.395
Women	89 (71)	66 (65)	–	–	–
Men	37 (29)	35 (35)	–	–	–
Living status, n (%) ^b			–	0.17 (2)	0.919
With someone	99 (79)	81 (80)	–	–	–
Alone	24 (19)	19 (19)	–	–	–
Other	2 (2)	1 (1)	–	–	–
Highest education, n (%) ^b			–	10.69 (4)	0.030
Unskilled	19 (15)	10 (10)	–	–	–
Skilled	21 (17)	12 (12)	–	–	–
Higher education (<4 years)	55 (45)	50 (50)	–	–	–
Higher education (>4 years)	15 (12)	25 (25)	–	–	–
Other	13 (11)	4 (4)	–	–	–
Employment, n (%) ^b			–	0.71 (3)	0.871
Employed or student	85 (68)	73 (72)	–	–	–
Unemployed	15 (12)	9 (9)	–	–	–
Disability pension or flexible work	10 (8)	8 (8)	–	–	–
Other (for example maternity leave)	15 (12)	11 (11)	–	–	–
Current sick leave, n (%) ^b			–	3.91 (3)	0.271
Full-time sick leave	19 (15)	7 (7)	–	–	–
Part-time sick leave	13 (10)	11 (11)	–	–	–
No absence	80 (64)	73 (72)	–	–	–
Not working	13 (10)	10 (10)	–	–	–

ACT-G, group-based acceptance and commitment therapy programme; iACT, internet-delivered acceptance and commitment therapy programme.
a. Eilenberg *et al.*¹⁹
b. Missing data in the ACT-G sample for living status (n = 1), highest education (n = 3), employment (n = 1) and current sick leave (n = 1).

in all, this suggests that self-referral can be an accurate method to recruit patients with health anxiety and possibly is non-inferior compared with clinician-referral.

Accuracy of patient self-referral

We found that 19% less clinician-referred patients were eligible for the diagnostic interview compared to the self-referred patients. This was partly because of more patients being excluded by clinicians at the research clinic (for example only presenting subclinical health anxiety symptoms) and partly because the clinician-referred patients more frequently failed to attend the diagnostic interview (i.e. declining participation without a reason or having improved in symptoms).

There can be several reasons for declining participation such as low motivation, lack of acceptance of the diagnosis and the treatment plan or barriers to assessment (for example transportation, going to a hospital). Self-referring patients may be more motivated when they initiate the treatment contact themselves, and barriers may be reduced when patients can undergo the diagnostic interview via video conferencing in their own home. However, many clinician-referred patients' symptoms also improved during the relatively longer waiting time for the diagnostic interview as compared with the faster procedure following self-referral. This might suggest that the more direct and faster pathway through self-referral may result in treatment initiation of some patients whose condition would have improved anyway. On the other hand, health anxiety tends to fluctuate but reoccur if left untreated.¹ Hence, self-referral may increase chances of timely treatment initiation.

There may be many possible explanations why self-referral seemed to be more accurate than clinician-referral in this study (for example motivation, less barriers to assessment and a faster procedure). As our results are based on only one trial applying a specific self-referral procedure with extensive patient educational material in a particular setting (Danish public healthcare system), we cannot draw any conclusions regarding self-referral for health

anxiety in general. Only one other trial relied solely on self-referral for recruiting patients with health anxiety.⁴⁰ In this internet-delivered cognitive-behavioural therapy trial, 495 patients commenced self-referring, 197 never completed the online procedure and only 102 patients were found eligible. This resulted in a total of 102/298 (34%) patients compared with 101/151 (67%) patients in our trial undertaking the treatment. Therefore, the accuracy of self-referral may depend on the individual procedure. Unfortunately, the self-referral procedure was not described by Newby and colleagues hampering a further comparison between the trials. However, it may well be that simply advertising is not sufficient. Our self-referral procedure was based on patient education material consisting both of clinical information and patient videos. This may have enhanced the accuracy in our study. Most clinical trials on health anxiety use a combination of self-referral and clinician-referrals from hospital settings or general practice.⁶ However, an efficient self-referral system has the potential to be cost-effective in that clinicians do not use resources related to referral, and therapists do not waste as much time waiting for unmotivated patients failing to attend an assessment.

Patient characteristics preceding recruitment

Self-referral has often been used as one recruitment method alongside clinician-referral in internet-delivered trials and results show that relevant patients tend to be captured across different psychiatric disorders.^{41–43} However, self-referred patients generally tend to be better educated,¹⁶ which was also the case in our study. We also found some support for the suggestion that self-referred patients may have a more distinct symptom profile in terms of less comorbidity and better physical-health-related quality of life compared with patients referred from primary care. This may seem contradictory since the self-referred patients also had higher levels of self-reported health anxiety symptoms and overall distress. However, the results also showed that self-referred patients were more reactive towards inner experiences compared with clinician-referred patients.³⁷ This is proposed to be a central mechanism in

Table 3 Clinical characteristics

Category	Clinician-referral, ACT-G ^a (n = 126)	Self-referral, iACT (n = 101)	Mean difference (95% CI)	t-test (d.f.)	Pearsons, χ^2 (d.f.)	z	P
Health anxiety							
Onset, years: mean (s.d.)	23.3 (10.6)	23.7 (12.0)	–	0.28 (223)	–	–	0.781
Duration, years: mean (s.d.)	13.0 (10.3)	15.9 (11.9)	–	–	–	–	–
Duration, years: median (IQR) ^b	11 (4–21)	12.5 (5.5–25)	–	–	–	1.79	0.073
Comorbidity, n (%)							
Anxiety disorder ^c	59 (47)	24 (24)	–	–	12.86 (1)	–	<0.001
Depressive disorder ^d	32 (25)	21 (21)	–	–	0.66 (1)	–	0.415
OCD ^e	9 (7)	3 (3)	–	–	–	–	0.234 ^f
Symptom profile, mean (s.d.) ^g							
Health anxiety							
Whiteley-7	69.3 (19.9)	74.9 (15.5)	5.6 (1.0 to 10.2)	–	–	–	0.018
Emotional distress							
SCL-8	52.2 (20.3)	57.5 (19.5)	5.3 (0.1 to 10.6)	–	–	–	0.047
Somatic symptoms							
SCL-SOM	36.4 (18.5)	41.5 (20.0)	5.1 (0.1 to 10.2)	–	–	–	0.047
OCD symptoms							
SCL-OCD	33.8 (18.3)	38.7 (20.4)	4.9 (–0.2 to 9.9)	–	–	–	0.061
Health-related quality of life							
SF-12 PCS	46.6 (10.6)	52.9 (12.2)	6.3 (3.1 to 9.4)	–	–	–	<0.001
SF-12 MCS	35.8 (9.7)	25.8 (11.1)	10.0 (7.1 to 12.9)	–	–	–	<0.001
Psychological flexibility							
AAQ-II	41.6 (21.7)	40.0 (17.7)	1.5 (–3.6 to 6.7)	–	–	–	0.560
Non-reactivity							
FFMQ non-react	31.3 (15.8)	25.2 (16.0)	6.0 (1.8 to 10.2)	–	–	–	0.005

ACT-G, group-based acceptance and commitment therapy programme; iACT, internet-delivered acceptance and commitment therapy programme; z, Wilcoxon rank-sum (Mann–Whitney); OCD, Obsessive-compulsive disorder; SCL, Symptom Checklist; SOM, Somatisation; SF-12, Short-Form Health Survey; PCS, Physical Component Summary; MCS, Mental Component Summary; AAQ, Acceptance and Action Questionnaire; FFMQ, Five Facet Mindfulness Questionnaire.

a. Eilenberg *et al.*¹⁹

b. Median and interquartile range reported because of a skewed distribution.

c. ICD-10 (F40.0–F40.2; F41.0–F41.1).

d. ICD-10 (F32.0–F32.2; F33.0–F33.2).

e. ICD-10 (F42.0–F42.1).

f. Fisher's exact test.

g. High score, more symptoms, except for health-related quality of life, psychological flexibility and non-reactivity.

health anxiety,^{38,39} and patients who are more reactive towards inner experiences might also be more inclined to search the internet for reassurance and seek treatment in a peak episode of health anxiety.⁴⁴ Therefore, the higher levels of distress could reflect that patients self-referred in a peak episode of health anxiety.

Self-referral may address patients with certain characteristics, or even require certain levels of illness recognition or health literacy to find and use a self-referral system.⁴⁵ Even though these speculations need further investigation, it is important to consider possible health inequalities if access is only granted through self-referral. On the other hand, only relying via clinician-referral may limit access to the more avoidant patients who tend to avoid health professionals and hospitals because of their fear of illness.

Strengths and limitations

The strengths of this study were the large number of participants from two RCTs carried out in the same research clinic. The amount of similar clinical data available in both trials enabled a comprehensive comparison. Furthermore, the empirically established diagnostic criteria and diagnostic interviews applied in the two trials ensured that the differences between the two patient groups were less likely to be caused by inclusion of different patient populations. The main limitation of this study was the different catchment areas entailing a risk that the patients may differ as a result of geographics rather than recruitment method. However, the catchment areas were both large (i.e. 2.5 to 5.7 million people), which reduces the risk considerably. Furthermore, the different recruitment periods with a few years in between might suggest that the two patient groups differ as a consequence of changing referral patterns over time rather than the different referral procedures and treatment preferences *per se*. However, ACT-G was also available in regular

care, during the period when iACT recruited patients through self-referral, and information about both treatments was available on the clinic's webpage and through the GPs. Thus, patients self-referring to iACT probably preferred this treatment format. Also, as for the clinician-referred patients, we do not know the relative proportion of GP and hospital clinician referrals. There could be a difference in accuracy between specialties, that is, GPs might be more literate in picking up health anxiety, which would mean the results do not generalise to all clinicians. This needs further investigation.

Finally, the slightly different eligibility criteria may have affected the characteristics of the included patients. The main difference in inclusion criteria between the trials was the age criteria, which turned out not to affect the results. Even though the two RCTs were not set up for this comparison, the large trials and overall comparability enabled an investigation of an important but largely unexplored area.


Clinical implications and future perspectives

Although the results of this study may primarily inform future study designs regarding the choice of recruitment method, they were based on two trials that were both embedded in a specialised hospital unit with a routine clinical function. Therefore, the results may also have some clinical implications. This study found that self-referral for psychological treatment seemed to work well in a patient population that most commonly seek medical reassurance or simply avoid the healthcare system. Patients with severe health anxiety were capable of preliminary self-assessment and self-referral to treatment. From a broader clinical perspective, self-referral and internet-delivered treatment ensure equal access to treatment for patients living in rural areas or patients simply not wishing to have face-to-face treatment. Instead of treating all patients similarly,

different ways of accessing treatment and modes of delivery enable a higher degree of individually tailored treatment. The increase in behavioural intervention technologies may have considerable implications for the delivery of future healthcare services.²²

National initiatives such as the English Improving Access to Psychological Therapy (IAPT) harness the internet to bridge the treatment gap by employing internet-delivered treatment and now self-referral for the treatment of depression and anxiety.⁴⁶ They have found that self-referred patients more accurately mirrored the general population including minority groups, who are often underrepresented among clinician-referrals. Harnessing technology in the dissemination of healthcare services has only just begun, and the future implications are numerous.⁴⁷ To start with, self-referral may generalise to other psychiatric disorders, and in the case of severe health anxiety, it may be a feasible supplement to clinician-referral. There is a need for studies directly comparing different recruitment methods in the same clinical trial to further investigate patient preferences and possible predictors for treatment outcome such as motivation.

In conclusion, patient self-referral and clinician-referral are both feasible and valid methods for recruiting patients with severe health anxiety although they may recruit patients with different characteristics. We found that the self-referred patients had higher levels of health anxiety and poorer mental health-related quality of life compared with the clinician-referred patients, underlining that self-referral is not primarily undertaken by people with mild symptoms and limited impairment. In addition, the self-referred patients were slightly older, better educated and had less comorbid anxiety disorders. Thus, allowing self-referral may lower barriers to treatment for a group of patients with considerable distress and a more distinct symptom profile.

Bitte Hoffmann , MSc, PhD, Postdoctoral Researcher, The Research Clinic for Functional Disorders and Psychosomatics, Aarhus University Hospital, Denmark; **Charlotte Ulrikka Rask**, MD, PhD, Professor, Child and Adolescent Psychiatric Centre, Risskov, Aarhus University Hospital, Denmark; **Erik Hedman-Lagerlöf**, MSc, PhD, Associate Professor, Department of Clinical Neuroscience, Karolinska Institute, Sweden; **Trine Eilenberg**, MSc, PhD, Clinical Psychologist, Department of Occupational Medicine, Aarhus University Hospital, Denmark; **Lisbeth Frostholt**, MSc, PhD, Associate Professor, The Research Clinic for Functional Disorders and Psychosomatics, Aarhus University Hospital, Denmark

Correspondence: Bitte Hoffmann, The Research Clinic for Functional Disorders and Psychosomatics, Aarhus University Hospital, Noerrebrogade 44, bldg. 2C, 1., 8000 Aarhus C, Denmark. Email: dittjese@rm.dk

First received 13 Nov 2018, final revision 28 Jun 2019, accepted 28 Jun 2019

Funding

This research was supported by the Danish foundation TrykFonden (grant numbers: 102644). The funding sources were not involved in designing and conducting the study or preparing this manuscript for publication.

Acknowledgements

We would like to thank PhD Tina Carstensen, psychologists Greta Lassen Lund and Katrine Ingeman Beck from the Research Clinic for Functional Disorders and Psychosomatics at Aarhus University Hospital for assisting with the diagnostic interviews and treatment. Also, we thank Jens S. Jensen for statistical assistance and Malene Skjøth for language revision. We would also like to thank all participants in our study.

References

- 1 Fink P, Ornbol E, Christensen KS. The outcome of health anxiety in primary care. A two-year follow-up study on health care costs and self-rated health. *PLoS One* 2010; **5**: e9873.
- 2 Tyrer P, Cooper S, Crawford M, Dupont S, Green J, Murphy D, et al. Prevalence of health anxiety problems in medical clinics. *J Psychosom Res* 2011; **71**: 392–4.

- 3 Fink P, Ornbol E, Toft T, Sparle KC, Frostholt L, Olesen F. A new, empirically established hypochondriasis diagnosis. *Am J Psychiatry* 2004; **161**: 1680–91.
- 4 Sunderland M, Newby JM, Andrews G. Health anxiety in Australia: prevalence, comorbidity, disability and service use. *Br J Psychiatry* 2013; **202**: 56–61.
- 5 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (5th edn) (DSM-5)*. American Psychiatric Association, 2013.
- 6 Cooper K, Gregory JD, Walker I, Lambe S, Salkovskis PM. Cognitive behaviour therapy for health anxiety: a systematic review and meta-analysis. *Behav Cogn Psychother* 2017; **45**: 110–23.
- 7 Perreault M, Lafortune D, Laverdure A, Chartier-Otis M, Belanger C, Marchand A, et al. Barriers to treatment access reported by people with anxiety disorders. *Can J Psychiatry* 2013; **58**: 300–5.
- 8 Marques L, LeBlanc NJ, Weingarden HM, Timpano KR, Jenike M, Wilhelm S. Barriers to treatment and service utilization in an internet sample of individuals with obsessive-compulsive symptoms. *Depress Anxiety* 2010; **27**: 470–5.
- 9 Rusch N, Angermeyer MC, Corrigan PW. Mental illness stigma: concepts, consequences, and initiatives to reduce stigma. *Eur Psychiatry* 2005; **20**: 529–39.
- 10 Soucy JN, Hadjistavropoulos HD. Treatment acceptability and preferences for managing severe health anxiety: perceptions of internet-delivered cognitive behaviour therapy among primary care patients. *J Behav Ther Exp Psychiatry* 2017; **57**: 14–24.
- 11 Kohn R, Saxena S, Levav I, Saraceno B. The treatment gap in mental health care. *Bull World Health Organ* 2004; **82**: 858–66.
- 12 Tyrer P, Eilenberg T, Fink P, Hedman E, Tyrer H. Health anxiety: the silent, disabling epidemic. *BMJ* 2016; **353**: i2250.
- 13 Pichlhofer O, Maier M. Unregulated access to health-care services is associated with overutilization—lessons from Austria. *Eur J Public Health* 2015; **25**: 401–3.
- 14 Hoffmann D, Rask CU, Hedman-Lagerlöf E, Ljótsson B, Frostholt L. Development and feasibility testing of internet-delivered acceptance and commitment therapy for severe health anxiety: pilot study. *JMIR Ment Health* 2018; **5**: e28.
- 15 Andrews G, Basu A, Cuijpers P, Craske MG, McEvoy P, English CL, et al. Computer therapy for the anxiety and depression disorders is effective, acceptable and practical health care: an updated meta-analysis. *J Anxiety Disord* 2018; **55**: 70–8.
- 16 Andersson G, Titov N. Advantages and limitations of internet-based interventions for common mental disorders. *World Psychiatry* 2014; **13**: 4–11.
- 17 Kim EH, Linker DT, Coumar A, Dean LS, Matsen FA, Kim Y. Factors affecting acceptance of a web-based self-referral system. *IEEE Trans Inf Technol Biomed* 2011; **15**: 344–7.
- 18 Brown JS, Boardman J, Whittinger N, Ashworth M. Can a self-referral system help improve access to psychological treatments? *Br J Gen Pract* 2010; **60**: 365–71.
- 19 Eilenberg T, Fink P, Jensen JS, Rief W, Frostholt L. Acceptance and commitment group therapy (ACT-G) for health anxiety: a randomized controlled trial. *Psychol Med* 2016; **46**: 103–15.
- 20 World Health Organization. *SCAN: Schedules for Clinical Assessment in Neuropsychiatry (Version 2.1 edn)*. World Health Organization, Division of Mental Health, 1998.
- 21 Dantoft TM, Ebstrup JF, Linneberg A, Skovbjerg S, Madsen AL, Mehlsen J, et al. Cohort description: the Danish study of Functional Disorders. *Clin Epidemiol* 2017; **9**: 127–39.
- 22 Mohr DC, Burns MN, Schueller SM, Clarke G, Klinkman M. Behavioral intervention technologies: evidence review and recommendations for future research in mental health. *Gen Hosp Psychiatry* 2013; **35**: 332–8.
- 23 Hajebi A, Motevalian A, Amin-Esmaeili M, Hefazi M, Radgoodarzi R, Rahimi-Movaghar A, et al. Telephone versus face-to-face administration of the Structured Clinical Interview for Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, for diagnosis of psychotic disorders. *Compr Psychiatry* 2012; **53**: 579–83.
- 24 Fink P, Ewald H, Jensen J, Sorensen L, Engberg M, Holm M, et al. Screening for somatization and hypochondriasis in primary care and neurological in-patients: a seven-item scale for hypochondriasis and somatization. *J Psychosom Res* 1999; **46**: 261–73.
- 25 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR)* 4th ed. text rev. APA, 2000.
- 26 World Health Organization, Collaborating Centres for Classification of Diseases. *ICD-10: International Statistical Classification of Diseases and Related Health Problems*. WHO, 2014.
- 27 Olsen LR, Mortensen EL, Bech P, Elsass P, Ivanou J, Poulsen S, et al. Symptom Checklist (SCL-90, SCL-90-R og SCL-92. In *Assessmentmetoder. Håndbog for psykologer og psykiatere* [Assessment Methods. Handbook for Psychologists and Psychiatrists] (1st edn) (eds LR Olsen, EL Mortensen, P Bech, P Elsass, J Ivanou, S Poulsen, et al.): 183. Dansk Psykologisk Forlag, 2006.

- 28 Fink P, Ornbol E, Huysse FJ, De Jonge P, Lobo A, Herzog T, et al. A brief diagnostic screening instrument for mental disturbances in general medical wards. *J Psychosom Res* 2004; **57**: 17–24.
- 29 Fink P, Ornbol E, Hansen MS, Sondergaard L, De Jonge P. Detecting mental disorders in general hospitals by the SCL-8 scale. *J Psychosom Res* 2004; **56**: 371–5.
- 30 Derogatis LR, Cleave PA. Confirmation of the dimensional structure of the SCL-90: a study in construct validation. *J Clin Psychol* 1977; **33**: 981–9.
- 31 Kim SW, Dysken MW, Kuskowski M. The symptom checklist-90: obsessive-compulsive subscale: a reliability and validity study. *Psychiatry Res* 1992; **41**: 37–44.
- 32 Bjorner JB, Damsgaard MT, Watt T, Groenvold M. Tests of data quality, scaling assumptions, and reliability of the Danish SF-36. *J Clin Epidemiol* 1998; **51**: 1001–11.
- 33 Ware Jr JE. SF-36 Health Survey Update. In *The Use of Psychological Testing for Treatment Planning and Outcome Assessment* (3rd edn) (ed ME Maruish): 693–718. Lawrence Erlbaum Associates Publishers, 2004.
- 34 Christensen LN, Ehlers L, Larsen FB, Jensen MB. Validation of the 12 item short form health survey in a sample from region central Jutland. *Soc Indic Res* 2013; **114**: 513–21.
- 35 Bond FW, Hayes SC, Baer RA, Carpenter KM, Guenole N, Orcutt HK, et al. Preliminary psychometric properties of the Acceptance and Action Questionnaire-II: a revised measure of psychological inflexibility and experiential avoidance. *Behav Therapy* 2011; **42**: 676–88.
- 36 Hayes SC, Strosahl K, Wilson KG, Bissett RT, Pistorello J, Toarmino D, et al. Measuring experiential avoidance: a preliminary test of a working model. *Psychol Rec* 2004; **54**: 553–78.
- 37 Baer RA, Smith GT, Lykins E, Button D, Krietemeyer J, Sauer S, et al. Construct validity of the five facet mindfulness questionnaire in meditating and nonmeditating samples. *Assessment* 2008; **15**: 329–42.
- 38 Eilenberg T, Hoffmann D, Jensen JS, Frostholm L. Intervening variables in group-based acceptance & commitment therapy for severe health anxiety. *Behav Res Ther* 2017; **92**: 24–31.
- 39 Hedman E, Hesser H, Andersson E, Axelsson E, Ljotsson B. The mediating effect of mindful non-reactivity in exposure-based cognitive behavior therapy for severe health anxiety. *J Anxiety Disord* 2017; **50**: 15–22.
- 40 Newby JM, Smith J, Uppal S, Mason E, Mahoney AEJ, Andrews G. Internet-based cognitive behavioral therapy versus psychoeducation control for illness anxiety disorder and somatic symptom disorder: a randomized controlled trial. *J Consult Clin Psychol* 2018; **86**: 89–98.
- 41 Titov N, Andrews G, Kemp A, Robinson E. Characteristics of adults with anxiety or depression treated at an internet clinic: comparison with a national survey and an outpatient clinic. *PLoS One* 2010; **5**: e10885.
- 42 Spence J, Titov N, Solley K, Dear BF, Johnston L, Wootton B, et al. Characteristics and treatment preferences of people with symptoms of posttraumatic stress disorder: an internet survey. *PLoS One* 2011; **6**: e21864.
- 43 Wootton BM, Titov N, Dear BF, Spence J, Kemp A. The acceptability of internet-based treatment and characteristics of an adult sample with obsessive compulsive disorder: an internet survey. *PLoS One* 2011; **6**: e20548.
- 44 Muse K, McManus F, Leung C, Meghreblian B, Williams JM. Cyberchondriasis: fact or fiction? A preliminary examination of the relationship between health anxiety and searching for health information on the Internet. *J Anxiety Disord* 2012; **26**: 189–96.
- 45 Bo A, Friis K, Osborne RH, Maindal HT. National indicators of health literacy: ability to understand health information and to engage actively with healthcare providers - a population-based survey among Danish adults. *BMC Public Health* 2014; **14**: 1095.
- 46 Clark DM. Realizing the mass public benefit of evidence-based psychological therapies: the IAPT program. *Annu Rev Clin Psychol* 2018; **14**: 159–83.
- 47 Holmes EA, Ghaderi A, Harmer CJ, Ramchandani PG, Cuijpers P, Morrison AP, et al. The Lancet Psychiatry Commission on psychological treatments research in tomorrow's science. *Lancet Psychiatry* 2018; **5**: 237–86.

