

Letter to the Editor

Visual scale for the quantification of hyperhidrosis

Escala visual para a quantificação da hiper-hidrose

Roberto de Menezes Lyra

To the Editor:

In most cases, patients with hyperhidrosis have unpleasant sensory and emotional experiences in the social sphere, experiences that are difficult to express to physicians in a simple way and to be understood by society. In contrast, assessing and quantifying

hyperhidrosis clinically is not an easy task; therefore, I have read with great enthusiasm the study by Sakiyama et al. describing a quantitative assessment of the intensity of palmar and plantar sweating in patients with primary palmoplantar hyperhidrosis.⁽¹⁾

Visual scale for the quantification of hyperhidrosis




Figure 1 – Visual scale for the quantification of hyperhidrosis.

A measuring device will certainly add to improving the quantification of hyperhidrosis during clinical assessment. This is because the clinical presentation of hyperhidrosis varies widely from person to person, although there is a certain pattern of presentation in localized hyperhidrosis. It is assumed that this is due in part to the large anatomical variation in the sympathetic chain and also to the fact that the sympathetic nervous system shows a metameric distribution that is not so precise, which makes it difficult to be mapped as dermatomes.^(2,3)

Formerly, in order to quantify hyperhidrosis, we had no other way but to measure it clinically with a visual scale by means of which the patient was asked to grade the intensity of hyperhidrosis on the basis of drawings in the scale, ranging from being normal to having the worst possible level of hyperhidrosis. Therefore, the intensity of hyperhidrosis at the initial assessment, as well as at each assessment and, if necessary, also in the postoperative period, can be recorded on the medical charts.⁽⁴⁾

In addition, we quantify the presentation of hyperhidrosis, whether it is generalized or localized, and the type of body distribution (Figure 1). We use hachures to demarcate the areas of excessive sweating on the drawings of the human body. Subsequently, using the visual scales shown in Figure 1, we determine the amount of sweat produced and the type of localized hyperhidrosis (palmar, axillary, plantar, or any combination of the three) presented by the patient.

Physiologically, there is great variability in sweat production, which depends on environmental

heat, seasons, physical exertion, stress, and even circadian cycle. Therefore, it is of the utmost importance that the patient guide us and quantify the areas that should be demarcated, thereby providing a strong foundation for reasoning about the choice of clinical or surgical treatment of hyperhidrosis.⁽⁵⁾

Roberto de Menezes Lyra
Thoracic Surgeon, *Hospital do Servidor*
***Público Estadual de São Paulo* – HSPE/**
SP, São Paulo Hospital for State Civil
Servants – São Paulo, Brazil

References

1. Sakiyama BY, Monteiro TV, Ishy A, de Campos JR, Kauffman P, Wolosker N. Quantitative assessment of the intensity of palmar and plantar sweating in patients with primary palmo-plantar hyperhidrosis. *J Bras Pneumol.* 2008;38(5):573-8. <http://dx.doi.org/10.1590/S1806-37132012000500006>
2. Zhang B, Li Z, Yang X, Li G, Wang Y, Cheng J, Tang X, Wang F. Anatomical variations of the upper thoracic sympathetic chain. *Clin Anat.* 2009;22(5):595-600. <http://dx.doi.org/10.1002/ca.20803> PMID:19418453
3. Schünke M, Schulte E, Schumacher U. *Dores viscerais.* In: Prometheus, atlas de anatomia: cabeça e neuroanatomia. Rio de Janeiro: Guanabara Koogan; 2007. p. 322-3
4. Lyra Rde M, Campos JR, Kang DW, Loureiro Mde P, Furian MB, Costa MG, et al. Guidelines for the prevention, diagnosis and treatment of compensatory hyperhidrosis. *J Bras Pneumol.* 2008;34(11):967-77. PMID:19099105
5. Boscardim PC, Oliveira RA, Oliveira AA, Souza JM, Carvalho RG. Thoracic sympathectomy at the level of the fourth and fifth ribs for the treatment of axillary hyperhidrosis. *J Bras Pneumol.* 2011;37(1):6-12. <http://dx.doi.org/10.1590/S1806-37132011000100003> PMID:21390426