

RESEARCH ARTICLE

# The Bacteriohopanepolyol Inventory of Novel Aerobic Methane Oxidising Bacteria Reveals New Biomarker Signatures of Aerobic Methanotrophy in Marine Systems

Darci Rush<sup>1</sup>\*, Kate A. Osborne<sup>1</sup>, Daniel Birgel<sup>2</sup>, Andreas Kappler<sup>3,4</sup>, Hisako Hirayama<sup>5</sup>, Jörn Peckmann<sup>2,6</sup>, Simon W. Poulton<sup>7</sup>, Julia C. Nickel<sup>8</sup>, Kai Mangelsdorf<sup>8</sup>, Marina Kalyuzhnaya<sup>9</sup>, Frances R. Sidgwick<sup>1</sup>, Helen M. Talbot<sup>1</sup>



**1** School of Civil Engineering & Geosciences, Newcastle University, Drummond Building, Newcastle upon Tyne, NE1 7RU, Newcastle-upon-Tyne, United Kingdom, **2** Institute of Geology, University of Hamburg, Hamburg, Germany, **3** Center for Applied Geoscience, University of Tübingen, Tübingen, Germany, **4** Center for Geomicrobiology, Department of Bioscience, Ny Munkegade 116, 8000, Aarhus C, Denmark, **5** Department of Subsurface Geobiological Analysis and Research, Japan Agency for Marine-Earth Science & Technology (JAMSTEC), Yokosuka, Japan, **6** Department of Geodynamics and Sedimentology, University of Vienna, 1090, Vienna, Austria, **7** School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, United Kingdom, **8** GFZ German Research Centre for Geosciences, Telegrafenberg, D-14473, Potsdam, Germany, **9** Faculty of Biology, San Diego State University, 5500 Campanile Drive, San Diego, 92182, United States of America

OPEN ACCESS

**Citation:** Rush D, Osborne KA, Birgel D, Kappler A, Hirayama H, Peckmann J, et al. (2016) The Bacteriohopanepolyol Inventory of Novel Aerobic Methane Oxidising Bacteria Reveals New Biomarker Signatures of Aerobic Methanotrophy in Marine Systems. PLoS ONE 11(11): e0165635. doi:10.1371/journal.pone.0165635

**Editor:** Zhe-Xue Quan, Fudan University, CHINA

**Received:** September 5, 2016

**Accepted:** October 15, 2016

**Published:** November 8, 2016

**Copyright:** © 2016 Rush et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The work was funded by a Starting Grant (No. 258734) awarded to HMT for project AMOPROX, from the European Research Council. KM and JCN thank Lundin Petroleum for financial support. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

\* These authors contributed equally to this work.

\* [darci.rush@nioz.nl](mailto:darci.rush@nioz.nl)

## Abstract

Aerobic methane oxidation (AMO) is one of the primary biologic pathways regulating the amount of methane (CH<sub>4</sub>) released into the environment. AMO acts as a sink of CH<sub>4</sub>, converting it into carbon dioxide before it reaches the atmosphere. It is of interest for (paleo)climate and carbon cycling studies to identify lipid biomarkers that can be used to trace AMO events, especially at times when the role of methane in the carbon cycle was more pronounced than today. AMO bacteria are known to synthesise bacteriohopanepolyol (BHP) lipids. Preliminary evidence pointed towards 35-aminobacteriohopane-30,31,32,33,34-pentol (aminopentol) being a characteristic biomarker for Type I methanotrophs. Here, the BHP compositions were examined for species of the recently described novel Type I methanotroph bacterial genera *Methylomarinum* and *Methylomarinovum*, as well as for a novel species of a Type I *Methylomicrobium*. Aminopentol was the most abundant BHP only in *Methylomarinovum caldicuralii*, while *Methylomicrobium* did not produce aminopentol at all. In addition to the expected regular aminotriol and aminotetrol BHPs, novel structures tentatively identified as methylcarbamate lipids related to C-35 amino-BHPs (MC-BHPs) were found to be synthesised in significant amounts by some AMO cultures. Subsequently, sediments and authigenic carbonates from methane-influenced marine environments were analysed. Most samples also did not contain significant amounts of aminopentol, indicating that aminopentol is not a useful biomarker for marine aerobic

**Competing Interests:** The authors have declared that no competing interests exist.

methanotrophic bacteria. However, the BHP composition of the marine samples do point toward the novel MC-BHPs components being potential new biomarkers for AMO.


## Introduction

Methane (CH<sub>4</sub>) is a potent greenhouse gas, and its atmospheric concentration has tripled since pre-industrial times (e.g. [1,2]). Global oceans hold large subsurface reservoirs of CH<sub>4</sub> in the form of gas hydrates. These stores are precariously dependent on temperature and pressure. A rapid destabilisation of gas hydrates has been proposed to have caused vast releases of marine CH<sub>4</sub> in the past [3]. Increased input of CH<sub>4</sub> into the atmosphere has been interpreted through records of excursions of significant δ<sup>13</sup>C depletion in the geological record, such as in the Palaeocene-Eocene Thermal Maximum (PETM) [4–7].

CH<sub>4</sub> release into the atmosphere is regulated by methanotrophy, which converts CH<sub>4</sub> into CO<sub>2</sub>, thereby playing a key role in the carbon biogeochemical cycle [8]. Although traditionally anaerobic archaea have been the most studied methanotrophs (e.g., AMNE-1 and ANME-2; cf. [9]), recent observations have highlighted the importance of bacteria performing aerobic CH<sub>4</sub> oxidation (AMO) in marine, estuarine, and riverine fan environments (e.g., [10–14]). For example, pelagic AMO activity rose significantly after the Macondo oil well blowout in 2010 [15]. However, this activity was short-lived, highlighting the complexity of natural community interactions in response to increased CH<sub>4</sub> [16]. It is thus important to recognise and trace methanotrophy during past extreme events in order to understand its potential to mitigate future CH<sub>4</sub> release.

AMO bacteria belong to two phyla, *Proteobacteria* and *Verrucomicrobia*. Most isolates of *Verrucomicrobia* are thermoacidophilic [17–20], and have been found primarily in acidic, geothermal environments [21]. Aerobic methanotrophic members of *Proteobacteria* belong to two distinct classes, separated based on their carbon assimilation pathways [8]. Type II methanotrophs, members of the *Alphaproteobacteria*, are associated with terrestrial settings ([8], and references therein), whereas Type I methanotrophs members of the *Gammaproteobacteria* are widespread in aquatic systems, although they are also found in terrestrial systems. Both Type I and Type II methanotrophs are known to synthesise bacteriohopanepolyol (BHP) lipids [22]. BHPs are precursors to hopanes, which are the most ubiquitous lipids in the geological record [23,24]. Therefore, being able to trace AMO using hopanoid biomarkers is of value to the study of the carbon cycle in the past.

Previous work exploring hopanoids as biomarker lipids found that methylation at the C-3 position (3-Me-BHPs; Fig 1, I<sup>3Me</sup>, II<sup>3Me</sup>, III<sup>3Me</sup>) was a possible indication of methanotrophic origin [25,26]. However, attributing C-3 methylation to AMO was challenged by the revelation that bacteria other than methanotrophs have the genes to methylate at C-3 [27]. C-3 methylation is more likely a requirement for cell survival in late stationary phase [27]. Moreover, not all methanotrophic bacteria have the gene to methylate at this position, [27,28], nor are 3-Me BHP precursors found in all CH<sub>4</sub>-influenced environments (Table 1). However, AMO bacteria are often considered to be the most likely source of 3-Me hopanoids in marine sediments due to their depleted carbon isotope signatures (e.g., [29]) and because they are frequently accompanied by 4-methylated steroids, which are also known as biomarkers of methanotrophic bacteria (e.g., [30,31]). In other studies, diplopterol and diploptene have been interpreted as biomarkers for methanotrophy, notably in anoxic environments [32–35]. However, neither diplopterol nor diploptene is source-specific to AMO [36], and these studies also relied on very



**Fig 1. Chemical structures of bacteriohopanepolyol lipids.** I, aminopentol; I<sup>3Me</sup>, 3-methyl-aminopentol; I<sup>MC</sup>, methylcarbamate-aminopentol; II, aminotetrol; II<sup>3Me</sup>, 3-methyl-aminotetrol; II<sup>MC</sup>, methylcarbamate-aminotetrol; III, aminotriol; III<sup>3Me</sup>, 3-methyl-aminotriol; III<sup>MC</sup>, methylcarbamate-aminotriol. The proposed structure of methylcarbamate(MC)-aminopentol (I<sup>MC</sup>), MC-aminotetrol (II<sup>MC</sup>), and MC-aminotriol (III<sup>MC</sup>) are tentatively based on mass spectral identification (S1 File).

doi:10.1371/journal.pone.0165635.g001

negative  $\delta^{13}\text{C}$  lipid values (e.g., ca. -61‰ for diplopterol; -61 to -74‰ for diploptene) as an indication of  $\text{CH}_4$  being the carbon source for the organism producing these lipids. Yet, recent work has emphasised that not all AMO-derived carbon shows a depleted isotope signature, especially in terrestrial systems where Type II methanotrophs tend to dominate. For example, only limited depletion in  $^{13}\text{C}$ , with values between -25‰ and -40‰, have been reported for hopanoids with an inferred methanotrophic origin from modern peat bogs [37,38] as well as other ancient lignite deposits [39]. Also, BHPs in Congo deep sea fan sediments, originating from low-latitude wetlands [40], had higher  $^{13}\text{C}$  values than expected (i.e., C<sub>30</sub> hopanol ~ -41‰; [14]). In marine  $\text{CH}_4$  seep carbonates from the Gulf of Mexico, BHPs, hopanoic acids, and 4-methylated sterols of aerobic methanotrophs were found with similar  $\delta^{13}\text{C}$  values as the  $\text{CH}_4$  source [41]. These observations can be the result of dilution from other heterotrophic bacterial sources that make it difficult to identify subordinate methanotroph contributions [37]. The metabolic pathways used by the AMO bacteria for  $\text{CH}_4$  assimilation can also have a profound effect on the level of isotopic depletion, with values for serine pathway methanotrophs (Type II) ranging from 12‰ depleted to 10‰ enriched relative to the  $\text{CH}_4$  substrate [42]. Furthermore, to analyse  $\delta^{13}\text{C}$  of intact BHPs, these must first be converted into primary alcohols by periodic acid/sodium borohydride cleavage [43]. For example, the  $\delta^{13}\text{C}$  value of 35-aminobacteriohopane-30,31,32,33,34-pentol (aminopentol; Fig 1, I) is measured on the C<sub>30</sub>-hopanol product, which includes all converted hexa-functionalised BHPs (i.e., not only aminopentol), as well as any free-hopanols that are present in a sample before BHP conversion. While the contamination of free-hopanols can be circumnavigated by column separations (e.g., [41]), measuring the  $\delta^{13}\text{C}$  values of intact BHPs is not currently possible.

Aminopentol is thought to be the most diagnostic BHP for AMO (see review in [14]). Aminopentol and its methylated and unsaturated homologues (i.e., I<sup>3Me</sup>,  $\Delta$ I) have been found

**Table 1. Presence and absence of aminopentol and related methylated and unsaturated homologues in previously investigated environmental settings.**

Sample	aminopentol			aminotetrol		aminotriol		Reference
	I	ΔI	I <sup>3Me</sup>	II	II <sup>3Me</sup>	III	III <sup>3Me</sup>	
<b>Soils</b>								
Pasture [manured] (UK)	+	-	-	+	-	+	+	[112]
Pasture [not manured] (UK)	-	-	-	+	-	+	+	[112]
Rice Paddy (Vietnam)	+	-	-	+	-	+	-	[112]
Woodland (North East England)	+	-	-	+	-	+	+	[112]
South West France	+	-	-	+	-	+	-	[116]
Amazon	+	-	-	+	-	+	-	[109]
Congo	+	+	-	+	-	+	+	[40]
Lean Delta Peamafrost [ice complex]	+	-	-	+	-	+	-	[110]
Human Sewage [treated]	-	-	-	+	-	+	-	[117]
Forest, Grassland Soils (Alberta, Canada)	-	-	-	+	-	+	-	[118]
<b>Peat and Lignite</b>								
River Tet Catchment (France)	+	-	-	+	-	+	-	[116]
Moorhouse (UK)	+	-	-	+	-	+	-	[46]
Misten Bog (Belgium)	+	-	-	+	-	+	-	[119]
Bisendorfer Moor (Germany)	+	-	-	+	-	+	+	[111]
The Cobham Lignite (UK)	+	-	-	+	-	+	-	[49]
<b>Geothermal Environments</b>								
Cyanobacterial mat (Surprise Valley, Nevada, USA)	+	-	-	+	-	+	-	[120]
Silica Sinter (Orakie Korako, Taupo Volcanic Zone, New Zealand)	+	-	+	+	-	+	-	[113]
Silica Sinter (Champagne Pool, Taupo Volcanic Zone, New Zealand)	-	-	-	-	-	+	-	[121]
Orange mat (Yellowstone, USA)	-	-	-	-	-	+	-	[122]
<b>Other Microbial mats</b>								
Mars Oasis (Antarctica)	-	-	-	-	-	+	-	[48]
Hypolith (Devon Island, Arctic)	-	-	-	-	-	+	-	[48]
Cyanobacterial Mat (Christmas Island, Kiribati)	-	-	-	-	-	+	-	[123]
<b>Lake Sediments</b>								
Lake Windermere (UK)	+	-	-	+	-	+	-	¶
Lake Cadagno (Switzerland)	+	-	-	+	-	+	-	[26]
Lake Holzmaar (Germany)	+	-	-	+	-	+	-	[26]
Lake Nkunga (Kenya)	+	-	-	+	-	+	-	[45]
Priest Pot (England)	+	-	-	+	-	+	-	[26,45]
La Piscina de Yuriria (Mexico)	+	-	+	+	+	+	-	[26,45]
Laguna de Zempoala (Mexico)	+	-	-	+	-	+	-	[26]
Loch Ness (UK)	+	-	-	+	-	+	-	[26,45]
Lake Druzby (Antarctica)	+	-	-	+	-	+	-	[26,45]
Sombre Lake (Signy Island, Southern Atlantic Ocean)	+	-	-	+	-	+	-	[26]
Heywood Lake (Signy Island, Southern Atlantic Ocean)	+	-	-	+	-	+	-	[26]
Ace Lake [Freshwater unit]	+	-	-	+	-	+	-	[104]
Ace Lake [meromictic unit]	-	-	-	+	-	+	-	[104]
Ace Lake [marine unit]	-	-	-	-	-	+	-	[104]
<b>Wetlands</b>								
Amazon	+	+	-	+	-	+	-	[109]
Congo	+	+	-	+	-	+	-	[40]
<b>Stream/River/Estuary Sediments</b>								

(Continued)

Table 1. (Continued)

Sample	aminopentol			aminotetrol		aminotriol		Reference
	I	ΔI	I <sup>3Me</sup>	II	II <sup>3Me</sup>	III	III <sup>3Me</sup>	
Arctic Rivers (Indigirka, Kolyma, Lena, Ob, Yenisei, Yukon, Mackenzie)	+	-	-	+	-	+	-	[124]
Glacial Outflow Stream (Svalbard)	+	-	-	+	-	+	-	[125]
Kalix River [surface sediment transect]	+	-	-	+	-	+	-	[126]
Kolyma River [surface sediment transect]	+	-	-	+	-	+	-	[127]
Yenisei River [mouth, and surrounding area]	+	-	-	+	-	+	-	[128]
Yangtze River [estuary, inner shelf]	+	-	-	+	-	+	-	[82,129]
Congo River [estuary]	+	+	-	+	-	+	-	[14,40]
<b>Water Column</b>								
Priest Pot [pond] (UK)	+	-	-	+	-	+	-	¶
River Water (Panama)	+	-	-	+	-	+	-	[130]
Yenisei River	+	-	-	+	-	+	-	[128]
Black Sea [oxic/anoxic transition zone]	+	-	+	+	-	+	-	[12]
Black Sea [oxic/anoxic transition zone]	+	-	-	+	-	+	-	[108]
Gotland Deep, Baltic Sea	+	-	-	+	-	+	-	[107]
Marine Water (off Panama coast)	-	-	-	-	-	+	-	[130]
Pelagic [sub-oxic and anoxic] (Arabian Sea, Peru Margin, Cariaco Basin)	-	-	-	-	-	+	-	[90]
Cariaco Basin	-	-	-	-	-	+	-	[131]
California Current	-	-	-	-	-	+	-	[91]
<b>Modern/Recent Marine Sediments</b>								
Gotland Deep, Baltic Sea	+	-	-	+	-	+	-	[107]
Congo River Deep-Sea Fan	+	+	-	+	-	+	-	[14]
Amazon Shelf and Fan	+	-	-	+	-	+	-	[109]
East Siberian Arctic Shelf	+	-	-	+	-	+	-	[79,126]
Black Sea	-	-	-	+	-	+	-	[88]
Southwest African Coast	-	-	-	+	-	+	(+)	[89]
Chukchi Sea	-	-	-	+	-	+	-	[132]
Alaskan Beaufort Sea	-	-	-	-	-	+	-	[132]
Baltic Sea	-	-	-	+	-	+	-	[133]
<b>Marine Carbonates</b>								
Carbonate, Gulf of Mexico	-	-	-	+	-	+	-	[72]
Authigenic carbonates, Gulf of Mexico	-	-	-	+	-	+	-	[41]
Seep Carbonate, Arabian Sea	-	-	-	+	-	+	-	[106]
<b>Other samples</b>								
Membrane Foulant [river water]	+	-	-	-	-	+	-	[134]
Membrane Foulant [brackish water]	+	-	-	+	-	+	-	[134]
Membrane Foulants [seawater]	-	-	-	-	-	+	-	[134]
Mussel Gill Tissue	-	-	-	+	-	+	-	[103]

+, detected in at least one of the samples; -, not detected in any of the samples

(+) indicates a methylated aminotriol was detected but the position of methylation was not identified

¶Talbot and Farrimond, unpublished data

doi:10.1371/journal.pone.0165635.t001

almost exclusively in Type I aerobic methanotrophs [22,44–46]. Moreover, aminopentol (I) has been found in a wide range of environments, which indicates potential as a biomarker for AMO (Table 1). Additionally, 35-aminobacteriohopane-31,32,33,34-tetrol (aminotetrol, II) and 35-aminobacteriohopane-32,33,34-triol (aminotriol, III) are also synthesised by Type I

and Type II AMO bacteria. However, **II** and **III** are less source-specific as both are synthesised by some species of sulfate reducing bacteria (SRB) of the genus *Desulfovibrio* [47], and **III** is synthesised by many other aerobic bacteria ([48], and references therein).

Aminopentol made up a very minor proportion of the BHP composition in a SRB culture (<0.1% of total BHPs in *Desulfovibrio salexigens*). The ratio of aminopentol to the more ubiquitous aminotriol was 1:1352 [47]. We can therefore discount SRB as the source of aminopentol in an environmental sample with a high ratio of aminopentol:aminotriol. A similar approach was recently used by [49] based on the ratio of aminotetrol:aminotriol which has been found in the range 1:20–100 in some species of *Desulfovibrio* SRB [47,50,51]. Interestingly, some of the species of SRB cultures analysed by Blumenberg et al. [47,50,51] also synthesised diploptol and diploptene, which could explain the enhanced presence of these lipids in CH<sub>4</sub>-influenced anoxic sites.

Representatives from only a small number of Type I methanotroph genera have been tested for BHP production (e.g., [22,36,46,52,53]). Many more recently described genera, including novel genera from marine and other (hyper)saline environments, have yet to be explored (e.g., [54–56]). Moreover, relatively few studies of BHP distributions have targeted marine environments (Table 1). It is important, therefore, to determine whether aminopentol, which is seen as a diagnostic marker for Type I methanotrophs, is present in methanotrophs isolated from marine and other saline environments, and whether we can find aminopentol in CH<sub>4</sub>-influenced marine sediments. This knowledge will have implications for the use of aminopentol as a biomarker to trace AMO in modern and ancient marine environments.

In this study, our goal was to develop an appropriate biomarker approach for AMO, which will allow high throughput analysis of sediment without the requirement for laborious chemical conversion steps prior to compound specific isotope analysis. To this end, we screened the BHP distributions of three genera of aerobic methanotrophs (i.e., *Methylomicrobium*, *Methylomarinum* and *Methylomarinovum*). *Methylomarinum* and *Methylomarinovum* have not previously been investigated for BHPs. Two species of *Methylomicrobium* have been described previously [52,53], however, we include an additional species *Methylomicrobium kenyense*. These data are combined with literature BHP distributions of other AMO genera, including the recently reported Type I genus, *Methylobacter* [28], in order to facilitate interpretation of BHP distributions in six selected modern marine sediments from CH<sub>4</sub>-influenced systems. Two marine sites not influenced by CH<sub>4</sub> were also investigated as controls for background marine BHP signatures.

## Methods

### Methanotroph pure cultures

**Cultivation of *Methylomarinum vadi* IT-4.** Previously described *Methylomarinum vadi* IT-4 was isolated from a microbial mat sample (in-situ temperature 30–40°C) collected at a shallow marine hydrothermal system (depth, ~23 m) in a coral reef off Taketomi Island, Okinawa, Japan [56,57]. Cultivation of this strain was performed at JAMSTEC, Japan, using MJmet medium at pH 6.6 at 37°C. A detailed site description and the enrichment and isolation procedures can be found in [57].

**Cultivation of *Methylomarinovum* spp.** Previously described *Methylomarinovum caldicuralii* IT-9 was isolated from the hot vent fluid (52°C) collected at the main vent site (depth, 23 m) in the shallow marine hydrothermal system where *Methylomarinum vadi* IT-4 was isolated [55,57]. *Methylomarinovum* sp. IN45 is a new isolate from a deep-sea hydrothermal field in Okinawa Trough, Japan (H. Hirayama, pers. comm.). The cultivation of strains IT-9 and IN45 was performed at JAMSTEC, Japan, using MJmet medium at 45°C, and at pH 6.2 and 6.6, respectively.

**Cultivation of *Methylomicrobium* spp.** *Methylomicrobium alcaliphilum* and *Methylomicrobium kenyense* were first isolated from highly alkaline soda lakes in Russia and Kenya, respectively [58]. Both *M. alcaliphilum* (DSM-number 19304) and *M. kenyense* (DSM-number 19305) are from the Leibniz Institute DSMZ (German Collection of Microorganisms and Cell Cultures). The cultivation of both strains was done at the Center for Applied Geosciences at the University of Tübingen, Germany, at pH 9.1 and at 28°C with a high salt NMS medium (1.5% NaCl).

## Marine sediment and carbonate samples

**Håkon Mosby Mud Volcano (HMMV).** HMMV has been extensively studied for both aerobic and anaerobic methanotrophy [13,59]. The flow of CH<sub>4</sub> in the center of the HMMV is restricted, and AMO was observed to be the most dominant process within the centre of the crater, performed by Type I methanotrophs [60]. Sediments from the HMMV were collected aboard the *RV Polarstern* (cruise ARK XXIV/2; 2009) and the *RV Maria S Merian* (cruise MSM 16/2; 2010) [61].

**Barents Sea carbonate crust (BSCC).** The Barents Sea is a well-studied area of active CH<sub>4</sub> seepage. The seafloor geology is marked by pockmarks [62], gas hydrates, gas flares [63], and patches of carbonate crusts [64], indicating active CH<sub>4</sub> seepage [10,64]. We analysed eight depths (each 2–3 cm thick) from a push core (P120020 PR4) of a cold seep carbonate crust sampled at Loppa High/Polheim Sub-Platform area in the SW Barents Sea (72° 34'02.07"N, 20° 52'05.96"E). The core penetrated to a depth of 19 cm below sea floor (cmbsf). These samples were taken aboard the *RV Fugro Meridian* in September 2012 by Lundin Petroleum Norway.

**Amon Mud Volcano (AMV).** The Amon mud volcano (AMV) is located in the Nile deep-sea fan, in the Eastern Basin of the Mediterranean Sea. Hydrocarbons, muds, and fluids are transported to the surface via one main feeding channel from the deep subsurface, creating a stark thermal gradient in the sediment. Oxidation of CH<sub>4</sub> in the water column directly above AMV has been attributed to AMO using <sup>13</sup>C and <sup>2</sup>H isotopic values [65]. Sediments from AMV were collected aboard the *RV Meteor* (cruise M70/2, BIONIL; 2006) and the *RV Maria S Merian* (cruise MSM 13/3; 2009) [66].

**New Zealand Seeps (NZS).** The Hikurangi continental margin, east of New Zealand's North Island has been described as a biogeographically "new" cold seep province, characterised by endemic faunal communities [67]. Surface (0–2 cmbsf) and subsurface (10–12 cmbsf) sediment samples from three New Zealand seep (NZS) sites were collected aboard the *RV Sonne* (cruise So-191; 2007) [68]. These were dominated by distinct biota: *Frenulata* (Omakere Ridge), *Ampharetidae* (Wairarapa Takahae), and sulfur-oxidising bacteria (Wairarapa Takahae).

**Golfo Dulce (GD) surface sediments.** A sill at 60 m water depth physically cuts off Golfo Dulce from the Pacific Ocean, which promotes anoxic conditions within the basin. Recently, GD was shown to contain authigenic carbonate formations at shallow (ca. 10 m) water depth [69]. GD sediments were collected along a transect from 10 to 140 m water depth in March 2008, as described in [70].

**Gulf of Mexico (GoM) cold seeps.** The Gulf of Mexico (GoM) holds an abundance of thermogenic gas. The venting of this gas from deep subsurface forms gas hydrates, free CH<sub>4</sub>, which are the sources of energy for microbial chemosynthetic communities, and authigenic carbonates [71]. Pancost et al., [72] described the BHP composition of material from five carbonate rock and nodule sites in GoM. Two of the carbonates were shown to contain low quantities of aminopentol. Here, we reinvestigated the BHP signatures of these sediments.

**GoM sediments.** The GoM also houses the outflow of the Mississippi River Delta, transporting terrestrial material into the Gulf. Three GoM sites to the best of our knowledge not

influenced by CH<sub>4</sub> (27°30'N, 87°20'W; 28°20'N, 89°38'W; 26°50'N, 92°40'W; two sediment depths at each site) were investigated in this study [73].

**Peru Margin (PM).** An intense upwelling regimes fertilises surface water productivity on the Peru Margin (PM). This lends to oxygen utilisation in the water column, causing the Eastern South Pacific Oxygen Minimum Zone (ESP OMZ) [74]. Three PM sediments were analysed (10–15, 20–25 and 40–45 cmbsf) from a core taken within the ESP OMZ, at 100 m water depth [73,75].

## Lipid extraction

**Total lipid extraction.** All freeze-dried bacterial cells and marine sediments, except the Barents Sea samples extracted at GFZ Potsdam, were extracted using a modified Bligh-Dyer method [76,77]. Briefly, freeze-dried material was extracted in 19 mL of a 10:5:4 (v:v:v) mixture of MeOH:chloroform:H<sub>2</sub>O in a 50 mL Teflon tube. This mixture was sonicated for 15 min at 40°C, and centrifuged for 10 min. The supernatant was transferred to a second tube, and the residue re-extracted twice more. The chloroform in the supernatant was separated from the aqueous phase by adding water until the H<sub>2</sub>O:MeOH ratio was 1:1 (v:v), and collected. This procedure was repeated for the subsequent extractions. The collected chloroform total lipid extract (TLE) was dried by rotary evaporation in a round-bottom flask. The extraction protocol at GFZ Potsdam was similar but used a mixture of MeOH:DCM:ammonium acetate buffer [78].

**Solid Phase Extraction.** In-house comparisons have shown that amino-BHPs are better detected after solid phase extraction (SPE). An aliquot of the TLE was separated over a 1 mg NH<sub>2</sub> solid phase extraction cartridge, as described in [79]. Briefly, the aliquot was dissolved and loaded onto a hexane-rinsed cartridge using 200 µL chloroform. Six mL of a 98:2 (v:v) diethyl ether:acetic acid solution was eluted. Residual material was dissolved with 200 µL 2:1 (v:v) chloroform:MeOH and loaded onto the cartridge, followed by elution with 10 mL of MeOH. BHPs were isolated from the MeOH fraction.

## Lipid analyses

**BHP preparation and HPLC/APCI-MSn analyses.** A known amount (ca. 5–10 µg/g dry sediment) of internal standard (5 $\alpha$ -pregnane-3 $\beta$ ,20 $\beta$ -diol) was added to SPE extracts of the TLE for BHP analysis. Samples were acetylated in 0.5 mL of a 1:1 (v:v) mixture of acetic anhydride and pyridine at 50°C for 1 h, then left to stand overnight at room temperature [80]. Solvent was dried under a stream of N<sub>2</sub> on a 50°C heating block. BHP samples were dissolved in MeOH:propan-2-ol (3:2; v:v), and filtered on 0.2 µm PTFE filters.

BHPs were analysed by high performance liquid chromatography coupled to positive ion atmospheric pressure chemical ionization mass spectrometry (HPLC/APCI-MS), using a data-dependent scan mode (2 events) on an HPLC system equipped with an ion trap MS, as described in [46,81]. Further structural information for novel BHPs was obtained by way of MS<sup>3</sup> spectra. BHP concentrations were (semi) quantitatively estimated based on the response factor of authentic standards (M. Rohmer; Strasbourg, France and [46,77]) relative to the internal standard.


## Results

In this study we investigated the BHP distributions in species of three AMO marine genera, and of eight marine environments, six of which were CH<sub>4</sub>-influenced.



### Novel nitrogen-containing BHP components

In addition to the ‘regular amino-BHPs’ (e.g., I, II, and III; Figs 1 and 2), a suite of novel compounds were found in the methanotrophs and screened marine samples. Identification of these compounds is described in detail in the Supplementary Information (S1 File). Briefly, these components were related to the 35-amino-BHPs but differ in their terminal groups at C-35, which are tentatively proposed to comprise a methylcarbamate rather than a simple amine on the basis of interpretation of their APCI MS<sup>2</sup> and MS<sup>3</sup> spectra. In each case, the novel compounds (I<sup>MC</sup>, I<sup>MC'</sup>, II<sup>MC</sup>, III<sup>MC</sup>; Figs 1 and 2) elute after their ‘regular’ amino-BHP analogues (i.e., I, I', II, III; Fig 2). This indicates that the tentatively-assigned terminal group structures are less polar than the regular terminal amines (after acetylation). The novel structures include: 35-methylcarbamate-bacteriohopane-32,33,34-triol (MC-triol herein; III<sup>MC</sup>), 35-methylcarbamate-bacteriohopane-31,32,33,34-tetrol (MC-tetrol herein; II<sup>MC</sup>), 35-methylcarbamate-bacteriohopane-30,31,32,33,34-pentol (MC-pentol herein; I<sup>MC</sup>) and an isomer of I<sup>MC</sup> (I<sup>MC'</sup>) akin



**Fig 2. Distribution of nitrogen-containing BHPs in novel Type I methanotroph cultures.** Partial mass chromatograms (15–25 min) showing relative abundances (%) of BHPs (shaded peaks) in the acetylated total lipid extracts of (a) *Methylo Marinum vadi* IT-4, (b) *Methylo Marinum caldicuralii* IT-9, (c) *Methylo Marinum* sp. IN45, (d) *Methylo Microbium alcaliphilum*, and (e) *Methylo Microbium kenyense*.

doi:10.1371/journal.pone.0165635.g002

to the early-eluting aminopentol isomer (I'), which was found, based on mass spectra, in a culture of *Methylovulum*-like strain M200 [46].

## Methanotroph BHP signatures

Four previously untested methanotrophs isolated from marine or saline, alkaline lacustrine environments, belonging to the three genera *Methylomarinum*, *Methylomarinovum*, and *Methylomicrobium*, were analysed for their BHP composition. An additional species *Methylomicrobium alcaliphilum*, the partial BHP composition of which was recently reported in [53], is also shown here in full for comparison with *Methylomicrobium kenyense*. All of the methanotroph cultures investigated only synthesised BHPs with a nitrogen atom at C-35 position (nitrogen-containing BHPs herein). The relative abundances of BHPs are indicated as the percentage of total BHPs in acetylated extracts, and are presented in Fig 2. The low starting mass of some of the dry cell material led to uncertainty in the calculations of absolute concentrations, which are therefore not reported.

**BHP inventory of *Methylomarinum vadi* IT-4.** The most abundant BHPs in *Methylomarinum vadi* IT-4 were aminotriol (III), 22.3% and MC-triol (III<sup>MC</sup>), 22.7% (Fig 2A). Aminopentol (I), MC-pentol (I<sup>MC</sup>), aminotetrol (II), and MC-aminotetrol (II<sup>MC</sup>) made up 15.4%, 14.0%, 11.1%, and 11.8% of total BHPs, respectively, with lower levels of the aminopentol isomers (I' and I'<sup>MC</sup>; Fig 2A). No C-3 methylated or unsaturated equivalents of aminotriol, aminotetrol and aminopentol were found in *Methylomarinum vadi* IT-4.

**BHP inventory of *Methylomarinovum* spp.** The most abundant BHP in *Methylomarinovum caldicuralii* IT-9 was aminopentol (I), 37.4% (Fig 2B). However, *Methylomarinovum* sp. IN45 had only 5.1% aminopentol (Fig 2C). *M. caldicuralii* IT-9 had 30.8% MC-pentol, 14.4% aminotetrol, and 13.3% MC-tetrol, and lower abundances of aminopentol isomer, MC-pentol isomer, aminotriol, and MC-triol (Fig 2B). The most abundant BHP in *Methylomarinovum* sp. IN45 was MC-tetrol (II<sup>MC</sup>), making up 51.1%. *Methylomarinovum* sp. IN45 also contained 12.7% aminotetrol, 15.6% MC-pentol, 12.1% MC-triol and lower levels of aminotriol, and the aminopentol isomers (Fig 2C). No C-3 methylated nor unsaturated equivalents of aminotriol, aminotetrol and aminopentol were present in either strains.

**BHP inventory of *Methylomicrobium* spp.** The *Methylomicrobium alcaliphilum* and *Methylomicrobium kenyense* cultures did not contain aminopentol (I) above detection limit (Fig 2D and 2E) although *M. kenyense* was found to contain minor abundance of 3-Me-aminopentol (I<sup>3Me</sup>; 1.0%; Fig 2E). The most abundant BHP in both *Methylomicrobium* cultures was aminotriol (III), making up ca. 65% of all BHPs in both species. The second most abundant BHP was 3-Me-aminotriol (III<sup>3Me</sup>) at 31.5% in *M. alcaliphilum* and slightly less in *M. kenyense* (23.9%). Both species also contained lower levels of aminotetrol (II) and 3-Me-aminotetrol (II<sup>3Me</sup>). *M. kenyense* also contained unsaturated compounds ( $\Delta$ III and  $\Delta$ III<sup>3Me</sup>; Fig 2E). The only MC compound identified in either *Methylomicrobium* sp. was MC-triol and then only at low levels (<3%).

## Marine sediment and carbonate BHP signatures

Eight marine settings were studied for their BHP signatures (Table 2). Six of these were known to be influenced by CH<sub>4</sub> (i.e., HHMV, BSCC, AMV, NZS, GoM cold seeps, GD) and two were used as comparison background marine levels (i.e., GoM sediments, PM).

**Håkon Mosby Mud Volcano (HMMV).** None of the HMMV sediment samples contained aminopentol (Table 2). The most abundant amino- and MC-BHPs in HMMV samples was either aminotriol (III) or MC-triol (III<sup>CME</sup>), making up 22–52% and 32–62% of total nitrogen-containing BHPs, respectively. Unsaturated aminotriol, unsaturated MC-triol, aminotetrol,

**Table 2. Concentrations (µg/g sediment) of amino-BHPs in marine sediment samples.**

		Amino-BHPs													Total nitrogen-containing BHPs (µg/g sediment)
		I	ΔI	I <sup>MC</sup>	I <sup>3Me</sup>	II	II <sup>MC</sup>	II <sup>3Me</sup>	III	ΔIII	III <sup>MC</sup>	III <sup>3Me</sup>	ΔIII <sup>MC</sup>	ΔIII <sup>3Me</sup>	
	<b>Base Peak<sup>a</sup></b>	<b>830</b>	<b>828</b>	<b>846</b>	<b>844</b>	<b>772</b>	<b>788</b>	<b>786</b>	<b>714</b>	<b>712</b>	<b>730</b>	<b>728</b>	<b>728</b>	<b>726</b>	
<b>Håkon Mosby Mud Volcano (HMMV)</b>															
new mud flow 0–1 cm	PS74/2 169–1 PUC3 229	bdl	bdl	bdl	bdl	0.02	0.01	bdl	0.17	0.02	0.14	bdl	0.01	bdl	0.38
aged flow 0–1 cm	ARK XXIV-2 PS74 172–1 237	bdl	bdl	bdl	bdl	0.01	0.01	bdl	0.29	0.05	0.81	bdl	0.13	bdl	1.3
center 0–1 cm	MSM16/2 847–1 MUC 53	bdl	bdl	bdl	bdl	0.01	0.01	bdl	0.63	0.09	1.26	bdl	0.19	bdl	2.19
center 10–12 cm	MSM16/2 HMMV 63	bdl	bdl	bdl	bdl	0.01	0.02	bdl	0.17	0.02	0.17	bdl	0.02	bdl	0.41
new mud flow 0–1 cm	MSM16/2 838–1 MUC 33	bdl	bdl	bdl	bdl	0.02	0.01	bdl	1.07	0.09	0.82	bdl	0.06	bdl	2.07
newer mud flow 0–1 cm	MSM16/2 855–1 MUC 73	bdl	bdl	bdl	bdl	0.01	0.01	bdl	1.1	0.13	1.24	bdl	0.15	bdl	2.64
newer mud flow 10–12 cm	MSM16/2 HMMV 83	bdl	bdl	bdl	bdl	0.02	0.01	bdl	0.17	0.02	0.11	bdl	0.01	bdl	0.33
aged flow 0–1 cm	MSM16/2 823–1 MUC 19	bdl	bdl	bdl	bdl	0.02	0.02	bdl	0.56	0.05	0.88	bdl	0.07	bdl	1.61
aged flow 10–12 cm	MSM16/2 HMMV 28	bdl	bdl	bdl	bdl	0.01	0.01	bdl	0.15	0.01	0.11	bdl	0.01	bdl	0.29
<b>Barents Sea Carbonate Crusts (BSCC) depth profile</b>															
0–2 cmbsf		bdl	bdl	bdl	bdl	1.36	bdl	bdl	4.45	0.12	bdl	bdl	bdl	bdl	5.93
2–4 cmbsf		bdl	bdl	bdl	bdl	0.21	bdl	bdl	0.93	0.06	bdl	bdl	bdl	bdl	1.2
4–6 cmbsf		bdl	bdl	bdl	bdl	0.04	bdl	bdl	0.27	0.03	0.02	bdl	bdl	bdl	0.36
6–8 cmbsf		bdl	bdl	bdl	bdl	0.03	bdl	bdl	0.19	0.02	0.02	bdl	bdl	bdl	0.26
8–10 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.07	bdl	0.02	bdl	bdl	bdl	0.09
10–13 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.03	bdl	0.01	bdl	bdl	bdl	0.04
13–16 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.08	bdl	bdl	bdl	bdl	bdl	0.08
16–19 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.11	bdl	bdl	bdl	bdl	bdl	0.11
<b>Amon Mud Volcano (AMV)</b>															
central dome 0–1 cm	M70/2a 760 PUC33 24	bdl	bdl	bdl	bdl	0.02	0.07	bdl	0.42	0.13	2.67	bdl	0.96	bdl	4.27
sulfur band 0–1 cm	M70/2a 765 PUC49+50 71	0.18	bdl	0.12	bdl	0.38	0.27	bdl	1.24	0.81	1.33	bdl	1.02	bdl	5.35
sulfur band 10–12 cm	M70/2 81	0.36	0.04	0.11	bdl	0.3	0.11	bdl	1.07	1.09	0.48	bdl	0.5	bdl	4.06
sulfur band 0–1 cm	M70/2a 790 PUC68 172	0.36	bdl	0.11	bdl	1.49	0.49	bdl	3.82	1.29	1.49	bdl	0.61	bdl	9.67
sulfur band, white mat	MSM13/3 947–1 PUC31 73D	0.09	bdl	0.11	bdl	0.31	0.24	bdl	0.64	0.36	0.71	bdl	0.44	bdl	2.9
sulfur band 0–1 cm	MSM13/3 968–1 PUC15 122	0.08	bdl	0.06	bdl	0.75	0.49	bdl	0.73	0.46	0.77	bdl	0.28	bdl	3.62
bacterial mats 0–1 cm	MSM13/3 929–1 PUC22 45D	0.05	bdl	0.01	bdl	1.13	0.17	bdl	1.11	0.43	0.5	bdl	0.16	bdl	3.56

(Continued)

Table 2. (Continued)

		Amino-BHPs													Total nitrogen-containing BHPs (ug/g sediment)
		I	ΔI	I <sup>MC</sup>	I <sup>3Me</sup>	II	II <sup>MC</sup>	II <sup>3Me</sup>	III	ΔIII	III <sup>MC</sup>	III <sup>3Me</sup>	ΔIII <sup>MC</sup>	ΔIII <sup>3Me</sup>	
	<b>Base Peak<sup>a</sup></b>	<b>830</b>	<b>828</b>	<b>846</b>	<b>844</b>	<b>772</b>	<b>788</b>	<b>786</b>	<b>714</b>	<b>712</b>	<b>730</b>	<b>728</b>	<b>728</b>	<b>726</b>	
bacterial mats 10–12 cm	MSM13/3 50D	bdl	bdl	bdl	bdl	0.01	bdl	bdl	0.03	bdl	0.02	bdl	bdl	bdl	0.07
gassy centre, 10–12 cm	MSM13/3 61D	bdl	bdl	bdl	bdl	0.01	bdl	bdl	0.04	0.01	0.02	bdl	0.01	bdl	0.08
<b>New Zealand Seeps (NZS)</b>															
0–2 cmbsf	Frenulata 45	bdl	bdl	bdl	bdl	0.09	0.06	bdl	0.21	0.03	0.29	bdl	0.05	bdl	0.73
10–12 cmbsf	Frenulata 45	bdl	bdl	bdl	bdl	0.13	0.03	bdl	0.26	0.04	0.15	bdl	0.02	bdl	0.63
0–2 cmbsf	Ampharetidae 309	0.06	bdl	bdl	bdl	1.11	0.17	bdl	1.32	1.17	2.59	bdl	0.26	bdl	6.67
10–12 cmbsf	Ampharetidae 309	bdl	bdl	bdl	bdl	0.01	0.02	bdl	0.08	0.02	0.13	bdl	0.02	bdl	0.28
0–2 cmbsf	Sulfur-oxidising bacteria 315	0.02	bdl	bdl	bdl	0.66	0.32	bdl	2.24	0.23	2.13	bdl	0.21	bdl	5.8
10–12 cmbsf	Sulfur-oxidising bacteria 315	bdl	bdl	bdl	bdl	0.03	bdl	bdl	0.14	0.02	0.1	bdl	bdl	bdl	0.28
<b>Golfo Dulce (GD) Surface Sediment</b>															
10 m water depth	SG 1	bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.01	bdl	bdl	bdl	bdl	bdl	0.01
24 m water depth	SG 2	0.16	bdl	bdl	bdl	0.03	bdl	bdl	0.36	bdl	0.02	bdl	bdl	bdl	0.57
77 m water depth	SG 4	0.02	bdl	bdl	bdl	0.01	bdl	bdl	0.15	0.03	0.01	bdl	bdl	bdl	0.23
90 m water depth	SG 5	0.04	bdl	bdl	bdl	0.02	bdl	bdl	0.17	0.05	0.01	bdl	bdl	bdl	0.29
110 m water depth	SG 6	bdl	bdl	bdl	bdl	0.01	bdl	bdl	0.08	0.07	0.02	bdl	bdl	bdl	0.18
120 m water depth	SG 7	0.04	bdl	bdl	bdl	0.02	bdl	bdl	0.36	0.2	0.02	bdl	bdl	bdl	0.63
140 m water depth	SG 8	0.01	bdl	bdl	bdl	<0.01	bdl	bdl	0.11	0.08	0.01	bdl	bdl	bdl	0.21
<b>Gulf of Mexico (GoM) Cold Seeps</b>															
GC 234 CNSTS 4434		bdl	bdl	bdl	bdl	0.01	bdl	bdl	0.09	bdl	bdl	bdl	bdl	bdl	0.1
234 4436		bdl	bdl	bdl	bdl	<0.01	bdl	bdl	0.04	bdl	bdl	bdl	bdl	bdl	0.04
GC 234 tube worm roots 4435		bdl	bdl	bdl	bdl	<0.01	bdl	bdl	0.02	bdl	bdl	bdl	bdl	bdl	0.02
GC 232		bdl	bdl	bdl	bdl	0.01	bdl	bdl	0.01	bdl	bdl	bdl	bdl	bdl	0.02
GC 185		bdl	bdl	bdl	bdl	0.76	bdl	bdl	0.2	bdl	bdl	bdl	bdl	bdl	0.95
<b>GoM Sediments</b>															
West Gulf	WG2-099/6	bdl	bdl	bdl	bdl	<0.01	bdl	bdl	0.01	bdl	bdl	bdl	bdl	bdl	0.01
26°50'N, 92°40'W	WG2-099/9	bdl	bdl	bdl	bdl	bdl	bdl	bdl	<0.01	bdl	<0.01	bdl	bdl	bdl	0.01
Eastern Gulf	88-C-1 DCS 128/17	bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.01	bdl	0.01	bdl	bdl	bdl	0.02
27°30'N, 87°20'W	DCS 128/13	bdl	bdl	bdl	bdl	bdl	bdl	bdl	<0.01	bdl	<0.01	bdl	bdl	bdl	0.01
	DCS 128/21	bdl	bdl	bdl	bdl	bdl	bdl	bdl	<0.01	bdl	0.01	bdl	bdl	bdl	0.01
Central Gulf	CGD-136/17	bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.01	bdl	0.01	bdl	bdl	bdl	0.01
28°20'N, 89°38'W	CGD-136/13	bdl	bdl	bdl	bdl	bdl	bdl	bdl	<0.01	bdl	0.01	bdl	bdl	bdl	0.01
<b>Peru Sediment</b>															
10–15 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.09	bdl	bdl	bdl	bdl	bdl	0.09
20–25 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	0.02	bdl	bdl	bdl	bdl	bdl	0.02

(Continued)

Table 2. (Continued)

	Amino-BHPs														Total nitrogen-containing BHPs (ug/g sediment)
	I	ΔI	I <sup>MC</sup>	I <sup>3Me</sup>	II	II <sup>MC</sup>	II <sup>3Me</sup>	III	ΔIII	III <sup>MC</sup>	III <sup>3Me</sup>	ΔIII <sup>MC</sup>	ΔIII <sup>3Me</sup>		
	<b>Base Peak<sup>a</sup></b>	<b>830</b>	<b>828</b>	<b>846</b>	<b>844</b>	<b>772</b>	<b>788</b>	<b>786</b>	<b>714</b>	<b>712</b>	<b>730</b>	<b>728</b>	<b>728</b>	<b>726</b>	
40–45 cmbsf		bdl	bdl	bdl	bdl	bdl	bdl	bdl	<b>0.09</b>	bdl	bdl	bdl	bdl	bdl	<b>0.09</b>

<sup>a</sup>base peak = [M + H-CH<sub>3</sub>COOH]<sup>+</sup>  
bdl—below detection limit

doi:10.1371/journal.pone.0165635.t002

and MC-tetrol all made up <10% of total nitrogen-containing BHPs. Some HMMV samples contained relatively high concentration of anhydrobacteriohopanetetrol (anhydro-BHT). Minor abundances of BHT and BHT isomer, BHT-cyclitol ether (BHT-CE), and BHT-glucosamine (BHT-G) were detected in some samples (S1 Table).

**Barents Sea carbonate crust (BSCC).** The 19 cm BSCC core contained a majority of aminotriol (III; 72–100% of nitrogen-containing BHPs; Table 2). MC-triol (III<sup>MC</sup>; 0–23%) peaked between 4 and 13 cm. Aminotetrol (II; 0–23%) was detected in the upper 8 cm. Minor contribution of unsaturated aminotriol (ΔIII; 0–7%) was detected in the upper sediments. Other BHPs detected included BHT, and low concentrations of anhydroBHT, BHT isomer, 2-methyl-BHT (2-Me-BHT), adenosylhopane, BHT-CE, and BHT-G (S1 Table).

**Amon Mud Volcano (AMV).** The most abundant nitrogen-containing BHPs in sediments from the Amon mud volcano were aminotriol (III; 9.8–48.9%) and MC-triol (III<sup>MC</sup>; 11.8–62.5%) (Table 2). Other nitrogen-containing BHPs in these sediments were aminotetrol (II; 0.5–31.8%), unsaturated aminotriol (ΔIII; 3.0–27.0%) and unsaturated MC-triol (ΔIII<sup>MC</sup>; 0–22.5%). Minor abundances of aminopentol (I; 0–8.9%), and MC-pentol (I<sup>MC</sup>; 0–3.8%) were found in some AMV sediments. Other BHPs in the AMV sediments were BHT, anhydroBHT, 2-Me-BHT, 3-Me-BHT, BHT-CE, BHT-G, BHT isomer, and adenosylhopane (S1 Table).

**New Zealand Seeps (NZS).** The most abundant nitrogen-containing BHP in the sediments from NZS were aminotriol (III; 28.5–72.7%), MC-triol (III<sup>MC</sup>; 13.2–46.4%) and aminotetrol (II; 5.1–20.6%) (Table 2). Unsaturated aminotriol (ΔIII), unsaturated MC-triol (ΔIII<sup>MC</sup>), and MC-tetrol (II<sup>MC</sup>) all made up <10%. Two sediment samples contained aminopentol, albeit in low abundance (<1% of nitrogen-containing BHPs). Both of these aminopentol positive samples also contained soil marker BHPs [77,82]. Other BHPs found in NZS samples were BHT, anhydroBHT, 2-methyl-BHT, 3-methyl-BHT, BHT isomer, adenosylhopane, and BHT-CE (S1 Table).

**Gulfo Dulce (GD).** The most abundant nitrogen-containing BHP in GD surface sediments was aminotriol (III; 44.3–100%; Table 2). Unsaturated aminotriol (ΔIII) made up 0–41.3%. MC-triol (III<sup>MC</sup>; 0–11.4%) and aminotetrol (II; 0–6.0%) were found in most samples. Aminopentol was found in all but two surface samples, one of which was the shallowest site. Aminopentol abundance decreased with increasing water depth with the highest abundance at 24 m water depth (27.6% of nitrogen-containing BHPs). BHT, anhydroBHT, BHT isomer, and 2-Me-BHT were also found in GD (S1 Table).

**Gulf of Mexico (GoM) cold seeps.** GoM cold seeps only contained aminotriol (III; 20.5–100%) and aminotetrol (II; 0–79.5%) nitrogen-containing BHPs (Table 2). Other BHPs included BHT, soil marker BHPs, 2-Me-BHT, BHT isomer, and BHT-CE (S1 Table).


**GoM sediments.** GoM sediments from near the outflow of Mississippi River Delta showed an abundance of aminotriol (**III**; 26.6–80.7% of total nitrogen-containing BHPs), MC-triol (**III<sup>MC</sup>**; 0–73.4%) (Table 2). One sample from the Western Basin had 19.3% aminotetrol (**II**) relative to total BHPs; however, the concentration of aminotetrol was low (<0.01 μg/g sediment). Other BHPs included BHT, BHT isomer, anhydroBHT, 2-methyl-BHT, BHT-CE, and soil marker BHPs (S1 Table).

**Peru Margin (PM).** The only nitrogen-containing BHP detected in sediments from the PM was aminotriol (**III**). Other BHPs found at this site included BHT, BHT isomer, anhydroBHT, 2-Me-BHT, adenosylhopane, and BHT-CE (S1 Table).

## Discussion

### BHP distributions in aerobic methanotrophs

**Previously reported BHP distributions in AMO bacteria.** Traditionally, Type I and Type II AMO bacteria had been distinguished by their different BHP signatures (e.g. [52]; see also review in [14]). Prior to the investigation of BHPs in the *Methylovulum*-like strain M200 [46], most screened Type I methanotrophs synthesised a high percentage of aminopentol (**I**) and lower contributions from aminotetrol (**II**) and in some cases aminotriol (**III**), and clustered in the left-hand corner of the amino-BHP ternary plot (Fig 3A). In contrast, Type II methanotrophs did not contain aminopentol, had varying contributions from **II** and **III**, and clustered along the right-hand axis of Fig 3A. The high relative abundance of **III** observed in *Methylovulum*-like strain M200 was, therefore, originally seen as an outlier [46]. Similarly, [52] showed that a culture of *Methylomicrobium album* did not contain aminopentol. At the time this was



**Fig 3. Relative composition (%) of 'regular' amino-BHPs.** Distributions of aminotriol, aminotetrol and aminopentol, including their C-3 methylated homologues, where present, are shown for (a) methanotroph cultures including literature data (circles) and data from new cultures of *Methylomicrobium*, *Methylomarinum*, and *Methylomarinovum* (diamonds; this study) and (b) sediments and microbial mats from methane-rich marine settings (HMMV = Håkon Mosby mud volcano; BSCC = Barents Sea carbonate crust; AMV = Amon mud volcano; NZ = New Zealand; GD Golfo Dulce; GoM = Gulf of Mexico; PM = Peru Margin). Methanotroph literature data from [36,42,46,52,83]. <sup>a</sup>Literature data from [36,42,83] was based on GC-MS hopanol quantifications after periodic acid treatment. Therefore, it is not possible to distinguish between amino- and non-amino-BHPs with the same number of functional groups. This is especially significant for the Type II aerobic methanotrophic bacteria that are known to synthesise bacteriohopanetetrol. For this reason, the axes of plot (a) are shown as functionality of the BHP-side chain.

doi:10.1371/journal.pone.0165635.g003

presumed to be a contaminated culture, however, [53] also did not report I synthesis in cultures of *Methylomicrobium alcaliphilum*. All of the recently analysed *Methylobacter* spp. [28] join the more typical Type I methanotrophs in the left-hand corner of the plot, however, *Methylobacter* sp. BB5.1 increased the spread of the cluster with almost 40% III content.

**Amino-BHP distributions in previously untested Type I AMO bacterial cultures.** It was assumed that the screened species of AMO Type I bacteria investigated in this study would display similar BHP distributions as those of previously reported Type I bacteria. All three bacterial genera screened do indeed only contain amino-BHPs (Fig 2). However, the relative distribution of specific nitrogen-containing BHPs varies between genera, as well as between species belonging to the same genus. To allow for a more accurate comparison with data from the literature (Fig 3A, circles), only aminopentol, aminotetrol, aminotriol, and their methylated equivalents were considered when producing the ternary plot of 'regular' nitrogen-containing BHPs of the novel Type I cultures (Fig 3A, diamonds).

Aminopentol is the most abundant BHP in the novel species *Methylomarinovum cadicuralii* IT-9, which is in agreement with literature BHP compositions of most other Type I methanotrophs (Fig 3A, circles), e.g., *Methylococcus capsulatus* [44,52], *Methylomonas* sp. [42,46,52], psychrotolerant isolate CEL 1923 [42], and *Methylocaldum tepidum* [22]. However, a species in the same genus (*Methylomarinovum* sp. IN45) has a much lower abundance of aminopentol (5.0% of total amino-BHPs; Fig 2C). *Methylomarinum vadi* IT-4 shows relatively high proportions of aminopentol, but it is not the most abundant BHP (Fig 2A). Moreover, in our screening of two species of *Methylomicrobium* spp., aminopentol was not detected, similar to reported cultures of *Methylomicrobium album* and *Methylomicrobium alcaliphilum* [52,53]. Our results seem to confirm the near-absence of aminopentol in all screened *Methylomicrobium* spp., which are the first Type I methanotrophs apparently unable to synthesise aminopentol. However, changes in BHP composition can occur at different growth stages and under different conditions (e.g. [27,83,84]), so further studies would be required to fully confirm this. It appears that the BHP distributions of *Methylomicrobium* and *Methylovulum*, which do not synthesise high amounts of aminopentol, should also no longer be considered outliers given the low levels of aminopentol in *M. vadi* IT-4 and *Methylomarinovum* spp. (Fig 2). This suggests a greater variance in the BHPs of Type I methanotrophs than previously thought. Furthermore, as *Methylomicrobium* has been isolated from a diverse range of marine environments [85–87], the absence of aminopentol in this genus might greatly affect its application as a marine aerobic methanotrophy biomarker. This, however, does not invalidate the use of aminopentol as a biomarker for methanotrophy.

There is also significant variation in the relative abundances of the other nitrogen-containing BHPs in Type I methanotrophs. A suite of novel BHPs identified as methylcarbamate (MC) BHPs are detected in all three genera screened in this study (Fig 2). These have not been reported in previous studies. Therefore, data available at Newcastle University from the analyses of *Methylococcus capsulatus* (Talbot et al., unpublished), *Methylovulum*-like strain M200, *Methylomonas methanica*, *Methylomonas*-like strain M5 [46], and *Methylobacter* spp. [28] were re-examined. III<sup>MC</sup> was identified retrospectively in *Methylomonas methanica*, and all species of *Methylobacter*. III<sup>MC</sup> constituted up 9.8% of total amino-BHPs in *Methylobacter* sp. BB 5.1. II<sup>MC</sup> was present in low abundance (<0.2%) in two of the three species of *Methylobacter* (*Methylobacter* sp. BBA 5.1 and *Methylobacter* sp. BB 5.1), but absent in *Methylobacter marinus* A 45. I<sup>MC</sup> was absent in *Methylococcus capsulatus* and *Methylovulum*-like strain M200, but was present in all species of *Methylobacter*. I<sup>MC</sup> was also present in *Methylomonas methanica* but absent in a different species of this genus (i.e., *Methylomonas*-like strain M5; [46]). These results indicate that MC-BHPs are not universally present when their regular

homologues are detected, and may be species specific and/or dependent on variations in growth conditions such as pH.

The two species of *Methylomarinovum* display significant variations in their BHP compositions (Fig 2B and 2C). *Methylomarinovum* sp. IN45 had a relatively low level of aminopentol. The most abundant BHP in *Methylomarinovum* sp. IN45 is  $\text{II}^{\text{MC}}$ . Although in lower abundances,  $\text{III}^{\text{MC}}$  and  $\text{I}^{\text{MC}}$  are also higher in comparison to their 'regular' homologues in this species compared to *M. caldicuralii* IT-9 (Fig 2B and 2C). In contrast, the most abundant BHP in *Methylomarinovum caldicuralii* IT-9 is aminopentol, followed by almost equal amounts of  $\text{I}^{\text{MC}}$ . The different relative BHP distributions between the *Methylomarinovum* spp. highlight that there can be significant variations within a genus. *Methylomarinovum* sp. IN45 was isolated from a deep-sea hydrothermal field and perhaps the high levels of methylcarbamate components observed are the result of a physiological adaptation to higher pressure in this environment. This may explain why the relative abundances of components in *Methylomarinovum caldicuralii* IT-9, the same genus but isolated from a shallow submarine hydrothermal environment, are quite different. Perhaps the complex functionality of the terminal group of the methylcarbamate components is more effective at stabilising the cell membrane and decreasing fluidity under these conditions.


3-methylaminotriol ( $\text{III}^{3\text{Me}}$ ) was observed in both *Methylomicrobium* spp. (23.9–31.5% of total BHPs; Fig 2D and 2E) in agreement with a recent report in [53]. This compound was accompanied by low levels of 3-methylaminotetrol ( $\text{II}^{3\text{Me}}$ ) in both species and trace amounts of 3-methylaminopentol in *M. kenyense* ( $\text{I}^{3\text{Me}}$ ). The absence of C-3 methylated structures in the previously investigated *Methylomicrobium album* strain BG8 [52] may appear inconsistent with the organisms investigated here; however, genomic investigations have revealed that *M. album* is separated from halo(alkali)philic representatives of the *Methylomicrobium* genus such as *M. alcaliphium* and *M. kenyense* [58], and perhaps specific environmental conditions influence the BHP composition of *Methylomicrobium* spp. as they seemingly do within the *Methylomarinovum* genus.

No C-3 methylated equivalents of aminotriol ( $\text{III}^{3\text{Me}}$ ), aminotetrol ( $\text{II}^{3\text{Me}}$ ), nor aminopentol ( $\text{I}^{3\text{Me}}$ ) were present in *Methylomarinum vadi* IT-4 or the *Methylomarinovum* spp., adding to examples of Type I methanotroph species that contain amino-BHPs, but not their C-3 methylated equivalents (e.g., see review in [14]). The most abundant BHPs in *Methylomarinum vadi* IT-4 were aminotriol ( $\text{III}$ ) and MC-triol ( $\text{III}^{\text{MC}}$ ), which were present in equal amounts (Fig 2A). Similar amounts of  $\text{II}$  and  $\text{II}^{\text{MC}}$ , and  $\text{I}$  and  $\text{I}^{\text{MC}}$  were also observed in this culture. A high proportion of  $\text{III}$  is unusual for a Type I methanotroph, but has been observed before in the *Methylovulum*-like strain M200. (Fig 4A; [46]). The new data reiterate that aminopentol is not always the most abundant BHP in Type I methanotrophs, nor necessarily the most appropriate biomarker for AMO.

## BHPs in marine environments

**Lack of BHP diversity in marine environments.** The screened marine sediments and authigenic carbonates do not show large diversity in their BHP signatures (S1 Table). The limited BHP distributions are also comparable to other reported marine sediment BHP signatures, all dominated by BHT and BHT isomer, from a number of locations including the Black Sea [88], the Benguela upwelling system [89], and the Arabian Sea [90]. More recently a similar pattern was also seen in water column samples from the California Current system, where the wide diversity observed in the gene responsible for hopane cyclisation (squalene-hopene cyclase) was not reflected by distinct BHP fingerprints related to this potential range of source-organisms [91]. However, genetic information is quickly lost, and we must strive to find lipid biomarkers to trace particular metabolisms in the geological record.





**Fig 4. Relative composition (%) of methylcarbamate(MC)- BHPs.** Distributions of MC-triol, MC-tetrol and MC-pentol are shown for (a) methanotroph Type I cultures, *Methylobacter*, *Methylomicrobium*, *Methylomarinum*, and *Methylomarinovum*, and (b) marine sediments (HMMV = Håkon Mosby mud volcano; BSCC = Barents Sea carbonate crust; AMV = Amon mud volcano; NZS = New Zealand; GD Golfo Dulce; GoM = Gulf of Mexico). *Methylobacter* data from [28].

doi:10.1371/journal.pone.0165635.g004

Non-nitrogen-containing BHP concentrations in the screened sediments do not show remarkable signatures (S1 Table). BHT and anhydro-BHT, thought to be a degradation product of BHT and other composite BHPs such as BHT cyclitol ether [92], were found at all sites. The presence of soil marker BHPs at some sites, particularly NZS, indicates that these sediments could be influenced by terrestrial input of organic matter (e.g., [77,82,93]). However, as adenosylhopane is an intermediate in the biosynthesis of all other side-chain extended BHPs [94], other sources cannot be entirely excluded. BHT isomer, a biomarker for anaerobic ammonium oxidation [70], was found in high concentrations in GD (previously reported in [70]), as well as in the PM sediments, which underlie the Peruvian OMZ, where anammox is known to be an important process [95], and where BHT isomer has previously been reported from the water column within the OMZ [90]. The most abundant of the three regular amino-BHPs in the CH<sub>4</sub>-influenced marine sediments was aminotriol, which is not source-specific (e.g., [48]).

**Aminopentol in marine sediments.** Although aminopentol was found in significant abundance in some of the reported and screened Type I methanotroph cultures (Fig 2), it was not found to be abundant in most of the CH<sub>4</sub>-influenced marine sites in this study (Table 2). In fact, it was only detected in AMV, GD surface sediments, and two NZS samples (Fig 3B; Table 2). The discrepancy between the distinct amino-BHP signatures of isolated Type I AMO bacteria and signatures of CH<sub>4</sub>-influenced marine sites is highlighted in the ternary plots of the relative composition of aminopentol, aminotetrol, and aminotriol (Fig 3A cf. Fig 3B). These differences could be due to the particular methanotrophic bacterial community responsible for methanotrophy in the CH<sub>4</sub>-influenced marine sediments. Ruff et al. [96] found that diversity in the global CH<sub>4</sub> seep microbiome was controlled by environmental factors such as

temperature and electron acceptor availability. Considering their findings, it is possible that the environmental conditions in most marine CH<sub>4</sub>-influenced sediments favour specific methanotroph communities. For example *Methylomicrobium* spp., found in saline environments [85–87] and saline, highly alkaline environments [58,97], and which do not produce aminopentol in significant amounts (Fig 2), could be present. However, the absence of C-3 methylated compounds is confounding for a *Methylomicrobium* source, pointing towards other methanotrophs that do not synthesise aminopentol. Yan et al. [98] found that 85% of the operational taxonomic units (OTUs) from the same sites as our GoM cold seeps did not group with known sequences of a subunit of particulate methane monooxygenase (*pmoA*). This would suggest the presence of novel methanotrophic species in GoM. In the same way, significant *pmoA* diversity has been observed in sediments from the North American margin [99], a shallow CH<sub>4</sub> seep [100], a marine estuary [101], and hydrocarbon seeps [102]. *pmoA* OTUs from the NZS sediments grouped with methanotrophic endosymbionts [68], including *Bathymodiolus* spp., which have been shown to contain neither aminopentol nor methylated BHPs [103]. Nevertheless, the absence of methylated amino-BHPs in the screened marine sediments (Table 2) may suggest *Methylomicrobium album*, or a related species that also does not synthesise methylated amino-BHPs, being the dominant methanotroph in CH<sub>4</sub>-influenced marine environments.

These are not the first reports of marine CH<sub>4</sub>-influenced environments not containing aminopentol (Table 1). For example, using methods targeting the functional gene *pmoA*, which is produced by most methanotrophs, Type I methanotrophs were detected in all three units of Ace Lake sediments. However, aminopentol was only detected in sediments deposited under freshwater conditions (unit III) despite the fact that the modern meromictic water column, containing relict seawater left behind after the sea level fell around 9000 years ago, hosts the Type I methanotroph *Methylosphaera hansonii* [104]. No aminopentol was detected in the methanotrophic symbionts in the gill tissue of a cold-seep mussel, despite other lipid-based evidence suggesting the presence of a Type I methanotroph [103,105]. Similarly, CH<sub>4</sub> seep carbonates from Alaminos Canyon, northern Gulf of Mexico [41] and the Northern Arabian Sea [106] were found to lack aminopentol. Conversely, aminopentol was detected in the water column of the Baltic Sea with supporting evidence for the presence of Type I methanotrophs from <sup>13</sup>C-depleted PLFAs [107]. Aminopentol was also detected in the water column of the Black Sea in the oxic-anoxic water transition, but not in the underlying sediment [12,88,108].

The presence of aminopentol in sediments from the AMV, located on the Nile deep-sea fan, in the Eastern Basin of the Mediterranean Sea (Table 2) may be explained by Nile River outflow carrying terrestrial wetland methanotrophy signatures into the Mediterranean, as seen in the Amazon and Congo River fans [14,109]. This would appear to indicate that aminopentol is still an excellent biomarker for terrestrial AMO. The near-absence of soil-marker BHPs in AMV (Table 2) may still point towards in-situ marine production of aminopentol. However, the relative abundance of soil-markers in terrestrial settings has recently been found to be strongly influenced by environmental factors; higher temperatures and low pH (in peatlands) can both strongly reduce the relative proportion of soil marker BHPs as a proportion of total BHPs [40,110,111]. Aminopentol was found in NZS sediments that also contained soil marker BHPs (Table 2). Therefore, aminopentol in sediments from NZS may have originated from terrestrial sources. Aminopentol in GD surface sediments may be the result of a distinct AMO community living in the specific environment prone to carbonate formation in GD. Unfortunately, samples were not properly preserved to be able to determine AMO diversity using genetic-based analyses of the *pmoA* gene in these sediments. The cumulative results of the studied marine sites do, however, indicate that an absence of aminopentol is not necessarily evidence for the absence of methanotrophs or aerobic methane oxidation.

## Alternative BHP biomarkers for AMO and implications for the marine sedimentary record of methanotrophy

**Regular amino-BHPs.** Screened Type I methanotrophs also produced varying amounts, depending on the genera, of aminotetrol (**II**) and aminotriol (**III**) (Fig 3A), both of which were found in CH<sub>4</sub>-influenced marine sediments (Fig 3B; Table 2). However, these two amino-BHPs are less source-specific to methanotrophic bacteria than aminopentol, and do not make ideal biomarker lipids for methanotrophy. Given that 3-Me-aminotriol (**III**<sup>3Me</sup>) made a significant contribution to the amino-BHP abundance in screened cultures of *Methylobacterium* spp. (23.9 and 31.5% of total amino-BHPs; Fig 2D and 2E) and 9.8% in *Methylobacter* sp. BB5.1 [28], it was expected that **III**<sup>3Me</sup> would be an important amino-BHP in CH<sub>4</sub>-influenced marine sediments. However, **III**<sup>3Me</sup> was not found in any of the screened sediments (Table 2). *Methylobacterium alcaliphilum* and *Methylobacterium kenyense* are adapted to high alkalinity, but not necessarily to high salinity [58]. This distinct lack of **III**<sup>3Me</sup> in marine sediment samples would seem to indicate that the *Methylobacterium* species we investigated are not the primary source of amino-BHPs in CH<sub>4</sub>-influenced marine environments.

**III**<sup>3Me</sup> has only occasionally been reported from environmental samples including some soils [82,112] and most recently in a peat core from Germany [111], but only at very low levels (Table 1). Other C-3 methylated amino-BHPs are even less common (Table 1). **I**<sup>3Me</sup> was first reported from a neo-volcanic, eutrophic and saline lake sediment (La Piscina de Yuriria, Mexico; [45]), and subsequently from a geothermal silica sinter (Opaheke Pool hot spring, New Zealand; [113]). The pentafunctionalised **II**<sup>3Me</sup> was also present in the Mexican lake sediment. Both of these compounds were reported in one study on the Black Sea water column [12], but were absent at another site [108]. The apparent discrepancy between the very limited occurrence of C-3 methylated BHPs (as measured using the periodic acid cleavage technique which converts polyfunctionalised BHPs into GC-amenable primary alcohols; e.g., [36,75]) and their wider occurrence in the form of 3-Me hopanes in ancient rocks and oils was first identified in [114]. These authors found 3-Me-BHPs to be abundant only in a very limited number of settings, under quite specific conditions (i.e., some alkaline lakes). The occurrence of 3-Me-hopanes in marine authigenic carbonates [31,115], which form under highly alkaline conditions are also consistent with a *Methylobacterium* source ([58], and references therein). It was further suggested that 3-Me-BHPs and hexafunctionalized BHPs appear to have different sources (possibly, but not necessarily restricted to, only Type I methanotrophs; [114]). Culture studies (on the moderately thermophilic Type I methanotroph *Methylococcus capsulatus*) have shown that production of C-3 methylated compounds may be related to growth stage. Higher relative proportions of methylated BHPs replaced the non-methylated equivalents during stationary phase growth [83], and appear to be necessary for maintaining intra-cellular membrane structures [27]. These important physiological roles for methylated BHPs are at odds with the very sparse occurrence of these compounds in modern settings (Table 1), and clearly our understanding of the factors controlling their biosynthesis and subsequent preservation in sediments is still limited, hampering interpretation of certain BHP signatures.

**Methylcarbamate-BHPs.** Most of the marine sediments influenced by CH<sub>4</sub> contained at least MC-triol, albeit at relatively low abundances (Fig 4B; Table 2). The fact that the MC-BHPs were found in all strains of methanotrophs analysed, though not all components in the suite were present in every strain, shows the biomarker potential of these BHPs for AMO (Fig 3A). MC-tetrol (**II**<sup>MC</sup>) was the most abundant component in *Methylomarinovum* sp. IN45, and MC-BHPs were found in higher abundance than the 'regular' 35-amino-BHP homologues, which may allow this particular hydrothermal vent species to be identified in environmental settings. Unsaturated MC-triol (**ΔIII**<sup>CME</sup>) was found in high abundance in AMV, HMMV, and

NZS, but was not found in any of the methanotroph cultures. This is possibly because the BHP signatures in most CH<sub>4</sub>-influenced marine sediments are sourced from AMO bacteria that have no cultured relatives or at least none which have been tested for BHP production.

Given the small diversity in BHPs found in marine sediments and the need for an AMO biomarker, there appear to be few BHPs that meet the criteria of being source-specific *and* abundant. This has significant implications for the development of a proxy using aminopentol to trace AMO in marine settings. Applying MC-BHPs combined with the traditional suite of amino-BHPs (e.g. aminopentol, aminotetrol, and aminotriol) seems to be the most appropriate biomarker course for AMO.

## Conclusions

Isolated methanotrophs from previously unexamined genera and species displayed marked differences in their relative abundances of amino-bacteriohopanepolyols (BHPs). Aminopentol (I) was the most abundant BHP in *Methylomarinovum caldicuralii* IT-9, which fits with the typical BHP signature of known Type I methanotrophs. However, the BHP signatures of *Methylomarinovum* sp. IN45 and *Methylomarinum vadi* IT-4 both did not show aminopentol as the most abundant BHP. Moreover, neither of the *Methylomicrobium* spp. contained aminopentol and only one contained a low level of 3-methyl-aminopentol showing that not all Type I methanotrophs synthesise aminopentol, agreeing with previous environmental studies. Considering *Methylomicrobium* can be prevalent in marine environments, this has implications for the use of aminopentol as a biomarker for marine methanotrophy. A suite of components related to amino-BHPs, but with methylcarbamate (MC) terminal groups, were detected for the first time, and were present in all Type I methanotroph strains tested. Marine sediments influenced by CH<sub>4</sub> did not contain significant amount of aminopentol, but did contain MC-BHPs. This study highlights the relatively low BHP diversity within marine sediments, and indicates that the combined use of MC-BHPs and amino-BHPs might be preferential to trace aerobic methane oxidation (AMO) in marine settings.

## Supporting Information

**S1 File. Identification of novel BHP compounds in methanotroph cultures.** (DOCX)

**S1 Table. Concentrations of other BHPs in marine sediment samples.** Concentration (μg/g sediment) of other BHPs in marine sediments samples presented in this study. (XLSX)

## Acknowledgments

We are deeply thankful to A. Boetius for supplying the sediments from HMMV, AMV, and NZS. Additional information about these sites can be found on the PANGAEA database (<http://www.pangaea.de/>). The PM sediments analysed in this study were originally provided by T. Eglinton and J. Repeta. R.D. Pancost is thanked for use of GoM cold seep material. The GoM sediments used in this study were provided by R. Sassen and S. Sweet of the Geochemical & Environmental Research Group (GERG) at Texas A&M University, U.S.A. We thank the captain and the crew of the *RV Kaiyo*, the operation team of the ROV Hyper-Dolphin, and the shipboard scientific party for the invaluable help in collecting deep-sea samples. We also thank M. Abe and K. Tanaka for their assistance in cultivation.

## Author Contributions

**Conceptualization:** KAO DR HMT.

**Formal analysis:** DR KAO HMT.

**Investigation:** KAO DR FRS.

**Methodology:** KAO DR HMT.

**Resources:** DB AK HH JP SWP JCN KM MK.

**Writing – original draft:** DR KO HMT.

**Writing – review & editing:** DB AK HH JP SWP JCN KM MK.

## References

1. Lelieveld J, Crutzen PJ, Dentener FJ (1998) Changing concentration, lifetime and climate forcing of atmospheric methane. *Tellus B* 50: 128–150.
2. IPCC (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (Eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
3. Dickens GR, O'Neil JR, Rea DK, Owen RM (1995) Dissociation of oceanic methane hydrate as a cause of the carbon isotope excursion at the end of the Paleocene. *Paleoceanography* 10: 965–971.
4. Dickens GR, Castillo MM, Walker JCG (1997) A blast of gas in the latest Paleocene: Simulating first-order effects of massive dissociation of oceanic methane hydrate. *Geology* 25: 259–262. PMID: [11541226](#)
5. Zachos JC, Rohl U, Schellenberg SA, Sluijs A, Hodell DA, Kelly DC, et al. (2005) Rapid acidification of the ocean during the Paleocene-Eocene thermal maximum. *Science* 308: 1611–1615. doi: [10.1126/science.1109004](#) PMID: [15947184](#)
6. Handley L, Pearson PN, McMillan IK, Pancost RD (2008) Large terrestrial and marine carbon and hydrogen isotope excursions in a new Paleocene/Eocene boundary section from Tanzania. *Earth Planet Sci Lett* 275: 17–25.
7. McInerney FA, Wing SL (2011) The Paleocene-Eocene thermal maximum: a perturbation of carbon cycle, climate, and biosphere with implications for the future. *Annu Rev Earth Planetary Sci* 39: 489–516.
8. Hanson RS, Hanson TE (1996) Methanotrophic bacteria. *Microbiol Rev* 60: 439–471. PMID: [8801441](#)
9. Sieburth JM, Johnson PW, Eberhardt MA, Sieracki ME, Lidstrom M, Laux D (1987) The first methane-oxidizing bacterium from the upper mixing layer of the deep ocean—*Methylomonas pelagica* sp. nov. *Curr Microbiol* 14: 285–293.
10. Hovland M, Svensen H, Forsberg CF, Johansen H, Fichler C, Fossa JH, et al. (2005) Complex pockmarks with carbonate-ridges off mid-Norway: Products of sediment degassing. *Mar Geol* 218: 191–206.
11. McDonald IR, Smith K, Lidstrom ME (2005) Methanotrophic populations in estuarine sediment from Newport Bay, California. *FEMS Microbiol Lett* 250: 287–293. doi: [10.1016/j.femsle.2005.07.016](#) PMID: [16085370](#)
12. Blumenberg M, Seifert R, Michaelis W (2007) Aerobic methanotrophy in the oxic-anoxic transition zone of the Black Sea water column. *Org Geochem* 38: 84–91.
13. Loesekann T, Knittel K, Nadalig T, Fuchs B, Niemann H, Boetius A, et al. (2007) Diversity and abundance of aerobic and anaerobic methane oxidizers at the Haakon Mosby mud volcano, Barents Sea. *Appl Environ Microbiol* 73: 3348–3362. doi: [10.1128/AEM.00016-07](#) PMID: [17369343](#)
14. Talbot HM, Handley L, Spencer-Jones CL, Dinga BJ, Schefuß E, Mann PJ, et al. (2014) Variability in aerobic methane oxidation over the past 1.2 Myrs recorded in microbial biomarker signatures from Congo fan sediments. *Geochim Cosmochim Acta* 133: 387–401.
15. Kessler JD, Valentine DL, Redmond MC, Du M, Chan EW, Mendes SD, et al. (2011) A persistent oxygen anomaly reveals the fate of spilled methane in the deep Gulf of Mexico. *Science* 331: 312–315. doi: [10.1126/science.1199697](#) PMID: [21212320](#)

16. Crespo-Medina M, Meile CD, Hunter KS, Diercks AR, Asper VL, Orphan VJ, et al. (2014) The rise and fall of methanotrophy following a deepwater oil-well blowout. *Nat Geosci* 7: 423–427.
17. Dunfield PF, Yuryev A, Senin P, Smirnova AV, Stott MB, Hou S, et al. (2007) Methane oxidation by an extremely acidophilic bacterium of the phylum *Verrucomicrobia*. *Nature* 450: 879–U18. doi: [10.1038/nature06411](https://doi.org/10.1038/nature06411) PMID: [18004300](https://pubmed.ncbi.nlm.nih.gov/18004300/)
18. Pol A, Heijmans K, Harhangi HR, Tedesco D, Jetten MSM, op den Camp HJM (2007) Methanotrophy below pH1 by a new *Verrucomicrobia* species. *Nature* 450: 874–U17. doi: [10.1038/nature06222](https://doi.org/10.1038/nature06222) PMID: [18004305](https://pubmed.ncbi.nlm.nih.gov/18004305/)
19. Islam T, Jensen S, Reigstad LJ, Larsen O, and Birkeland N- K (2008) Methane oxidation at 55 degrees C and pH 2 by a thermoacidophilic bacterium belonging to the *Verrucomicrobia* phylum. *P Natl Acad Sci USA* 105: 300–304.
20. van Teeseling MCF, Pol A, Harhangi HR, van der Zwart S, Jetten MSM, Op den Camp HJM (2014) Expanding the verrucomicrobial methanotrophic world: description of three novel species of *Methylocidimicrobium* gen. nov. *Appl Environ Microbiol* 80: 6782–6791. doi: [10.1128/AEM.01838-14](https://doi.org/10.1128/AEM.01838-14) PMID: [25172849](https://pubmed.ncbi.nlm.nih.gov/25172849/)
21. Sharp CE, Smirnova AV, Graham JM, Stott MB, Khadka R, Moore TR, et al. (2014) Distribution and diversity of *Verrucomicrobia* methanotrophs in geothermal and acidic environments. *Environ Microbiol* 16: 1867–1878. doi: [10.1111/1462-2920.12454](https://doi.org/10.1111/1462-2920.12454) PMID: [24650084](https://pubmed.ncbi.nlm.nih.gov/24650084/)
22. Cvejic JH, Bodrossy L, Kovacs KL, Rohmer M (2000) Bacterial triterpenoids of the hopane series from the methanotrophic bacteria *Methylocaldum* spp.: phylogenetic implications and first evidence for an unsaturated aminobacteriopanepolyol. *FEMS Microbiol Lett* 182: 361–365. PMID: [10620693](https://pubmed.ncbi.nlm.nih.gov/10620693/)
23. Ourisson G, Albrecht P (1992) Hopanoids 1. Geohopanoids—the most abundant natural-products on Earth. *Acc Chem Res* 25: 398–402.
24. Farrimond P, Fox PA, Innes HE, Miskin IP, Head IM (1998) Bacterial sources of hopanoids in recent sediments: improving our understanding of ancient hopane biomarkers. *Ancient Biomolecules* 2: 147–166.
25. Zundel M, Rohmer M (1985) Prokaryotic triterpenoids. 1. 3-beta-methylhopanoids from *Acetobacter* species and *Methylococcus-capsulatus*. *Eur J Biochem* 150: 23–27. PMID: [3926494](https://pubmed.ncbi.nlm.nih.gov/3926494/)
26. Talbot HM, Watson DF, Pearson EJ, Farrimond P (2003) Diverse bihopanoid compositions of non-marine sediments. *Org Geochem* 34: 1353–1371.
27. Welander PV, Summons RE (2012) Discovery, taxonomic distribution, and phenotypic characterization of a gene required for 3-methylhopanoid production. *P Natl Acad Sci USA* 109: 12905–12910.
28. Osborne KA (2015) Environmental controls on bacteriohopanepolyol signatures in estuarine sediments. PhD Thesis. Newcastle University, UK.
29. Collister JW, Summons RE, Lichtfouse E, Hayes JM (1992) An isotopic biogeochemical study of the Green River oil-shale. *Org Geochem* 19: 265–276. PMID: [11540064](https://pubmed.ncbi.nlm.nih.gov/11540064/)
30. Bouvier P, Rohmer M, Benveniste P, Ourisson G (1976)  $\Delta 8(14)$ -steroids in the bacterium *Methylococcus capsulatus*. *Biochem J* 159: 267–271. PMID: [999649](https://pubmed.ncbi.nlm.nih.gov/999649/)
31. Birgel D, Peckmann J (2008) Aerobic methanotrophy at ancient marine methane seeps: A synthesis. *Org Geochem* 39: 1659–1667.
32. Elvert M, Suess E, Greinert J, Whiticar MJ (2000) Archaea mediating anaerobic methane oxidation in deep-sea sediments at cold seeps of the eastern Aleutian subduction zone. *Org Geochem* 31: 1175–1187.
33. Pancost RD, Sinninghe Damsté JS, de Lint S, van der Maarel M, Gottschal JC, Medinaut Shipboard Scientific Party (2000) Biomarker evidence for widespread anaerobic methane oxidation in Mediterranean sediments by a consortium of methanogenic archaea and bacteria. *Appl Environ Microbiol* 66: 1126–1132.
34. Hinrichs KU (2001) A molecular recorder of methane hydrate destabilization. *Geochem Geophys Geosy* 2: art. no.-2000GC000118.
35. Thiel V, Peckmann J, Richnow HH, Luth U, Reitner J, Michaelis W (2001) Molecular signals for anaerobic methane oxidation in Black Sea seep carbonates and a microbial mat. *Mar Chem* 73: 97–112.
36. Rohmer M, Bouviernave P, Ourisson G (1984) Distribution of hopanoid triterpenes in prokaryotes. *J Gen Microbiol* 130: 1137–1150.
37. Pancost RD, van Geel B, Baas M, Sinninghe Damsté JS (2000)  $\delta^{13}C$  values and radiocarbon dates of microbial biomarkers as tracers for carbon cycling in peat deposits. *Geology* 28: 663–666.
38. van Winden JF, Kip N, Reichart G- J, Jetten MSM, Op den Camp HJM, Sinninghe Damsté JS (2010) Lipids of symbiotic methane-oxidizing bacteria in peat moss studied using stable carbon isotopic labelling. *Org Geochem* 41: 1040–1044.

39. Inglis GN, Collinson ME, Riegel W, Wilde V, Robson BE, Lenz OK, et al. (2015) Ecological and biogeochemical change in an early Paleogene peat-forming environment: Linking biomarkers and palynology. *Palaeogeogr Palaeoclimatol* 438: 245–255.
40. Spencer-Jones CL, Wagner T, Dinga BJ, Schefuß E, Mann PJ, Poulsen JR, et al. (2015) Bacteriohopanepolyols in tropical soils and sediments from the Congo River catchment area. *Org Geochem* 89: 1–13.
41. Birgel D, Feng D, Roberts HH, Peckmann J (2011) Changing redox conditions at cold seeps as revealed by authigenic carbonates from Alaminos Canyon, northern Gulf of Mexico. *Chem Geol* 285: 82–96.
42. Jahnke LL, Summons RE, Hope JM, Des Marais DJ (1999) Carbon isotopic fractionation in lipids from methanotrophic bacteria II: the effects of physiology and environmental parameters on the biosynthesis and isotopic signatures of biomarkers. *Geochim Cosmochim Acta* 63: 79–93. PMID: [11541777](#)
43. Crossman ZM, McNamara N, Parekh N, Ineson P, Evershed RP (2001) A new method for identifying the origins of simple and complex hopanoids in sedimentary materials using stable isotope labelling with  $^{13}\text{C}(\text{CH}_4)$  and compound specific stable isotope analyses. *Org Geochem* 32: 359–364.
44. Neunlist S, Rohmer M (1985) Novel hopanoids from the methylotrophic bacteria *Methylococcus capsulatus* and *Methylomonas methanica*—(22S)-35-aminobacteriohopane-30,31,32,33,34-pentol and (22S)-35-amino-3-beta-methylbacteriohopane-30,31,32,33,34-pentol. *Biochem J* 231: 635–639. PMID: [3935106](#)
45. Talbot HM, Farrimond P (2007) Bacterial populations recorded in diverse sedimentary biohopanoid distributions. *Org Geochem* 38: 1212–1225.
46. van Winden JF, Talbot HM, Kip N, Reichart GJ, Pol A, McNamara NP, et al. (2012) Bacteriohopanepolyol signatures as markers for methanotrophic bacteria in peat moss. *Geochim Cosmochim Acta* 77: 52–61.
47. Blumenberg M, Hoppert M, Krueger M, Dreier A, Thiel V (2012) Novel findings on hopanoid occurrences among sulfate reducing bacteria: Is there a direct link to nitrogen fixation? *Org Geochem* 49: 1–5.
48. Talbot HM, Summons RE, Jahnke LL, Cockell CS, Rohmer M, Farrimond P (2008) Cyanobacterial bacteriohopanepolyol signatures from cultures and natural environmental settings. *Org Geochem* 39: 232–263.
49. Talbot HM, Bischoff J, Inglis GN, Collinson ME, Pancost RD (2016) Polyfunctionalised bio- and geohopanoids in the Eocene Cobham Lignite. *Org Geochem* 96: 77–92.
50. Blumenberg M, Krüger M, Nauhaus K, Talbot HM, Oppermann BI, Seifert R, et al. (2006) Biosynthesis of hopanoids by sulfate-reducing bacteria (genus *Desulfovibrio*). *Environ Microbiol* 8: 1220–1227. doi: [10.1111/j.1462-2920.2006.01014.x](#) PMID: [16817930](#)
51. Blumenberg M, Oppermann BI, Guyoneaud R, Michaelis W (2009) Hopanoid production by *Desulfovibrio bastinii* isolated from oilfield formation water. *FEMS Microbiol Lett* 293: 73–78. doi: [10.1111/j.1574-6968.2009.01520.x](#) PMID: [19222571](#)
52. Talbot HM, Watson DF, Murrell JC, Carter JF, Farrimond P (2001) Analysis of intact bacteriohopanepolyols from methanotrophic bacteria by reversed-phase high-performance liquid chromatography-atmospheric pressure chemical ionisation mass spectrometry. *J Chromatogr A* 921: 175–185. PMID: [11471801](#)
53. Banta AB, Wei JH, Welander PV (2015) A distinct pathway for tetrahymanol synthesis in bacteria. *P Natl Acad Sci USA* 112: 13478–13483.
54. Dedysh SN (2009) Exploring methanotroph diversity in acidic northern wetlands: Molecular and cultivation-based studies. *Microbiology* 78: 655–669.
55. Hirayama H, Abe M, Miyazaki M, Nunoura T, Furushima Y, Yamamoto H, et al. (2014) *Methylomarinovum caldicuralii* gen. nov., sp. nov., a moderately thermophilic methanotroph isolated from a shallow submarine hydrothermal system, and proposal of the family Methylothermaceae fam. nov. *Int J Syst Evol Microbiol* 64: 989–999. doi: [10.1099/ijs.0.058172-0](#) PMID: [24425820](#)
56. Hirayama H, Fuse H, Abe M, Miyazaki M, Nakamura T, Nunoura T, et al. (2013) *Methylomarinum vadi* gen. nov., sp. nov., a methanotroph isolated from two distinct marine environments. *Int J Syst Evol Microbiol* 63: 1073–1082. doi: [10.1099/ijs.0.040568-0](#) PMID: [22707534](#)
57. Hirayama H, Sunamura M, Takai K, Nunoura T, Noguchi T, Oida H, et al. (2007) Culture-dependent and -independent characterization of microbial communities associated with a shallow submarine hydrothermal system occurring within a coral reef off Taketomi Island, Japan. *Appl Environ Microbiol* 73: 7642–7656. doi: [10.1128/AEM.01258-07](#) PMID: [17921273](#)

58. Kalyuzhnaya MG, Khmelenina V, Eshinimaev B, Sorokin D, Fuse H, Lidstrom M, et al. (2008) Classification of halo(alkali)philic and halo(alkali)tolerant methanotrophs provisionally assigned to the genera *Methylomicrobium* and *Methylobacter* and emended description of the genus *Methylomicrobium*. *Int J Syst Evol Microbiol* 58: 591–596. doi: [10.1099/ijs.0.65317-0](https://doi.org/10.1099/ijs.0.65317-0) PMID: [18319461](https://pubmed.ncbi.nlm.nih.gov/18319461/)
59. Elvert M, Niemann H (2008) Occurrence of unusual steroids and hopanoids derived from aerobic methanotrophs at an active marine mud volcano. *Org Geochem* 39: 167–177.
60. Niemann H, Losekann T, de Beer D, Elvert M, Nadalig T, Knittel K, Amann et al. (2006) Novel microbial communities of the Haakon Mosby mud volcano and their role as a methane sink. *Nature* 443: 854–858. doi: [10.1038/nature05227](https://doi.org/10.1038/nature05227) PMID: [17051217](https://pubmed.ncbi.nlm.nih.gov/17051217/)
61. Felden J, Wenzhoefer F, Feseker T, and Boetius A (2010) Transport and consumption of oxygen and methane in different habitats of the Hakon Mosby Mud Volcano (HMMV). *Limnol Oceanogr* 55: 2366–2380.
62. Nickel JC, di Primio R, Mangelsdorf K, Stoddart D, Kallmeyer J (2012) Characterization of microbial activity in pockmark fields of the SW-Barents Sea. *Mar Geol* 332: 152–162.
63. Chand S, Thorsnes T, Rise L, Brunstad H, Stoddart D, Bøe R, et al. (2012) Multiple episodes of fluid flow in the SW Barents Sea (Loppa High) evidenced by gas flares, pockmarks and gas hydrate accumulation. *Earth Planet Sci Lett* 331: 305–314.
64. Crémière A, Lepland A, Chand S, Sahy D, Condon DJ, Noble SR, et al. (2016) Timescales of methane seepage on the Norwegian margin following collapse of the Scandinavian Ice Sheet. *Nat Commun* 7: 1–10.
65. Mastalerz V, de Lange GJ, Dählmann A (2009) Differential aerobic and anaerobic oxidation of hydrocarbon gases discharged at mud volcanoes in the Nile deep-sea fan. *Geochim Cosmochim Acta* 73: 3849–3863.
66. Felden J, Lichtschlag A, Wenzhoefer F, de Beer D, Feseker T, Ristova PP, et al. (2013) Limitations of microbial hydrocarbon degradation at the Amon mud volcano (Nile deep-sea fan). *Biogeosciences* 10: 3269–3283.
67. Baco AR, Rowden AA, Levin LA, Smith CR, Bowden DA (2010) Initial characterization of cold seep faunal communities on the New Zealand Hikurangi margin. *Mar Geol* 272: 251–259.
68. Ruff SE, Arnds J, Knittel K, Amann R, Wegener G, Ramette A, et al. (2013) Microbial Communities of Deep-Sea Methane Seeps at Hikurangi Continental Margin (New Zealand). *PLOS ONE* 8(9): e72627. doi: [10.1371/journal.pone.0072627](https://doi.org/10.1371/journal.pone.0072627) PMID: [24098632](https://pubmed.ncbi.nlm.nih.gov/24098632/)
69. Wild C, Rixen T, Sánchez-Noguera C, Merico A, Jiménez C, Cortés J, et al. (2015) A carbonate platform associated with shallow cold methane seeps in Golfo Dulce, Pacific Costa Rica. *Galaxea, Journal of Coral Reef Studies* 17: 13–14.
70. Rush D, Sinninghe Damsté JS, Poulton SW, Thamdrup B, Garside AL, Acuña González J, et al. (2014) Anaerobic ammonium-oxidising bacteria: A biological source of the bacteriohopanetetrol stereoisomer in marine sediments. *Geochim Cosmochim Acta* 140: 50–64.
71. Sassen R, Roberts HH, Carney R, Milkov AV, DeFreitas DA, Lanoil B, et al. (2004) Free hydrocarbon gas, gas hydrate, and authigenic minerals in chemosynthetic communities of the northern Gulf of Mexico continental slope: relation to microbial processes. *Chem Geol* 205: 195–217.
72. Pancost RD, Zhang CLL, Tavacoli J, Talbot HM, Farrimond P, Schouten S, et al. (2005) Lipid biomarkers preserved in hydrate-associated authigenic carbonate rocks of the Gulf of Mexico. *Palaeogeogr Palaeoclimatol* 227: 48–66.
73. Watson DF (2002) Environmental distribution and sedimentary fate of hopanoid biological marker compounds. PhD Thesis. Newcastle University, UK.
74. Paulmier A, Ruiz-Pino D (2009) Oxygen minimum zones (OMZs) in the modern ocean. *Prog Oceanogr* 80: 113–128.
75. Farrimond P, Head IM, Innes HE (2000) Environmental influence on the biohopanoid composition of recent sediments. *Geochim Cosmochim Acta* 64: 2985–2992.
76. Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Phys* 37: 911–917.
77. Cooke MP, Talbot HM, Farrimond P (2008) Bacterial populations recorded in bacteriohopanepolyol distributions in soils from Northern England. *Org Geochem* 39: 1347–1358.
78. Zink K-G, Mangelsdorf K (2004) Efficient and rapid method for the extraction of intact phospholipids from sediments combined with molecular structure elucidation using LC-ESI-MS/MS analysis. *Anal Bioanal Chem* 380: 798–812. doi: [10.1007/s00216-004-2828-2](https://doi.org/10.1007/s00216-004-2828-2) PMID: [15480579](https://pubmed.ncbi.nlm.nih.gov/15480579/)
79. Bischoff J, Sparkes RB, Doğrul Selver A, Spencer RGM, Gustafsson Ö, Semiletov IP et al. (2016) Source, transport and fate of soil organic matter inferred from microbial biomarker lipids on the East Siberian Arctic Shelf. *Biogeosci Discuss*: doi: [10.5194/bg-2016-128](https://doi.org/10.5194/bg-2016-128)



80. Spencer-Jones CL (2016) Novel concepts derived from microbial biomarkers in the Congo System: Implications for continental methane cycling. PhD Thesis. Newcastle University, UK.
81. Talbot HM, Rohmer M, Farrimond P (2007) Rapid structural elucidation of composite bacterial hopanoids by atmospheric pressure chemical ionisation liquid chromatography/ion trap mass spectrometry. *Rapid Commun Mass Spectrom* 21: 880–892. doi: [10.1002/rcm.2911](https://doi.org/10.1002/rcm.2911) PMID: [17294511](https://pubmed.ncbi.nlm.nih.gov/17294511/)
82. Zhu C, Talbot HM, Wagner T, Pan J- M, Pancost RD (2011) Distribution of hopanoids along a land to sea transect: Implications for microbial ecology and the use of hopanoids in environmental studies. *Limnol Oceanogr* 56: 1850–1865.
83. Summons RE, Jahnke LL, Roksandic Z (1994) Carbon isotopic fractionation in lipids from methanotrophic bacteria: Relevance for interpretation of the geochemical record of biomarkers. *Geochim Cosmochim Acta* 58: 2853–286. PMID: [11540111](https://pubmed.ncbi.nlm.nih.gov/11540111/)
84. Joyeux C, Fouchard S, Llopiz P, Neunlist S (2004) Influence of the temperature and the growth phase on the hopanoids and fatty acids content of *Frateuria aurantia* (DSMZ 6220). *FEMS Microbiol Ecol* 47: 371–379. doi: [10.1016/S0168-6496\(03\)00302-7](https://doi.org/10.1016/S0168-6496(03)00302-7) PMID: [19712325](https://pubmed.ncbi.nlm.nih.gov/19712325/)
85. Fuse H, Ohta M, Takimura O, Murakami K, Inoue H, Yamaoka Y, et al. (1998) Oxidation of trichloroethylene and dimethyl sulfide by a marine *Methylomicrobium* strain containing soluble methane monooxygenase. *Biosci Biotech Bioch* 62: 1925–1931.
86. Jensen S, Neufeld JD, Birkeland N- K, Hovland M, Murrell JC (2008) Methane assimilation and trophic interactions with marine *Methylomicrobium* in deep-water coral reef sediment off the coast of Norway. *FEMS Microbiol Ecol* 66: 320–330. doi: [10.1111/j.1574-6941.2008.00575.x](https://doi.org/10.1111/j.1574-6941.2008.00575.x) PMID: [18811651](https://pubmed.ncbi.nlm.nih.gov/18811651/)
87. Sauter LM, Latypova E, Smalley NE, Lidstrom ME, Hallam S, Kalyuzhnaya MG (2012) Methanotrophic communities of Saanich Inlet: A microcosm perspective. *Syst Appl Microbiol* 35: 198–203. doi: [10.1016/j.syapm.2011.10.006](https://doi.org/10.1016/j.syapm.2011.10.006) PMID: [22406542](https://pubmed.ncbi.nlm.nih.gov/22406542/)
88. Blumenberg M, Seifert R, Kasten S, Bahlmann E, Michaelis W (2009) Euphotic zone bacterioplankton sources major sedimentary bacteriohopanepolyols in the Holocene Black Sea. *Geochim Cosmochim Acta* 73: 750–766.
89. Blumenberg M, Mollenhauer G, Zabel M, Reimer A, Thiel V (2010) Decoupling of bio- and geohopanooids in sediments of the Benguela Upwelling System (BUS). *Org Geochem* 41: 1119–1129.
90. Sáenz JP, Wakeham SG, Eglinton TI, Summons RE (2011) New constraints on the provenance of hopanoids in the marine geologic record: Bacteriohopanepolyols in marine suboxic and anoxic environments. *Org Geochem* 42: 1351–1362.
91. Kharbush JJ, Ugalde JA, Hogle SL, Allen EE, Aluwihare LI (2013) Composite bacterial hopanoids and their microbial producers across oxygen gradients in the water column of the California Current. *Appl Environ Microbiol* 79: 7491–7501. doi: [10.1128/AEM.02367-13](https://doi.org/10.1128/AEM.02367-13) PMID: [24077702](https://pubmed.ncbi.nlm.nih.gov/24077702/)
92. Schaeffer P, Schmitt G, Adam P, Rohmer M (2008) Acid-catalyzed formation of 32,35-anhydrobacteriohopanetetrol from bacteriohopanetetrol. *Org Geochem* 39: 1479–1482.
93. Cooke MP, Talbot HM, and Wagner T (2008) Tracking soil organic carbon transport to continental margin sediments using soil-specific hopanoid biomarkers: A case study from the Congo fan (ODP site 1075). *Org Geochem* 39: 965–971.
94. Bradley AS, Pearson A, Sáenz JP, Marx CJ (2010) Adenosylhopane: The first intermediate in hopanoid side chain biosynthesis. *Org Geochem* 41: 1075–1081.
95. Hamersley MR, Lavik G, Woebken D, Rattray J E, Lam P, Hopmans EC, et al. (2007) Anaerobic ammonium oxidation in the Peruvian oxygen minimum zone. *Limnol Oceanogr* 52: 923–933.
96. Ruff SE, Biddle JF, Teske AP, Knittel K, Boetius A, Ramette A (2015) Global dispersion and local diversification of the methane seep microbiome. *P Natl Acad Sci USA* 112: 4015–4020.
97. Lidstrom ME (1988) Isolation and characterization of marine methanotrophs. *A Van Leeuw J Microb* 54: 189–199.
98. Yan T, Zhou J, Zhang CL (2006) Diversity of functional genes for methanotrophs in sediments associated with gas hydrates and hydrocarbon seeps in the Gulf of Mexico. *FEMS Microbiol Ecol* 57: 251–259. doi: [10.1111/j.1574-6941.2006.00122.x](https://doi.org/10.1111/j.1574-6941.2006.00122.x) PMID: [16867143](https://pubmed.ncbi.nlm.nih.gov/16867143/)
99. Tavormina PL, Ussler W III, Orphan VJ (2008) Planktonic and sediment-associated aerobic methanotrophs in two seep systems along the North American margin. *Appl Environ Microbiol* 74: 3985–3995. doi: [10.1128/AEM.00069-08](https://doi.org/10.1128/AEM.00069-08) PMID: [18487407](https://pubmed.ncbi.nlm.nih.gov/18487407/)
100. Wasmund K, Kurtböke DI, Burns KA, Bourne DG (2009) Microbial diversity in sediments associated with a shallow methane seep in the tropical Timor Sea of Australia reveals a novel aerobic methanotroph diversity. *FEMS Microbiol Ecol* 68: 142–151. doi: [10.1111/j.1574-6941.2009.00667.x](https://doi.org/10.1111/j.1574-6941.2009.00667.x) PMID: [19573197](https://pubmed.ncbi.nlm.nih.gov/19573197/)

101. Moussard H, Stralis-Pavese N, Bodrossy L, Neufeld JD, Murrell JC (2009) Identification of active methylotrophic bacteria inhabiting surface sediment of a marine estuary. *Environ Microbiol Rep* 1: 424–433. doi: [10.1111/j.1758-2229.2009.00063.x](https://doi.org/10.1111/j.1758-2229.2009.00063.x) PMID: [23765896](https://pubmed.ncbi.nlm.nih.gov/23765896/)
102. Redmond MC, Valentine DL, Sessions AL (2010) Identification of novel methane-, ethane-, and propane-oxidizing bacteria at marine hydrocarbon seeps by stable isotope probing. *Appl Environ Microbiol* 76: 6412–6422. doi: [10.1128/AEM.00271-10](https://doi.org/10.1128/AEM.00271-10) PMID: [20675448](https://pubmed.ncbi.nlm.nih.gov/20675448/)
103. Kellermann MY, Schubotz F, Elvert M, Lipp JS, Birgel D, Prieto-Mollar X, et al. (2012) Symbiont-host relationships in chemosynthetic mussels: A comprehensive lipid biomarker study. *Org Geochem* 43: 112–124.
104. Coolen MJL, Talbot HM, Abbas BA, Ward C, Schouten S, Volkman JK, et al. (2008) Sources for sedimentary bacteriohopanepolyols as revealed by 16S rDNA stratigraphy. *Environ Microbiol* 10: 1783–1803. doi: [10.1111/j.1462-2920.2008.01601.x](https://doi.org/10.1111/j.1462-2920.2008.01601.x) PMID: [18397311](https://pubmed.ncbi.nlm.nih.gov/18397311/)
105. Jahnke LL, Summons RE, Dowling LM, Zahiralis KD (1995) Identification of methanotrophic lipid biomarkers in cold-seep mussel gills—chemical and isotopic analysis. *Appl and Environ Microbiol* 61: 576–582.
106. Himmler T, Birgel D, Bayon G, Pape T, Ge L, Bohrmann G, et al. (2015) Formation of seep carbonates along the Makran convergent margin, northern Arabian Sea and amolecular and isotopic approach to constrain the carbon isotopic composition of parent methane. *Chem Geol* 415: 102–117.
107. Berndmeyer C, Thiel V, Schmale O, Blumenberg M. (2013) Biomarkers for aerobic methanotrophy in the water column of the stratified Gotland Deep (Baltic Sea). *Org Geochem* 55: 103–111.
108. Wakeham SG, Amann R, Freeman KH, Hopmans EC, Jorgensen BB, Putnam IF, et al. (2007) Microbial ecology of the stratified water column of the Black Sea as revealed by a comprehensive biomarker study. *Org Geochem* 38: 2070–2097.
109. Wagner T, Kallweit W, Talbot HM, Mollenhauer G, Boom A, Zabel M (2014) Microbial biomarkers support organic carbon transport from methane-rich Amazon wetlands to the shelf and deep sea fan during recent and glacial climate conditions. *Org Geochem* 67: 85–98.
110. Höfle S, Kusch S, Talbot HM, Mollenhauer G, Zubrzycki S, Burghardt S, et al. (2015) Characterization of bacterial populations in Arctic permafrost soils using bacteriohopanepolyols. *Org Geochem* 88: 1–16.
111. Talbot HM, McClymont E, Inglis GN, Evershed R, Pancost RD (2016) Origin and preservation of bacteriohopanepolyol and geohopanoid signatures in Sphagnum peat from Bissendorfer Moor (Germany). *Org Geochem* 97: 95–110.
112. Cooke MP (2011) The role of bacteriohopanepolyols as biomarkers for soil bacterial communities and soil derived organic matter. PhD Thesis. Newcastle University, UK.
113. Gibson RA, Talbot HM, Kaur G, Pancost RD, Mountain B (2008) Bacteriohopanepolyol signatures of cyanobacterial and methanotrophic bacterial populations recorded in a geothermal vent sinter. *Org Geochem* 39: 1020–1023.
114. Farrimond P, Talbot HM, Watson DF, Schulz LK, Wilhelms A (2004) Methylhopanoids: Molecular indicators of ancient bacteria and a petroleum correlation tool. *Geochim Cosmochim Acta* 68: 3873–3882.
115. Birgel D, Meister P, Lundberg R, Horath TD, Bontognali TRR, Bahniuk AM, De Rezende CE, Vasconcelos C, McKenzie JA (2015) Methanogenesis produces strong <sup>13</sup>C enrichment in stromatolites of Lagoa Salgada, Brazil: a modern analogue for Palaeo-/Neoproterozoic stromatolites? *Geobiology* 13: 245–266. doi: [10.1111/gbi.12130](https://doi.org/10.1111/gbi.12130) PMID: [25773379](https://pubmed.ncbi.nlm.nih.gov/25773379/)
116. Kim J-H, Talbot HM, Zarzycka B, Bauersachs T, Wagner T (2011) Occurrence and abundance of soil-specific bacterial membrane lipid markers in the Tet watershed (southern France): Soil-specific BHPs and branched GDGTs. *Geochem Geophys Geosy* 12: Q02008, doi: [10.1029/2010GC003364](https://doi.org/10.1029/2010GC003364)
117. Redshaw CH, Cooke MP, Talbot HM, McGrath S, Rowland SJ (2008) Low biodegradability of fluoxetine HCl, diazepam and their human metabolites in sewage sludge-amended soil. *J Soil Sediment* 8: 217–230.
118. Xu Y, Cooke MP, Talbot HM, Simpson MJ (2009) Bacteriohopanepolyol signatures of bacterial populations in Western Canadian soils. *Org Geochem* 40: 79–86.
119. van Winden JF, Talbot HM, De Vleeschouwer FD, Reichart G-J, Sinninghe Damsté JS (2012) Variation in methanotroph-related proxies in peat deposits from Misten Bog, Hautes-Fagnes, Belgium. *Org Geochem* 53: 73–79.
120. Zhang CL, Huang Z, Li Y-L, Romanek CS, Mills GL, Gibson RA, et al. (2007) Lipid biomarkers, carbon isotopes, and phylogenetic characterization of bacteria in California and Nevada hot springs. *Geomicrobiol J* 24: 519–534.

121. Gibson RA, Sherry A, Kaur G, Pancost RD, Talbot HM (2014) Bacteriohopanepolyols preserved in silica sinters from Champagne Pool (New Zealand) indicate a declining temperature gradient over the lifetime of the vent. *Org Geochem* 69: 61–69.
122. Ricci JN, Coleman ML, Welander PV, Session AL, Summons RE, Spear JR et al. (2014) Diverse capacity for 2-methylhopanoid production correlates with a specific ecological niche. *ISME J* 8:675–684. doi: [10.1038/ismej.2013.191](https://doi.org/10.1038/ismej.2013.191) PMID: [24152713](https://pubmed.ncbi.nlm.nih.gov/24152713/)
123. Blumenberg M, Arp G, Reitner J, Schneider D, Daniel R, Thiel V (2013) Bacteriohopanepolyols in a stratified cyanobacterial mat from Kiritimati (Christmas Island, Kiribati). *Org Geochem* 55: 55–62.
124. Cooke MP, van Dongen BE, Talbot HM, Semiletov I, Shakhova N, Guo L, et al. (2009) Bacteriohopanepolyol biomarker composition of organic matter exported to the Arctic Ocean by seven of the major Arctic rivers. *Org Geochem* 40: 1151–1159.
125. Rethemeyer J, Schubotz F, Talbot HM, Cooke MP, Hinrichs K-U, Mollenhauer G (2010) Distribution of polar membrane lipids in permafrost soils and sediments of a small high Arctic catchment. *Org Geochem* 41: 1130–1145.
126. Doğrul Selver A, Talbot HM, Gustafsson Ö, Boulton S, van Dongen BE (2012) Soil organic matter transport along a sub-Arctic river–sea transect. *Org Geochem* 51: 63–72.
127. Doğrul Selver A, Sparkes RB, Bischoff J, Talbot HM, Gustafsson Ö, Semiletov IP, et al. (2015) Distributions of bacterial and archaeal membrane lipids in surface sediments reflect differences in input and loss of terrestrial organic carbon along a cross-shelf Arctic transect. *Org Geochem* 84: 16–26.
128. De Jonge C, Talbot HM, Bischoff J, Stadnitskaia A, Cherkashov G, Sinninghe Damsté JS (2016) Bacteriohopanepolyol distribution in Yenisei River and Kara Sea suspended particulate matter and sediments traces terrigenous organic matter input. *Geochim Cosmochim Acta* 174: 85–101.
129. Zhu C, Talbot HM, Wagner T, Pan JM, Pancost RD (2010) Intense aerobic methane oxidation in the Yangtze Estuary: A record from 35-aminobacteriohopanepolyols in surface sediments. *Org Geochem* 41: 1056–1059.
130. Sáenz JP, Eglinton TI, Summons RE (2011) Abundance and structural diversity of bacteriohopanepolyols in suspended particulate matter along a river to ocean transect. *Org Geochem* 42: 774–780.
131. Wakeham SG, Turich C, Schubotz F, Podlaska A, Li XN, Varela R, et al. (2012) Biomarkers, chemistry and microbiology show chemoautotrophy in a multilayer chemocline in the Cariaco Basin. *Deep-Sea Res Pt I* 63: 133–156.
132. Taylor KA, Harvey HR (2011) Bacterial hopanoids as tracers of organic carbon sources and processing across the western Arctic continental shelf. *Org Geochem* 42: 487–497.
133. Blumenberg M, Berndmeyer C, Moros M, Muschalla M, Schmale O, Thiel V (2013) Bacteriohopanepolyols record stratification, nitrogen fixation and other biogeochemical perturbations in Holocene sediments of the central Baltic Sea. *Biogeosciences* 10: 2725–2735.
134. Mondamert L, Labanowski J, N'Goye F, Talbot HM, Croué JP (2011) High pressure membrane foulants of seawater, brackish water and river water: origin assessed by sugar and bacteriohopanepolyol signatures. *Biofouling* 27: 21–32. doi: [10.1080/08927014.2010.536614](https://doi.org/10.1080/08927014.2010.536614) PMID: [21104476](https://pubmed.ncbi.nlm.nih.gov/21104476/)