



Frailty index and risk of cardiovascular diseases: a mendelian randomization study

Jun Li^{1#}, Heng Chen^{1#}, Wei He^{2#}, Limin Luo³, Xiaogang Guo¹

¹Department of Cardiology, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China; ²Division of Gastroenterology and Hepatology, Key Laboratory of Gastroenterology and Hepatology, Ministry of Health, State Key Laboratory for Oncogenes and Related Genes, Renji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai, China; ³Department of Dermatology, The First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, China

Contributions: (I) Conception and design: J Li, H Chen, W He; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: J Li, H Chen, W He; (V) Data analysis and interpretation: J Li, H Chen, W He, L Luo; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work and should be considered as co-first authors.

Correspondence to: Xiaogang Guo. Department of Cardiology, The First Affiliated Hospital, Zhejiang University School of Medicine, 79 Qingchun Road, Hangzhou 310003, China. Email: gxx22222@zju.edu.cn.

Background: Previous epidemiological evidence has suggested that frailty status might be associated with cardiovascular diseases (CVDs). However, the exact causality remains unestablished. In this study, we employed Mendelian randomization and sought to investigate the potential causality in association of frailty index (FI) with cardiovascular outcomes [coronary artery disease (CAD), myocardial infarction (MI), atrial fibrillation (AF), and heart failure (HF)].

Methods: Independent single nucleotide polymorphisms (SNPs) at genome-wide significance for FI were obtained from a recent genome-wide association study (GWAS) meta-analysis of European descent (n=175,226). The association of these SNPs with CVDs was examined in summary statistics from corresponding GWASs of European descent (CAD: 184,305 cases and 60,801 controls; MI: 184,305 cases and 43,676 controls; AF: 1,030,836 cases and 60,620 controls; and HF: 977,323 cases and 47,309 controls). Replication analyses were performed using GWAS datasets from FinnGen.

Results: In the meta-analysis of inverse-variance weighted estimates from different data sources, genetically determined higher FI conferred an odds ratio (OR) of 1.46 [95% confidence interval (CI): 1.13 to 1.87; P=0.003] for CAD, 1.62 (95% CI: 1.21 to 2.17, P=0.001) for MI, and 1.46 (95% CI: 1.24 to 1.72; P=4.89×10⁻⁶) for HF. However, FI failed to be potentially influential on AF risk (OR, 1.43; 95% CI: 0.93 to 1.66; P=0.107). Several complementary analyses also received broadly concordant results.

Conclusions: We have provided genetic evidence of a causal association between FI and the risk of CAD, MI, and HF. Further studies are warranted to clarify whether FI is causally related to AF risk.

Keywords: Causal association; cardiovascular diseases (CVDs); frailty index (FI); Mendelian randomization; genome-wide association study (GWAS)

Submitted Aug 12, 2022. Accepted for publication Sep 08, 2022.

doi: 10.21037/atm-22-4239

View this article at: <https://dx.doi.org/10.21037/atm-22-4239>

Introduction

Frailty describes a syndrome in a growing number of older adults, characterized by deteriorative function of multiple systems and increased vulnerability to endogenous as well

as exogenous exposures (1). In the clinical setting, frailty can be defined as a reduction in physical strength and endurance, with or without a decline in cognitive ability. Individuals with frailty experience a higher risk of adverse

outcomes including falls, disability, and hospitalization (1,2). To assess this complex condition, Searle *et al.* introduced the frailty index (FI), which is based on the proportion of age-related deficits among a list of 30 physical parameters (3).

Cardiovascular diseases (CVDs) are commonly known as a group of diseases including coronary artery disease (CAD), myocardial infarction (MI), atrial fibrillation (AF), and heart failure (HF). An increased prevalence of CVDs among the aging population has led to elevated morbidity and mortality worldwide (4). Identification of modifiable risk factors including hypertension, diabetes mellitus, and smoking, has facilitated the management of CVDs. Notably, observational studies have suggested a pattern of associations between frailty and CVDs. Frailty is more prevalent in patients with CVDs than in those without it (5,6). A longitudinal cohort study enrolling 4,211 community-dwellers showed that experiencing baseline frailty was correlated with an increased risk of CVDs over an 8-year follow-up (7). Besides, a previous study has further revealed that the FI might have a more pivotal value than traditional CVD risk factors to discriminate CVD events (8). Regarding the casual effect of frailty on CVDs, true relationships may be distorted by reverse causation or residual confounders (9). Well-designed large-scale cohort studies can to some extent overcome these obstacles and shed light into this issue, but at a relatively high cost.

Therefore, exploring the potential causal link could facilitate enhanced management of CVDs. Mendelian randomization (MR) is an application of genetic variants to plausibly infer the causal associations between phenotypic traits (exposures) and health-related outcomes. The upside of this approach lies in leveraging a large sample size from a genome-wide association study (GWAS) and minimizing bias caused by reverse causation and confounding factors. There are 3 basic principles when performing MR analysis. First, the instrumental variables (IVs) should be associated with exposure of interest at a genome-wide significance level; Second, the IVs are irrelevant to any confounders that may affect the exposure or outcome; Third, the IVs do not directly lead to the outcome, except through its association with the exposure (10). This approach has been applied to the fields of genetics, epidemiology, statistics, econometrics, and bioinformatics (11). In the present study, we used instrumental variables (IVs) identified from a recent GWAS meta-analysis for FI, and performed a 2-sample MR to decipher whether genetically determined higher FI causally leads to increased CVDs risk. The robustness of the results was tested by replicating the main analyses using different

outcome datasets. We present the following article in accordance with the STREGA reporting checklist (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-4239/rc>).

Methods

Study design

The potential causal effect of genetic liability to FI on CVDs was assessed using a 2-sample MR study based on summary statistics from published GWAS meta-analyses and the FinnGen consortium (<https://www.finnngen.fi/en>). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Data sources

Summary-level GWAS data related to CAD/MI, AF, and HF were drawn from the Coronary ARtery Disease Genome-wide Replication and Meta-analysis plus The Coronary Artery Disease Genetics (CARDIoGRAMplusC4D) consortium (12), a GWAS meta-analysis conducted by Nielsen *et al.* (13), and the Heart Failure Molecular Epidemiology for Therapeutic Targets (HERMES) consortium (14), respectively. Replication analyses were performed using data from FinnGen consortium (15). Detailed information for these outcome datasets (sample size, ethnicity, case definition, adjustment, etc.) is listed in the [Table S1](#). No ethical permission or informed consent was necessary given that this MR study was based on publicly available summary statistics.

IVs selection

A total of 14 FI-related single nucleotide polymorphisms (SNPs) were provided by a large GWAS meta-analysis comprising European participants from UK Biobank (n=164,610) and TwinGene (n=10,616) (16). The FI was defined based on the accumulation of deficits as described previously (3). All these SNPs reached a genome-wide association ($P < 5 \times 10^{-8}$) upon adjustment for age, gender, and 10 principal components, and were not in linkage disequilibrium with each other ($r^2 < 0.001$ across a 10,000 kb window) according to the European 1000 genomes panel (17). The details of the associations of genetic variants with the exposure and outcomes are displayed in [Table S2](#). The phenotypic variance explained by

these SNPs (R^2) was calculated using the method described by Shim *et al.* (18). To attain the first assumption of MR design, SNPs with F statistics [$F = R^2(n - k - 1)/k(1 - R^2)$] higher than 10 were considered valid IVs. The rs9275160 was not available in the CARDIoGRAMplusC4D dataset; no suitable proxy was found by searching an online website tool (<https://snipa.helmholtz-muenchen.de/snipa3/>).

Statistical analysis

Effect estimates of each SNP on outcomes were pooled using multiplicative random-effects inverse-variance weighted (IVW) as the primary statistical method (19). The IVW method confers convincing results when all 3 MR assumptions are satisfied. However, it was susceptible to horizontal pleiotropy. Therefore, a set of complementary analyses were used, including the simple median, the weighted median (20), MR-Egger regression (21), and MR pleiotropy residual sum and outlier (MR-PRESSO) (22) methods. These approaches make distinct assumptions on the presence of invalid IVs, and were applied to test the robustness of the results. Heterogeneity was evaluated using Cochrane's Q test and I^2 index. Significant heterogeneity was considered when $P_{\text{Cochrane's Q}} < 0.05$ and $I^2 > 50\%$ for IVW estimates (23). The MR-Egger regression can evaluate horizontal pleiotropy bias by employing its intercept as an indicator ($P_{\text{intercept}} < 0.05$ suggests pleiotropy) (24). In addition, potential pleiotropic outlying IVs were detected by MR-PRESSO approach (22).

The effect size for each SNP was scaled to genetically determined 1 total point increase in FI. The IVW results based on GWAS datasets and the FinnGen consortium were combined using meta-analysis in a fixed-effect model if no heterogeneity was found, otherwise, a random-effect model was applied. Post hoc statistical power was calculated using sample size and proportion of cases of outcome datasets, Type-I error rate (0.05), odds ratio (OR), and percentage of variation explained by IVs (Table S3) (25). Considering the multiple tests, associations with a Bonferroni-corrected P value of < 0.0125 were considered significant. All analyses were performed using R packages TwoSampleMR (26) and MR-PRESSO (22) within software R (version 4.1.0; The R Foundation for Statistical Computing, Vienna, Austria).

Results

For all IVs considered, the F statistics ranged from 28.5 to 113.6, suggesting that these IVs exhibited sufficient strength

for the present MR. All together they explained ~0.3% of phenotypic variation of FI (Table S2).

The IVW analyses showed that a genetically determined 1 point increment in FI conferred an OR of 1.47 [95% confidence interval (CI): 1.10 to 1.96; $P=0.009$] for CAD, 1.62 (95% CI: 1.15 to 2.29, $P=0.006$) for MI, 1.15 (95% CI: 0.92 to 1.44; $P=0.222$) for AF, and 1.42 (95% CI: 1.19 to 1.71; $P=1.34 \times 10^{-4}$) for HF in the GWAS datasets (Figure 1). The causal associations remained broadly consistent in the analyses based on the FinnGen consortium (Figure 1). The meta-analysis combining different data sources was in further support of the causal effect of FI on CAD (OR, 1.46; 95% CI: 1.13 to 1.87; $P=0.003$), MI (OR, 1.62; 95% CI: 1.21 to 2.17; $P=0.001$), and HF (OR, 1.46; 95% CI: 1.24 to 1.72; $P=4.89 \times 10^{-6}$) (Figure 1). However, the results showed that the association for AF may not be causal (OR, 1.43; 95% CI: 0.93 to 1.66; $P=0.107$) (Figure 1).

We had sufficient power ($>90\%$) in detecting the OR in all cases (except for AF) by applying both GWAS meta-analyses and FinnGen consortium (Table S3). Complementary analyses including weighted median, simple median, and MR-PRESSO methods were in accord with prior results, albeit with a smaller magnitude with wide CIs in several analyses (Figure 2). Cochrane's Q test and I^2 index suggested a modest heterogeneity for FI-CAD and FI-MI associations in the FinnGen consortium, whereby no evidence was found for horizontal pleiotropy based on the P value for MR Egger intercept (Table S4). In addition, MR-PRESSO detected no pleiotropic outlying SNPs for all results considered (except for CAD in the FinnGen consortium: rs4146140) (Table S4). There was no observed substantial difference when we recalculated MR estimates by excluding this SNP (Figure 2).

Discussion

The present MR study yielded strong evidence indicating causal relationships between genetically predicted FI and risk of CAD, MI, and HF, with sufficient statistical power. The results were broadly consistent in replication analyses and several complementary analyses. In consideration of heterogeneous results from the different data sources and insufficient power, having a higher FI may not put a person at a higher risk of AF.

The genetic determinants of FI include gene loci associated with traits such as BMI, CVDs, smoking, HLA proteins, depression and neuroticism (16). Higher educational attainment and lower BMI are associated with




Figure 1 IVW analysis estimates of genetically determined FI with CVDs risk. *, the rs9275160 was not available in the CARDIoGRAMplusC4D dataset; no suitable proxy was found. IVW, inverse-variance weighted; FI, frailty index; CVDs, cardiovascular diseases; CARDIoGRAMplusC4D, Coronary ARtery DIsease Genome-wide Replication and Meta-analysis plus The Coronary Artery Disease Genetics; HERMES, Heart Failure Molecular Epidemiology for Therapeutic Targets; SNPs, single-nucleotide polymorphisms; OR, odds ratio; CI, confidence interval.

decreased FI. Besides, mental health plays a pivotal role in the biological mechanisms of frailty (16). The degree of frailty is expected to increase along with aging, and FI has been considered an indicator of biological age that could even outperform DNA methylation age (27). However, frailty acts as a modifiable variable that could be adjusted by controlling the risk factors (1). Investigators in recent years have successfully decoded the underlying the associations between frailty and other diseases and health-related outcomes. The causal association between FI and risk of CAD, MI, and HF found in this MR study was in line with observations from several traditional epidemiology studies (7,28,29). Notably, the clinical data also showed that frailty was an essential independent predictor of prognosis for patients with acute coronary syndrome (30,31). Despite adjusting for some clinical and biochemical variables, current conclusions from observational studies also call for further validation due to limited sample size and biases such as residual confounders and reverse causality. In the present MR study, we managed to provide genetic evidence for the causal relationships between FI and CVDs outcomes. Our conclusion strengthens the conceptual framework that FI increases the risk of CAD, MI, and HF, and that effort to prevent FI may have substantial cardiovascular benefits.

Life style changes, such as regular exercise and appropriate food intake have been considered to curb the progression of frailty (1). Importantly, identification of frailty using FI could aid clinicians in providing better primary, secondary, and tertiary prevention of these CVDs. Besides, frailty index performs better than traditional cardiovascular risk factors in predicting the risk of CVD events (8). Frailty index, together with traditional cardiovascular risk factors, will provide greater prognostic value when clinicians identify those with higher CVDs risks.

Previous studies have suggested some molecular and cellular pathways through which frailty leads to CVDs. First, frail patients often present with higher levels of oxidative stress (32), resulting in an accumulation of cellular damage that impairs endothelium function and further triggers the onset of atherosclerotic CVD (33). Second, the Cardiovascular Health Study reported that frailty status was characterized by elevated inflammatory marker (C-reactive protein) and blood clotting markers (factor VIII and D dimer) (34). High inflammation levels and a hypercoagulable state perpetuate CVDs (35,36). Furthermore, cross-sectional studies have demonstrated that frail people are placed at a higher risk of IGF-1 and sex hormone deficiency (37), which are positively related




Figure 2 Complementary analyses of genetically determined FI with CVDs risk. *, MR-PRESSO instrumental variable outlier detected: rs4146140. FI, frailty index; CVDs, cardiovascular diseases; CARDIoGRAMplusC4D, Coronary ARtery DIsease Genome-wide Replication and Meta-analysis plus The Coronary Artery Disease Genetics; HERMES, Heart Failure Molecular Epidemiology for Therapeutic Targets; MR-PRESSO, MR-pleiotropy residual sum and outlier; OR, odds ratio; CI, confidence interval.

to higher CVDs risk (38,39). A recent cross-sectional study found a pattern of association between frailty and high waist circumference and high body fat mass, and low skeletal muscle mass (40). Meanwhile, these body composition changes have been recognized as risk factors as predictors of CVD outcomes (41,42). Therefore, body composition changes may present one of the biologic mechanisms of how frailty leads to higher CVDs risks.

The frailty state was more pronounced among those developed with AF (43). A longitudinal study from Ireland showed that the frailty phenotype in people with AF may be useful in detecting early deterioration and accelerated aging (44). Besides, patients with AF and FI had a greater tendency to experience stroke, bleeding, and mortality (45). However, prevalent frailty status may not significantly affect AF incidence as reported by the

Framingham Heart Study Offspring cohort study (hazard ratio, 1.22; 95% CI: 0.95 to 1.55) (46), which corroborated the results from the present MR study. This may be due to the fact that the mechanism of AF is to some extent different from that of atherosclerotic CVDs. Nonetheless, it is still plausible that frailty may increase the risk of AF given the limited sample size and insufficient statistical power of this MR study. Further well-designed clinical trials are needed to shed light on this important issue.

GWAS is an observational study which is sought to find genetic variants associated with a trait at genome-wide scale. GWASs provide genome-wide significant variants that can be utilized as instrumental variables in the MR studies (11). The biggest advantage of GWAS is its increased scale and scope in the past decades. Thereby, investigators are enabled to perform MR studies more efficiently (11). The MR framework, by applying hitherto largest GWAS meta-analyses, is one of the major strengths of this study. Using genetic variants randomly assorted and constant after conception, MR analysis minimized the influence of the reverse causation and confounding factors. We had sufficient statistical power to assess the causal association of FI with CAD, MI, and HF. Importantly, the MR estimates were broadly in accordance with across replication analyses using the FinnGen dataset and complementary analyses by other MR methods with no overt horizontal pleiotropy. Taken together, the MR estimates by multiple means reinforced the causal effect of FI on CVDs.

There were several limitations to our study. Firstly, study samples were confined to European cohorts, which may limit the generalizability of conclusions to different populations. Secondly, we had limited statistical power for the FI-AF association in both Nielsen *et al.* GWAS dataset and the FinnGen dataset. This may be ascribed to the small sample size as well as the low percentage of phenotypic variance explained by IVs. Therefore, we should take prudent steps when properly interpreting the corresponding results. Third, moderate heterogeneity was detected in several analyses. However, here we used IVW in multiplicative random effects, which are known to be applicable in the case of heterogeneity (47).

Conclusions

This MR study ascertained a contribution to the causal associations between genetically predicted FI and the risk of CAD, MI, and HF. However, FI may not be causative in AF incidence. Programs aimed at curbing frailty may be of

benefit in the prevention of atherosclerotic CVDs and HF.

Acknowledgments

The authors thank the GWAS meta-analysis of atrial fibrillation (Nielsen *et al.*), CARDIoGRAMplusC4D (Coronary ARtery Disease Genome-wide Replication and Meta-analysis plus The Coronary Artery Disease Genetics), HERMES (Heart Failure Molecular Epidemiology for Therapeutic Targets) consortium, frailty index (Atkins *et al.*), and the FinnGen consortium for providing summary-level data.

Funding: This work was supported by grants from the National Natural Science Foundation of China (Nos. 82170331, U21A20337, and 82003372); and the Key Research and Development Plan of Zhejiang Province (No. 2020C03017).

Footnote

Reporting Checklist: The authors have completed the STREGA reporting checklist. Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-4239/rc>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-4239/coif>). All authors report that this work was supported by grants from the National Natural Science Foundation of China (Nos. 82170331, U21A20337, and 82003372); and the Key Research and Development Plan of Zhejiang Province (No. 2020C03017). The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the

formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Hoogendijk EO, Afilalo J, Ensrud KE, et al. Frailty: implications for clinical practice and public health. *Lancet* 2019;394:1365-75.
2. Clegg A, Young J, Iliffe S, et al. Frailty in elderly people. *Lancet* 2013;381:752-62.
3. Searle SD, Mitnitski A, Gahbauer EA, et al. A standard procedure for creating a frailty index. *BMC Geriatr* 2008;8:24.
4. Lennon RP, Claussen KA, Kuersteiner KA. State of the Heart: An Overview of the Disease Burden of Cardiovascular Disease from an Epidemiologic Perspective. *Prim Care* 2018;45:1-15.
5. Frisoli A Jr, Ingham SJ, Paes ÂT, et al. Frailty predictors and outcomes among older patients with cardiovascular disease: Data from Fragicor. *Arch Gerontol Geriatr* 2015;61:1-7.
6. Afilalo J, Alexander KP, Mack MJ, et al. Frailty assessment in the cardiovascular care of older adults. *J Am Coll Cardiol* 2014;63:747-62.
7. Veronese N, Koyanagi A, Smith L, et al. Multidimensional frailty increases cardiovascular risk in older people: An 8-year longitudinal cohort study in the Osteoarthritis Initiative. *Exp Gerontol* 2021;147:111265.
8. Farooqi MAM, Gerstein H, Yusuf S, et al. Accumulation of Deficits as a Key Risk Factor for Cardiovascular Morbidity and Mortality: A Pooled Analysis of 154 000 Individuals. *J Am Heart Assoc* 2020;9:e014686.
9. Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology contribute to understanding environmental determinants of disease? *Int J Epidemiol* 2003;32:1-22.
10. Burgess S, Scott RA, Timpson NJ, et al. Using published data in Mendelian randomization: a blueprint for efficient identification of causal risk factors. *Eur J Epidemiol* 2015;30:543-52.
11. Burgess S and Thompson SG Mendelian Randomization: Methods for Causal Inference Using Genetic Variants (Second Edition). CRC Press 2015.
12. Nikpay M, Goel A, Won HH, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet* 2015;47:1121-30.
13. Nielsen JB, Thorolfsdottir RB, Fritsche LG, et al. Biobank-driven genomic discovery yields new insight into atrial fibrillation biology. *Nat Genet* 2018;50:1234-9.
14. Shah S, Henry A, Roselli C, et al. Genome-wide association and Mendelian randomisation analysis provide insights into the pathogenesis of heart failure. *Nat Commun* 2020;11:163.
15. Access results | FinnGen. n.d. Available online: https://www.finnngen.fi/en/access_results. (Accessed 20 April 2022).
16. Atkins JL, Jylhävä J, Pedersen NL, et al. A genome-wide association study of the frailty index highlights brain pathways in ageing. *Aging Cell* 2021;20:e13459.
17. 1000 Genomes Project Consortium; Abecasis GR, Altshuler D, et al. A map of human genome variation from population-scale sequencing. *Nature* 2010;467:1061-73.
18. Shim H, Chasman DI, Smith JD, et al. A multivariate genome-wide association analysis of 10 LDL subfractions, and their response to statin treatment, in 1868 Caucasians. *PLoS One* 2015;10:e0120758.
19. Burgess S, Bowden J, Fall T, et al. Sensitivity Analyses for Robust Causal Inference from Mendelian Randomization Analyses with Multiple Genetic Variants. *Epidemiology* 2017;28:30-42.
20. Bowden J, Davey Smith G, Haycock PC, et al. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol* 2016;40:304-14.
21. Burgess S, Thompson SG. Interpreting findings from Mendelian randomization using the MR-Egger method. *Eur J Epidemiol* 2017;32:377-89.
22. Verbanck M, Chen CY, Neale B, et al. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* 2018;50:693-8.
23. Greco M FD, Minelli C, Sheehan NA, et al. Detecting pleiotropy in Mendelian randomisation studies with summary data and a continuous outcome. *Stat Med* 2015;34:2926-40.
24. Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 2015;44:512-25.
25. Brion MJ, Shakhbazov K, Visscher PM. Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol* 2013;42:1497-501.
26. Hemani G, Zheng J, Elsworth B, et al. The MR-Base platform supports systematic causal inference across the human phenome. *Elife* 2018;7:34408.
27. Kim S, Myers L, Wyckoff J, et al. The frailty index

- outperforms DNA methylation age and its derivatives as an indicator of biological age. *Geroscience* 2017;39:83-92.
28. Veronese N, Siggeirsdottir K, Eiriksdottir G, et al. Frailty and Risk of Cardiovascular Diseases in Older Persons: The Age, Gene/Environment Susceptibility-Reykjavik Study. *Rejuvenation Res* 2017;20:517-24.
 29. Arahamian I, Petrella M, Robello EC, et al. The association between cardiovascular risk factors and major cardiovascular diseases decreases with increasing frailty levels in geriatric outpatients. *Exp Gerontol* 2021;153:111475.
 30. Ekerstad N, Swahn E, Janzon M, et al. Frailty is independently associated with short-term outcomes for elderly patients with non-ST-segment elevation myocardial infarction. *Circulation* 2011;124:2397-404.
 31. Alonso Salinas GL, Sanmartin M, Pascual Izco M, et al. Frailty is an independent prognostic marker in elderly patients with myocardial infarction. *Clin Cardiol* 2017;40:925-31.
 32. Mulero J, Zafrilla P, Martinez-Cacha A. Oxidative stress, frailty and cognitive decline. *J Nutr Health Aging* 2011;15:756-60.
 33. Soccio M, Toniato E, Evangelista V, et al. Oxidative stress and cardiovascular risk: the role of vascular NAD(P) H oxidase and its genetic variants. *Eur J Clin Invest* 2005;35:305-14.
 34. Walston J, McBurnie MA, Newman A, et al. Frailty and activation of the inflammation and coagulation systems with and without clinical comorbidities: results from the Cardiovascular Health Study. *Arch Intern Med* 2002;162:2333-41.
 35. Buckley DI, Fu R, Freeman M, et al. C-reactive protein as a risk factor for coronary heart disease: a systematic review and meta-analyses for the U.S. Preventive Services Task Force. *Ann Intern Med* 2009;151:483-95.
 36. Zhang H, Yao J, Huang Z, et al. Prognostic Value of Baseline d-Dimer Level in Patients With Coronary Artery Disease: A Meta-Analysis. *Angiology* 2022;73:18-25.
 37. Cappola AR, Xue QL, Fried LP. Multiple hormonal deficiencies in anabolic hormones are found in frail older women: the Women's Health and Aging studies. *J Gerontol A Biol Sci Med Sci* 2009;64:243-8.
 38. Ren J, Anversa P. The insulin-like growth factor I system: physiological and pathophysiological implication in cardiovascular diseases associated with metabolic syndrome. *Biochem Pharmacol* 2015;93:409-17.
 39. Corona G, Rastrelli G, Monami M, et al. Hypogonadism as a risk factor for cardiovascular mortality in men: a meta-analytic study. *Eur J Endocrinol* 2011;165:687-701.
 40. Xu L, Zhang J, Shen S et al. Association Between Body Composition and Frailty in Elder Inpatients. *Clin. Interv Aging* 2020;15:313-20.
 41. Garg PK, Biggs ML, Kizer JR et al. Associations of body size and composition with subclinical cardiac dysfunction in older individuals: the cardiovascular health study. *International Journal of Obesity* 2021;45:2539-45.
 42. Yoo JH, Park SW, Jun JE et al. Relationship between low skeletal muscle mass, sarcopenic obesity and left ventricular diastolic dysfunction in Korean adults. *Diabetes/ metabolism Research and Reviews* 2021;37:e3363.
 43. Polidoro A, Stefanelli F, Ciacciarelli M, et al. Frailty in patients affected by atrial fibrillation. *Arch Gerontol Geriatr* 2013;57:325-7.
 44. Richard G, O'Halloran AM, Doody P, et al. Atrial fibrillation and acceleration of frailty: findings from the Irish Longitudinal Study on Ageing. *Age Ageing* 2022;51:afab273.
 45. Wilkinson C, Clegg A, Todd O, et al. Atrial fibrillation and oral anticoagulation in older people with frailty: a nationwide primary care electronic health records cohort study. *Age Ageing* 2021;50:772-9.
 46. Orkaby AR, Kornej J, Lubitz SA, et al. Association Between Frailty and Atrial Fibrillation in Older Adults: The Framingham Heart Study Offspring Cohort. *J Am Heart Assoc* 2021;10:e018557.
 47. Bowden J, Del Greco M F, Minelli C, et al. A framework for the investigation of pleiotropy in two-sample summary data Mendelian randomization. *Stat Med* 2017;36:1783-802.
- (English Language Editor: J. Jones)

Cite this article as: Li J, Chen H, He W, Luo L, Guo X. Frailty index and risk of cardiovascular diseases: a mendelian randomization study. *Ann Transl Med* 2022;10(18):1007. doi: 10.21037/atm-22-4239