

Cite this: *Chem. Sci.*, 2024, 15, 6770

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 21st February 2024

Accepted 28th March 2024

DOI: 10.1039/d4sc01234h

rsc.li/chemical-science

Easy access to polyhalogenated biaryls: regioselective (di)halogenation of hypervalent bromines and chlorines†

Daniel Carter Martos,^{ID} ^a Maxime de Abreu,^{ID} ^a Pascal Hauk,^{ID} ^a Philipp Fackler^b and Joanna Wencel-Delord^{ID} ^{†*}

Polyhalogenated biaryls are unique motifs offering untapped potential as versatile building blocks for the expedient synthesis of complex biaryl compounds. Overcoming the limitations of conventional syntheses, we introduce a novel, metal-free, operationally simple and one-pot approach to regioselectively (di)halogenate biaryl compounds under mild conditions using cyclic biaryl hypervalent bromine and chlorine substrates as masked arynes. Through chemoselective post-functionalizations, these valuable products can expand the toolbox for synthesizing biaryl-containing scaffolds, addressing a critical gap in the field.

Introduction


Polyhalogenated biaryls are valuable motifs with applications in many fields such as agrochemicals,¹ pharmaceuticals,² natural products,³ and organic materials⁴ (Scheme 1A). Moreover, polyhalogenated biaryls are uniquely promising as privileged building blocks for the assembly of complex biaryl structures through well-established chemoselective modifications, streamlining the synthesis of value-added compounds.

Despite their importance, constructing polyhalogenated biaryls remains a significant challenge. Conventional transition metal-catalyzed cross-couplings are, by far, the most common approach to prepare such compounds. However, not only do these methods require harsh conditions, expensive catalysts, and pre-installed functionalities, but they typically face significant chemo- and regioselectivity issues when multiple halogens are present, leading to undesired side products, over-functionalizations and polymerizations (Scheme 1B).⁵


Attempts to tackle these challenges through late-stage halogenations, such as Sandmeyer reactions,⁶ pyrylium C–N activations,⁷ electrophilic aromatic substitutions (S_EAr),⁸ or C–H

functionalizations,^{9,10} have their own distinct disadvantages and result in extended and complex synthetic sequences. Alternative approaches to construct polyhalogenated biaryls


A. Natural products, drugs and materials containing polyhalogenated biaryls




B. Conventional cross-coupling methods and drawbacks (generic example)



C. Preparation of cyclic biaryl hypervalent Br and Cl compounds (Lanzi et al.)



D. This work: regioselective (di)halogenation of cyclic biaryl hypervalent Br and Cl compounds.



^aLaboratoire d'Innovation Moléculaire et Applications (LIMA, UMR CNRS 7042), Université de Strasbourg/Université de Haute Alsace, ECPM, 67087 Strasbourg, France. E-mail: wenceldelord@unistra.fr

^bMerck Electronics KGaA, 64293 Darmstadt, Germany

† Electronic supplementary information (ESI) available: Detailed experimental procedures and spectroscopic data for all new compounds. X-ray data analysis for compounds 5b, 5c, 5e and 16. CCDC 2322393–2322396. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4sc01234h>

‡ Present address: Institut für Organische Chemie, Universität Würzburg, Am Hubland, 97074 Würzburg, Germany. Email: joanna.wencel-delord@uni-wuerzburg.de

Scheme 1 Importance and synthesis of polyhalogenated biaryl compounds.



using lithium bases have also been explored, but are frequently substrate-specific and utilize harsh conditions or hazardous reagents.¹¹

Notably, the aforementioned methods are incompatible with the synthesis of biaryls bearing highly reactive iodine and bromine substituents at specific positions. As a result, numerous polyhalogenated biaryls are commercially unavailable, prohibitively expensive, or yet undiscovered. This highlights the need for more innovative, versatile and efficient protocols to selectively introduce halogen motifs on biaryl scaffolds.

Recently, our group has delved into investigating the chemistry of bench-stable cyclic biaryl hypervalent bromine and chlorine reagents, which remain rare and poorly explored. These unusual reagents, which are safe to handle and easy to prepare (Scheme 1C), undergo unexpected metal-free cycloadditions and *meta*-functionalizations with C-, O- and N-nucleophiles though *in situ* aryne formation at room temperature.¹²

Based on this, we hypothesized that harnessing the unique reactivity of such hypervalent bromine and chlorine compounds as masked biaryl arynes provides a conceptually different strategy to prepare a wide array of polyhalogenated biaryl compounds in a mild and straightforward manner.

It is worth noting that several copper-catalyzed methodologies for the halogenation of analogous cyclic diaryliodonium salts have been reported.¹³ Nevertheless, this approach diverges in its mechanism, yielding selectively *ortho*-functionalized products, while *meta*-functionalization of hypervalent iodines remains extremely rare and requires high reaction temperatures and drastic conditions.¹⁴

In clear contrast, the use of hypervalent bromine and chlorine precursors, which are accessible either through standard Suzuki coupling or metal-free conditions, opens up the possibility of generating biaryl aryne intermediates, which have the potential to undergo vicinal difunctionalization through the concerted addition of both nucleophilic and electrophilic species under extremely mild reaction conditions (Scheme 1C). Such a double functionalization presents however an important challenge. Indeed, while 1,2-difunctionalization of arynes is widely reported, mainly through cycloadditions or reactions with ambiphilic reagents,¹⁵ versatile 3-component strategies remain uncommon.¹⁶ Furthermore, to the best of our knowledge, the 1,2-dihalogenation of arynes is mostly limited to diiodo, dibromo or dichlorinations, and is scarcely reported with mixed halogens.¹⁷ Moreover, the electrophilic intermolecular trapping of arynes originating from cyclic diaryl-bromonium or chloronium salts remains unaccomplished.¹⁸ In this work, we report the first example of a concerted nucleophilic and electrophilic intermolecular 1,2-difunctionalization of our biaryl aryne precursors, allowing the direct, mild and metal-free installation of customized halogen motifs in a fully regioselective manner (Scheme 1D).

Results and discussion

Meta-selective monohalogenations

Our investigation began with the bromination of hypervalent compound **1a** to synthesize product **3b**. Addition of Cs₂CO₃ as mild base prompted simultaneous deprotonation and ring


opening of **1a** to generate aryne intermediate **Int1**. *In situ* attack of **Int1** with a nucleophilic bromide source should thus achieve the targeted *meta*-selective bromination.

Preliminary experiments using alkali metal bromides (KBr, NaBr, and LiBr) showed promising results, particularly when employed alongside the corresponding crown ether additives (Table 1, entries 1–4). Increased temperature did not lead to higher yields (entry 5), and swapping solvents (entries 6, and 7) or reagents (entry 8, and 9) led to varying results, frequently producing mixtures of *meta* and *ortho* bromination. Remarkably, the most favourable outcomes were obtained using tetra-*n*-butylammonium bromide (TBAB), furnishing the desired product in quantitative yields and full *meta*:*ortho* selectivities (entries 11–13).

To broaden the generality of this methodology, other halogenation reactions, namely iodination, chlorination and fluorination, were subsequently investigated. Re-evaluation of the most promising conditions in Table 1 showed that tetra-*n*-butylammonium halide salts in THF provided the optimal global reaction conditions for all types of halogenations (see ESI†).


With the optimized conditions in hand, the scope of this transformation was investigated (Scheme 2). We initially explored all possible halogen permutations, achieving quantitative yields for most combinations except fluorination products **3d** and **4d** (82% and 88%, respectively), and iodination product **4a** (50%). Electron donating substituents, including methyl (**3e**, **4e**), methoxy (**3h**), and *tert*-butyl (**3j**), afforded the corresponding products in moderate to excellent yields (49–90%). Likewise, electron withdrawing groups such as –CF₃ (**3k**)

Table 1 Optimization of *meta*-selective bromination^a



Entry	[Br ⁻]	Additive	Solvent	Yield, % ^b
1	KBr	—	DCM	(42)/(54) ^c
2	KBr	18-Crown-6	DCM	(71)
3	NaBr	15-Crown-5	DCM	56
4	LiBr	12-Crown-4	DCM	99 (87)
5	LiBr	—	DCM	67/43 ^d
6	LiBr	—	MeCN	99 (77)
7	LiBr	—	MeCN/H ₂ O 1 : 1	40 + 40 ^e
8	NH ₄ Br	—	DCM/H ₂ O 1 : 1	3 + 34 ^e
9	TBAB	—	DCM	30 + 24 ^e
10	TBAB	—	MeCN	72
11	TBAB	—	MeCN/H ₂ O 1 : 1	99 (99)
12	TBAB	—	THF	99 (99)
13	TBAB	—	AcOEt	99

^a Standard conditions: **1a** (0.10 mmol), Cs₂CO₃ (2.0 equiv.), [Br⁻] nucleophile (2.0 equiv.), additive (2.0 equiv.), solvent (2.0 mL), argon, room temperature, 16 h. ^b Yields determined by ¹H NMR analysis using CH₂Br₂ as internal standard; isolated yields in parentheses. ^c 10 equiv. KBr. ^d 40 °C. ^e Undesired *ortho*-bromination.




Scheme 2 Scope of *meta*-selective halogenations. ^aStandard conditions: hypervalent bromine **1** or chlorine **2** (0.10 mmol), Cs₂CO₃ (2.0 equiv.), tetra-*n*-butylammonium halide TBAY (2.0 equiv.), THF (2.0 mL), argon, room temperature, 16 h. Isolated yields are reported unless otherwise stated. ^bYield determined by ¹H NMR analysis using CH₂Br₂ as internal standard. Volatile product, isolated yield: 30%. ^cThe isolated yield encompasses the total yield of both isomers, irrespective of their regiochemical pathway A/B. The subsequent ratio indicates the proportion of this total obtained through pathway A relative to pathway B, and was determined *via* ¹H NMR analysis.

and -CO₂Me (**3l**) led to good yields, and push-pull type structures (**3f**, **3g**, **3i**) yielded the respective halogenations in moderate to high yields (55–92%).

While the nucleophilic attack on the aryne occurs selectively at the *meta* position, the initial deprotonation and generation of the aryne species itself is less regioselective. In non-symmetrical substrates where the aryne formation was possible on both aromatic rings, mixtures of regioisomers resulted from their subsequent *meta*-selective halogenations. Despite this, a preference towards deprotonation and the resulting halogenation on the most electron-poor aromatic ring was observed (Scheme 2, path A vs. path B, compounds **3h–l**).

Extension to 1,2-dihalogenations

With the monohalogenation conditions established, we aimed to further expand the methodology to 1,2-dihalogenations by treating the aryne intermediate with both nucleophilic and



Scheme 3 Predicted aryl carbanion species and subsequent electrophilic trapping.

electrophilic halogen sources in a one-pot procedure. We hypothesized that a *meta*-selective nucleophilic attack on the aryne moiety would generate a highly reactive aryl carbanion species on the adjacent *ortho* position, which, under appropriate conditions, could be trapped by suitable halogen electrophiles (Scheme 3). It must be highlighted that, for the monohalogenation protocol, this reactive carbanion is likely quenched by deprotonating another molecule of starting material in an autocatalytic fashion.

To develop such a rare dihalogenation protocol, we began by screening various electrophilic iodine reagents alongside the optimized monobromination conditions of TBAB in THF (Table 2). Initial attempts yielded limited success with common iodine electrophiles such as NIS, I₂ or I-Cl (entries 1–5). However, the use of nonafluoro-1-iodobutane resulted in a comparatively remarkable outcome, furnishing the desired dihalogenated biaryl **5b** in 86% yield (entry 6).

Table 2 Optimization of regioselective dihalogenation^a

Entry	Z	[Z ⁺] (equiv.)	T, °C	Yield ^b , %
1	I (5b)	I ₂ (1.2)	rt	< 5
2		NIS (3.0)	rt	< 5
3		I-Cl (3.0)	rt	< 5
4		CHI ₃ (3.0)	rt	nr ^c
5		<i>n</i> C ₆ F ₁₃ -I (3.0)	rt	nr
6		<i>n</i> C ₄ F ₉ -I (5.0)	rt	(86)
7	Br (5e)	CBr ₄ (5.0)	rt	< 5
8		NBS (5.0)	rt	< 5
9		<i>n</i> C ₆ F ₁₃ -Br (5.0)	rt	6
10		C ₆ F ₅ -Br (20)	rt	17
11		<i>n</i> C ₈ F ₁₇ -Br (15)	50	(16)
12		C ₆ F ₅ -Br (15)	50	(91)
13	Cl (5h)	CCl ₄ (5.0)	rt	< 5
14		CCl ₄ (1.0 mL)	rt	nr
15		NCS (5.0)	rt	nr
16		C ₆ F ₅ -Cl (5.0)	rt	< 5
17		C ₆ F ₅ -Cl (15)	50	(30)
18		CCl ₄ (15)	50	(53)

^a Standard conditions: **1a** (0.10 mmol), Cs₂CO₃ (2.0 equiv.), TBAB (2.0 equiv.), [Z⁺] electrophile, THF (2.0 mL), argon, Temperature, 16 h. ^b Yields determined by ¹H NMR analysis using CH₂Br₂ as internal standard; isolated yields in parentheses. ^c No reaction.

Motivated by this breakthrough, we set out to expand the protocol to bromane and chlorane reagents. However, the screening of Br and Cl electrophiles proved more challenging due to their decreased reactivity compared to their iodine counterparts. While some modest improvements were made using perfluoroalkyl and perfluoroaryl halides as the electrophile sources, the yields remained low (Table 2, entries 9, 10 and 16). Ultimately, temperature was found to be a key factor, as the best results for trapping with Br (entry 12), and Cl (entry 18), were observed when performing the reaction at 50 °C.


Having established the conditions for the regioselective dihalogenations, the scope of the reaction was surveyed for both hypervalent bromine and chlorine substrates (Scheme 4). Biaryls containing halogens in several different arrangements were synthesized in moderate to excellent yields, including *ortho*-iodinated products **5a–d** and **6a–b** (45–98%), *ortho*-brominated products **5e–f** (88–91%), and *ortho*-chlorinated products **5g–h** and **6c** (53–74%). We likewise examined the compatibility of the procedure with electron donating groups such as –Me (**5j**, **5k**, **6d**) and –OMe (**6e**); and electron withdrawing groups such as –CO₂Me **5l**, all of which resulted in modest to high yields (56–75%). We were also able to dihalogenate binaphthyl-derived substrate **5i** in 69% yield, a particularly interesting result

given the potential of such compounds as building blocks for non-C₂ symmetrical privileged ligands. Crucially, the potential for this methodology to be extended to other electrophiles was demonstrated *via* the electrophilic trapping with MeOD to furnish deuterated product **5m** in 87% yield and 92% deuterium incorporation. Furthermore, compound **5m** offered additional proof that our electrophiles were indeed functionalizing the desired *ortho* position. X-ray structures were also obtained for selected dihalogenated products, unequivocally confirming the predicted regioselectivity of this transformation.


Chemoselective post-functionalizations

The potential of **5** and **6** as new and unique building blocks was then investigated *via* chemoselective and sequential post-functionalizations on products **5c** and **5g** (Scheme 5).

Compound **5g** was first subjected to several iodine-selective transformations. Grignard formation with *i*-PrMgCl followed by treatment with DMF furnished benzaldehyde **7** in moderate but non-optimized 42% yield, and Pd-catalyzed Sonogashira, Buchwald-Hartwig and Suzuki–Miyaura cross-couplings yielded



Scheme 4 Scope of the regioselective dihalogenations. Standard conditions: hypervalent bromine **1** or chlorine **2** (0.10 mmol), Cs₂CO₃ (2 equiv.), tetra-*n*-butylammonium halide TBAY (2.0 equiv.), [Z⁺] electrophile [Z=I: nonafluoro-1-iodobutane nC₄F₉-I (5.0 equiv.); Z=Br: bromopentafluorobenzene C₆F₅-Br (15 equiv.); Z=Cl: carbon tetrachloride CCl₄ (15 equiv.)], THF (2.0 mL), argon, temperature [Z=I: room temperature, Z=Br/Cl: 50 °C], 16 h. Isolated yields are reported unless otherwise stated. ^aFollowing standard conditions, electrophile: MeOD (30 equiv.), temperature: 50 °C.



Scheme 5 Chemoselective post-functionalizations of **5g** and **5c**. Conditions: (a) *i*-PrMgCl, THF, 0 °C then DMF, 0 °C to rt. (b) *p*-(trifluoromethyl)phenylacetylene, Pd(PPh₃)₂Cl₂, K₂CO₃, CuI, DME/H₂O, 100 °C. (c) *p*-TolNHMe, Pd(OAc)₂, DPEphos, NaOtBu, toluene, 100 °C. (d) *p*-TolB(OH)₂, Pd(PPh₃)₂Cl₂, K₂CO₃, DME/H₂O, 100 °C. (e) 3-pyridylB(OH)₂, Pd(PPh₃)₄, K₂CO₃, dioxane/H₂O, 100 °C. (f) cyclopropylB(OH)₂, Pd(OAc)₂, PCy₃, K₃PO₄, toluene/H₂O, 100 °C. (g) *n*-BuLi, THF, –78 °C then PhCHO, –78 °C to rt. (h) *p*-anisidine, Pd(dppf)Cl₂, NaOtBu, dioxane, 120 °C. (i) PhB(OH)₂, Pd(OAc)₂, PPh₃, K₂CO₃, THF/H₂O, 80 °C. (j) *t*-BuLi, THF, –78 °C then DMF, –30 °C. (k) *m*-CPBA, TFOH, DCM, 0 °C to rt.

the respective functionalized biaryls **8–10** in moderate to quantitative yields, with little to no presence of over-functionalization on bromine.

Selective functionalization on bromine over chlorine can also be achieved, as illustrated with compound **10** *via* Suzuki–Miyaura cross-couplings with 3-pyridylboronic acid and cyclopropylboronic acid, affording products **11** and **12** respectively in moderate to high yields. Additionally, selective lithium–bromine exchange on **10** was realized, followed by trapping with benzaldehyde, which furnished atropostable biaryl **13** in 61% yield as a mixture of atropodiastereomers.

Compound **5c** proved more challenging to functionalize selectively due to the increased steric hindrance about the biaryl axis. Buchwald–Hartwig amination using *p*-anisidine led to substituted carbazole **14** in 63% yield as a single product, while Suzuki–Miyaura arylation successfully furnished **15** in 58% yield. Interestingly, fluorene derivative **16** was obtained in 54% yield by treatment of **5c** with *tert*-butyllithium, followed by addition of DMF, and its structure was confirmed by X-ray analysis. Finally, we were also able to achieve a selective oxidation on iodine to form appealing cyclic diaryliodonium compound **17** in a considerable 44% yield, which could serve as a unique platform for subsequent transformations.

Conclusions

In conclusion, we have developed a mild, metal-free and highly efficient regioselective (di)halogenation procedure by harnessing the reactivity of cyclic biaryl hypervalent bromine and chlorine compounds. By integrating mild base-mediated aryne formation with *in situ* nucleophilic attack, employing readily available and environmentally benign tetra-*n*-butylammonium halides, we successfully furnished a wide variety of *meta*-halogenated 2-chloro- and 2-bromobiaryls in up to quantitative yields at room temperature. More significantly, we expanded the scope of our methodology to vicinal dihalogenations by achieving an unprecedented electrophilic intermolecular trapping the reactive biaryl carbanion intermediates, effortlessly preparing a range of rare 2,2',3'-trihalogenated biaryl compounds with perfect site selectivity in a simple, mild, one-pot process. Through our innovative and conceptually distinct (di)halogenation protocol, we unlock the opportunity to efficiently and flexibly synthesize a diverse library of tailor-made polyhalogenated biaryls stemming from only a handful of common hypervalent compounds as precursors. This approach thus circumvents the need for complex, multi-step *de novo* syntheses which would be required to produce such a wide variety of rare and valuable compounds. Furthermore, some compounds synthesized utilizing our method may be altogether inaccessible through traditional approaches, underscoring the importance of our protocol in expanding the space of polyhalogenated biaryls. Moreover, the unique potential of these polyhalogenated biaryls as powerful synthetic building blocks was subsequently demonstrated through several chemoselective post-functionalization reactions, thus providing a valuable toolkit for the streamlined synthesis of polyfunctionalized biaryl scaffolds.

Data availability

Detailed experimental procedures, characterization data and crystallographic data are available in the ESI.†

Author contributions

J. W.-D. and P. F. conceived and directed the project. D. C. M. and M. d. A. developed the method and performed the synthetic experiments. D. C. M. and P. H. carried out the post-functionalization studies. D. C. M. and J. W.-D. wrote the manuscript with assistance from all authors.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

J. W.-D., D. C. M., P. H. and P. F. thank the European Union's H2020 research and innovation programme for funding this project under the Marie Skłodowska-Curie Grant Agreement No. 860762. This work was financially supported by the CNRS (Centre National de la Recherche Scientifique), the Université de Strasbourg, the Ministère de l'Éducation Nationale et de la Recherche (France) and Merck Electronics KGaA (Darmstadt, Germany). J. W. D. and M. d. A. are grateful to the European Commission for the ERC-Starting Grant "ALCHIMIE" No. 949804. The authors thank Dr Emeric Wasielewski (NMR service) and Matthieu Chessé (analytical facilities) from LIMA (UMR CNRS 7042); Dr Lydia Karmazin, Dr Nathalie Gruber and Dr Corinne Bailly for X-ray diffraction analyses (Service de Radiocristallographie, Fédération de Chimie "Le Bel" FR2010, Université de Strasbourg); and Dr Hélène Nierengarten and Dr Stéphanie Coutin for High Resolution Mass Spectrometry analyses (Service de Spectrométrie de Masse, Fédération de Chimie "Le Bel" FR2010, Université de Strasbourg). The authors are also thankful to Prof. Françoise Colobert, Dr Sabine Choppin, Dr Mark M. Maturi and Dr Tomas J. Saiegh for helpful discussions.

Notes and references

- 1 Illustrative examples may be found in: P. Jeschke, *Eur. J. Org. Chem.*, 2022, 2022, e202101513, DOI: [10.1002/ejoc.202101513](https://doi.org/10.1002/ejoc.202101513).
- 2 For selected examples, see: (a) D. Benedetto Tiz, L. Bagnoli, O. Rosati, F. Marini, L. Sancineto and C. Santi, *Molecules*, 2022, 27, 1643; (b) Z.-W. Zhang, H. Tu, M. Jiang, S. Vanan, S. Y. Chia, S.-E. Jang, W.-T. Saw, Z.-W. Ong, D.-R. Ma, Z.-D. Zhou, J. Xu, K.-H. Guo, W.-P. Yu, S.-C. Ling, R. A. Margolin, D. G. Chain, L. Zeng and E.-K. Tan, *Sci. Signaling*, 2022, 15, eabk3411; (c) Y. Wang, S. Dana, H. Long, Y. Xu, Y. Li, N. Kaplaneris and L. Ackermann, *Chem. Rev.*, 2023, 123, 11269–11335; (d) M. Bruncko, T. K. Oost, B. A. Belli, H. Ding, M. K. Joseph, A. Kunzer, D. Martineau, W. J. McClellan, M. Mitten, S.-C. Ng,

- P. M. Nimmer, T. Oltersdorf, C.-M. Park, A. M. Petros, A. R. Shoemaker, X. Song, X. Wang, M. D. Wendt, H. Zhang, S. W. Fesik, S. H. Rosenberg and S. W. Elmore, *J. Med. Chem.*, 2007, **50**, 641–662; (e) T. S. Haque, R. L. Martinez, V. G. Lee, D. G. Riexinger, M. Lei, M. Feng, B. Koplowitz, C. Mapelli, C. B. Cooper, G. Zhang, C. Huang, W. R. Ewing and J. Krupinski, *Peptides*, 2010, **31**, 1353–1360; (f) R. R. Shah, J. M. Redmond, A. Mihut, M. Menon, J. P. Evans, J. A. Murphy, M. A. Bartholomew and D. M. Coe, *Bioorg. Med. Chem.*, 2020, **28**, 115326.
- 3 Selected examples: (a) A. Isnansetyo and Y. Kamei, *Antimicrob. Agents Chemother.*, 2003, **47**, 480–488; (b) S. Adak and B. S. Moore, *Nat. Prod. Rep.*, 2021, **38**, 1760–1774; (c) C. Wagner, M. El Omari and G. M. König, *J. Nat. Prod.*, 2009, **72**, 540–553.
- 4 For illustrative examples, see: (a) M. L. Tang and Z. Bao, *Chem. Mater.*, 2011, **23**, 446–455; (b) B. Park, H. Bae, J.-W. Ha, C. Lee, J. Lee, Y. Heo, B. Kim, S. C. Yoon, H. Choi and S.-J. Ko, *Org. Electron.*, 2023, **113**, 106717; (c) H. Shudo, M. Kuwayama, M. Shimasaki, T. Nishihara, Y. Takeda, N. Mitoma, T. Kuwabara, A. Yagi, Y. Segawa and K. Itami, *Nat. Commun.*, 2022, **13**, 3713; (d) Y.-M. Liu, H. Hou, Y.-Z. Zhou, X.-J. Zhao, C. Tang, Y.-Z. Tan and K. Müllen, *Nat. Commun.*, 2018, **9**, 1901; (e) H. Tamaoki, R. Katoono, K. Fujiwara and T. Suzuki, *Angew. Chem., Int. Ed.*, 2016, **55**, 2582–2586; (f) Y. Sakamoto, T. Suzuki, A. Miura, H. Fujikawa, S. Tokito and Y. Taga, *J. Am. Chem. Soc.*, 2000, **122**, 1832–1833; (g) S. Komatsu, Y. Sakamoto, T. Suzuki and S. Tokito, *J. Solid State Chem.*, 2002, **168**, 470–473; (h) M. Ikai, S. Tokito, Y. Sakamoto, T. Suzuki and Y. Taga, *Appl. Phys. Lett.*, 2001, **79**, 156–158; (i) Y. Liu, Y. Huang, S. Liu, D. Chen, C. Tang, Z. Qiu, J. Zhu and Y. Tan, *Angew. Chem., Int. Ed.*, 2019, **58**, 13276–13279; (j) M. Mahl, K. Shoyama, J. Rühle, V. Grande and F. Würthner, *Chem.–Eur. J.*, 2018, **24**, 9409–9416.
- 5 Selected reviews: (a) C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot and V. Snieckus, *Angew. Chem., Int. Ed.*, 2012, **51**, 5062–5085; (b) A. Biffis, P. Centomo, A. del Zotto and M. Zecca, *Chem. Rev.*, 2018, **118**, 2249–2295; (c) J. Pérez Sestelo and L. A. Sarandeses, *Molecules*, 2020, **25**, 4500.
- 6 Representative examples: (a) F.-C. Favre-Besse, O. Poiré, T. Bersot, E. Kim-Grellier, S. Daumas, S. El Mestikawy, F. C. Acher and N. Pietrancosta, *Eur. J. Med. Chem.*, 2014, **78**, 236–247; (b) V. Akhmetov, M. Feofanov, S. Troyanov and K. Amsharov, *Chem.–Eur. J.*, 2019, **25**, 7607–7612; (c) K. Kantarod, D. Soorukram, C. Kuhakarn, P. Surawatanawong, W. Wattanathana, V. Reutrakul and P. Leowanawat, *Chem. Commun.*, 2022, **58**, 9468–9471.
- 7 C. Ghiazza, T. Faber, A. Gómez-Palomino and J. Cornella, *Nat. Chem.*, 2022, **14**, 78–84.
- 8 (a) P. B. D. de la Mare, *Electrophilic Halogenation: Reaction Pathways Involving Attack by Electrophilic Halogens on Unsaturated Compounds; Cambridge Chemistry Texts*, Cambridge University Press, Cambridge, 1976; (b) B. Galabov, D. Nalbantova, P. V. R. Schleyer and H. F. Schaefer, *Acc. Chem. Res.*, 2016, **49**, 1191–1199; (c) W. Wang, X. Yang, R. Dai, Z. Yan, J. Wei, X. Dou, X. Qiu, H. Zhang, C. Wang, Y. Liu, S. Song and N. Jiao, *J. Am. Chem. Soc.*, 2022, **144**, 13415–13425.
- 9 Selected reviews on the topic: (a) J. Wang, C. Liu, Q. Zheng and G. Rao, *Eur. J. Org. Chem.*, 2020, **2020**, 3737–3765; (b) R. Das and M. Kapur, *Asian J. Org. Chem.*, 2018, **7**, 1524–1541; (c) D. A. Petrone, J. Ye and M. Lautens, *Chem. Rev.*, 2016, **116**, 8003–8104.
- 10 Additional examples: (a) C. K. Hazra, Q. Dherbassy, J. Wencel-Delord and F. Colobert, *Angew. Chem., Int. Ed.*, 2014, **53**, 13871–13875; (b) S. T. Linde, V. Corti, V. H. Lauridsen, J. N. Lamhauge, K. A. Jørgensen and N. M. Rezayee, *Chem. Sci.*, 2023, **14**, 3676–3681.
- 11 (a) V. Mamane, E. Aubert, P. Peluso and S. Cossu, *J. Org. Chem.*, 2013, **78**, 7683–7689; (b) F. Leroux and M. Schlosser, *Angew. Chem., Int. Ed.*, 2002, **41**, 4272–4274; (c) L. Bonnafoux, R. Gramage-Doria, F. Colobert and F. R. Leroux, *Chem.–Eur. J.*, 2011, **17**, 11008–11016; (d) F. R. Leroux, L. Bonnafoux, C. Heiss, F. Colobert and D. A. Lanfranchi, *Adv. Synth. Catal.*, 2007, **349**, 2705–2713.
- 12 For benchmark chemistry on synthesis and functionalization of hypervalent bromine and chlorine reagents from our research group, see: (a) M. Lanzi, Q. Dherbassy and J. Wencel-Delord, *Angew. Chem., Int. Ed.*, 2021, **60**, 14852–14857; (b) M. Lanzi, R. A. Ali Abdine, M. De Abreu and J. Wencel-Delord, *Org. Lett.*, 2021, **23**, 9047–9052; (c) M. Lanzi, T. Rogge, T. S. Truong, K. N. Houk and J. Wencel-Delord, *J. Am. Chem. Soc.*, 2023, **145**, 345–358; (d) M. de Abreu, T. Rogge, M. Lanzi, T. Saiegh, K. N. Houk and J. Wencel-Delord, *Angew. Chem., Int. Ed.*, 2024, e202319960, DOI: [10.1002/anie.202319960](https://doi.org/10.1002/anie.202319960); (e) M. Lanzi and J. Wencel-Delord, *Chem. Sci.*, 2024, **15**, 1557–1569.
- 13 (a) B. Wu and N. Yoshikai, *Angew. Chem., Int. Ed.*, 2015, **54**, 8736–8739; (b) J. Ke, B. Zu, Y. Guo, Y. Li and C. He, *Org. Lett.*, 2021, **23**, 329–333; (c) K. Zhu, Z. Song, Y. Wang and F. Zhang, *Org. Lett.*, 2020, **22**, 9356–9359.
- 14 H. Jiang, M. Liu, Z. Ye, Z. Song, Y. Wu and F. Zhang, *Org. Chem. Front.*, 2023, **10**, 3856–3860.
- 15 Selected reviews and articles: (a) X. Yang and G. Chit Tsui, *Chem. Sci.*, 2018, **9**, 8871–8875; (b) A. Nilova, P. A. Sibbald, E. J. Valente, G. A. González-Montiel, H. C. Richardson, K. S. Brown, P. H. Cheong and D. R. Stuart, *Chem.–Eur. J.*, 2021, **27**, 7168–7175; (c) B. Baire, D. Niu, P. H. Willoughby, B. P. Woods and T. R. Hoye, *Nat. Protoc.*, 2013, **8**, 501–508; (d) P. M. Tadross and B. M. Stoltz, *Chem. Rev.*, 2012, **112**, 3550–3557; (e) S. Cho and Q. Wang, *Org. Lett.*, 2020, **22**, 1670–1674; (f) J. He, D. Qiu and Y. Li, *Acc. Chem. Res.*, 2020, **53**, 508–519; (g) C. Ni, L. Zhang and J. Hu, *J. Org. Chem.*, 2008, **73**, 5699–5713; (h) D. R. Stuart, *Synlett*, 2016, **28**, 275–279; (i) A. Nilova, B. Metze and D. R. Stuart, *Org. Lett.*, 2021, **23**, 4813–4817; (j) O. Smith, M. J. Hindson, A. Sreenithya, V. Tataru, R. S. Paton, J. W. Burton and M. D. Smith, *Nat. Synth.*, 2024, **3**, 58–66.
- 16 Selected examples: (a) J. Mindner, S. Rombach and D. B. Werz, *Org. Lett.*, 2024, **26**(10), 2124–2128; (b) J.-A. García-López, M. Çetin and M. F. Greaney, *Angew.*

- Chem., Int. Ed.*, 2015, **54**, 2156–2159; (c) S. Cho and Q. Wang, *Org. Lett.*, 2020, **22**, 1670–1674; (d) Y. Zeng, L. Zhang, Y. Zhao, C. Ni, J. Zhao and J. Hu, *J. Am. Chem. Soc.*, 2013, **135**, 2955–2958; (e) Y. Zeng and J. Hu, *Synthesis*, 2016, **48**, 2137–2150; (f) J. L. Henderson, A. S. Edwards and M. F. Greaney, *J. Am. Chem. Soc.*, 2006, **128**, 7426–7427; (g) H. Yoshida, Y. Asatsu, Y. Mimura, Y. Ito, J. Ohshita and K. Takaki, *Angew. Chem., Int. Ed.*, 2011, **50**, 9676–9679; (h) C. E. Hendrick, S. L. McDonald and Q. Wang, *Org. Lett.*, 2013, **13**, 3444–3447; (i) K.-P. Wang, S. Y. Yun, P. Mamidipalli and D. Lee, *Chem. Sci.*, 2013, **4**, 3205–3211; (j) S. S. Bhojgude, T. Roy, R. G. Gonnade and A. T. Biju, *Org. Lett.*, 2016, **18**, 5424–5427; (k) C. Ni, L. Zhang and J. Hu, *J. Org. Chem.*, 2008, **73**, 5699–5713.
- 17 Precedents for the 1,2-dihalogenation of arynes: (a) Y. Zeng, G. Li and J. Hu, *Angew. Chem., Int. Ed.*, 2015, **54**, 10773–10777; (b) D. Niu, T. Wang, B. P. Woods and T. R. Hoye, *Org. Lett.*, 2014, **16**, 254–257; (c) L. Friedman and F. M. Logullo, *Angew. Chem., Int. Ed. Engl.*, 1965, **4**, 239–240; (d) M. A. Birkett, D. W. Knight, P. B. Little and M. B. Mitchell, *Tetrahedron*, 2000, **56**, 1013; (e) D. Rodríguez-Lojo, A. Cobas, D. Peña, D. Pérez and E. Guitán, *Org. Lett.*, 2012, **14**, 1363–1365.
- 18 Prior attempts in our group to functionalize biaryl arynes derived from cyclic diarylbromonium and chloronium species with electrophiles have failed, despite the variety of strategies to do so which are reported with other aryne precursors, notably Kobayashi-type arynes. We suspect that the observed difficulty is attributed to steric hindrance inflicted by the adjacent aryl substituent.