

THE ROLE OF LIPOXIN A4 IN CYSTIC FIBROSIS LUNG DISEASE

Valérie Urbach^{a,b,*}, Gerard Higgins^a, Paul Buchanan^a, Fiona Ringholz^a

Abstract: In Cystic Fibrosis (CF), mutations of the CFTR gene result in defective Cl⁻ secretion and Na⁺ hyperabsorption by epithelia which leads to airway lumen dehydration and mucus plugging and favours chronic bacterial colonization, persistent inflammation and progressive lung destruction. Beyond this general description, the pathogenesis of CF lung disease remains obscure due to an incomplete understanding of normal innate airway defense. This mini-review aims to highlight the role of the pro-resolution lipid mediator, Lipoxin A₄, which is inadequately produced in CF, on several aspects of innate immunity that are altered in CF airway disease.

3RD ANNUAL WORLD CONGRESS OF MOLECULAR & CELL BIOLOGY

Cystic Fibrosis airway disease

Cystic Fibrosis (CF) is a lethal genetic disorder which results from a mutation of the gene coding for the Cystic Fibrosis Transmembrane conductance Regulator (CFTR), a cyclic AMP-dependent Cl⁻ channel [1]. Cystic Fibrosis affects various organs in which the CFTR protein is normally expressed. The major clinical features of CF are chronic pulmonary disease, exocrine pancreatic insufficiency and male infertility, however, the lung disease is the main cause of morbidity and mortality in CF [2-4]. Healthy airways are lined by an epithelial layer that plays a major role in defense against inhaled pathogens involving several specialized epithelial functions including: mechanical barrier, adequate surface hydration due to an intricate regulation of ions and water transport, mucus secretion, production of antimicrobial peptides, expression of receptors that recognize pathogen associated molecular patterns (PAMPs), secretion of cytokines that control the local immune responses in the airway lumen. In CF mutations of the CFTR gene results in defective Cl⁻ secretion and Na⁺ hyperabsorption by airway epithelia [5,6]. This contributes to reduction of the periciliary fluid volume, the airway lumen dehydration, reduction of the periciliary fluid volume and mucus plugging [7]. This results in an impaired mucociliary clearance of pathogens from the lung, favouring chronic bacterial colonization, persistent inflammation and progressive destruction of the lung [8]. In addition to the abnormality of epithelial ion transport, other epithelial dysfunctions have been described in chronically inflamed and infected CF airways, intrinsic pro-inflammatory properties, amplified inflammatory responses to infections and reduced bacterial clearance. However, beyond this general description, the pathogenesis of the CF lung disease remains obscure.

Anti-inflammatory therapy in Cystic Fibrosis

Whilst the field continues to celebrate the success, for a minority of people with CF, in achieving therapeutic benefits via CFTR modulation strategies, reduction of lung inflammation and restoration of airway hydration / muco-ciliary clearance remain core goals of CF therapy for the majority. Several anti-inflammatory approaches have been examined in CF, however, the ideal anti-inflammatory drug is not yet available [9]. A recent systematic review of the risks and benefits of Inhaled corticosteroids (ICS) in CF, examining evidence from 13 trials, concluded that there is insufficient evidence to establish whether ICS are beneficial in CF, but withdrawal in those already taking them has been shown to be safe [10]. It is established that ICS use can have adverse effects on growth. A systematic review of the efficacy of non-steroidal anti-inflammatory drugs in CF concluded that treatment with high-dose ibuprofen was associated with a significantly lower annual rate of decline in lung function (especially in children), however, the adoption of ibuprofen into therapy has not been universally accepted [11,12]. Redressing the imbalance in fatty acid metabolism described in CF, by supplementation of Docosahexaenoic Acid may be helpful, and efforts are ongoing to evaluate the potential therapeutic benefit [13].

The Specialized Pro-resolving Mediators

New perspectives have emerged in inflammation research with the discovery of new classes of biologically active lipid mediators playing specialised roles in the active resolution of inflammation – the “specialized pro-resolving mediators” (*SPM*) [14]. Furthermore, the acute inflammatory response is a protective mechanism that evolved to eliminate invading organisms and yet be self-limited with an active resolution phase designed to restore tissue homeostasis. The resolution phase is carried out by the actions of *SPM* which are non-immunosuppressive [14]. The temporal evolution of acute inflammation toward its active resolution is directed by the sequential expression and activity of characteristic classes of eicosanoid mediator in a process termed “class switching” [15]. Prostaglandins are biosynthesised early, initiating the inflammatory response. Leukotrienes follow, typified by Leukotriene B₄ (LTB₄) which plays its role in amplification and propagation of inflammation [15] acting in concert with the cytokine Interleukin 8 (IL-8) as a potent

^aNational Children’s Research Centre, Crumlin, Dublin 12, Ireland

^bInstitut National de la Santé et de la Recherche Médicale, U845, Faculté de Médecine Paris Descartes - Site Necker - 156 rue Vaugirard 75015, Paris, France

* Corresponding author.

E-mail address: valerie.urbach@nrcr.ie (Valérie Urbach)

neutrophil chemo-attractant [16,17]. Lipoxin A₄ (LXA₄) is the first eicosanoid of the *SPM* family to be expressed in the active resolution phase of inflammation. LXA₄ production is followed by the biosynthesis of Resolvins and Protectins at the inflammatory site. These *SPM* are biosynthesized *in vivo* in inflammatory exudates from essential fatty acids; Lipoxins (LX) from arachidonic acid; E-series resolvins (Rv) from Omega-3 Eicosapentaenoic acid (EPA); D-series resolvins and protectins (PD) from Docosahexaenoic acid (DHA) [14].

Anti-inflammatory properties of Lipoxin A₄

The anti-inflammatory properties of LXA₄ have been reported in a wide variety of tissues. LXA₄ inhibits nuclear factor-kappaB activation, which results in inhibition of pro-inflammatory cytokine release and inhibition of inflammatory responses in microglial cells, astrocytoma cells, macrophages, peripheral blood mononuclear cells (PBMC), polymorphonuclear leukocytes (PMN) and intestinal epithelial cells [18-20]. LXA₄ inhibits neutrophil functions, most notably inhibiting LTB₄ induced neutrophil chemotaxis, neutrophil adherence and transmigration across intestinal epithelium and endothelium and inhibiting superoxide anion and peroxynitrite generation [14] [21-23]. LXA₄ facilitate neutrophil apoptosis [24] and stimulates phagocytosis of apoptotic neutrophils by macrophages [25][26]. This is a critical point since delayed neutrophil apoptosis appears to be a component of the pathophysiology in patients with inflammatory diseases, including cystic fibrosis [27,28] and frequently correlates with disease severity and outcome. In the airways, *in vitro* and *in vivo* studies also report that LXA₄ displays diverse and potent anti-inflammatory actions [19,29,30]. In human airways, LXA₄ suppresses IL-8 production by leukocytes and bronchial epithelial cells [30,31]. LXA₄ was shown to arrest neutrophilic inflammation and decrease infection in a mouse model of chronic airway inflammation and infection [32]. LXA₄ has been proposed as a novel regulators of adaptive immunity and may have therapeutic potential in chronic immune disorders [33]. The pro-resolution properties of LXA₄ are mediated by the ALX/FPR2 receptor. The ALX/FPR2 is a G protein-coupled receptor of seven-transmembrane domains that is expressed mainly by mammalian phagocytic leukocytes. The known effect of ALX/FPR2 receptor in the in host defense and inflammation is not only triggered by LXA₄ since other pro-resolution mediators such as resolving D1 also mediate their effects through this receptor [34] [35].

Abnormal eicosanoid class switching in Cystic Fibrosis

LXA₄ is produced by lipoxygenase (LO) interactions resulting from trans-cellular cooperations [36] of neutrophils [37], eosinophils [38], alveolar macrophages [39], platelets [40] or airway epithelial cells [41], each expressing different Lipoxygenase (LO) enzymes which act in sequence in LXA₄ biosynthesis [16,42]. The 5-LO expressed by neutrophils can utilise the 15(S)-hydroxyeicosatetraenoic acid released by epithelial cells as substrate to synthesize lipoxins [37]. The platelet 12-LO [40], the macrophages or epithelial 15-LO [39] [19] are each able to transform Leukotriene A₄, released by neutrophils, into LXA₄. Up-regulation of 15-LO activity generates an inverse relationship between LTB₄ and LXA₄ biosynthesis. The activity of 15-LO promotes LXA₄ biosynthesis and blocks leukotriene biosynthesis, both as a result of 15-LO products competing for flux at the 5-LO level, and by diversion of the

intermediate Leukotriene A₄ away from LTB₄ towards LXA₄ biosynthesis [15,41,42]. The levels of LXA₄ have been reported to be decreased in chronic airway inflammatory disease such as asthma, chronic obstructive pulmonary disease and CF [29,32,43,44]. A decreased proportion of pro-resolving compounds (LXA₄) compared to pro-inflammatory (LTB₄) is associated with decrease of lung function [45]. The absolute content of LXA₄ concentration in CF BAL fluid from patients with CF is not significantly different from controls [46]. However, a significant suppression in LXA₄/neutrophil ratios in BAL fluid of patients with CF compared with pulmonary inflammatory controls was reported [32,46]. More specifically, *in vitro* studies support a role for CFTR in LXA₄ production. CFTR inhibition reduced LXA₄ synthesis by 50% during platelets/ PMN co-incubation by inhibiting the lipoxin synthase activity of platelets 12-LO. Platelets from patients with CF generated 40% less LXA₄ compared to healthy subject [47]. The decreased LXA₄ production in CF provides a mechanistic explanation of the failure to actively resolve acute airway inflammation seen in these patients.




Figure 1. LXA₄ restores the Airway Surface Liquid (ASL) layer in CF bronchial epithelium. Live cell imaging using confocal microscopy of bronchial epithelium in primary culture from a child with CF before and after treatment with LXA₄. The bronchial epithelial cells are stained in green using *calcein green* and the ASL in red using dextran coupled to *texas-red*.

Lipoxin A₄ regulates ion transport and the airway surface liquid height

LXA₄ stimulates a rapid and transient intracellular Ca²⁺ increase and induces Cl⁻ secretion through human bronchial epithelial cells by Ca²⁺-activated Cl⁻ channel and not CFTR [48]. Furthermore, LXA₄ effects on ion transport lead into an increase of the airway surface liquid (ASL) layer height in models of fully differentiated bronchial epithelia derived from primary culture of bronchial brushings from patients with CF (Figure 1). LXA₄ exerts this effect on the ASL dynamics via the ALX/FPR2 receptor which is expressed in the apical membrane of airway epithelial cells. The sustained increase in ASL height induced by LXA₄ in non-CF and CF bronchial epithelial results from stimulation of an intracellular calcium signal and Ca²⁺-activated Cl⁻ secretion via NPPB sensitive Cl⁻ channels [49]. LXA₄ thus restores Cl⁻ secretion and adequate ASL height which are affected in CF airways, highlighting a role for LXA₄ in the control of innate immune defence. The inadequate endogenous LXA₄ biosynthesis in CF contributes to the reduce ASL volume and impaired mucociliary clearance in addition to alter resolution of inflammation in the airway, thus amplifying the vicious circle of airway dehydration, chronic infection and inflammation.

Lipoxin A₄ regulates airway epithelial integrity

Epithelial repair is a key process required to maintain epithelial barrier integrity and respiratory function, however in CF, repeated

infections and inflammatory insults result in damage to the airways, triggering the repair process [50]. Epithelial repair initially involves cell spreading and migration, followed by proliferation to repopulate the denuded area that has been created by injury [51-53]. This is then followed by differentiation of the epithelium [54]. Recent research suggests that epithelial repair and differentiation of the CF airway epithelium is down-regulated or delayed [55-59]. More specifically, cell migration and proliferation appear to both be reduced during repair in CF bronchial epithelial cells compared to non-CF [60]. This delay in repair of the CF epithelium renders the lung more susceptible to ongoing bacterial infection and thus may lead to more epithelial damage [61]. It was recently reported that LXA₄ can trigger epithelial cell migration and proliferation and thus play a role in repair of corneal and bronchial epithelia [62-64] [65-67]. LXA₄ triggers an increase in migration, proliferation, and wound repair of non-CF and CF bronchial epithelia. These responses to LXA₄ are mediated by the ALX/FPR2 receptor via the downstream activation of K_{ATP} channels and ERK MAP kinase phosphorylation [60]. This effect of LXA₄ both ion transport and repair are consistent with the role of ion channels in two key processes of repair, migration and proliferation [68]. In particular potassium channels have been shown in numerous cell types to be involved in cell migration and proliferation [57,58,69-72]. Furthermore, LXA₄ enhances airway epithelial tight junction formation. LXA₄ stimulates ZO-1, claudin-1 and occludin expression and trafficking at the apical membrane resulting in enhanced transepithelial electrical resistance in human airway epithelia [73]. Therefore the reduced levels of LXA₄ in the CF airways [74] may account for the reduced capacity for epithelial repair in the CF epithelium.




Figure 2. Pleiotropic effects of LXA₄. LXA₄ which is abnormally produced in Cystic Fibrosis controls various airway physiological functions. LXA₄ regulates bronchial epithelium ion transport, enhances the airway surface liquid (ASL) height, protects epithelial barrier integrity and reduces inflammation.

Therapeutic potential for LXA₄ in cystic fibrosis treatment

Clearance of airway secretions has been a first line therapy for patients with CF and a variety of airway clearance therapies have been developed [75,76]. One of the greatest challenges into reversing the CF defect in the airways is to design strategies to overcome the absence of functional CFTR by stimulating chloride secretion via alternative pathways, thus restoring airway hydration and mucociliary clearance. This can be achieved via the activity of calcium-activated

chloride channels stimulated by agents that raise the intracellular calcium concentration. This strategy has been plagued by the attendant side effects of the amplification of the calcium dependant pro-inflammatory response resulting in the undesirable activation of NFκB [26]. Thus identification of agents, particularly natural endogenous biologicals, that stimulate alternative non-CFTR Cl⁻ secretory pathways and promote ASL hydration and optimal ASL height recovery are likely to be of therapeutic benefit in improving mucociliary clearance in patients with CF. The effects of LXA₄ inhalation has been evaluated in a pilot study of eight asthmatic and healthy adult subjects. The challenge was tolerated, had no adverse effect on pulse or blood pressure and demonstrated favourable effects on specific airway conductance [77].

Conclusion and future prospect

Taken together, the discovery of the multiple LXA₄ functions in restoring bronchial epithelium ion transport in enhancing ASL height, in restoring epithelial barrier function and in reducing inflammation might provide significant advance in improving quality of life and longevity for CF patients (Figure 2).

Citation

Urbach V, Higgins G, Buchanan P, Ringholz F (2013) The role of Lipoxin A4 in Cystic Fibrosis Lung Disease. *Computational and Structural Biotechnology Journal*. 6 (7): e201303018. doi: <http://dx.doi.org/10.5936/csbj.201303018>

References

- Riordan JR, Rommens JM, Kerem B, Alon N, Rozmahel R, et al. (1989) Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* 245: 1066-1073.
- Boat TF, Cheng PW (1989) Epithelial cell dysfunction in cystic fibrosis: implications for airways disease. *Acta Paediatr Scand Suppl* 363: 25-29; discussion 29-30.
- Mickle JE, Cutting GR (1998) Clinical implications of cystic fibrosis transmembrane conductance regulator mutations. *Clin Chest Med* 19: 443-458, v.
- Mickle JE, Macek M, Jr., Fulmer-Smentek SB, Egan MM, Schwiebert E, et al. (1998) A mutation in the cystic fibrosis transmembrane conductance regulator gene associated with elevated sweat chloride concentrations in the absence of cystic fibrosis. *Hum Mol Genet* 7: 729-735.
- Kunzelmann K, Kiser GL, Schreiber R, Riordan JR (1997) Inhibition of epithelial Na⁺ currents by intracellular domains of the cystic fibrosis transmembrane conductance regulator. *FEBS Lett* 400: 341-344.
- Stutts MJ, Canessa CM, Olsen JC, Hamrick M, Cohn JA, et al. (1995) CFTR as a cAMP-dependent regulator of sodium channels. *Science* 269: 847-850.
- Davis PB, Drumm M, Konstan MW (1996) Cystic fibrosis. *Am J Respir Crit Care Med* 154: 1229-1256.
- Pillarsetti N, Linnane B, Ranganathan S (2010) Early bronchiectasis in cystic fibrosis detected by surveillance CT. *Respirology* 15: 1009-1011.
- Balfour-Lynn IM (2007) Anti-inflammatory approaches to cystic fibrosis airways disease. *Curr Opin Pulm Med* 13: 522-528.

10. Balfour-Lynn IM, Welch K (2012) Inhaled corticosteroids for cystic fibrosis. *Cochrane Database Syst Rev* 11: CD001915.
11. Lands LC, Stanojevic S (2013) Oral non-steroidal anti-inflammatory drug therapy for lung disease in cystic fibrosis. *Cochrane Database Syst Rev* 6: CD001505.
12. Bush A, Davies J (2007) Non-steroidal anti-inflammatory therapy for inflammatory lung disease in cystic fibrosis (at least at the moment). *J Pediatr* 151: 228-230.
13. Van Biervliet S, Van Biervliet JP, Robberecht E, Christophe A (2005) Docosahexaenoic acid trials in cystic fibrosis: a review of the rationale behind the clinical trials. *J Cyst Fibros* 4: 27-34.
14. Serhan CN, Chiang N, Van Dyke TE (2008) Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol* 8: 349-361.
15. Levy BD, Clish CB, Schmidt B, Gronert K, Serhan CN (2001) Lipid mediator class switching during acute inflammation: signals in resolution. *Nat Immunol* 2: 612-619.
16. Haeggstrom JZ, Funk CD (2011) Lipoxygenase and leukotriene pathways: biochemistry, biology, and roles in disease. *Chem Rev* 111: 5866-5898.
17. Mukaida N, Okamoto S, Ishikawa Y, Matsushima K (1994) Molecular mechanism of interleukin-8 gene expression. *J Leukoc Biol* 56: 554-558.
18. Wu SH, Liu B, Dong L, Wu HJ (2010) NF-kappaB is involved in inhibition of lipoxin A4 on dermal inflammation and hyperplasia induced by mezerein. *Exp Dermatol* 19: e286-288.
19. Vachier I, Chanez P, Bonnans C, Godard P, Bousquet J, et al. (2002) Endogenous anti-inflammatory mediators from arachidonate in human neutrophils. *Biochem Biophys Res Commun* 290: 219-224.
20. Gewirtz AT, McCormick B, Neish AS, Petasis NA, Gronert K, et al. (1998) Pathogen-induced chemokine secretion from model intestinal epithelium is inhibited by lipoxin A4 analogs. *J Clin Invest* 101: 1860-1869.
21. Bonnans C, Fukunaga K, Levy MA, Levy BD (2006) Lipoxin A(4) regulates bronchial epithelial cell responses to acid injury. *Am J Pathol* 168: 1064-1072.
22. Colgan SP, Serhan CN, Parkos CA, Delp-Archer C, Madara JL (1993) Lipoxin A4 modulates transmigration of human neutrophils across intestinal epithelial monolayers. *J Clin Invest* 92: 75-82.
23. Takano T, Clish CB, Gronert K, Petasis N, Serhan CN (1998) Neutrophil-mediated changes in vascular permeability are inhibited by topical application of aspirin-triggered 15-epi-lipoxin A4 and novel lipoxin B4 stable analogues. *J Clin Invest* 101: 819-826.
24. El Kebir D, József L, Khreiss T, Pan W, Petasis NA, et al. (2007) Aspirin-triggered lipoxins override the apoptosis-delaying action of serum amyloid A in human neutrophils: a novel mechanism for resolution of inflammation. *J Immunol* 179: 616-622.
25. Godson C, Mitchell S, Harvey K, Petasis NA, Hogg N, et al. (2000) Cutting edge: lipoxins rapidly stimulate nonphlogistic phagocytosis of apoptotic neutrophils by monocyte-derived macrophages. *J Immunol* 164: 1663-1667.
26. El Kebir D, Filep JG (2010) Role of neutrophil apoptosis in the resolution of inflammation. *ScientificWorldJournal* 10: 1731-1748.
27. Soleti R, Porro C, Martínez MC (2013) Apoptotic process in cystic fibrosis cells. *Apoptosis* 18: 1029-1038.
28. McKeon DJ, Condliffe AM, Cowburn AS, Cadwallader KC, Farahi N, et al. (2008) Prolonged survival of neutrophils from patients with Delta F508 CFTR mutations. *Thorax* 63: 660-661.
29. Bonnans C, Vachier I, Chavis C, Godard P, Bousquet J, et al. (2002) Lipoxins are potential endogenous antiinflammatory mediators in asthma. *Am J Respir Crit Care Med* 165: 1531-1535.
30. Bonnans C, Gras D, Chavis C, Mainprice B, Vachier I, et al. (2007) Synthesis and anti-inflammatory effect of lipoxins in human airway epithelial cells. *Biomed Pharmacother* 61: 261-267.
31. Jozsef L, Zouki C, Petasis NA, Serhan CN, Filep JG (2002) Lipoxin A4 and aspirin-triggered 15-epi-lipoxin A4 inhibit peroxynitrite formation, NF-kappa B and AP-1 activation, and IL-8 gene expression in human leukocytes. *Proc Natl Acad Sci U S A* 99: 13266-13271.
32. Karp CL, Flick LM, Park KW, Softic S, Greer TM, et al. (2004) Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway. *Nat Immunol* 5: 388-392.
33. Parkinson JF (2006) Lipoxin and synthetic lipoxin analogs: an overview of anti-inflammatory functions and new concepts in immunomodulation. *Inflamm Allergy Drug Targets* 5: 91-106.
34. Brink C, Dahlén SE, Drazen J, Evans JF, Hay DW, et al. (2003) International Union of Pharmacology XXXVII. Nomenclature for leukotriene and lipoxin receptors. *Pharmacol Rev* 55: 195-227.
35. Krishnamoorthy S, Recchiuti A, Chiang N, Yacoubian S, Lee CH, et al. (2010) Resolvin D1 binds human phagocytes with evidence for proresolving receptors. *Proc Natl Acad Sci U S A* 107: 1660-1665.
36. Fierro IM, Serhan CN (2001) Mechanisms in anti-inflammation and resolution: the role of lipoxins and aspirin-triggered lipoxins. *Braz J Med Biol Res* 34: 555-566.
37. Chavis C, Vachier I, Chanez P, Bousquet J, Godard P (1996) 5(S),15(S)-dihydroxyicosatetraenoic acid and lipoxin generation in human polymorphonuclear cells: dual specificity of 5-lipoxygenase towards endogenous and exogenous precursors. *J Exp Med* 183: 1633-1643.
38. Serhan CN, Hirsch U, Palmblad J, Samuelsson B (1987) Formation of lipoxin A by granulocytes from eosinophilic donors. *FEBS Lett* 217: 242-246.
39. Levy BD, Romano M, Chapman HA, Reilly JJ, Drazen J, et al. (1993) Human alveolar macrophages have 15-lipoxygenase and generate 15(S)-hydroxy-5,8,11-cis-13-trans-icosatetraenoic acid and lipoxins. *J Clin Invest* 92: 1572-1579.
40. Serhan CN, Sheppard KA (1990) Lipoxin formation during human neutrophil-platelet interactions. Evidence for the transformation of leukotriene A4 by platelet 12-lipoxygenase in vitro. *J Clin Invest* 85: 772-780.
41. Claria J, Lee MH, Serhan CN (1996) Aspirin-triggered lipoxins (15-epi-LX) are generated by the human lung adenocarcinoma cell line (A549)-neutrophil interactions and are potent inhibitors of cell proliferation. *Mol Med* 2: 583-596.
42. Serhan CN., guest editor (2005) Special Issue: The lipoxins and the aspirin-triggered lipoxins. *Prostaglandins Leukot Essent Fatty Acids* 73: 139-321.
43. Planaguma A, Kazani S, Marigowda G, Haworth O, Mariani TJ, et al. (2008) Airway lipoxin A4 generation and lipoxin A4 receptor expression are decreased in severe asthma. *Am J Respir Crit Care Med* 178: 574-582. Epub 2008 Jun 2026.
44. Balode L, Strazda G, Jurka N, Kopeika U, Kislina A, et al. (2012) Lipoxygenase-derived arachidonic acid metabolites in chronic obstructive pulmonary disease. *Medicina (Kaunas)* 48: 292-298.
45. Carlo T, Kalwa H, Levy BD (2013) 15-Epi-lipoxin A4 inhibits human neutrophil superoxide anion generation by regulating polyisoprenyl diphosphate phosphatase 1. *FASEB J*.
46. Starosta V, Ratjen F, Rietschel E, Paul K, Griesse M (2006) Anti-inflammatory cytokines in cystic fibrosis lung disease. *Eur Respir J* 28: 581-587.
47. Mattoscio D, Evangelista V, De Cristofaro R, Recchiuti A, Pandolfi A, et al. (2010) Cystic fibrosis transmembrane conductance regulator

- (CFTR) expression in human platelets: impact on mediators and mechanisms of the inflammatory response. *FASEB J* 24: 3970-3980.
48. Bonnans C, Mainprice B, Chanez P, Bousquet J, Urbach V (2003) Lipoxin A4 stimulates a cytosolic Ca²⁺ increase in human bronchial epithelium. *J Biol Chem* 278: 10879-10884.
 49. Verriere V, Higgins G, Al-Alawi M, Costello RW, McNally P, et al. (2012) Lipoxin A4 stimulates calcium-activated chloride currents and increases airway surface liquid height in normal and cystic fibrosis airway epithelia. *PLoS ONE* 7: e37746.
 50. Regamey N, Jeffery PK, Alton EW, Bush A, Davies JC (2011) Airway remodelling and its relationship to inflammation in cystic fibrosis. *Thorax* 66: 624-629.
 51. Stripp BR, Reynolds SD (2008) Maintenance and repair of the bronchiolar epithelium. *Proc Am Thorac Soc* 5: 328-333.
 52. Zahm JM, Chevillard M, Puchelle E (1991) Wound repair of human surface respiratory epithelium. *Am J Respir Cell Mol Biol* 5: 242-248.
 53. Zahm JM, Kaplan H, Hérard AL, Doriot F, Pierrot D, et al. (1997) Cell migration and proliferation during the in vitro wound repair of the respiratory epithelium. *Cell Motil Cytoskeleton* 37: 33-43.
 54. Erjefält JS, Erjefält I, Sundler F, Persson CG (1995) In vivo restitution of airway epithelium. *Cell Tissue Res* 281: 305-316.
 55. Trinh NT, Bardou O, Privé A, Maillé E, Adam D, et al. (2012) Improvement of defective cystic fibrosis airway epithelial wound repair after CFTR rescue. *Eur Respir J*.
 56. Maillé E, Trinh NT, Privé A, Bilodeau C, Bissonnette E, et al. (2011) Regulation of normal and cystic fibrosis airway epithelial repair processes by TNF- α after injury. *Am J Physiol Lung Cell Mol Physiol* 301: L945-955.
 57. Trinh NT, Privé A, Maillé E, Noël J, Brochiero E (2008) EGF and K⁺ channel activity control normal and cystic fibrosis bronchial epithelia repair. *Am J Physiol Lung Cell Mol Physiol* 295: L866-880.
 58. Trinh NT, Privé A, Kheir L, Bourret JC, Hijazi T, et al. (2007) Involvement of KATP and KvLQT1 K⁺ channels in EGF-stimulated alveolar epithelial cell repair processes. *Am J Physiol Lung Cell Mol Physiol* 293: L870-882.
 59. Hajj R, Lesimple P, Nawrocki-Raby B, Birembaut P, Puchelle E, et al. (2007) Human airway surface epithelial regeneration is delayed and abnormal in cystic fibrosis. *J Pathol* 211: 340-350.
 60. Buchanan PJ, McNally P, Harvey BJ, Urbach V (2013) Lipoxin A4-mediated KATP potassium channel activation results in cystic fibrosis airway epithelial repair. *Am J Physiol Lung Cell Mol Physiol* 305: L193-201.
 61. Roger P, Puchelle E, Bajolet-Laudinat O, Tournier JM, Debordeaux C, et al. (1999) Fibronectin and α 5 β 1 integrin mediate binding of *Pseudomonas aeruginosa* to repairing airway epithelium. *Eur Respir J* 13: 1301-1309.
 62. Bonnans C, Fukunaga K, Levy MA, Levy BD (2006) Lipoxin A(4) regulates bronchial epithelial cell responses to acid injury. *Am J Pathol* 168: 1064-1072.
 63. Gronert K (2005) Lipoxins in the eye and their role in wound healing. *Prostaglandins Leukot Essent Fatty Acids* 73: 221-229.
 64. Kenchegowda S, Bazan NG, Bazan HE (2011) EGF stimulates lipoxin A4 synthesis and modulates repair in corneal epithelial cells through ERK and p38 activation. *Invest Ophthalmol Vis Sci* 52: 2240-2249.
 65. Wang SB, Hu KM, Seamon KJ, Mani V, Chen Y, et al. (2012) Estrogen negatively regulates epithelial wound healing and protective lipid mediator circuits in the cornea. *FASEB J* 26: 1506-1516.
 66. Kenchegowda S, Bazan NG, Bazan HE (2011) EGF stimulates lipoxin A4 synthesis and modulates repair in corneal epithelial cells through ERK and p38 activation. *Invest Ophthalmol Vis Sci* 52: 2240-2249.
 67. Gronert K, Maheshwari N, Khan N, Hassan IR, Dunn M, et al. (2005) A role for the mouse 12/15-lipoxygenase pathway in promoting epithelial wound healing and host defense. *J Biol Chem* 280: 15267-15278.
 68. Kirk KL (2010) CFTR channels and wound healing. Focus on "Cystic fibrosis transmembrane conductance regulator is involved in airway epithelial wound repair". *Am J Physiol Cell Physiol* 299: C888-890.
 69. Rao JN, Platoshyn O, Li L, Guo X, Golovina VA, et al. (2002) Activation of K(+) channels and increased migration of differentiated intestinal epithelial cells after wounding. *Am J Physiol Cell Physiol* 282: C885-898.
 70. Pardo LA (2004) Voltage-gated potassium channels in cell proliferation. *Physiology (Bethesda)* 19: 285-292.
 71. Kessler W, Budde T, Gekle M, Fabian A, Schwab A (2008) Activation of cell migration with fibroblast growth factor-2 requires calcium-sensitive potassium channels. *Pflugers Arch* 456: 813-823.
 72. Lotz MM, Wang H, Song JC, Pories SE, Matthews JB (2004) K⁺ channel inhibition accelerates intestinal epithelial cell wound healing. *Wound Repair Regen* 12: 565-574.
 73. Grumbach Y, Quynh NV, Chiron R, Urbach V (2009) LXA4 stimulates ZO-1 expression and transepithelial electrical resistance in human airway epithelial (16HBE14o-) cells. *Am J Physiol Lung Cell Mol Physiol* 296: L101-108.
 74. Karp CL, Flick LM, Park KW, Softic S, Greer TM, et al. (2004) Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway. *Nat Immunol* 5: 388-392.
 75. Pisi G, Chetta A (2009) Airway clearance therapy in cystic fibrosis patients. *Acta Biomed* 80: 102-106.
 76. Tarran R, Grubb BR, Parsons D, Picher M, Hirsh AJ, et al. (2001) The CF salt controversy: in vivo observations and therapeutic approaches. *Mol Cell* 8: 149-158.
 77. Christie PE, Spur BW, Lee TH (1992) The effects of lipoxin A4 on airway responses in asthmatic subjects. *Am Rev Respir Dis* 145: 1281-1284.

Competing Interests:

The authors have declared that no competing interests exist.



© 2013 Urbach et al.

Licensee: Computational and Structural Biotechnology Journal.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are properly cited.

What is the advantage to you of publishing in *Computational and Structural Biotechnology Journal (CSBJ)* ?

- ✚ Easy 5 step online submission system & online manuscript tracking
- ✚ Fastest turnaround time with thorough peer review
- ✚ Inclusion in scholarly databases
- ✚ Low Article Processing Charges
- ✚ Author Copyright
- ✚ Open access, available to anyone in the world to download for free

WWW.CSBJ.ORG