



Circular RNA as a prospective molecular tool for the study of neuroprotection in cerebral ischemia

Svetlana A. Limborska, Ivan B. Filippenkov

Institute of Molecular Genetics, Russian Academy of Sciences, Moscow, Russia

Correspondence to: Svetlana A. Limborska. Institute of Molecular Genetics, Russian Academy of Sciences, Kurchatov Sq. 2, Moscow, Russia.

Email: limbor@img.ras.ru.

Comment on: Han B, Zhang Y, Zhang Y, *et al.* Novel insight into circular RNA HECTD1 in astrocyte activation via autophagy by targeting MIR142-TIPARP: implications for cerebral ischemic stroke. *Autophagy* 2018;14:1164-84.

Submitted Oct 30, 2018. Accepted for publication Nov 30, 2018.

doi: 10.21037/tcr.2018.12.03

View this article at: <http://dx.doi.org/10.21037/tcr.2018.12.03>

We recently read the article by Han *et al.* [2018]: “Novel insight into circular RNA *HECTD1* in astrocyte activation via autophagy by targeting *MIR142-TIPARP*: implications for cerebral ischemic stroke” published in *Autophagy* (1). This study reported that circular RNA of *HECTD1* gene (*circHECTD1*) is upregulated in the blood of patients with acute ischemic stroke (AIS) and in animal and cell line ischemia model conditions. Simultaneously, knockdown of *circHECTD1* reduced infarct areas and attenuated neuronal deficits and astrocyte activation in a mouse model of transient middle cerebral artery occlusion (tMCAO). Han *et al.* [2018] showed that *circHECTD1* functions as an endogenous microRNA sponge to inhibit *MIR142* activity, resulting in the inhibition of TCDD-inducible poly[ADP-ribose] polymerase (*TIPARP*) gene expression with subsequent inhibition of astrocyte activation via autophagy. In conclusion, the authors hypothesized that *circHECTD1* can serve as a novel biomarker and therapeutic target for stroke (1).

Here, we want to emphasize the significance of the investigation by Han *et al.* [2018] as an important step towards the development of new approaches in understanding the pathogenesis of stroke. We also discuss the role of circular RNAs in the diagnosis and treatment of cerebral ischemia and comment on a number of research articles that preceded Han *et al.* [2018] in this field of study.

Undoubtedly, the problem of vascular diseases of the brain and, in particular, ischemic stroke, retains its medical and social significance due to the high rates of morbidity, disability and mortality of this disease in the world (2). To date, much attention has been paid to the study of specific

types of RNAs, as markers in diagnosis and as targets in the treatment of ischemic stroke. It is well known that cerebral ischemia causes a cascade of biochemical and transcriptome changes in brain tissues (3). It has been shown that reperfusion after ischemia causes additional damage in brain cells, including the destruction of the endothelial cells of the microvascular brain, disturbance of the blood-brain barrier, accumulation of excess oxygen radicals and activation of apoptosis (4,5). Recently, in a rat model of tMCAO, we revealed the activation of a large number of genes involved in inflammation, the immune response, apoptosis and the stress response (6). Simultaneously, a massive downregulation of genes that ensure the functioning of neurotransmitter systems was observed in tMCAO conditions.

To date, it has been shown that not only messenger RNA (mRNA) but also various types of regulatory RNA are involved in the response to the pathological effects (7-10). Of particular interest are circular RNAs (circRNAs), which have a covalently closed structure and are often formed in protein-coding genes during back-splicing (11,12). circRNAs demonstrate an increased resistance to the action of exonucleases and a predominantly brain-specific expression pattern (13-15). However, the functional significance of circRNA is not well understood. Currently, great attention is being paid to the study of the functioning of circRNAs as microRNA (miRNA) sponges. CircRNA acting as competitive endogenous RNA (ceRNA) competes with mRNA for binding to miRNA and diminishing the effect of miRNA on transcriptional and post-transcriptional levels of regulation of gene expression (16,17). The

Table 1 Circular RNAs associated with ischemia conditions

Circular RNA	Ischemia conditions	Confirmation of specific expression	Confirmation of miRNA-circRNA interaction	Neuroprotective effect	Reference
<i>circHECTD1</i>	tMCAO in mice, plasma of AIS patients, human glioblastoma A172 cell line treated with OGD-R	Microarrays and real-time RT-PCR	Luciferase activity assays	Knockdown are associated with neuroprotection	Han <i>et al.</i> [2018]
<i>circDLGAP4</i>	tMCAO in mice, plasma of AIS patients, mouse brain endothelial bEnd.3 cells treated with OGD-R	Real-time RT-PCR	Luciferase activity assays, miRNA pull-down and FISH assays	Upregulation are associated with neuroprotection	Bai <i>et al.</i> [2018]
<i>circ_016128, circ_007362, circ_006839, circ_000113, circ_002664, circ_008018, circ_011381, circ_015350, circ_006696, circ_001729, circ_000741, circ_016423, circ_009396, circ_017370, circ_010383, circ_016289</i>	tMCAO in mice	Microarrays and real-time RT-PCR	Computational prediction	–	Mehta <i>et al.</i> [2017]
<i>mmu_circRNA_40001, mmu_circRNA_013120, mmu_circRNA_40806</i>	tMCAO in mice	Microarrays and real-time RT-PCR	Computational prediction	–	Liu <i>et al.</i> [2017]
<i>mmu-circRNA-015947</i>	HT22 hippocampal cell culture treated with OGD-R	Microarrays and real-time RT-PCR	Computational prediction	–	Lin <i>et al.</i> [2016]

tMCAO, transient middle cerebral artery occlusion.

functions of several circRNAs as miRNA sponges were investigated in various pathologies. In particular, the role of circRNA CIRS-7 in preventing neuropsychiatric disorders in mice, associated with its functioning as a ceRNA, was recently established (17). In addition, in Alzheimer's disease (18) and various types of cancer (19-21), results showing that circRNA-miRNA-mRNA competition may be associated with pathogenesis regulation were obtained.

In the last 2 years, studies have investigated the transcriptional profile and functional properties of circRNAs under conditions simulating cerebral ischemia (see *Table 1*). Recently, in HT22 hippocampal cell culture under conditions of oxygen glucose deprivation-reoxygenation (OGD-R), which simulated damage after cerebral ischemia and reperfusion, Lin *et al.* [2016] showed that circRNAs may function as miRNA sponges (22). In the OGD-R model, it was shown that circRNA expression is associated with the metabolic pathways of apoptosis and immune response. In tMCAO model conditions, Mehta *et al.* [2017] showed that the differentially expressed circRNAs might be controlled by a set of transcription factors in stroke.

The bioinformatics analysis of biological and molecular functions showed that circRNAs altered their expression after tMCAO controlled biological regulation, the metabolic process, cell communication, and binding to proteins, ions and nucleic acids (23). Simultaneously, it was shown that 16 circRNAs contain binding sites for many miRNAs. In addition, in the tMCAO model conditions in mice, Liu *et al.* [2017] (24) detected the changes in the expression of more than one thousand circRNAs using microarray analysis. These circRNAs were associated with signalling pathways regulating the processes of cell survival and death. Moreover, Liu *et al.* [2017] predicted possible interactions between circRNAs and miRNAs, which can provide potential information for revealing the mechanisms of brain damage in ischemic stroke. Thus, in the studies by Lin *et al.* [2016], Liu *et al.* [2017] and Mehta *et al.* [2017], the transcriptional profile of circRNAs was studied in detail under the conditions of the individual ischemia model. CircRNAs that most significantly altered the expression during ischemia were identified. However, the association with metabolic pathways and biological processes in the cell, as well as with miRNA interaction, was

mainly predictive and computational. Thus, further studies of circRNAs functioning as miRNA sponges in ischemia conditions are needed for specific examples.

A recent study published new important information on the functioning of circRNAs in ischemia. Bai *et al.* [2018] showed that the circRNA of *DLGAP4* gene (*circDLGAP4*) functions as an miRNA sponge to inhibit MIR143 activity, resulting in the inhibition of homologous to the E6-AP C-terminal domain E3 ubiquitin protein ligase 1 expression (25). In contrast, *circDLGAP4* levels were significantly decreased in the plasma of AIS patients and in a mouse stroke model. It has been shown that upregulation of *circDLGAP4* expression significantly attenuated neurological deficits, and decreased infarct areas and blood-brain barrier damage in the tMCAO mouse stroke model. From our perspective, several important conclusions can be immediately drawn from this study. First, blood is widely used in human diagnostic procedures, and the confirmation that transcription pattern in peripheral blood cells can reflect what occurs in the brain is of great significance. Second, the study clearly indicates that the regulation of the regenerative mechanisms of brain cells during ischemia occurs with the participation of circRNA, and in this case, is associated with upregulation of *circDLGAP4* expression. Third, the study by Bai *et al.* emphasizes the role of circRNA-miRNA-mRNA interactions in ensuring the regulation of pathogenesis and regeneration in ischemia.

In studying the functioning of circRNAs as miRNA sponges in cerebral ischemia, we next considered the work by Han *et al.* [2018] on novel insights into *circHECTD1* functioning in cerebral ischemia (1). In this paper, the authors hypothesized that upregulated *circHECTD1* directly binds to miRNA MIR142 and acts as an endogenous sponge to inhibit MIR142 activity, which results in astrocyte activation and thus contributes to cerebral infarction. First, the authors convincingly showed that *circHECTD1* is upregulated in mouse brain in tMCAO model conditions and in human glioblastoma A172 cell line treated with OGD-R. Han *et al.* [2018] clearly demonstrated that the expression pattern of *circHECTD1* in the blood of patients with AIS and in ischemia model conditions is similar. Second, they showed that the knockdown of *circHECTD1* was associated with the reduction of brain infarction in tMCAO mice. Consequently, *circHECTD1* participates in the regulation of the injury and regenerative mechanisms of brain cells in conditions of ischemia. Third, it was shown that in interaction with MIR142, which negatively affects the level of mRNA of *TIPARP* gene, *circHECTD1* diminished miRNA activity. As a result, *circHECTD1*-MIR142-*TIPARP* competition leads to modulation of astrocyte

activation via autophagy in cerebral ischemia conditions.

In conclusion, we want to emphasize the significance of the studies by Han *et al.* [2018] and Bai *et al.* [2018] as important steps towards understanding the role of circRNAs in cerebral ischemia. It should be noted that a considerable number of recent studies show the complex nature of the regulation of pathogenesis and regeneration processes in ischemia, including the functioning of circRNAs. Han *et al.* [2018] and Bai *et al.* [2018] conclude that *circHECTD1* or *circDLGAP4* can serve as a novel biomarker and therapeutic target for stroke (1,25). We are optimistic about such conclusions because their work contributes to stimulating further research. Subsequent analysis of circRNA-miRNA-mRNA interactions is an important step in fully elucidating the mechanisms of damage and regeneration in cerebral ischemia and in determining strategies for achieving a neuroprotective effect in the conditions of this disease.

Acknowledgments

Funding: This work was supported by the program of the Presidium of the Russian Academy of Sciences “Fundamental researches for biomedical technologies”.

Footnote

Provenance and Peer Review: This article was commissioned and reviewed by the Section Editor Chunlin Ou (Cancer Research Institute of Central South University, Changsha, China).

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr.2018.12.03>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the

formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Han B, Zhang Y, Zhang Y, et al. Novel insight into circular RNA HECTD1 in astrocyte activation via autophagy by targeting MIR142-TIPARP: implications for cerebral ischemic stroke. *Autophagy* 2018;14:1164-84.
- Mukherjee D, Patil CG. Epidemiology and the global burden of stroke. *World Neurosurg* 2011;76:S85-90.
- Wang C, Liu M, Pan Y, et al. Global gene expression profile of cerebral ischemia-reperfusion injury in rat MCAO model. *Oncotarget* 2017;8:74607-22.
- Rosenberg GA, Estrada EY, Dencoff JE. Matrix metalloproteinases and TIMPs are associated with blood-brain barrier opening after reperfusion in rat brain. *Stroke* 1998;29:2189-95.
- Lochhead JJ, McCaffrey G, Quigley CE, et al. Oxidative stress increases blood-brain barrier permeability and induces alterations in occludin during hypoxia-reoxygenation. *J Cereb Blood Flow Metab* 2010;30:1625-36.
- Dergunova LV, Filippenkov IB, Stavchansky VV, et al. Genome-wide transcriptome analysis using RNA-Seq reveals a large number of differentially expressed genes in a transient MCAO rat model. *BMC Genomics* 2018;19:655.
- Yang P, Qiu Z, Jiang Y, et al. Silencing of cZNF292 circular RNA suppresses human glioma tube formation via the Wnt/ β -catenin signaling pathway. *Oncotarget* 2016;7:63449-55.
- Li Y, Dong Y, Huang Z, et al. Computational identifying and characterizing circular RNAs and their associated genes in hepatocellular carcinoma. *PLoS One* 2017;12:e0174436.
- Xia W, Qiu M, Chen R, et al. Circular RNA hsa_circ_0067934 is upregulated in esophageal squamous cell carcinoma and promoted proliferation. *Sci Rep* 2016;6:35576.
- Xu Y, Yao Y, Zhong X, et al. Downregulated circular RNA hsa_circ_0001649 regulates proliferation, migration and invasion in cholangiocarcinoma cells. *Biochem Biophys Res Commun* 2018;496:455-61.
- Lasda E, Parker R. Circular RNAs: diversity of form and function. *RNA* 2014;20:1829-42.
- Filippenkov IB, Sudarkina OY, Limborska SA, et al. Multi-step splicing of sphingomyelin synthase linear and circular RNAs. *Gene* 2018;654:14-22.
- Chen BJ, Yang B, Janitz M. Region-specific expression of circular RNAs in the mouse brain. *Neurosci Lett* 2018;666:44-7.
- Rybak-Wolf A, Stottmeister C, Glažar P, et al. Circular RNAs in the Mammalian Brain Are Highly Abundant, Conserved, and Dynamically Expressed. *Mol Cell* 2015;58:870-85.
- Filippenkov IB, Sudarkina OY, Limborska SA, et al. Circular RNA of the human sphingomyelin synthase 1 gene: Multiple splice variants, evolutionary conservatism and expression in different tissues. *RNA Biol* 2015;12:1030-42.
- Memczak S, Jens M, Elefsinioti A, et al. Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature* 2013;495:333-8.
- Piwecka M, Glažar P, Hernandez-Miranda LR, et al. Loss of a mammalian circular RNA locus causes miRNA deregulation and affects brain function. *Science* 2017;357:eaam8526.
- Akhter R. Circular RNA and Alzheimer's Disease. In: Xiao J, editor. *Circular RNAs. Advances in Experimental Medicine and Biology*, vol 1087. Singapore: Springer, 2018:239-43.
- Li F, Zhang L, Li W, et al. Circular RNA ITCH has inhibitory effect on ESCC by suppressing the Wnt/ β -catenin pathway. *Oncotarget* 2015;6:6001-13.
- Liu J, Kong F, Lou S, et al. Global identification of circular RNAs in chronic myeloid leukemia reveals hsa_circ_0080145 regulates cell proliferation by sponging miR-29b. *Biochem Biophys Res Commun* 2018;504:660-5.
- Chen B, Luo M, Liang J, et al. Surface modification of PGP for a neutrophil-nanoparticle co-vehicle to enhance the antidepressant effect of baicalein. *Acta Pharm Sin B* 2018;8:64-73.
- Lin SP, Ye S, Long Y, et al. Circular RNA expression alterations are involved in OGD/R-induced neuron injury. *Biochem Biophys Res Commun* 2016;471:52-6.
- Mehta SL, Pandi G, Vemuganti R. Circular RNA Expression Profiles Alter Significantly in Mouse Brain After Transient Focal Ischemia. *Stroke* 2017;48:2541-8.
- Liu C, Zhang C, Yang J, et al. Screening circular RNA expression patterns following focal cerebral ischemia in mice. *Oncotarget* 2017;8:86535-47.
- Bai Y, Zhang Y, Han B, et al. Circular RNA DLGAP4 Ameliorates Ischemic Stroke Outcomes by Targeting miR-143 to Regulate Endothelial-Mesenchymal Transition Associated with Blood-Brain Barrier Integrity. *J Neurosci* 2018;38:32-50.

Cite this article as: Limborska SA, Filippenkov IB. Circular RNA as a prospective molecular tool for the study of neuroprotection in cerebral ischemia. *Transl Cancer Res* 2019;8(Suppl 2):S126-S129. doi: 10.21037/tcr.2018.12.03