



REVIEW

Edible Insects as a Protein Source: A Review of Public Perception, Processing Technology, and Research Trends

Tae-Kyung Kim^{1,†}, Hae In Yong^{1,†}, Young-Boong Kim¹, Hyun-Wook Kim², and Yun-Sang Choi^{1,*}

¹Research Group of Food Processing, Korea Food Research Institute, Wanju 55365, Korea

²Department of Animal Science & Biotechnology, Gyeongnam National University of Science and Technology, Jinju 52725, Korea

OPEN ACCESS

Received May 17, 2019

Revised July 8, 2019

Accepted July 10, 2019

*Corresponding author : Yun-Sang Choi
Research Group of Food Processing, Korea
Food Research Institute, Wanju 55365, Korea
Tel: +82-63-219-9387
Fax: +82-63-219-9076
E-mail: keys0517@kfri.re.kr

*ORCID
Tae-Kyung Kim
<https://orcid.org/0000-0002-6349-4314>
Hae In Yong
<https://orcid.org/0000-0003-0970-4496>
Young-Boong Kim
<https://orcid.org/0000-0001-8441-198X>
Hyun-Wook Kim
<https://orcid.org/0000-0002-4397-9664>
Yun-Sang Choi
<https://orcid.org/0000-0001-8060-6237>

† These authors contributed equally to this work.

Abstract This review summarizes the current trends related to insect as food resources among consumers, industry, and academia. In Western societies, edible insects have a greater potential as animal feed than as human food because of cultural biases associated with harmful insects, although the abundant characteristics of edible insects should benefit human health. Nevertheless, many countries in Asia, Oceania, Africa, and Latin America utilize insects as a major protein source. Using insects can potentially solve problems related to the conventional food-supply chain, including global water, land, and energy deficits. Academic, industry, and government-led efforts have attempted to reduce negative perceptions of insects through developing palatable processing methods, as well as providing descriptions of health benefits and explaining the necessity of reducing reliance on other food sources. Our overview reveals that entomophagy is experiencing a steady increase worldwide, despite its unfamiliarity to the consumers influenced by Western eating habits.

Keywords edible insects, animal feed, food resources, entomophagy

Introduction

Several projections have suggested that world population will reach over 9 billion by 2050 (Grafton et al., 2015; Park and Yun, 2018). This increase in population requires approximately double the current food production (Belluco et al., 2013). Alarmingly, global warming is gradually decreasing the areas used for food production worldwide (Dobermann et al., 2017). The climate change and the environmental destruction from industrial development also negatively affect the food productivity (van Huis and Oonincx, 2017). In light of worsening resource shortage, several foods have been proposed as alternatives, with insects receiving the most attention (Patel et al., 2019).

Insects are institutionally accepted as a food in many regions and historically consumed (Murefu et al., 2019), providing sufficient nutritional value for humans (Zielinska et al., 2018). However, the rapid increase in food production through technological advancement has largely eliminated insects from our diets (Gao et al., 2018).

The reappearance of insects as a viable food group can be attributed to their nutritional, environmental, and economic value (Nongonierma and FitzGerald, 2017). The increased scrutiny of edible insects is part of a multifaceted strategies for achieving global food security (van Huis, 2015). In general, insects have high protein content and excellent production efficiency compared with other conventional food groups (Kohler et al., 2019; Nongonierma and FitzGerald, 2017; Zielinska et al., 2015). This characteristic is particularly valuable given that future protein consumption is expected to increase, but food supply declines (Gao et al., 2018; Kohler et al., 2019; Patel et al., 2019). Currently, Europe and the United States have the fastest growing edible-insect industry, where a trend is associated with high meat intake (Mlcek et al., 2014). Moreover, it is recognized that a steady increase in the global market size of the insect industry, with applications reaching beyond food into material and drug development (van Thielen et al., 2019).

Despite the numerous practical advantages, many barriers remain in insect-food development because the concept of insect foods is not similar with conventional Western eating habit (Murefu et al., 2019). Currently, insect foods are in a transitional stage, and a promising new technique involves developing new food products via combining protein-processing technology with insects (Kohler et al., 2019; Patel et al., 2019). This review aims to provide a much needed overview of current research on insect-food development. As we will demonstrate, the insect industry is on track to become a successful protein resource that will lead the global market.

Industrial Applications of Insects as Food and Feed Sources

Food industry

For hundreds of years, native cultures in Asia, South America, Africa, and Europe include the consumption of various species of insects (Dobermann et al., 2017; Raheem et al., 2018). Approximately 2,000 insect species are consumed in at least 113 countries (Yen, 2009). A survey of markets in Bangkok, Thailand, for instance, identified 164 insect species being sold for food (Yhoun-aree, 2010). The most commonly eaten insects are beetles, caterpillars, bees, ants, crickets, grasshoppers, and locusts (Raheem et al., 2018). In Zimbabwe, Zambia, and Nigeria, edible insects are commonly available in school cafeterias and open markets, forming a profitable business (Mutungi et al., 2019). Indeed, up to 50% of dietary protein is derived from insects, and they actually have higher market value than other protein sources (Dobermann et al., 2017). Some insects are appreciated for their organoleptic characteristics and consumed in high-class restaurants (DeFoliart, 1999). For example, *escamoles* (ant eggs) are considered a delicate gourmet dish in Mexico, Laos, Cambodia, and Europe (Ramos-Elorduy, 1997).

An interest in edible insects has increased rapidly because the Food and Agriculture Organization (FAO) has begun promoting insects as viable dietary options for humans (van Huis et al., 2013). Globally, the edible-insect market is expected to exceed USD 522 million by 2023 (Han et al., 2017). Since 2012, the Korean edible-insect market (especially for human consumption) has made major progress with government support and successful research endeavors.

However, lingering negative perceptions of insects hamper global market expansion and limit insects as a mainstream dining option, which may be related to the fact that people are skeptical to novel foods due to general neophobic tendencies (Dobermann et al., 2017). Thus, inexperienced consumers perceive insects as a source of fear or disgust, have strong rejection

of insects as a normal food in their diet, and totally neglect their high nutritive value (van Huis, 2016). Overcoming such intrinsic attitudes are one of the main challenges facing the edible-insect industry (Yen, 2009). Fortunately, the positive social perception of novel foods does lead to its consumption (Yen, 2010), as consumer attitudes are driven multiple factors (e.g., food-choice motives) beyond neophobia (van Thielen et al., 2019). Thus, initiatives promoting edible insects should emphasize their practical value, which can create consumer demand (Sun-Waterhouse et al., 2016). Governments and NGOs can provide the information about benefits related to nutrition and environmental sustainability (Ruby et al., 2015). Tasting events or educational workshops can also provide opportunities to learn about edible insects (Han et al., 2017). Another method for improving consumer perception is the creation of cookbooks with insect recipes (Raheem et al., 2018). Overall, the repeated positive exposure to edible insects raises awareness and could encourage consumption (van Thielen et al., 2019).

Moreover, an increase in accessibility to edible insects is to develop insect-based ingredients instead of final food products showing their original appearance (Mishyna et al., 2019; Nongonierma and FitzGerald, 2017; Purschke et al., 2018; Yi et al., 2016). Incorporating edible insects in already-familiar foods may be more acceptable for an insect-phobic culture than providing insects directly as a food option, and using insects as food ingredients is beneficial for the formation of sustainable business models (Han et al., 2017).

A major hurdle in the edible insect industry is the lack of systematic work to guarantee safety and shelf-life (van Huis, 2016). Insect farming also requires standardization and quality control, a goal that requires government legislation and regulations (Han et al., 2017).

Feed industry

High nutritional value, minimal space requirements, and low environmental impact combine to make insects an appealing option for animal feed. Another major advantage is that insects are already used for the natural part of many animal diets (Rumpold and Schlüter, 2013b; Veldkamp and Bosch, 2015). Insect-based animal feeds are particularly attractive when considering the cost of standard feeds, currently accounting for 70% of livestock-production expenses (van Huis et al., 2013).

The most promising, well-studied candidates for industrial feed production are black soldier flies, larvae, yellow mealworms, silkworms, grasshoppers, and termites (Dobermann et al., 2017). Such previous research has revealed that insect meal can partially replace commercial meal in broiler feed, particularly protein sources. For example, housefly-larvae meal can replace 4% of the fish meal in broiler diets without negative effects on carcass weight and feed efficiency (Awoniyi et al., 2003). Another report indicated that broiler diets containing 31%, 26%, and 20% soybean meal can be successfully modified to contain 0%, 5%, and 10% dried yellow mealworm, respectively (Ramos-Elorduy et al., 2002). Furthermore, growth performance and carcass quality could be improved when broiler chicks were fed a diet containing 10%–15% housefly larvae (Hwangbo et al., 2009). As compared with commercial corn/soy-based diets, a housefly-larvae-based diet significantly increased the carcass weight, total feed intake, and average daily gain of broiler chickens (Pretorius, 2011). However, a more recent study found that replacing soybean oil with black-soldier-fly-larvae meal has no impacts on the growth performance of broilers (Schiavone et al., 2017). In addition to the nutritional value, insect-based feed could have a further advantage in improving the taste of final meat products. In the Philippines, for example, consumers prefer the taste of pasture-grown chickens fed with grasshoppers, resulting in higher price compared with chickens on commercial feed (Litton, 1993).

Insect-based feeds have also been tested in egg-producing poultry. The replacing fish meal with dried mealworm increased egg production by 2.4% (Wang et al., 1996). Fully replacing the protein content with larvae meal in a laying-hen diet did not negatively affect feed intake, feed conversion efficiency, egg production hen health, and immune status (Marono et al., 2017).

Smallholder farms in Asia and Africa commonly use insects as fish feed (Dobermann et al., 2017). Housefly-larvae meal can replace up to 75% of fishmeal in Nile tilapia diets without any adverse effects (Wang et al., 2017). Replacing fish meal with black-soldier-fly meal in diets does not alter the odor, flavor, or texture of Atlantic salmon (*Salmo salar*) (Lock et al., 2016). Likewise, mealworms can successfully replace 40%–80% of the standard catfish (*Ameiurus melas* Raf.) diet without adverse effects on growth performance (Roncarati et al., 2015). Another viable alternative to fish meal is silkworm pupa, which was tested successfully for African catfish (*Clarias gariepinus*) fingerling diets (Kurbanov et al., 2015). In carp (*Cyprinus carpio*), a silkworm-meal diet was superior to a leaf-meal diet for improving nutrient digestibility, nutrient retention, and feed conversion efficiency (Swamy and Devaraj, 1994).

In summary, existing studies clearly demonstrate that insects are a promising protein source for animal feeds and has the potential to meet increasing global demand (Rumpold and Schlüter, 2013b). Before the mass-production of such insect-based feeds, however, governments and companies should aim to address health and safety concerns related to edible insects, such as the presence of anti-nutrient properties (Dobermann et al., 2017).

Value of Edible Insects as Food Resources

Nutritional value of edible insects

Insect nutritional value varies with diet, developmental stage, sex, species, growth environment, and measurement methods (van Huis and Oonincx, 2017). Nevertheless, researchers generally agree that insects are extremely rich in protein, fat, and vitamins (Rumpold and Schlüter, 2013a). Nutrient compositions of edible insects as published in literatures (based on dry matter) is summarized in Table 1. Nitrogen is a critical nutrient, and proteins directly involved in N supply comprise 16.5% of an adult human body (Melo et al., 2011). On average, the protein content of edible insects ranges 35%–60% dry weight or 10%–25% fresh weight (Melo et al., 2011; Schlüter et al., 2017), which are higher than plant protein sources, including cereal, soybeans, and lentils (Bukkens, 1997). At the upper range, insects provide more protein than even meat and chicken eggs (Mlcek et al., 2014). Edible insects in Orthoptera (crickets, grasshoppers, locusts) are particularly protein-rich (Rumpold and Schlüter, 2013a). However, insect protein digestibility is highly variable due to the presence of a hard exoskeleton (van Huis, 2016). Exoskeletons with high proportion of chitin component are especially difficult to digest (Schlüter et al., 2017). Indeed, we currently do not know whether humans are capable of digesting chitin (Muzzarelli et al., 2012). Of course, the removal of exoskeleton through a part of processing is a viable option (Rumpold and Schlüter, 2013b). Some studies have found that insect protein digestibility is 77%–98% without after the exoskeleton (Defoliart, 1992).

The second largest component of insect nutrient composition is fat (Mlcek et al., 2014). Various factors such as species, sex, reproduction stage, season, diet, and habitat all combine to influence insect fat content (Schlüter et al., 2017). Orthoptera, Lepidoptera (caterpillars), cockroaches (blattodea), Isoptera (termites), Hemiptera, and Coleoptera (beetles, grubs) have the averaged fat content of 13.41%, 27.66%, 29.90%, 32.74%, 30.26%, and 33.40%, respectively (Rumpold and Schlüter, 2013a). Larvae and pupae have more fat than adult insect (Mlcek et al., 2014). In addition, females are fatty than males (de Castro et al., 2018; Mlcek et al., 2014). The fatty acid profiles of insects are also dependent on species and diet (Schlüter et al., 2017), although insects generally have more unsaturated fatty acids (UFA) compared to saturated fatty acids (SFA) (de Castro et al., 2018). On average, Hymenoptera (ants, bees, and wasps) and 41.97% for Isoptera (termites) have 30.83% and 41.97% SFA, respectively (Rumpold and Schlüter, 2013a). A notable exception occurs in *Imbrasia ertli* caterpillars, which possess SFA comprising 38% arachidic acid (C20:0) (Bukkens, 1997; Rumpold and Schlüter, 2013a).

Table 1. Nutrition composition of edible insects (based on dry matter)

Type of insects	Scientific name	Protein content (%)	Fat content (%)	Carbohydrates content (%)	Reference
Larvae	<i>Allomyrina dichotoma</i>	54.18	20.24	- ¹⁾	Ghosh et al. (2017)
	<i>Anaphe infracta</i>	20.00	15.20	-	Banjo et al. (2006)
	<i>Anaphe recticulata</i>	23.00	10.20	-	Banjo et al. (2006)
	<i>Anaphe venata</i>	25.70	23.21	-	Banjo et al. (2006)
	<i>Gonimbrasia belina</i>	56.95	10.00	7.80	Siulapwa et al. (2012)
	<i>Gynanisa maja</i>	55.92	12.10	10.70	Siulapwa et al. (2012)
	<i>Protaetia brevitarsis</i>	44.23	15.36	-	Ghosh et al. (2017)
	<i>Rhynchophorus phoenicis</i>	22.06	66.61	5.53	Ekpo and Onigbinde (2005)
	<i>Tenebrio molitor</i>	46.44	32.70	-	Ravzanaadii et al. (2012)
Beetle	<i>Heteroligus meles</i>	38.10	32.01	20.10	Jonathan (2012)
	<i>Oryctes boas</i>	26.00	1.50	-	Banjo et al. (2006)
	<i>Rhynchophorus phoenicis</i>	50.01	21.12	20.23	Jonathan (2012)
	<i>Rhynchophorus phoenicis</i>	28.42	31.40	-	Banjo et al. (2006)
Grasshopper	<i>Ruspolia differens</i>	44.59	49.00	8.40	Siulapwa et al. (2012)
	<i>Zonocerus variegatus</i>	26.80	3.80	-	Banjo et al. (2006)
Cricket	<i>Brachytrypes</i> spp.	6.25	2.34	-	Banjo et al. (2006)
	<i>Gryllus bimaculatus</i>	58.32	11.88	-	Ghosh et al. (2017)
	<i>Teleogryllus emma</i>	55.65	25.14	-	Ghosh et al. (2017)
Termites	<i>Macrotermes bellicosus</i>	20.10	28.20	-	Banjo et al. (2006)
	<i>Macrotermes falciger</i>	43.26	43.00	32.80	Siulapwa et al. (2012)
	<i>Macrotermes notalensis</i>	22.10	22.50	-	Banjo et al. (2006)
Bee	<i>Apis mellifera</i>	21.00	12.30	-	Banjo et al. (2006)
Dragonfly	<i>Aeschna multicolor</i>	54.24	16.72	-	Ramos-Elorduy et al. (1998)
	<i>Anax</i> sp.	26.22	22.93	-	Ramos-Elorduy et al. (1998)

¹⁾ Do not observed in the reference.

Carbohydrates in insects mainly exist in two forms of chitin and glycogen. The former is a polymer of N-acetyl-D-glucosamine that is the primary component of exoskeleton (Bukkens, 1997; van Huis et al., 2013), while the latter is an energy source stored in cells and muscle tissues (Schluter et al., 2017). The averaged carbohydrate content of edible insects ranges from 6.71% (stink bug) to 15.98% (cicada) (Mlcek et al., 2014).

Some insects (e.g., grasshoppers, crickets, termites, and mealworms) are rich in iron, zinc, calcium, copper, phosphorus, magnesium, and manganese. Invertebrates without a mineralized skeleton have very little calcium content (de Castro et al., 2018; Mlcek et al., 2014). Most edible insects have similar iron content to beef (Bukkens, 1997), but we currently know little about mineral bioavailability of insects (de Castro et al., 2018). A rare study found that consuming insects can provide the high proportions of daily mineral recommendations for humans, particularly in terms of iron (Latunde-Dada et al., 2016). The investigations of vitamin content are also insufficient, but available data indicate that edible insects contain carotene, vitamin B₁, B₂, B₆, C, D, E, and K (Mlcek et al., 2014). In particular, Orthoptera and Coleoptera are rich in folic acid (Rumpold and

Schlüter, 2013a).

Food processing of edible insects

As traditional animal-protein intake methods are called into question, insects are increasingly viewed as the food of the future (Sogari, 2015). In the Netherlands, the efforts to promote entomophagy include studies aimed at customizing insects for Western tastes, resulting in the sale of some insects (e.g., locusts, mealworms) at retail markets (Schösler et al., 2012). Insects have been processed into powder or meal to minimize visual associations and increase palatability (Bubler et al., 2016). In addition, researchers have been investigating the functional properties of insect proteins, including gelling capacity, foam capacity, emulsion capacity, and solubility in various buffers or solvents. In Table 2, previous studies regarding the food processing properties (functional properties) of edible insects are presented. Appropriate extraction methods (e.g., defatting, pH change, zymolysis, or sonication) are also being investigated (Mishyna et al., 2019; Nongonierma and FitzGerald, 2017; Purschke et al., 2018; Yi et al., 2016). These studies will help the use of edible insects as a food ingredient for producing traditional foods.

Medicinal applications

Cultures that consume insects also tend to associate them with various health benefits beyond nutrition (Raheem et al., 2018). For example, caterpillar fungus supposedly has immunostimulatory and anti-cancer properties (Chen et al., 2009). In traditional Chinese medicine, male *Antheraea pernyi* is considered aphrodisiac (Chen et al., 2009), while ant-alcohol is also thought to enhance immunity and libido (Liu et al., 2004). Some evidence exists to suggest that termites (*Macrotermes annandalei*) may have immunostimulatory effects (Chen et al., 2009). Another insect historically considered to have beneficial health effects is the silkworm (*Bombyx mori* L.) (Kim et al., 2008). Recent analyses have identified a blood-glucose-lowering agent, resulting in the development of silkworm powder as a diabetic medicine in Korea, and such health benefit was also reported by the Chinese Ministry of Health and State Food and Drug Administration (Belluco et al., 2013; Nongonierma and FitzGerald, 2017).

The traditional claims of medicinal properties have resulted in multiple studies aiming to empirically determine the properties of edible insects. The several analyses of insect enzymatic hydrolysates have identified antioxidant and antidiabetic properties, as well as the ability to inhibit angiotensin converting enzyme (ACE) (de Castro et al., 2018; Nongonierma and FitzGerald, 2017). The hydrolysates of *S. littoralis* larvae exhibit 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) and 2,2-diphenyl-1-picrylhydrazyl scavenging activity that indicates a powerful antioxidant activity (Vercruyssen et al., 2009).

In summary, the discovery of physiologically important substances contained in edible insects makes medicinal use a potentially important application (Han et al., 2017). However, given the current lack of scientific data, more researches are warranted to confirm the benefits of consuming insects as health foods or medicine.

Food safety of edible insects

Despite the obvious nutritional value of edible insect, the food safety issues of edible insects such as microorganism, allergic reaction, and toxicity may catch up with latent consumer (Rumpold and Schlüter, 2013a). Harmful factors affect food safety of edible insect is summarized in Table 3. In the perspective on microbial safety, insect gut is considered as a main habitat, and body surface and mouthparts are also included. Microbial colonization is reached vertically through ovary, egg capsule, spawning, and horizontally through feed and environment (Dematheis et al., 2012; Yun et al., 2014). Therefore,

Table 2. Previous studies regarding the food processing properties of edible insects

Type of insects	Scientific name	Observation	Reference
Larvae	<i>Cirina forda</i>	<ul style="list-style-type: none"> • Protein solubility • Oil absorption capacity • Foaming stability 	Omotoso (2006)
Larvae	<i>Grylloides sigillatus</i>	<ul style="list-style-type: none"> • Protein solubility 	Zielinska et al. (2018)
Grasshopper	<i>Schistocerca gregaria</i>	<ul style="list-style-type: none"> • Water holding capacity 	
Cricket	<i>Tenebrio molitor</i>	<ul style="list-style-type: none"> • Oil holding capacity • Foaming capacity • Emulsion capacity 	
Larvae	<i>Cirina forda</i>	<ul style="list-style-type: none"> • Protein solubility • Oil absorption capacity • Water absorption capacity • Foaming capacity • Emulsion capacity 	Osasona and Olaofe (2010)
Larvae	<i>Oryctes owariensis</i>	<ul style="list-style-type: none"> • Oil absorption capacity 	Assielou et al. (2015)
Larvae	<i>Bombyxmori L.</i>	<ul style="list-style-type: none"> • Water absorption capacity • Foaming capacity • Foaming stability • Emulsion capacity • Emulsion stability 	Omotoso (2015)
Larvae	<i>Tenebrio molitor</i>	<ul style="list-style-type: none"> • Water binding capacity 	Bubler et al. (2016)
Larvae	<i>Hermetia illucens</i>	<ul style="list-style-type: none"> • Fat binding capacity • Protein solubility 	
Larvae	<i>Tenebrio molitor</i>	<ul style="list-style-type: none"> • Foaming capacity 	Yi et al. (2013)
Larvae	<i>Zophobas morio</i>	<ul style="list-style-type: none"> • Foaming stability 	
Larvae	<i>Alphitobius diaperinus</i>	<ul style="list-style-type: none"> • Gel formation 	
Cricket	<i>Acheta domesticus</i>		
Cockroach	<i>Blaptica dubia</i>		

the use of insect as food sources may have a potential hazard as a pathogenic vector. In addition, the microbiota ratio in insect gut can be an important part being used as food sources, because the total amount of biomass from gut microbiota could account for 1%–10% of the body weight of insect (Douglas, 2015). In addition, it is practically difficult to remove gut part from edible insects (Douglas, 2015). Although insect-specific pathogenic microorganisms had no risk to human health, they can colonize only cells or tissues of insect. However, the contamination of human pathogenic microorganisms can occur, and this contamination may be regulated by controlled breeding condition with special care (Schluter et al., 2017). In allergic reactions, some edible insects had allergic cross-reactive proteins with arachnids and crustaceans, which are known as arthropods (Ribeiro et al., 2018). Furthermore, carmine used for food dye is obtained from the bodies of female cochineal insects and this dye could induce allergic reaction to some patients (Belluco et al., 2013). Considering that only a few studies

Table 3. Harmful factors affecting the food safety of edible insects

Issues	Derived hazard sources from edible insect		Note			
Microorganism	Gut and surface	<i>Enterococcus</i>	Insect-specific pathogenic microorganism is not harmful to human			
		<i>Streptococcus</i>				
		<i>Staphylococcus</i>				
		<i>Pseudomonas</i>				
		<i>Bacillus</i>				
		<i>Clostridium</i>				
		<i>Enterobacteriaceae</i>				
		<i>Aspergillus</i>				
		<i>Penicillium</i>				
		<i>Alternaria</i>				
		<i>Candida</i>				
		Allergene		Pan-allergenic structures	Tropomyosin	Cross-reactivity and co-sensitisation were observed with other crustaceans
					Myosin	
Actin						
Troponin C						
Tubulin						
Hemocyanin						
Defensin						
Triosephosphate isomerase						
α -Amylase						
Trypsin						
Phospholipase A						
Hyaluronidase						
Arginine kinase						
Chitin						
Pathogenic molds	<i>Aspergillus</i>					
	<i>Penicillium</i>					
Pathogenic yeast	<i>Candida</i>					
Chemical or toxin	Synthesis toxin		Oxalate		There were no reports that edible insect had toxin	
			Tannin			
		Phytate				
		Thiaminases				
		Cantharidin				
		Toluene				
		Quinones				
		Alkanes				
		Cyanogenic glycosides				
		Microorganism toxin	<i>Bacillus</i>			
			<i>Clostridium</i>			
			<i>Aspergillus</i>			

Sources: Belluco et al. (2013), Dematheis et al. (2012), Ribeiro et al. (2018), Schluter et al. (2017), Yun et al. (2014).

regarding the allergic reaction of edible insects have been carried out, we need to pay attention to the unknown potential allergens contained in edible insects. Further study and investigation determining the hazard substances including allergens and toxic substances and their impacts on pathological symptom in human body should be performed for guaranteeing the safe use of edible insects as feed and food resources.

Historical and Current Consumption Patterns of Edible Insects

Historical insect intake

The lack of sophisticated tools and well-developed hunting ability mean that ancient cultures likely consumed insects frequently (Valadez, 2003). Upon the development of agriculture and domestication of livestock, however, insect-eating habits have been disappearing in the many regions (Abrams, 1987). The resultant abundance of food naturally could also cause a decrease in the need for insect consumption. With changing cultural mores, insects have evolved from a primary food source to snacks, luxury ingredients, and bait (Smeathman, 1781). However, some countries have still continued to use insects as ordinary food resources to this day. Below, we will introduce and discuss detailed examples of current patterns of insect consumption.

Edible insects in Asia

In Korea, canned silkworm pupae are sold in retail markets and processed as snacks. Additionally, grasshoppers (*Oxya velox*) are eaten in rural areas (Pemberton, 1994). In Japan, a dish called *inago* involves fried grasshoppers seasoned with soy sauce. Moreover, bee or wasp larvae/adults are considered expensive delicacies; they are eaten raw, boiled with soy sauce, or served with rice (DeFoliart, 1999). Entomophagy has existed in China for over 2000 years; approximately 324 species from 11 orders are consumed possibly (Feng et al., 2018). India has the developed multiple uses for insects, including silk, fertilizer, food, and medicine. Notably, eri silkworm (*Samia ricini*) pupae is a delicacy in northeastern India (Peigler, 1993). Overall, about 255 species are used as food, although the intensity varies depending on seasonal or regional differences in culture (Chakravorty, 2014). In Thailand, insects are an important source of protein, fat, and other nutrients; the villages of Ubon alone consume 20–60 g of insects daily (Sungpuag and Puwastien, 1983). Over 80 species are considered edible food resources (DeFoliart, 1999). Furthermore, Thailand's Ministry of Public Health recommends insect consumption for rural communities to obtain necessary nutrients (DeFoliart, 1999), in addition, it has also spread from rural to urban markets (Yhounng-aree, 2010). The information on major edible insects consumed in Asia is summarized in Table 4.

Edible insects in Oceania

Sago grub (*Rhynchophorus ferrugineus*) is a popular edible insect in Papua New Guinea and a main part of an annual grub festival (Mercer, 1993). In local regions, locusts, crickets, mole-crickets, mantises, and even spiders are consumed (Meyer-Rochow and Changkija, 1997). Aboriginal tribes consumed a wide variety of insects from Cossidae, Noctuidae, Cerambycidae, and bees (Macfarlane, 1978). In Australia, entomophagy is low among European-derived populations (Yen, 2010). However, the edible-insect market has increased dramatically alongside an interest in bushfood, and insects are now available as restaurant menus (Irvine, 1989). More information about the edible insects in Oceania are given in Table 5.

Edible insects in Africa

Table 4. Most consumed insects as food resources traditionally in Asia

Country	Scientific and common name of consumed insects		Consumed type
Korea	<i>Bombyx mori</i>	Silkworm	Pupae
	<i>Oxya velox</i>	Grasshopper	Adult
Japan	<i>Oxya yezoensis</i>	Grasshopper	Adult
	<i>Oxya japonica</i>	Grasshopper	Adult
	<i>Vespula lewisii</i>	Bee	Larvae, pupae
	<i>Bombyx mori</i>	Silkworm	Pupae, female adult
China	<i>Bombyx mori</i>	Silkworm	Pupae
	<i>Antheraea pernyi</i>	Tussah silkworm	Pupae
	<i>Tenebrio molitor</i>	Mealworm	Larvae
	<i>Polyrhachis dives</i>	Weaver ants	Adult
	<i>Apis cerana</i>	Honey bee	Larvae, pupae
	<i>Locusta migratoria</i>	Locust	Adult
	<i>Coccoidea</i>	Scale insect	Adult
	<i>Cicadoidea</i>	Cicadas	Adult
	<i>Apongopus chinensis</i>	Stinkbugs	Adult
	<i>Odontotermes formosanus</i>	Termites	Adult
	<i>Anax parthenope</i>	Dragonfly	Nymphs
India	<i>Samia cynthia</i>	Eri silkworm	Larvae, pupae, adult
	<i>Corizus hyoscyami</i>	Cinnamon bug	Adult
	Termites	Termites	Adult (winged, queen)
	<i>Schistocerca gregaria</i>	Locust	Adult
	Honey bee		Adult
	<i>Udonga montana</i>	Pentatomid bug	Adult
Thailand	<i>Oecophylla smaragdina</i>	Weaver ant	Adult
	<i>Meimuna opalifera</i>	Walker's cicada	Adult
	<i>Allonemobius fasciatus</i>	Ground cricket	Adult
	<i>Anurogryllus arboreus</i>	Short tailed cricket	Adult
	<i>Belostomatidae</i>	Giant water bug	Adult
	<i>Bombyx mori</i>	Silkworm	Pupae
	<i>Cotinis nitida</i>	June beetle	Adult
	<i>Omphisa fuscidentalis</i>	Bamboo caterpillar	Larvae

Sources: Chakravorty (2014), DeFoliart (1999), Feng et al. (2018), Pemberton (1994), Raheem et al. (2018).

Edible insects play an important role in the food culture of Africa (Mutungi et al., 2019). Caterpillars and termites are the most popular insects, but the insects from other orders are also widely eaten (DeFoliart, 1999), including Lepidoptera, Orthoptera, and Coleoptera (Kelemu et al., 2015). In total, approximately 470 species are consumed, and eaten insects are summarized in Table 6. Some examples from specific countries include grasshoppers, considered a source of energy and

Table 5. Most consumed insects as food resources traditionally in Oceania

Country	Scientific and common name of consumed insects	Consumed type
Papua New Guinea	<i>Rhynchophorus ferrugineus</i>	Sago grub
	Locusts	Larvae
	Crickets	Adult
	Mantis	Adult
Australia	Cossidae	Larvae
	Noctuidae	Larvae
	Cerambycids	Larvae
	Bees	Adult, honey

Sources: Macfarlane (1978), Mercer (1993), Meyer-Rochow and Changkija (1997).

protein in Uganda (Mbabazi, 2011). Additionally, edible insects could account for over 20% of the animal protein produced in the Democratic Congo, where more than 65 species are consumed (DeFoliart, 1999). Overall, entomophagy is widespread and important to human nutrition in Africa (Mutungi et al., 2019).

Edible insects in the Americas

Major edible insects consumed in the Americas are shown in Table 7. Edible insects are a major protein source for Amazonian tribes, with women and children actively foraging for them (Dufour, 1987). In particular, *Rhynchophorus palmarum* and Atta ants are popular and mass-produced in South America (DeFoliart, 1999). In Mexico, edible insects are traditionally eaten in both rural and urban regions. However, increasing westernization of cities after Spanish conquest eventually has limited entomophagy largely to rural areas (Ramos-Elorduy, 1997). Nevertheless, Mexican restaurants often served *escamol*, an insect dish fried with aromatic spices (DeFoliart, 1999). In Colombia, the Yukpa people prefer insects over meat, but it has been forced to decrease insect consumption owing to massive deforestation (Ruddle, 1973).

Negative attitudes toward edible insects

The adoption of Western values among countries that traditionally consumed insects is correlated with a decrease in this behavior (DeFoliart, 1999; Ramos-Elorduy, 1997; Yen, 2010). Therefore, Western attitudes toward edible insects must be adjusted to successfully expand the global market. Several explanations are available for why Western cultures generally did not develop entomophagy, including insect size, dispersed distribution, and seasonal unavailability (van Huis, 2016). Existing cultural differences were then used by Western cultures to denigrate traditional peoples, leading to an association of disgust with insect consumption (Looy et al., 2014). Furthermore, despite the fact that only 0.2% of insects are actually detrimental to human life, the persistent perceptions of harmful insects contribute to the reluctance in accepting entomophagy (van Huis, 2016). Further systemic efforts must be made to alter cultural attitudes through cooperation between academia, government, and industry.

Prospect of Insects as Food Resources

Despite the numerous advantages of insect consumption, the future of insect industry is unfavorable in Western societies (Sogari, 2015; van Huis, 2016), given that existing cultural distaste cannot be changed rapidly (DeFoliart, 1999). Fortunately,

Table 6. Most consumed insects as food resources traditionally in Africa

Country	Scientific and common name of consumed insects		Consumed type
Nigeria	<i>Macrotermes natalensis</i>	Termite	Adult
	<i>Microtermes bellicosus</i>	Termite	Adult
	<i>Brachytrupes membranaceus</i>	Locust	Adult
	<i>Nezara viridula</i>	Stinkbug	Adult
	<i>Anaphe reticulata</i>	Reticulate bagnet	Larvae
	<i>Cirina forda</i>	Pallid emperor moth	Larvae
	<i>Anaphe venata</i>	African silkworm	Larvae
	<i>Oryctes monocerus</i>	Palm beetle	Larvae
	<i>Rhynchophorus phoenicis</i>	Palm weevil	Larvae
	<i>Analeptes trifasciata</i>	Stem girdler	Larvae
	<i>Oryctes boas</i>	Rhinoceros beetle	Larvae
	<i>Oryctes monoceros</i>	Rhinoceros beetle	Larvae
	<i>Aphodius rufipes</i>	Dung beetle	Larvae
	<i>Heteroligus meles</i>	Yam beetle	Larvae
	<i>Apis mellifera</i>	Honey bee	Egg, larvae, pupa, honey
	<i>Brachytrupes membranaceus</i>	Giant African cricket	Adult
	<i>Gymnogryllus lucens</i>	Cricket	Adult
	<i>Cytacanthacris naeruginosus</i>	Short horned grasshopper	Adult
	<i>Zonocerus variegatus</i>	Grasshopper	Adult
	<i>Gryllotalpa africana</i>	Mole cricket	Adult
Uganda	<i>Macrotermes</i> spp.	Termites	Adult
	<i>Ruspolia nitidula</i>	Grasshopper	Adult
	<i>Chaoborus edulis</i>	Lake fly	Adult
	<i>Ruspolia differens</i>	Grasshopper	Adult
Congo	<i>Oryctes owariensis</i>	Rhinoceros beetle	Larvae
	<i>Anaphe panda</i>	Silkworm	Larvae
	Termites		Adult
	Bees		Egg, larvae, pupa, honey
Kenya	<i>Pseudacanthotermes militaris</i>	Winged termite	Adult
	<i>Macrotermes bellicosus</i>	Winged termite	Adult
	<i>Macrotermes subhyllanus</i>	Winged termite	Adult
	<i>Pseudacanthotermes spiniger</i>	Winged termite	Adult
	<i>Carebara vidua</i>	Black ant	Adult
	<i>Ruspolia differens</i>	Grasshopper	Adult

Sources: DeFoliart (1999), Mutungi et al. (2019), Raheem et al. (2018).

the increased consumer knowledge about edible insects also increases the willingness to pay for insect food (Piha et al., 2018). Continuous promotional efforts to increase exposure, coupled with development to enhance taste and appearance

Table 7. Most consumed insects as food resources traditionally in America

Country	Scientific and common name of consumed insects		Consumed type
Mexico	<i>Rhynchophorus palmarum</i>	Palm weevil	Larvae
	<i>Sphenarium</i>	Grasshopper	Adult
	<i>Schistocerca</i>	Grasshopper	Adult
	<i>Atizies taxcoensis</i>	Stink bug	Adult
	<i>Liometopum apiculatum</i>	Ant	Larvae, pupae
	<i>Liometopum occidentale</i>	Ant	Larvae, pupae
Colombia	<i>Aidemona azteca</i>	Grasshopper	Adult
	<i>Tropidacris latreillei</i>	Grasshopper	Adult
	<i>Podischnus agenor</i>	Beetles	Larvae
	<i>Caryobruchus scheelae</i>	Beetles	Larvae
	<i>Corydalus</i> spp.	Dobsonfly	Larvae
	<i>Atta</i>	Ant	Adult
	<i>Eumenes canaliculata</i>	Potter wasps	Larvae
	<i>Polybia ignobilis</i>	Wasps	Egg, larvae
	<i>Trigona trinidadensis</i>	Bee	Honey
<i>Trigona clavipes</i>	Bee	Honey	

Sources: DeFoliart (1999), Ramos-Elorduy (1997), Ruddle (1973).

(Looy et al., 2014; van Huis et al., 2013), has successfully improved the negative perceptions in some Western countries. For example, consumers in Belgium increasingly accept insects as an excellent food source (van Thielen et al., 2019). Furthermore, the insect industry in the Netherlands has been successfully marketed, and the freeze-dried insect powder is sold as a meat replacement (Raheem et al., 2018). To increase acceptance, social, practical, and contextual factors affecting food consumption must be emphasized to consumers (House, 2016). Such efforts involve continuous education and promotion regarding the potential for edible insects to solve environmental, population, and land-availability problems today and in the future.

Here, we introduce available patents related to edible insects. The aim of this examination is to provide the objective information on the technological status of insect consumption, which will then be used for establishing strategic research and development. The patent applications related to edible insects have been filed yearly, with a notable uptick since mid-2010. Currently, 167 patents have been filed (Fig. 1). Thus, the research and development on entomophagy is being actively pursued. A sharp decrease in patent applications is observed starting from 2018, because it is probably that patents are released 18 months after application. In terms of the raw materials from edible insects, the most common patent applications are related to pupa, silkworm, brown goat, and grasshopper, in that order. Our analysis shows that most patents have been applying to protect food processing methods using edible insects.

To better understand technological growth, we compared the number of applicants and applications across various time intervals (Fig. 2). We employed a time-series analysis based on patent number, using a technical life-cycle method (Little, 1981). The degree of competition is defined as applicant number in the analysis section. Production efficiency is a means of analyzing trends in time series by replacing application number or the extent of increase in applications.




Fig. 1. Patent application trend by edible insect technology year. Bars indicate the number of patent application each year, and a dotted line indicate the cumulative count of patent application.




Fig. 2. Technology growth stage of edible insect patent application.

In general, the technology life cycle consists of introduction, growth, maturity, and decay. Applicants and applications begin to appear during the introduction period. During the growth period, application numbers increase. During maturity, the number of applications received, and the number of continuing applications start to decrease. During decline, the number of applications remains steadily low. In our analysis of total effective patent applications filed, we divided the technical life cycle trends into four sections, approximately a four-year term: 1998–2002, 2003–2007, 2008–2012, and 2013–2018. We observed a continuous increase in applicant and application number from section one section to four. This pattern indicates that edible-insect-related technology may enter the stage of growth. Taken together, these analysis of patent trends suggest that the technologies related to edible insects are experiencing rapid and continuous growth.

Conclusions

Entomophagy could be the solution to an increasingly urgent food-security problem confronting the world. As we have described, numerous countries already use insects as alternative food protein and feed sources. Existing researches confirm

the considerable nutritional and medicinal values of edible insects. Various strategies have been deployed to expand the edible-insect market and to counteract existing animus among Western societies toward entomophagy. As a result, patent numbers have increased continuously. In conclusion, this review shows the current status of edible insect remains insufficient to replace traditional animal foods worldwide, although edible insects have strong potential to become a primary source of nutrients.

Conflict of Interest

The authors declare no potential conflict of interest.

Acknowledgements

This research was supported by Main Research Program (E0193118-01) of the Korea Food Research Institute (KFRI) funded by the Ministry of Science and ICT (Korea).

Author Contributions

Conceptualization: Choi YS. Data curation: Kim TK, Yong HI. Formal analysis: Kim YB, Kim HW. Methodology: Kim TK, Yong HI. Software: Choi YS, Kim HW. Validation: Kim YB, Kim HW. Investigation: Choi YS. Writing - original draft: Choi YS, Kim TK, Yong HI. Writing - review & editing: Kim TK, Yong HI, Kim YB, Kim HW, Choi YS.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abrams HL. 1987. The preference for animal protein and fat: A cross-cultural survey. Temple University Press, Philadelphia, PA, USA.
- Assielou B, Due EA, Koffi MD, Dabonne S, Kouame PL. 2015. *Oryctes owariensis* larvae as good alternative protein source: Nutritional and functional properties. *Annu Res Rev Biol* 8:1-9.
- Awoniyi TAM, Aletor VA, Aina JM. 2003. Performance of broiler-chickens fed on maggot meal in place of fishmeal. *Int J Poult Sci* 2:271-274.
- Azua RV. 2003. La domesticación animal. Mexico City: Universidad Nacional Autónoma de Mexico/Instituto de Investigaciones Antropológicas.
- Banjo AD, Lawal OA, Songonuga EA. 2006. The nutritional value of fourteen species of edible insects in southwestern Nigeria. *Afr J Biotechnol* 5:298-301.
- Belluco S, Losasso C, Maggioletti M, Alonzi CC, Paoletti MG, Ricci A. 2013. Edible insects in a food safety and nutritional perspective: A critical review. *Compr Rev Food Sci Food Saf* 12:296-313.
- Bubler S, Rumpold BA, Jander E, Rawel HM, Schluter OK. 2016. Recovery and techno-functionality of flours and proteins

- from two edible insect species: Meal worm (*Tenebrio molitor*) and black soldier fly (*Hermetia illucens*) larvae. *Heliyon* 2:e00218.
- Bukkens SG. 1997. The nutritional value of edible insects. *Ecol Food Nutr* 36:287-319.
- Chakravorty J. 2014. Diversity of edible insects and practices of entomophagy in India: An overview. *J Biodivers Biopros Dev* 1:124.
- Chen X, Feng Y, Chen ZY. 2009. Common edible insects and their utilization in China. *Entomol Res* 39:299-303.
- de Castro RJS, Ohara A, dos Santos Aguilar JG, Domingues MAF. 2018. Nutritional, functional and biological properties of insect proteins: Processes for obtaining, consumption and future challenges. *Trends Food Sci Technol* 76:82-89.
- Defoliart GR. 1992. Insects as human food: Gene DeFoliart discusses some nutritional and economic aspects. *Crop Protection* 11:395-399.
- DeFoliart GR. 1999. Insects as food: Why the western attitude is important. *Annu Rev Entomol* 44:21-50.
- Dematheis F, Kurtz B, Vidal S, Smalla K. 2012. Microbial communities associated with the larval gut and eggs of the western corn rootworm. *PLOS ONE* 7:e44685.
- Dobermann D, Swift JA, Field LM. 2017. Opportunities and hurdles of edible insects for food and feed. *Nutr Bull* 42:293-308.
- Douglas AE. 2015. Multiorganismal insects: Diversity and function of resident microorganisms. *Annu Rev Entomol* 60:17-34.
- Dufour DL. 1987. Insects as food: A case study from the northwest Amazon. *American Anthropol* 89:383-397.
- Ekpo KE, Onigbinde AO. 2005. Nutritional potentials of the larva of *Rhynchophorus phoenicis* (F). *Pak J Nutr* 4:287-290.
- Feng Y, Chen XM, Zhao M, He Z, Sun L, Wang CY, Ding WF. 2018. Edible insects in China: Utilization and prospects. *Insect Sci* 25:184-198.
- Gao Y, Wang D, Xu ML, Shi SS, Xiong JF. 2018. Toxicological characteristics of edible insects in China: A historical review. *Food Chem Toxicol* 119:237-251.
- Ghosh S, Lee SM, Jung C, Meyer-Rochow VB. 2017. Nutritional composition of five commercial edible insects in South Korea. *J Asia-Pac Entomol* 20:686-694.
- Grafton RQ, Daugbjerg C, Qureshi ME. 2015. Towards food security by 2050. *Food Secur* 7:179-183.
- Han R, Shin JT, Kim J, Choi YS, Kim YW. 2017. An overview of the South Korean edible insect food industry: Challenges and future pricing/promotion strategies. *Entomol Res* 47:141-151.
- House J. 2016. Consumer acceptance of insect-based foods in the Netherlands: Academic and commercial implications. *Appetite* 107:47-58.
- Hwangbo J, Hong EC, Jang A, Kang HK, Oh JS, Kim BW, Park BS. 2009. Utilization of house fly-maggots, a feed supplement in the production of broiler chickens. *J Environ Biol* 30:609-614.
- Irvine G. 1989. Putting insects on the Australian menu. *Food Aust* 41:565-566.
- Jonathan AA. 2012. Proximate and anti-nutritional composition of two common edible insects: Yam beetle (*Heteroligus meles*) and palm weevil (*Rhynchophorus phoenicis*). *Food Sci* 49:9782-9786.
- Kelemu S, Niassy S, Torto B, Fiaboe K, Affognon H, Tonnang H, Maniania NK, Ekesi S. 2015. African edible insects for food and feed: Inventory, diversity, commonalities and contribution to food security. *J Insect Food Feed* 1:103-119.
- Kim SA, Kim KM, Oh BJ. 2008. Current status and perspective of the insect industry in Korea. *Entomol Res* 38:S79-S85.
- Kohler R, Kariuki L, Lambert C, Biesalski HK. 2019. Protein, amino acid and mineral composition of some edible insects

- from Thailand. *J Asia Pac Entomol* 22:372-378.
- Kurbanov AR, Milusheva RY, Rashidova SS, Kamilov BG. 2015. Effect of replacement of fish meal with silkworm (*Bombyx mori*) pupa protein on the growth of *Clarias gariepinus* Fingerling. *Int J Fish Aquat Stud* 2:25-27.
- Latunde-Dada GO, Yang W, Vera Aviles M. 2016. *In vitro* iron availability from insects and sirloin beef. *J Agric Food Chem* 64:8420-8424.
- Little AD. 1981. The strategic management of technology. European management forum. Davos, Swiss.
- Litton E. 1993. Grasshopper consumption by humans and freerange chickens reduces pesticide use in the Philippines. *Food Insects Newsletter* 6:3.
- Liu DW, Sun QS, Li T. 2004. The anti-fatigue activity of polyhachis vicina roger extract in mice. *Chin J Food Hyg* 16:334-343.
- Lock ER, Arsiwalla T, Waagbo R. 2016. Insect larvae meal as an alternative source of nutrients in the diet of Atlantic salmon (*Salmo salar*) postsmolt. *Aquac Nutr* 22:1202-1213.
- Looy H, Dunkel FV, Wood JR. 2014. How then shall we eat? Insect-eating attitudes and sustainable foodways. *Agric Human Values* 31:131-141.
- Macfarlane WV. 1978. Aboriginal desert hunter-gatherers in transition. Symposium on the Nutrition of Aborigines, Commonwealth Scientific and Industrial Research Organisation, Melbourne. pp 49-62.
- Marono S, Loponte R, Lombardi P, Vassalotti G, Pero ME, Russo F, Gasco L, Parisi G, Piccolo G, Nizza S, Di Meo C, Attia YA, Bovera F. 2017. Productive performance and blood profiles of laying hens fed *Hermetia illucens* larvae meal as total replacement of soybean meal from 24 to 45 weeks of age. *Poult Sci* 96:1783-1790.
- Mbabazi M. 2011. Dietary contribution of grasshoppers (*Ruspolia nitidula*) and white ants (*Macrotermes bellicosus*) and influence of processing methods on their nutrient composition. Ph. D. Dissertation, Makerere Univ., Kampala, Uganda.
- Melo V, Garcia M, Sandoval H, Jimenez HD, Calvo C. 2011. Quality proteins from edible indigenous insect food of Latin America and Asia. *Emir J Food Agric* 23:283-289.
- Mercer CWL. 1993. Insects as food in Papua New Guinea. In Proceedings of Invertebrates Farming Seminar. Institute of Tropical Medicine, Antwerp, Belgium. pp 157-162.
- Meyer-Rochow VB, Changkija S. 1997. Uses of insects as human food in Papua New Guinea, Australia, and North-East India: Cross-cultural considerations and cautious conclusions. *Ecol Food Nutr* 36:159-185.
- Mishyna M, Martinez JJI, Chen J, Benjamin O. 2019. Extraction, characterization and functional properties of soluble proteins from edible grasshopper (*Schistocerca gregaria*) and honey bee (*Apis mellifera*). *Food Res Int* 116:697-706.
- Mlcek J, Rop O, Borkovcova M, Bednarova M. 2014. A comprehensive look at the possibilities of edible insects as food in Europe—a review. *Pol J Food Nutr Sci* 64:147-157.
- Murefu TR, Macheke L, Musundire R, Manditsera FA. 2019. Safety of wild harvested and reared edible insects: A review. *Food Control* 101:209-224.
- Mutungi C, Irungu FG, Nduko J, Mutua F, Affognon H, Nakimbugwe D, Ekesi S, Fiaboe KKM. 2019. Postharvest processes of edible insects in Africa: A review of processing methods, and the implications for nutrition, safety and new products development. *Crit Rev Food Sci Nutr* 59:276-298.
- Muzzarelli RAA, Boudrant J, Meyer D, Manno N, DeMarchis M, Paoletti MG. 2012. Current views on fungal chitin/chitosan, human chitinases, food preservation, glucans, pectins and inulin: A tribute to Henri Braconnot, precursor of the carbohydrate polymers science, on the chitin bicentennial. *Carbohydr Polym* 87:995-1012.

- Nongonierma AB, FitzGerald RJ. 2017. Unlocking the biological potential of proteins from edible insects through enzymatic hydrolysis: A review. *Innov Food Sci Emerg Technol* 43:239-252.
- Omotoso OT. 2006. Nutritional quality, functional properties and anti-nutrient compositions of the larva of *Cirina forda* (Westwood) (Lepidoptera: Saturniidae). *J Zhejiang Univ Sci B* 7:51-55.
- Omotoso OT. 2015. An evaluation of the nutrients and some anti-nutrients in silkworm, *Bombyxmori* L. (Bombycidae: Lepidoptera). *Jordan J Biol Sci* 8:45-50.
- Osasona AI, Olaofe O. 2010. Nutritional and functional properties of *Cirina forda* larva from Ado-Ekiti, Nigeria. *Afr J Food Sci* 4:775-777.
- Park S, Yun E. 2018. Edible insect food: Current scenario and future perspectives. *Food Sci Anim Resour Ind* 7:12-20.
- Patel S, Suleria HAR, Rauf A. 2019. Edible insects as innovative foods: Nutritional and functional assessments. *Trends Food Sci Technol* 86:352-359.
- Peigler RS. 1993. Wild silks of the world. *Am Entomol* 39:151-162.
- Pemberton RW. 1994. The revival of rice-field grasshoppers as human food in South Korea. *Pan-Pac Entomol* 70:323-327.
- Piha S, Pohjanheimo T, Lahteenmaki-Uutelala A, Kreckova Z, Otterbring T. 2018. The effects of consumer knowledge on the willingness to buy insect food: An exploratory cross-regional study in Northern and Central Europe. *Food Qual Prefer* 70:1-10.
- Pretorius Q. 2011. The evaluation of larvae of *Musca domestica* (common house fly) as protein source for broiler production. Ph. D. Dissertation, Stellenbosch University, Stellenbosch, Republic of South Africa.
- Purschke B, Meinschmidt P, Horn C, Rieder O, Jager H. 2018. Improvement of techno-functional properties of edible insect protein from migratory locust by enzymatic hydrolysis. *Eur Food Res Technol* 244:999-1013.
- Raheem D, Carrascosa C, Oluwole OB, Nieuwland M, Saraiva A, Millan R, Raposo A. 2018. Traditional consumption of and rearing edible insects in Africa, Asia and Europe. *Crit Rev Food Sci Nutr* 59:2169-2188.
- Ramos-Elorduy J, Gonzalez EA, Hernandez AR, Pino JM. 2002. Use of *Tenebrio molitor* (Coleoptera: Tenebrionidae) to recycle organic wastes and as feed for broiler chickens. *J Econ Entomol* 95:214-220.
- Ramos-Elorduy J, Pino-M JM, Correa SC. 1998. Edible insects of the state of Mexico and determination of their nutritive values. *An Inst Biol Univ Nac Auton Mex Ser Zool* 69:65-104.
- Ramos-Elorduy BJ. 1997. The importance of edible insects in the nutrition and economy of people of the rural areas of Mexico. *Ecol Food Nutr* 36:347-366.
- Ravzanaadii N, Kim SH, Choi WH, Hong SJ, Kim NJ. 2012. Nutritional value of mealworm, *Tenebrio molitor* as food source. *Int J Indust Entomol* 25:93-98.
- Ribeiro JC, Cunha LM, Sousa-Pinto B, Fonseca J. 2018. Allergic risks of consuming edible insects: A systematic review. *Mol Nutr Food Res* 62:1700030.
- Roncarati A, Gasco L, Parisi G, Terova G. 2015. Growth performance of common catfish (*Ameiurus melas* Raf.) fingerlings fed mealworm (*Tenebrio molitor*) diet. *J Insects Food Feed* 1:233-240.
- Ruby MB, Rozin P, Chan C. 2015. Determinants of willingness to eat insects in the USA and India. *J Insects Food Feed* 1:215-225.
- Ruddle K. 1973. The human use of insects: Examples from the Yukpa. *Biotropica* 5:94-101.
- Rumpold BA, Schluter OK. 2013a. Nutritional composition and safety aspects of edible insects. *Mol Nutr Food Res* 57:802-823.

- Rumpold BA, Schluter OK. 2013b. Potential and challenges of insects as an innovative source for food and feed production. *Innov Food Sci Emerg Technol* 17:1-11.
- Schiavone A, Cullere M, De Marco M, Meneguz M, Biasato I, Bergagna S, Dezzutto D, Gai F, Dabbou S, Gasco L, Zotte AD. 2017. Partial or total replacement of soybean oil by black soldier fly larvae (*Hermetia illucens* L.) fat in broiler diets: Effect on growth performances, feed-choice, blood traits, carcass characteristics and meat quality. *Ital J Anim Sci* 16:93-100.
- Schluter O, Rumpold B, Holzhauser T, Roth A, Vogel RF, Quasigroch W, Vogel S, Heinz V, Jager H, Bandick N, Kulling A, Knorr D, Steinberg P, Engel KH. 2017. Safety aspects of the production of foods and food ingredients from insects. *Mol Nutr Food Res* 61:1600520.
- Schosler H, De Boer J, Boersema JJ. 2012. Can we cut out the meat of the dish? Constructing consumer-oriented pathways towards meat substitution. *Appetite* 58:39-47.
- Siulapwa N, Mwambungu A, Lungu E, Sichilima W. 2014. Nutritional value of four common edible insects in Zambia. *Int J Sci Res* 3:876-884.
- Smeathman H. 1781. Some account of the termites, which are found in Africa and other hot climates. In a letter from Mr. Henry Smeathman, of Clement's Inn, to Sir Joseph Banks, Bart. *P. R. S. Philos Trans R Soc Lond* 71:17810033.
- Sogari G. 2015. Entomophagy and Italian consumers: An exploratory analysis. *Prog Nutr* 17:311-316.
- Sun-Waterhouse D, Waterhouse GIN, You L, Zhang J, Liu Y, Ma L, Gao J, Dong Y. 2016. Transforming insect biomass into consumer wellness foods: A review. *Food Res Int* 89:129-151.
- Sungpuag P, Puwastien P. 1983. Nutritive value of unconventional protein source: Insect. *Pochanagan Sarn* 1:5-12.
- Swamy HV, Devaraj KV. 1994. Nutrient utilization by common carp (*Cyprinus carpio* Linn) fed protein from leaf meal and silkworm pupae meal based diets. *Indian J Anim Nutr* 11:67-71.
- van Huis A, Ooninx DGAB. 2017. The environmental sustainability of insects as food and feed. A review. *Agron Sustain Dev* 37:43.
- van Huis A, van Itterbeeck J, Klunder H, Mertens E, Halloran A, Muir G, Vantomme P. 2013. Edible insects: Future prospects for food and feed security. Food and Agriculture Organization of the United Nations, Rome.
- van Huis A. 2015. Edible insects contributing to food security? *Agric Food Secur* 4:20.
- van Huis A. 2016. Edible insects are the future? *Proc Nutr Soc* 75:294-305.
- van Thielen L, Vermuyten S, Storms B, Rumpold B, van Campenhout L. 2019. Consumer acceptance of foods containing edible insects in Belgium two years after their introduction to the market. *J Insects Food Feed* 5:35-44.
- Veldkamp T, Bosch G. 2015. Insects: A protein-rich feed ingredient in pig and poultry diets. *Anim Front* 5:45-50.
- Vercruyse L, Smaghe G, Beckers T, van Camp J. 2009. Antioxidative and ACE inhibitory activities in enzymatic hydrolysates of the cotton leafworm, *Spodoptera littoralis*. *Food Chem* 114:38-43.
- Wang L, Li J, Jin JN, Zhu F, Roffeis M, Zhang XZ. 2017. A comprehensive evaluation of replacing fishmeal with housefly (*Musca domestica*) maggot meal in the diet of Nile tilapia (*Oreochromis niloticus*): Growth performance, flesh quality, innate immunity and water environment. *Aquac Nutr* 23:983-993.
- Wang Y, Chen Y, Li X, Xia J, Du Q, Zhi C. 1996. Study on the rearing larvae of *Tenebrio molitor* Linne and the effects of its processing and utilizing. *Acta Agriculture Universitatis Henensis* 30:288-292.
- Yen AL. 2009. Edible insects: Traditional knowledge or western phobia? *Entomol Res* 39:289-298.
- Yen AL. 2010. Edible insects and other invertebrates in Australia: Future prospects. In *Forest insects as food: Humans bite*

- back. Durst PB, Johnson DV, Leslie RN, Shono K (ed). FAO, Bangkok, Thailand. pp 65-84.
- Yhoung-aree J. 2010. Edible insects in Thailand: Nutritional values and health concerns. In Forest insects as food: Humans bite back. Durst PB, Johnson DV, Leslie RN, Shono K (ed). Food and Agriculture Organization of the United Nations, Bangkok, Thailand. pp 201-216.
- Yi L, Lakemond CMM, Sagis LMC, Eisner-Schadler V, van Huis A, van Boekel MAJS. 2013. Extraction and characterisation of protein fractions from five insect species. *Food Chem* 141:3341-3348.
- Yi L, van Boekel MAJS, Boeren S, Lakemond CMM. 2016. Protein identification and *in vitro* digestion of fractions from *Tenebrio molitor*. *Eur Food Res Technol* 242:1285-1297.
- Yun JH, Roh SW, Whon TW, Jung MJ, Kim MS, Park DS, Yoon C, Nam YD, Kim YJ, Choi JH, Kim JY, Shin NR, KIM SH, Lee WJ, Bae JW. 2014. Insect gut bacterial diversity determined by environmental habitat, diet, developmental stage, and phylogeny of host. *Appl Environ Microbiol* 80:5254-5264.
- Zielinska E, Baraniak B, Karas M, Rybczynska K, Jakubczyk A. 2015. Selected species of edible insects as a source of nutrient composition. *Food Res Int* 77:460-466.
- Zielinska E, Karas M, Baraniak B. 2018. Comparison of functional properties of edible insects and protein preparations thereof. *LWT-Food Sci Technol* 91:168-174.