



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

10

Adenovirus Vaccines

Gregory C. Gray and Dean D. Erdman

HISTORY OF DISEASE

In the 1950s, multiple researchers^{1,2} identified new viruses as the cause of acute respiratory disease (ARD), pharyngitis, conjunctivitis, and pneumonitis. Soon afterward, these viruses were recognized to be a related group of viruses and given their present name: adenoviruses.³

WHY THE DISEASE IS IMPORTANT

The adenoviruses are often the cause of respiratory illness among U.S. military trainees^{4,5} and children.^{6–13} In the 1960s, adenoviruses were recognized to infect as many as 80% of trainees, with 20% requiring hospitalization.⁵ Median attack rates for trainees ranged from 6 to 16.7/100 per month at the most affected northern and Midwestern U.S. military bases, and from 2.3 to 2.6/100 per month for posts in the South and West.¹⁴ Up to 40% of the trainees in a unit were lost to illness within a 2-week period, and many who were hospitalized had to restart training.¹⁵ Adenoviruses were also recognized to cause as much as 15% of instances of gastroenteritis in infants and children, and up to 10% of instances of pneumonia among hospitalized children.^{16–18} Adenovirus morbidity can be severe and lead to death, especially in young children, transplantation recipients, and other immunocompromised patients.

The search for vaccines against adenoviruses was driven by adenovirus morbidity among military trainees, their disruption to military training, and their medical care costs.¹⁵ In 1971, after a series of misstarts, military recruits began routinely receiving live oral enteric-coated vaccines for adenovirus types 4 and 7 (Ad4 and Ad7), which were safe and effective.¹⁹ With few exceptions, adenovirus morbidity was markedly controlled. However, in 1996, the sole manufacturer of these vaccines ceased production. As vaccine stores were depleted, U.S. military trainees again experienced large outbreaks of adenovirus disease, with some deaths reminding public health officials of the impact of adenoviruses.^{20–22} In one comprehensive 2006 study, Russell and colleagues documented 98% Ad4 attack rates among 180 susceptible military recruits during 4 weeks of training.²³ The importance of these pathogens has been further emphasized through the recent emergence of multiple novel Ad3, Ad7, and Ad14 strains and the subsequent epidemics and deaths they caused.^{24–29}

BACKGROUND**Clinical Description**

Human adenoviruses have been classically grouped into seven species (A through G) containing as many as 68 unique types (Ad1 through Ad68) and multiple more genotypes

often determined by restriction enzyme digests (Ad7a, Ad7d2, etc). However, the nomenclature and classification system is undergoing change. In 2013, the International Committee of Taxonomy of Viruses added a genus prefix (eg, human adenovirus A has become human mastadenovirus A).³⁰ A number of investigators have additionally argued that it is time to move away from neutralization and hemagglutination assays in determining serotype and to chiefly use a genetic classification approach.³¹ Such virologists have used large DNA sequence comparisons to describe as many as 68 novel adenovirus types,^{32,33} although this approach has met with some debate.^{34–37}

No matter which adenovirus nomenclature or classification system is agreed upon, it is clear that adenoviruses affect most organ systems. Individual adenovirus types often have different tissue tropisms that lead to specific signs and symptoms (Table 10.1).^{39,40}

Endemic Respiratory Adenovirus in Children

Although most children become infected with many of the common adenoviruses early in life, only approximately 50% of these infections result in disease.^{39,41,42} Isolation studies have indicated that Ad1, Ad2, Ad5, Ad3, and Ad6 are most prevalent types detected,^{39,43} and often manifest by pharyngitis, bronchitis, bronchiolitis, croup, or pneumonia.^{43,44} The incidence of adenovirus disease is higher in late winter, spring, and early summer, and both sexes are equally affected.^{41,45}

Epidemic Respiratory Adenovirus in Children

Occasionally, epidemics occur in daycare facilities, neonatal intensive care units, chronic care facilities, and orphanages, especially with Ad3 and Ad7.^{46–49} In addition to causing outbreaks in closed populations, Ad7 can cause large epidemics affecting children in open communities.^{50,51}

Acute Respiratory Disease of Military Recruits

In young adults who live in closed communities such as boarding schools and military recruit camps, adenoviruses may cause epidemics of illnesses similar to influenza, including tracheobronchitis and pneumonia severe enough to require hospitalization.^{2,4,52,53} Adenovirus pneumonia may be fatal, especially when it is associated with Ad7 strains.^{16,54–56} The incubation period of the disease is 4 to 5 days.⁵³ Before the use of vaccines in the U.S. military, Ad4 and Ad7 accounted for 90% of all respiratory illnesses among recruits who were hospitalized, whereas Ad3, Ad14, and Ad21 were less frequently observed.⁴ At some northern basic training sites, rates of 6 to 8 per 100 trainees per week translated into 600 to 800 ARD hospital admissions per week.¹⁵ Recently, the newly

TABLE 10.1 Clinical Syndromes Associated With Human Adenovirus Infections

Clinical Syndrome	Common Types	Population at Risk
Endemic respiratory	1, 2, 3, 5, 6	Infants, children
Epidemic respiratory	7, 14, 55	Children
Acute respiratory disease	3, 4, 7, 14, 21	Military recruits
Pharyngoconjunctival fever	1, 3, 4, 7, 14	School-age children, young adults
Keratoconjunctivitis	8, 11, 19, 37, 53, 54	All age groups
Hemorrhagic cystitis	11, 34, 35	Immunocompromised patients, children
Gastroenteritis	31, 40, 41, 52	Children, immunocompromised patients
Other syndromes	2, 4, 7, 12, 19, 32, 37	Children, adults
Immune deficiency	34, 35, 43–49, 50, 51	Transplantation recipients, persons with AIDS, and immunocompromised patients

There is agreement regarding the classification of adenoviruses 1 to 51, which have been classically typed with neutralization and hemagglutination assays. However, the classification of adenoviruses 52 to 68 remains controversial as these have been chiefly identified through genetic sequencing. In particular, adenovirus 55 is a recombination of 11 and 14.^{126,148}

recognized Ad55 (Ad11 and Ad14 recombinant) has been documented to have caused significant ARD morbidity among military trainees in China.⁵⁷

Typical ARD is a febrile disease with symptoms of sore throat, fever, cough, coryza, rhinorrhea, headache, and chest pain.^{2,39,58} With extension into the lungs, physical examinations can reveal rales and rhonchi with little evidence of consolidation, and chest radiography often shows patchy interstitial infiltrates, principally in the lower lung fields.^{2,53,59} Symptoms last 3 to 10 days.³⁹ The infection is often self-limited, but occasionally deaths occur.^{2,60} Routes of transmission are thought to include direct contact or aerosolized virus inhaled into the lung. The virus has been isolated from the oropharynx more than 2 weeks after exposure.^{2,61} In a 2006 report, Russell and colleagues demonstrated viable adenovirus was ubiquitous on surfaces in the military trainee barracks.²³

Pharyngoconjunctival Fever

This syndrome is characterized by pharyngitis, conjunctivitis, and spiking fever.⁶² First described in the 1920s as associated with swimming, the cause has been subsequently linked to insufficient chlorination.^{63–65} One or both eyes are affected, and diarrhea, coryza, tonsillar exudates, and lymphadenopathy may be observed. The most frequently identified types have been Ad3 and Ad7, but other types, such as Ad1, Ad4, and Ad14, are also associated with pharyngoconjunctival fever.⁶² The disease is associated with summer camps, swimming pools, and lakes; occurs in children and young adults; and often spreads to other family members.^{62,66,67} The incubation period is 6 to 9 days, and the virus may be isolated from pool water.^{62,65} There is little bacterial superinfection and no permanent damage to the eye.⁵⁹

Epidemic Keratoconjunctivitis

Epidemics of conjunctivitis in adults caused by adenoviruses were first described as “shipyard eye.”^{68,69} The disease was observed at industrial settings where shipbuilding took place and was probably transmitted because of inadequate infection control practices when workers sought care for chemical irritation and minor trauma from paint and rust chips.⁶⁸ The disease has an incubation time of 8 to 10 days and is characterized by conjunctivitis, edema of the eye, pain, photophobia, and lacrimation. Superficial erosions and subepithelial infiltrates of the cornea may occur.³⁹ Preauricular lymph gland swelling and involvement of cervical and submaxillary lymph

glands may be observed.³⁹ Epidemic keratoconjunctivitis (EKC) is associated with Ad8, Ad19, Ad37, and, rarely, other types.^{70–74} In addition, Ad19 and Ad37 have been isolated from the genital tract of young adults with EKC, and the possibility of sexual transmission has been considered.^{75,76} Numerous epidemics of EKC have been associated with ophthalmology practices, where spread from contaminated ophthalmic solutions, fingers, and instruments has been implicated.^{70,72,74,77,78}

Hemorrhagic Cystitis

Hemorrhagic cystitis syndrome in children is caused by adenovirus infections in 23% to 51% of cases in the United States and Japan.⁷⁹ Although the specific route of spread is unknown, Ad11 and Ad21 were the most frequently isolated adenoviruses.⁷⁹ Boys were two to three times more often affected than girls. Clinical findings included gross hematuria of 3 days' duration. Dysuria, microscopic hematuria, and urinary frequency lasted a few days longer.^{79,80} No viremia or structural abnormalities were found. Adenoviral antigen in exfoliated bladder epithelial cells can be demonstrated by immunofluorescence.⁷⁹ Cases of acute hemorrhagic cystitis after renal and bone marrow transplantation are increasingly being reported.^{81–83} Ad34 and Ad35, as well as species B adenoviruses, were first isolated from renal transplantation recipients, but neither was associated with symptoms of hemorrhagic cystitis.⁸⁴

Gastroenteritis

Adenoviruses are also recognized to cause watery diarrhea with fever. Ad40 and Ad41 are the most prevalent strains, but Ad50, Ad51, and Ad52 have been detected among the immunocompromised. AdV52 has been associated with one outbreak cluster of acute gastroenteritis.¹²⁴ These strains often require special cells lines for culture and they may not demonstrate cytopathic effect.⁸⁵ Electron microscopy or special assays are often necessary for their study.⁸⁵ Numerous outbreaks have been described, and adenoviruses may account for up to 12% of all infant diarrhea.^{86–88}

Rare Acute Manifestations

A number of case or outbreak reports document other, more unusual manifestations of human adenovirus infection. These occur most frequently among children. Although the studies

are inconsistent, adenovirus is implicated as a possible fetal pathogen.^{89,90} Adenovirus infections among children are associated with sudden infant death,⁹¹ encephalitis, meningoencephalitis,^{92,93} cerebral edema,⁹⁴ acute flaccid paralysis,⁹⁵ pertussis-like syndromes,^{96,97} mononucleosis-like syndromes,⁹⁸ and neonatal disseminated infections.⁹⁹ Among adults, adenovirus infections have been associated with a toxic shock–like syndrome,¹⁰⁰ encephalitis, genital lesions, orchitis, urethritis, and cervicitis.¹⁰¹ Nosocomial transmission to susceptible healthcare workers and patients has been reported. This is likely related to the long periods of viral shedding among adenovirus-infected immunocompromised hosts, the possible aerosolization of the virus, and fomite transmission in the hospital setting.^{40,95,102}

Complications

Adenovirus Infections in Immunocompromised Patients

Adenoviruses have been implicated as opportunistic agents in patients with immune deficiency states, such as patients with AIDS, patients receiving cancer chemotherapy, bone marrow transplant recipients, and patients undergoing solid organ transplantation.^{11,12,82,103–106} The patients are prone to pneumonia and disseminated adenoviral infection. They have also developed parotitis¹⁰⁷ and urinary tract disease.¹⁰⁸ A number of the more recently described types, such as Ad43 through Ad51, have been recovered from people with AIDS.^{108,109}

A review of 201 bone marrow recipients over a 4-year period indicated that adenovirus infections occurred in 20.9% of patients, with a higher incidence in children than in adults (31.3% vs 13.6%).¹¹⁰ Ad35 was the most common type identified.¹¹⁰ In 1992, Hierholzer reported that adenovirus infections in immunocompromised patients with pneumonia were associated with case fatality rates as high as 60%, compared with only 15% among immunocompetent hosts with pneumonia.⁹ Radiographs of adenovirus-infected patients often demonstrate patchy interstitial infiltrates, usually in the lower lung fields.¹¹¹ Fig. 10.1 shows the chest radiograph of a 20-year-old recipient of an autologous bone marrow transplant for diffuse large cell lymphoma (B-cell type) with systemic Ad11




Figure 10.1. Pneumonia caused by adenovirus type 11 in a 20-year-old recipient of an autologous bone marrow transplant. (Courtesy Stuart Ray, MD, Johns Hopkins University, Baltimore, MD.)

infection, which was obtained on day 45 posttransplantation. He died on day 80. Adenovirus was recovered from conjunctival, urine, and bronchoalveolar lavage cultures. A case fatality rate of 50% occurred in immunocompromised people with hepatitis and associated adenovirus infections, compared with 10% in similarly infected immunocompetent patients with hepatitis.⁹ Positive tests of serum by polymerase chain reaction (PCR) for adenovirus has been reported to predict severe disseminated adenovirus infection in immunocompromised patients.¹¹² Nephritis and kidney failure caused by adenovirus infections have been reported in adults and children receiving bone marrow transplants.^{113,114} A study of 532 hematopoietic stem cell transplant recipients between 1986 and 1997 found a 12% incidence of adenovirus infections, with children being more likely than adults to have a positive culture (23% vs 9%).¹¹⁵

Chronic Diseases Caused by Adenoviruses

Thanks often to molecular techniques, human adenoviruses have been associated with a number of chronic disease conditions, including chronic airway obstruction,¹⁰¹ pulmonary dysplasia,¹¹⁶ myocarditis, and cardiomyopathy.¹¹⁷ Although the results of studies are mixed, an intriguing series of human and animal studies associate a number of human adenovirus types with obesity.^{118–121}

Virology

Human adenoviruses are nonenveloped, double-stranded DNA viruses belonging to the genus Mastadenovirus, family Adenoviridae.¹²² They are classified by species (A–G) through hemagglutination properties and by serotype (1–51) with horse or rabbit antisera (Table 10.2).^{40,59,123} Different strains within serotypes are further distinguished by whole-genome restriction-enzyme digest patterns. However, as typing serum samples are difficult to acquire and sequencing costs are decreasing, virologists are considering full genome sequencing as a more modern and accurate method to detect unique strains and have described as many as 68 unique adenovirus types.^{124–128}

Adenoviruses are considered large viruses, with an estimated diameter of approximately 920 Å.¹²⁹ The viruses have an icosahedral capsid shell that is made up of 240 hexon bases, 12 penton bases, and 12 fibers that are associated with the penton bases (Fig. 10.2). Four minor proteins (IIIa, VI, VIII,

TABLE 10.2 Human Adenovirus Species and Types

Adenovirus Species	Types
A	12, 18, 31, 61
B	3, 7, 11, 14, 16, 21, 34, 35, 50, 55, 66
C	1, 2, 5, 6, 57
D	8–10, 13, 15, 17, 19, 20, 22–30, 32, 33, 36–39, 42–49, 51, 53, 54, 56, 58–60, 63–67
E	4
F	40, 41
G	52

Some information for the recently described adenovirus types is not yet available.

Modified from Lion T. Adenovirus infections in immunocompetent and immunocompromised patients. Clin Microbiol Rev. 2014;27(3):441–462.




Figure 10.2. Image of adenovirus virion determined by cryoelectron microscopy showing newly understood interactions of the major components (hexon, penton, and fiber proteins) and minor proteins (IIIa, VIII, and IX). The proteins interact in groups of nine (GON) and groups of six (GOS) capsomers. (Courtesy Hong Zhu, University of California, Los Angeles. Modified from Harrison SC. *Looking inside adenovirus*. *Science*. 2010;329:1026–1027.)

and IX) add to the complexity of the capsid. The pentons and hexons are each derived from different viral polypeptides.

Adenovirus hexons are both type-specific and species-specific antigens, primarily inducing species-specific complement-fixing antibodies, whereas the pentons are especially active in hemagglutination.¹³⁰ The fibers also evoke type-specific antibodies, vary in length among human strains, and are sometimes absent in particular animal strains.^{131–134} The genome core of the virus is composed of five more proteins (V, VII, u, Iva2, and terminal protein) and a single molecule of linear, double-stranded DNA of 26×106 to 45×106 molecular weight. The G + C base compositions of the human virus genomes range from 47% to 60%.^{110,122,135,136}

Adenoviruses are unusually stable to physical and chemical agents, as well as adverse pH, and, thus, survive for long periods outside the host, making them available for transmission to others. They can be destroyed by heat at 56°C for 30 minutes, UV irradiation, 0.25% sodium dodecyl sulfate, chlorine at $0.5 \mu\text{g}/\text{mL}$, and formalin, but are resistant to ether and chloroform.

Adenoviruses replicate in the cell nucleus and tend to be very host-specific. However, there is mounting evidence that humans and animals have exchanged adenoviruses. In 2009, a simian adenovirus epidemic occurred among nonhuman primates at a U.S. research center and subsequently infected the primate caretakers with some evidence of human-to-human transmission.^{137,138} A 2011 report documented the recent detection of 45 distinct adenovirus in nonhuman primates most of which mapped phylogenetically very closely to human adenovirus species.¹³⁹ In 2012, human infections with a novel avian adenovirus were associated with avian chlamydiosis outbreak in Hong Kong.¹⁴⁰ Finally, it is also important to recognize that Ad4 and Ad52 are thought to have

originated from nonhuman primates.^{124,141,142} Hence, some virologists have recently reviewed the probability that adenoviruses may be zoonotic.¹⁴³

Some adenovirus types have been determined to be oncogenic in animals and to transform cell lines, but oncogenicity has not been observed in humans. Hybridization or genetic recombination of multiple strains of adenovirus may occur. Sometimes, these recombinant strains or other emergent novel adenovirus strains lead to epidemics. A number of recent examples have been reported in the medical literature.^{26,27,29,144–148}

Pathogenesis as It Relates to Prevention

Depending on the route of inoculation, the type of virus, and the immune state of the host, adenoviruses can cause diseases or asymptomatic infections. Respiratory infection is presumed to result from inhalation of aerosolized virus, whereas ocular infection, gastroenteritis, and nosocomial infections may arise from fomites, water, or fecal–oral contact. Reactivation of latent adenovirus is also believed to occur.⁵³ Some 50% to 80% of tonsils removed surgically may have adenoviruses isolated from the tissue, suggesting that these viruses may remain in a latent state for years.^{149,150} Virus has been also isolated from lymphocytes, kidney, blood, cerebrospinal fluid, and most body organs.^{9,11,12,54,55,93,151–153} In the lungs, extensive pathology has been found with microscopic necrosis of the tracheal and bronchial epithelium.³⁹ Acidophilic intranuclear inclusions are seen in bronchial epithelial cells in addition to the basophilic masses of cells surrounded by clear halos, which may indicate aggregations of viral material.⁵⁴ A mononuclear infiltrate, rosette formation, and focal necrosis of mucous glands are characteristically seen.

Three types of interaction of adenovirus with infected cells can occur. A lytic infection may take place during which the virus completes an entire replicative cycle.⁵⁹ Between 10^5 to 10^6 progeny viruses per cell are produced, of which only 1% to 5% are actually infectious.⁵⁹ The second type of interaction is chronic or latent infection, in which small amounts of virus are produced, and an inapparent infection results. Viral shedding from the gastrointestinal tract may occur for years.³⁹ In fact reactivation of such latent infection likely explains much adenovirus disease in the severely compromised. Correspondingly, monitoring viral loads in the stool of transplant patients has predicted disseminated disease and been useful in guiding antiviral therapy.¹⁵⁴

In addition to aerosolization, intestinal shedding of respiratory virus is an important factor to consider in the prevention of nosocomial spread in hospitals and chronic care homes.^{39,145,155} Persistent infection has been reported in epithelial cells from monkeys.¹⁵⁶ Lymphoid cells are thought to be the reservoir for these persistent infections.^{157,158} The third type of interaction is oncogenic transformation, whereby the viral DNA is integrated into the host genome and replicated with the cellular host DNA, but only the early steps in the viral cycle occur and no infectious virions are produced.¹⁵⁹

The genes from adenoviruses are expressed in the cell nucleus in two phases: “early” (E), which precedes viral DNA replication, and “late.”¹⁵⁷ Early genes encode proteins that function to thwart immunosurveillance, especially those from the E3 transcription unit.¹⁵⁷ The late genes primarily encode viral structural proteins. The functions of the E1 proteins include the induction of DNA synthesis in quiescent cells, immortalization of primary cells in cooperation with activated *ras* or with the E1B proteins, transactivation of delayed early genes, induction or repression of several cellular genes, and induction of apoptosis. These proteins modulate the sensitivity of adenovirus-infected cells to tumor necrosis factor (TNF),

a key inflammatory cytokine with antiviral properties.¹⁵⁷ None of the E3 genes are required for adenovirus replication in cultured cells, but several of the E3-coded proteins (10.4 K, 14.5 K, and 14.7 K) inhibit TNF cytotoxicity.^{157,160,161} Because a major function of TNF may be to prevent viral replication, the inhibition of TNF by these viral proteins may be a significant mechanism of pathogenesis.

Another significant E3-coded protein is Gp 19 K. This glycoprotein is located in the endoplasmic reticulum and forms a complex with class I antigens of the major histocompatibility complex (MHC), preventing cells from being killed by cytotoxic T lymphocytes.¹⁵⁷ A cotton rat animal model was used by Ginsberg and colleagues and Ginsberg and Prince to study the pathogenesis of Ad2 and Ad5, which cause pneumonia similar to that seen in humans.^{160,162} Two phases of infection were seen; the initial phase, characterized by the infiltration of monocytes and neutrophils, and a later phase associated with the infiltration of lymphocytes. The pathology seemed to reflect the response by host immune defenses to viral infection. The Gp 19 K markedly reduced the transport of the class I MHC to the surface of the infected cells and impeded the attack of cytotoxic T cells.^{157,160,161} It is now known that only the early genes are required to induce the complete pathogenesis of adenovirus infection in cotton rats.¹⁶¹ Although several cytokines, such as TNF- α , interleukin-1, and interleukin-6, were elaborated during the first 2 to 3 days of the infection in the cotton rat model, only TNF- α had a major role in pathogenesis.¹⁶¹ Steroids almost completely eliminated the pneumonic inflammatory response to infection.¹⁶¹

Pathology caused by latent infection with adenoviruses has been linked to chronic obstructive pulmonary disease (COPD).^{158,162} Some have suggested that childhood viral diseases represent an independent risk factor for COPD.¹⁶³ The adenoviral E1A proteins can stimulate the transcription of many heterologous viral and cellular genes. These proteins have the ability to interact with the DNA binding domains of several cellular transcription factors and activate a wide variety of genes.^{164,165} The adenoviral genome has been found to be present in the lungs of more patients with COPD than in control subjects.¹⁶⁶ E1A proteins are expressed in epithelial cells of human lung tissue, and by increasing the expression of several genes important in controlling the inflammatory process, these may contribute to the pathogenesis of COPD. The events described may amplify the airway inflammation associated with cigarette smoking.¹⁶¹

The isolation and cloning of a 46-kDa protein adenovirus receptor, which mediates attachment and infection of group B viruses, may facilitate the development of new strategies to limit diseases caused by adenoviruses.¹⁶⁷ The more common receptor is referred to as CAR. This protein has been identified as the receptor for the coxsackie B virus and adenoviruses. Recently, another receptor, desmoglein 2, was identified for Ad3, Ad7, Ad11, and Ad14 strains.¹⁶⁸

Laboratory Diagnosis

Adenoviral infections generally cannot be diagnosed on clinical grounds alone because the clinical signs and symptoms of these infections are variable and often resemble those caused by other microorganisms.¹⁰¹ Traditionally, laboratory support and qualified personnel are necessary to accurately diagnose adenovirus infections. Considerations include specimen type and timing, collection and storage procedures, types of laboratory tests performed, including serological assays, and availability of newer types of diagnostic assays.

Specimen Collection. The optimal specimen for diagnosis of adenovirus infection depends on the clinical presentation

and the suspected type.¹⁰¹ Adenoviruses can be detected in a variety of specimens, including nasal and throat secretions, conjunctival scrapings and swabs, stool, blood, cerebrospinal fluid, and biopsied tissue specimens. Specimens should be collected early in the illness and shipped promptly at 4°C or on dry ice if prompt testing is not possible. Swab and fresh tissue specimens must be transported in appropriate viral transport media^{40,58,84,101} whereas fluid samples like urine, stool, and cerebrospinal fluid should not be diluted in transport media. For culture, nonsterile specimens should be treated with antibiotics prior to inoculation. Stool specimens are brought to 10% to 20% suspensions with buffered saline and clarified by low-speed centrifugation. Blood for culture or molecular testing should be collected with an anticoagulant to prevent clotting. Serum separated from clotted blood samples can be used for serological diagnosis (see below).

Cell Culture. Through biological amplification, cell culture offers a sensitive method for adenovirus detection by monitoring for cytopathic effect (CPE). Because adenoviruses are host-specific, isolation of human adenoviruses is most easily accomplished in human cells.¹⁶⁹ Most human adenoviruses have been successfully isolated in cell culture. Several diploid and continuous cell lines, including A549, HeLa, HEP-2, KB, MRC-5 and commercial mixed cell lines with A549 cells, reportedly give good overall recovery and produce typical CPE.^{170,171} However, the fastidious enteric adenoviruses Ad40 and Ad41 are an exception. For efficient recovery, most Ad40 and Ad41 strains require the Graham-293 Ad5-transformed secondary HEK (human embryo kidney) cell line for primary isolation.

CPE may develop slowly in monolayers of inoculated cells necessitating several subpassages before becoming visible. Infected cells become rounded, enlarged, and refractile and aggregate into irregular "grape-like" clusters.⁴⁰ A 4-week incubation with blind passage is recommended. A rapid procedure for culture identification of adenovirus is the shell vial technique. Here, cell monolayers prepared in glass vials or multi-well plastic culture plates are inoculated with the clinical specimen, centrifuged, and stained after 2 to 3 days incubation with commercial monoclonal antibodies.¹⁷² Cell culture is declining as a routine diagnostic method for detection and identification of human adenoviruses as a result of cost and delays. Antigen and molecular detection techniques have supplanted culture in many laboratories (see below).¹⁶⁹ However, while direct detection methods offer important diagnostic benefits, culture is the only means available for obtaining sufficient virus for immunotyping, assessing virus susceptibility to antivirals, and identifying infectious virus in clinical and environmental specimens.

Direct Visualization of Virus Particles

Electron and immunoelectron microscopy¹⁷³ were once commonly used to detect and identify the fastidious adenoviruses in stool specimens from young children, thereby establishing their association with acute gastroenteritis.^{40,102,174-177} These methods are still used to localize virus in cells from biopsy and autopsy specimens to confirm disease association and to study disease pathology.

Antigen Assays. Detection of adenovirus antigenic proteins can be accomplished by reacting with virus-specific polyclonal or monoclonal antibodies labeled with various reporter systems. Examples include enzyme immunoassay, immunofluorescence assay, time-resolved fluoroimmunoassay, and latex agglutination.¹⁷⁸⁻¹⁸¹ These assays can be used to detect adenovirus proteins directly in clinical specimens or to

identify virus isolates. Immunoassays are simple to perform, less costly and provide results more quickly than culture or molecular tests, but are generally less sensitive.^{181,182} The immunofluorescence assay is often used in the clinical diagnostic laboratory for rapid identification of adenovirus in cells from unfrozen specimens. Point-of-care antigen immunoassays^{181,183} for rapid adenovirus detection and immunohistological staining for adenovirus antigens in biopsy and autopsy specimens are still in common use^{182,184}

Molecular Assays. Molecular methods are rapidly replacing classical culture and antigen detection for diagnosis of adenovirus infection. The PCR assay in particular has become a popular diagnostic alternative for these viruses, offering the potential for rapid and sensitive detection and being easily tailored for species-specific and type-specific identification as described below. PCR assays¹⁸⁵ using adenovirus group-specific primers individually,^{185,186} or combined in assays for multiple human pathogens¹⁸⁷ have proven comparable or better than cell culture or immunodiagnostic methods for detection of adenoviruses in clinical samples. Several commercial multiplex assays for respiratory pathogens including adenovirus have been cleared by the U.S. Food and Drug Administration (FDA) for clinical diagnostic use.¹⁸⁸ Use of quantitative real-time PCR assays have the added benefit of allowing monitoring of adenovirus levels that can be used to predict patient prognosis, manage chemotherapy and monitor efficacy of antiviral therapy.¹⁸⁹

Adenovirus Typing Assays. Once detected, further identification of adenoviruses to species and type can be accomplished by a variety of immunologic and molecular methods to support outbreak investigations and patient clinical management. For example, adenovirus infections in bone marrow or lung transplant recipients can be rapidly fatal. It is in the patients' interest that healthcare providers determine the species and type of any infecting adenovirus as some are more sensitive to specific antiviral therapy.¹⁹⁰ It is also important to learn if the adenovirus represents a community-acquired strain, a healthcare-acquired strain, a donor-associated strain, or the reactivation of a latent virus. Efforts to further identify adenoviruses have led to the discovery of cross-species infections with simian adenoviruses acquired from nonhuman primate reservoirs.^{138,191}

Serum neutralization (SN) using defined hyperimmune animal antisera is the gold standard method used for classifying human adenovirus types.⁴⁰ SN targets type-specific neutralizing epitopes on the adenovirus hexon and fiber proteins and to a lesser extent on the penton protein. Hemagglutination inhibition (HI) has also been used of immunotyping adenoviruses. HI targets type-specific epitopes on the adenovirus fiber protein, and when used in combination with SN, can identify hexon/fiber recombinant viruses. Because immunotyping methods require reference antisera that are difficult to obtain and standardize and often difficult to interpret, molecular methods have generally replaced immunotyping for rapid routine adenovirus typing.

Numerous molecular methods for species and type-specific identification of adenoviruses have been described, including conventional^{192–194} and real-time PCR^{195,196} assays using species-specific and type-specific primers and probes, PCR amplification coupled with restriction endonuclease analysis,^{197,198} microarrays,¹⁹⁹ and mass spectrometry²⁰⁰ and, increasingly, PCR-coupled sequence-based typing strategies that target hypervariable regions of the adenovirus hexon or fiber genes or both.^{194,201–208} Several commercial assays have been marketed that use PCR to identify human adenovirus to species.^{209–211} As molecular typing strategies often focus only

on specific hypervariable regions of the virus genome, potentially important virus strains resulting from recombinations or mutations can be missed. McCarthy and colleagues proposed a molecular typing algorithm that combined examinations of the hexon and fiber genes.²¹²

Restriction enzyme analysis (REA) of whole-virus genomes can be used to identify genetic variants of a serotype, aiding epidemiologic studies and outbreak investigations.^{213,214} Extensive studies of Ad3, Ad4, and Ad7 using REA have been described^{20,215–218} and a number of novel viruses have thus been identified.²⁶ However, REA is work intensive, dependent on virus isolation to recover genomic DNA and requires expertise to perform and interpret, and is therefore restricted to a few experienced laboratories.

Classical and next generation sequencing technologies are increasingly being used to obtain full genomic sequences of human adenoviruses. Genome sequences are available for all human adenovirus prototype types and sequences of newly identified genomic variants are now routinely placed in public domain providing reference data for adenovirus identification.¹²⁶ A new whole genome-based characterization system for adenoviruses is under consideration to replace immunotyping for designating new adenovirus types.^{31,36}

Serological Assays. Acute adenovirus infection can also be diagnosed by serological demonstration of a fourfold or greater rise in antibody titer between paired acute-phase and convalescent-phase sera. The first acute specimen should be collected as early as possible after onset of illness and the convalescent specimen should be collected 2 to 4 weeks later. Serologic tests for adenoviruses have included complement fixation (CF), HI, SN, indirect immunofluorescence, and enzyme immunoassay (EIA).¹⁶⁹ The classical CF test measures adenovirus-group specific antibodies, and though once widely used, lacks sensitivity. The EIA also measures adenovirus group-specific antibodies, but is more sensitive than the CF test and is readily automated and therefore more widely used today.^{40,101,219} HI and SN assays also are more sensitive than CF, but because they measure type-specific antibodies, they are not suitable for routine diagnosis and most reagents are not commercially available.⁴⁰ Although valuable for outbreak investigations and essential for seroprevalence and vaccine efficacy studies, serology is rarely used today for clinical diagnosis because of the inconvenience of serum collection and the 2- to 4-week delay needed to obtain convalescent serum.

Treatment and Prevention With Antimicrobials

Currently, there are no FDA-approved treatment or prophylaxis therapies against adenovirus infections. There are numerous off-label therapeutic case or case series reports, but their results are mixed. There are few, if any, controlled, prospective therapeutic studies.²²⁰ In the 1960s, γ -globulin was studied as prophylaxis against adenovirus infections without good effect.²²¹ However, Maisch and colleagues found a positive effect of intravenous immunoglobulin therapy in patients with adenovirus-associated inflammatory cardiomyopathy.²²² Some studies suggest that interferon- β may have a beneficial effect on EKC and viral cardiomyopathies.^{223,224} More recently, adoptive transfer of adenovirus-specific T-cells has been explored with some success but this is now reserved for immunocompromised patients who do not respond to antiviral therapy.²²⁵

Off-label antiviral use reports of the antivirals ribavirin, cidofovir, ganciclovir, acyclovir, and vidarabine^{220,226} have also been disappointing. Only ribavirin and cidofovir have been used often enough to be worthy of discussion. Nebulized ribavirin was used with some success for two children with pneumonia caused by adenovirus.²²⁷ Intravenous ribavirin has

been reportedly used in immunocompromised patients and in patients with hemorrhagic cystitis and disseminated disease.^{219,228–231} Intravenous ribavirin was given to 12 adult blood and marrow transplant recipients without appreciable benefit.²³² The mixed reports may be explained by a recent finding that only human adenoviruses of species C are susceptible to ribavirin.¹⁹⁰ In contrast, all species of human adenovirus appear to be susceptible to cidofovir.¹⁹⁰ Perhaps this finding explains the better results with this antiviral drug. In a retrospective study of 45 bone marrow transplant recipients treated with cidofovir for adenovirus infections, 31 (69%) had successful outcomes; however, in 18 (40%) of the 45 patients, cidofovir-associated toxicity developed.²³³

A new experimental oral antiviral therapy against DNA viruses, brincidofovir (formerly CMX001), shows great promise against adenoviruses.^{225,234,235} An analog of cidofovir, it does not have cidofovir's renal toxicity profile. Currently, brincidofovir is being evaluated against DNA viruses including adenoviruses. It has been awarded "Fast Track Designation" by the FDA.

EPIDEMIOLOGY

Incidence and Prevalence Data

Because only approximately 50% of childhood adenovirus infections result in disease, prevalence as detected by antibody studies is high.^{42,236,237} Those in species C (Ad1, Ad2, Ad5, and Ad6) are usually endemic and acquired in early childhood, often before 2 years of age.³⁹ By school age, most children have been exposed to several types of adenoviruses. Infections caused by Ad4, Ad7, Ad14, and Ad21 may occur at a later age. Many of the other types occur sporadically or in epidemics. Many adenovirus infections are subclinical or asymptomatic, especially those in species A and D. Conversely, Ad4, Ad7, and Ad21 usually cause symptomatic respiratory disease.³⁹

Most types of adenoviruses are thought to be endemic worldwide.⁴⁵ In 1983, Schmitz and colleagues⁴⁵ summarized 10 years of adenovirus reports sent to the World Health Organization. They noted an increasing prevalence of Ad7, Ad8, and Ad19 and a declining prevalence of Ad3 and Ad4.⁴⁵ Age predilections were highly significant for infants for species A (Ad12, Ad18, and Ad31); for infants and small children for species C (Ad1, Ad2, Ad5, and Ad6); for school children for Ad3; for school children and adults for Ad7; and for adults for Ad4, Ad8, and other types of species B, D, and E.⁴⁵ A predilection for males was observed for all types in species B and C and in Ad4 and Ad19.⁴⁵

A summary of 2237 clinical adenovirus submissions to 22 U.S. laboratories during a 25-month period (2004–2006) revealed that Ad3 (prevalence: 34.6%), Ad2 (prevalence: 24.3%), Ad1 (prevalence: 17.7%), and Ad5 (prevalence: 5.3%) explained most illnesses among children. The authors noted an increasing prevalence of Ad21 over time.¹⁷⁷

Even though most types of adenovirus can be found throughout the world, sometimes marked differences are observed in the geographical distribution of specific genotypes. Of particular note are the distributions of Ad3, Ad4, and Ad7 genotypes, with anecdotal suggestions that some genotypes may be more virulent and have competitive advantage.²³⁸ As yet, no clear markers of increased adenovirus virulence have been identified.

In 2009, Lebeck and colleagues²⁶ reported on 516 clinical Ad3 isolates from 15 U.S. laboratories with restriction enzyme digests. The most prevalent types were Ad3a2 (36.9%), Ad3a50 (27.1%), Ad3a51, (18.0%), and Ad3a17 (4.6%). Ad3a50 and Ad3a51 were newly described strains that became more prevalent in 2006. Multivariable modeling revealed that children

younger than 2 years; persons with chronic disease; and persons infected with Ad3a2, Ad3a50, or multiple or rare strains were at increased risk of severe Ad3 clinical disease. Similarly, beginning in 1986, Chile, Uruguay, and Argentina experienced the emergence of a novel Ad7h strain that supplanted the endemic Ad7c strains.^{239,240} Among pediatric patients, the Ad7h strain was associated with longer hospitalizations, higher patient temperatures, greater oxygen requirements, and 55% secondary attack rates among siblings, and it explained 94% of adenovirus mortality.^{241,242} Similarly, first detected in 1992 in Israel, Ad7d2 spread to the United States in 1996 and since then has been associated with three of the five documented U.S. civilian outbreaks of Ad7²⁴ and at least one military outbreak.²⁰ These Ad7d2 outbreaks led to at least 19 deaths and the suggestion that this new genotype was a more virulent strain. In a 2005 report from Iowa, Ad7d2 had displaced other Ad7 strains.²⁴³

Most recently, a novel genotype of a seldom-detected Ad14 strain (classified as Ad14p1) emerged in the United States and has spread to other countries causing considerable morbidity and death.^{28,146,147,244–249} It differs from the Ad14p prototype strain by markedly greater epithelial cell binding, cell cytotoxicity, plaque morphology, and minor mutations in the fiber knob and E1A genes.^{250,251,251a}

High-Risk Groups

As noted, virtually all children have adenovirus infections, and, occasionally, these infections are severe and lead to death. Additionally, adenovirus infections are quite common among the immunocompromised and often associated with significant morbidity. Adenovirus epidemics often occur at intensive care units or long-term care facilities. For instance, in 1999 a novel Ad7b variant caused 84% of the 50 residents (age range: 1–46 years) of a New York long-term care facility to become ill, 26 were hospitalized, and 7 died.²⁵² In a 2007 national U.S. adenovirus study, the authors found risk factors for severe clinical adenovirus disease included age younger than 7 years, chronic disease, recent transplantation, and infection with Ad5 or Ad21.¹⁷⁷

Epidemics among military populations, especially when adenovirus vaccines were not available, have been particularly well documented.⁴ Before the use of vaccines in the U.S. military, Ad4 and Ad7 accounted for 60% of all ARD among hospitalized recruits. Ad3, Ad14, and Ad21 also occurred, but less commonly.¹⁴ Up to 80% of recruits became infected with adenovirus, while seasoned military personnel experienced lower rates.⁴ Such outbreaks have also been reported among military recruits in China, Finland, The Netherlands, The United Kingdom, Turkey, Singapore, and South Korea.^{253–261}

Modes of Transmission and Reservoirs of Infection

The disease characteristics of the adenoviruses vary by species and type (see Table 10.1).³⁹ The reasons for these differences are not well understood. Adenoviruses are transmitted by direct contact, aerosolized virus, the fecal–oral route, and via water. Although recent data suggest that human Ad4 and Ad52 may have simian origins,^{124,141} and a novel adenovirus may have crossed species between humans and animals,^{137,262} humans are thought to be by far the primary reservoir for human adenovirus.²⁶³

Significance as a Health Burden

The most common environment for adenovirus transmission is the home. However, high rates of adenovirus transmission

occur wherever large groups of susceptible people gather. Transmission rates are higher in children's institutions and daycare centers, and in lower socioeconomic groups.^{46,47} Enteric adenoviruses may be an important pathogen in the daycare setting.²⁶⁴ Epidemics associated with Ad3 often occur in association with swimming activities.⁶⁶ Ad8 has been associated with transmission in physicians' offices.⁷⁰

Nosocomial outbreaks of adenovirus keratoconjunctivitis have been reported from the accident and emergency department of a major eye hospital in the United Kingdom.^{265–267} Nosocomial conjunctivitis, pharyngitis, and pneumonia caused by adenoviruses have been noted in hospital intensive care units.^{17,268,269}

With increasing numbers of immunocompromised patients in hospitals, adenovirus infections have become of more frequent cause of severe disease.⁹ In addition, reactivation of a latent infection could possibly initiate a nosocomial outbreak. Persons with deficient cell-mediated immunity are at greatest risk for adverse outcomes.³⁹ Bone marrow transplant recipients are especially susceptible to adenovirus infections.^{82,110} The death rate among immunodeficient patients with pneumonia may be as high as 60%.⁹

Chronic diarrhea in patients with AIDS is often a diagnostic problem. A prospective study using extensive diagnostic techniques, such as duodenal, jejunal, and rectal biopsies, demonstrated that 6.5% of the patients had adenovirus infections.²⁷⁰

Although the civilian experience has not established a requirement for a vaccine, the epidemic nature and extensive morbidity experienced by military recruits during the 1960s demonstrated an overwhelming need for adenovirus vaccines for military use. One well-studied and typical epidemic at Fort Dix, NJ, exemplified this requirement.¹⁵ A platoon of 48 men was followed up prospectively for their 8 weeks of basic training. Of 92 episodes of respiratory illness, 24 required hospitalization. The documented hospitalization rate for ARD due to adenovirus Ad4 was 5/100 soldiers per week.¹⁵ At large basic training posts, this rate translated to approximately 500 to 800 ARD admissions per week, which had a devastating impact on military hospitals.^{15,19} Excess medical costs and the fact that soldiers had to be "recycled" owing to lost training time resulted in significant economic loss to the military. Serious disruptions to training schedules led to administrative attempts to control epidemics, such as sleeping head-to-foot and keeping military units separated (cohorting).¹⁹ The impact of adenovirus infections in the military led to the development of adenovirus vaccines.

PASSIVE IMMUNIZATION

While years ago immunoglobulin therapy was not found to be particularly useful in preventing adenovirus infections among military trainees,²²⁰ modern passive immunization strategies may prove helpful. In particular, recently isolating Ad-specific T cells from stem cell donors have been effective in treating hematopoietic stem cell transplantation recipients who develop severe adenovirus that is nonresponsive to antivirals.³⁷

ACTIVE IMMUNIZATION

The first adenovirus vaccine prototype was an injectable bivalent Ad4 and Ad7 vaccine grown in monkey kidney cells and formalin-inactivated.¹⁹ After safety tests, a small trial in 1957 reduced admissions for adenoviruses by 98%, and a large trial of 8238 soldiers was 90% effective in reducing hospitalization.^{271,272} A trivalent vaccine, which also included Ad3, was also tested.²⁷³ Large-scale production of vaccine lots led to variation in antigenicity, resulting in a lowering of protection

rates, with only a 52% reduction in hospital illnesses caused by adenoviruses.²⁷⁴ Even this low rate of protection was reported in 1965 to have saved the Army approximately \$5 million a year.²⁷⁴ Adenovirus seed lots were found to be contaminated with simian virus 40 (SV40), a virus oncogenic in animals, and in 1963, the license for the vaccine was rescinded.¹⁵ Although this caused considerable concern at the time, an epidemiologic study of 1959–1961 era U.S. Army male veterans showed no evidence of an increased risk of various cancers that plausibly could be the result of an SV40 exposure.²⁷⁵

Use of a live vaccine for ARD caused by Ad7 was investigated, and oral administration was reported to induce high antibody levels in 1960.²⁷⁶ Couch and colleagues²⁷⁷ and Chanock and colleagues²⁷⁸ demonstrated that some adenoviruses infect the gastrointestinal tract but do not produce symptoms in adults. This finding led to the administration of the virus as a vaccine in an enteric-coated capsule, which produced an asymptomatic, intestinal infection and neutralizing antibodies.^{277,278} Many safety studies were performed in human volunteers, including the administration of Ad4 and Ad7 vaccines separately and simultaneously.²⁷⁷ Neutralizing antibodies were elicited, and asymptomatic infections were evidenced by the recovery of virus from rectal specimens. Initial vaccine virus production using HEK cells was later modified to use human diploid fibroblast strains WI-26 and WI-38 because the primary cells were not suitable for large-scale production and there was concern of their contamination with other human pathogens.²⁷⁸ Because some adenoviruses are oncogenic in animals, much attention was given to safety studies in hamsters and the transformation of cell lines by adenoviruses.^{279,280}

Additional field trials for the Ad4 vaccine were conducted at Parris Island, SC, and Great Lakes, IL, indicating that the vaccine was highly protective, safe, and antigenic and showed no evidence of transmission among military trainees. Vaccine use led to reductions in ARD rates by 50%.^{281,282} Intrinsic vaccine efficacy was as high as 82%.²⁸²

Use of the monotypic Ad4 vaccine was followed by the appearance of ARD caused by Ad7 at Fort Dix, NJ.¹⁵ The Army Adenovirus Surveillance Program was initiated to identify the agents of ARD and to assess fluctuations in disease patterns.¹⁴

Before an Ad7 vaccine could be developed and licensed, numerous studies addressed the oncogenicity potential of Ad7 to humans.^{283–286} A trial of the Ad7 vaccine demonstrated safety, infectivity, antigenicity, and lack of communicability similar to that observed for the Ad4 vaccine.²⁸⁷ Additional trials indicated that when the Ad4 and Ad7 vaccines were given simultaneously, no decrease in immunogenicity occurred, and with mass immunization, there was 95% suppression of Ad7-associated disease.^{287–289} Numerous outbreaks of Ad21 in military populations in the United States and Europe led to the initiation of studies using an Ad21 prototype vaccine in safety and immunogenicity trials.^{290–294} Trials of Ad1, Ad2, and Ad5 vaccines were also conducted with the goal of protecting children.²⁹⁵ However, Ad1, Ad2, Ad5, and Ad21 vaccine production was never pursued.

Beginning in 1971, live, enteric-coated, oral vaccines for Ad4 and Ad7 were routinely administered, and adenovirus morbidity was greatly reduced.¹⁹ Ad4 and Ad7 vaccines were licensed by the FDA in 1980 and produced by Wyeth Laboratories. In 1994, the Department of Defense was notified that the sole producer of Ad4 and Ad7 vaccines, Wyeth Laboratories Inc, Marietta, PA, would stop production permanently. Beginning in 1984, Wyeth repeatedly notified the Department of Defense of the need to negotiate a contract that would consider the renovation of the existing vaccine production facility.²⁹⁶ The contract was not renegotiated.²⁹⁶ Bulk vaccine production ceased in 1995, and the last vaccine was placed in

tablet form in 1996. Through requests for extensions of expiration dates and use of vaccines only during September through March, rather than year-round, the U.S. Department of Defense attempted to obtain maximum benefit from the remaining supply. By 1999, all vaccine supplies were depleted.

After the loss of the manufacture of the Ad4 and Ad7 vaccines, there were many documented outbreaks of ARD due to adenoviruses in U.S. military training centers. The first recorded epidemic took place during April-May 1995, at Fort Jackson, SC, during a lapse of vaccine administration that occurred because of a logistical error that temporarily interrupted vaccine production.²¹ Ad4 was identified as the etiologic agent in unvaccinated soldiers who experienced hospitalization rates of 11.6% during basic training.²¹ Numerous additional outbreaks occurred at the other U.S. military camps,^{20,22,297} some causing large numbers of hospitalizations^{22,297,298} and some spreading to more advanced training camps.²⁹⁹ Results of a nationwide seroprevalence survey among unimmunized U.S. Army trainees confirmed the lack of protective neutralization antibodies to Ad4 and Ad7, with nearly 90% being susceptible to at least one type.^{299,300} Evaluation of the epidemic, clinical, and immunological risk factors for adenovirus infection was performed for a subset of patients, demonstrating that anti-Ad4 immunity was low in new recruits, risk of illness was higher in smokers, and 81% of patients from whom paired serum samples were collected demonstrated a fourfold or higher increase in anti-Ad4 titer after infection.³⁰¹

Gray and colleagues provided a concise overview of population-based surveillance for respiratory disease at four U.S. military training centers as the last of the stores of vaccines were depleted.³⁰² Between October 1996 and June 1998, 53.1% of 1,814 throat cultures from symptomatic trainees yielded adenoviruses.³⁰² Ad4, Ad7, Ad3, and Ad21 accounted for 57%,

25%, 9%, and 7% of the isolates,³⁰² respectively (although the prevalence of Ad7 detections has markedly declined in recent years).³⁰³ Russell and colleagues³⁰⁴ examined an estimated 73,748 clinical cases of adenovirus infection that occurred during the period of 1999 to 2004 among U.S. military trainees at eight training sites. They observed that adenovirus explained from 52.3% to 76.4% of detected febrile respiratory illnesses at these sites.³⁰⁴ The aggregate number of adenovirus infections among U.S. military trainees has fluctuated with the availability of vaccines and season (Fig. 10.3). In some winter months, more than 2200 preventable clinical infections were documented. Realizing that the detected adenovirus illnesses greatly underestimated true clinical disease, Russell and colleagues³⁰⁴ posited that the true count of clinical adenovirus disease occurring in these camps was an estimated 45,000 cases per year, 90% of which would be prevented were the adenovirus vaccines available. This realization of preventable morbidity was not lost to public health officials inside and outside the U.S. Department of Defense, with many calling for reacquisition and use of the vaccines.³⁰⁵⁻³⁰⁸

On September 25, 2001, 7 years after Wyeth announced permanent cessation of production of the adenovirus vaccines, two widely publicized adenovirus-associated deaths in non-immunized recruits,³⁰⁹ and strongly worded letters from the Institute of Medicine and the Armed Forces Epidemiological Board to expedite contract efforts, induced the Department of Defense to contract for production of adenovirus vaccines.^{296,306,310,311} The decision to resume production was based on a continuing low prevalence of antibodies to Ad4 and Ad7 in people coming into the military,³⁰⁰ the occurrence of documented outbreaks when the vaccines were not given,^{20-22,302} and almost 30 years of experience of not having ARD outbreaks when potent adenovirus vaccines were being




Figure 10.3. Febrile respiratory illness (FRI) and adenovirus (Adv) rates per 100 person-weeks per month with estimated monthly number of adenovirus-associated clinical encounters among trainees at eight U.S. military training sites, June 1998 through December 2014. (Courtesy Anthony Hawksworth, Naval Health Research Center, San Diego, CA.)

administered.¹⁹ A genomic analysis of the Wyeth Ad4 and Ad7 vaccine strains show no evidence of attenuation.³¹² In 2001 the contract was awarded and seed viruses strains from Wyeth were transferred to Barr Laboratories Inc. (now part of Teva Pharmaceuticals Industries, Inc.).

In the fall of 2004, the Teva Ad4 and Ad7 vaccines were studied in a Phase I double-blind, placebo-controlled trial involving 58 Army medic trainees who had just completed U.S. Army basic training.³¹³ Among the subjects who had been previously exposed to adenovirus during basic training and before enrollment, 79% and 78% were seropositive to Ad4 or Ad7, respectively. For the trial, 58 initially seronegative subjects were selected. Both vaccines were given to 30, and 28 received placebo tablets. For 32 subjects, final follow-up was permitted at day 180. The participants were closely followed up with cultures, symptom diaries, and clinical examinations during a 56-day period for evidence of vaccine reactions and adenoviral disease. The most commonly reported adverse events in the vaccine recipients were nasal congestion (33%) and dry cough (33%), sore throat (27%), headache (20%), abdominal pain (17%), arthralgia (13%), nausea (13%), and diarrhea (13%). Symptom reporting did not differ between vaccine and placebo groups. Vaccine virus was shed in the stools of 73% and 71% of volunteers seronegative to Ad4 and Ad7, respectively, for as long as 21 days after vaccination. By day 28 after vaccination, 73% of vaccine recipients seroconverted to Ad4, while 63% seroconverted to Ad7. A randomized, double-blind, placebo-controlled Phase II/III trial (NCT00382408) of these vaccines was conducted in 2006–2007 at two military training facilities. Ad4 and Ad7 vaccine or placebo was administered to 4040 U.S. military trainees in a 3:1 ratio. As described above the vaccine was found to have excellent efficacy (99.3% for Ad4) and safety profiles similar to the previously manufactured Wyeth vaccine.

In March 2011, after an arduous 10-year vaccine development process, eight likely preventable adenovirus-associated deaths,³¹⁴ thousands of hospitalizations, and more than \$100 million in funding,^{315,316} Teva was awarded FDA licensure for their Ad4 and Ad7 vaccines. The vaccines are indicated for military personnel ages 17 to 50 years of age and contraindicated for persons who are pregnant, who cannot swallow the vaccine tablets without chewing, or who have a severe reaction to the vaccines.

Constituents Including Antibiotics, Preservative, Adjuvants, Etc.

Wyeth vaccine tablets contained live viruses, materials added for the growth and maintenance of viruses and cells, and other pharmaceutical materials. Human-diploid fibroblast cells (strain WI-38) were used for virus preparation, and growth was maintained in Minimal Essential Medium, Eagle solution, antibiotics (neomycin sulfate, gentamicin sulfate, and amphotericin B), fetal bovine serum, and sodium bicarbonate. After harvesting, the viral growth was freed of particulate material by filtration and dried by lyophilization. During drying, additives were used to preserve viability. Before processing into tablets, the virus preparation was diluted with lactose powder.³¹⁷

Using the same Wyeth master seed viruses and a manufacturing process similar to the process Wyeth used, Teva is now producing the Ad4 and Ad7 vaccines. Teva's tablets consist of three layers: a central core containing at least 4.5 log₁₀ tissue culture infective dose (TCID₅₀) of lyophilized formulated adenovirus mixed with anhydrous lactose, microcrystalline cellulose, polacrillin potassium, and magnesium stearate, an




Figure 10.4. Teva Pharmaceuticals Industries Ltd. enteric-coated adenovirus type 4 and type 7 vaccine packaging and tablets. The adenovirus type 7 (b7) tablet is yellow. The enteric coat is designed to resist stomach acid. (Courtesy John Shaw, Teva Pharmaceuticals, Inc.)

outer layer of inactive excipients, and a protective enteric coating consisting of cellulose acetate phthalate, alcohol, acetone, and castor oil (Fig. 10.4).³¹⁸ The Ad7 vaccine tablet also contains FD&C Yellow #6 aluminum lake dye to give it a distinguishing yellow color.

Preparations Available, Including Combinations

The Teva adenovirus vaccines contain live nonattenuated Ad4 or Ad7 in the form of enteric-coated tablets. They are produced similarly to how the Wyeth vaccines (described above) were produced. The final vaccine is composed of two tablets (one tablet of Ad4 and one tablet of Ad7) designed to pass intact through the stomach and release the live virus in the intestine. Each enteric-coated tablet contains an inner core tablet containing anhydrous lactose, microcrystalline cellulose, polacrillin potassium, magnesium stearate, and live adenovirus, either Type 4 or Type 7, at a potency of no fewer than 32,000 tissue-culture infective doses (4.5 log₁₀ TCID₅₀) per tablet. The outer tablet layer contains microcrystalline cellulose, magnesium stearate, and anhydrous lactose, with an enteric coating consisting of cellulose acetate phthalate, alcohol, acetone, and castor oil. The Ad7 tablet also contains FD&C Yellow #6 aluminum lake dye.

The Ad4 vaccine (white tablets) and the Ad7 vaccine (yellow tablets) are packaged in separate bottles and the bottles are combined in a single package.³¹⁸ The tablets can be administered simultaneously but have to be swallowed without chewing. Vomiting and diarrhea may interfere with vaccine effectiveness. These vaccines are indicated for military populations shown to be at risk of ARD from the specific adenovirus types represented in the tablets.

The adenovirus vaccines are routinely given to military recruits and U.S. Coast Guard cadets along with numerous other vaccines in the first 8 days after entering training. Type-specific neutralizing antibodies to the adenovirus vaccines are produced in the vaccinated trainees and were associated with protection against ARD. Interference with immune responses when the adenovirus vaccines were given with other vaccines has not been identified.³¹⁹

DOSAGE AND ROUTE: INTRAMUSCULAR, SUBCUTANEOUS, AND ORAL

The initial oral Ad4 vaccine was developed using dried virus contained in enteric-coated capsules and found to be effective in preventing infections.²⁷⁸ Original trials for Ad7 vaccine were conducted using 0.05 mL of a 1:10 dilution of the Ad7 pool (10^6 TCID₅₀) within a hard gelatin capsule and given orally.²⁷⁷ Time of disintegration within the intestine was assayed roentgenographically using a barium sulfate-containing capsule and varied between 1 and 5 hours.²⁷⁷ Two small dose ranging studies using vaccine capsules suggested that for both types, nearly 100% seroconversion was achieved by approximately 10^4 TCID₅₀ per capsule.^{289,313} A 1969 dose-ranging study used three Ad7 doses: $10^{6.8}$ TCID₅₀, $10^{4.8}$ TCID₅₀, and less than 10 TCID₅₀ per capsule.^{277,289} There was 100% antibody response with the highest dose, 95% response with the intermediate dose, and 56% response with the lowest dose. Another study evaluated the response when both vaccines were given individually at doses of $10^{5.4}$ TCID₅₀ for Ad7 and 10^4 TCID₅₀ for Ad4, and simultaneously for both vaccines at the same doses.²⁸⁷ There was no decrease in the immunogenicity of the Ad4 vaccine when it was given with the Ad7 vaccine.²⁸⁹ In the mid 1960s, Wyeth Laboratories took over production of the adenovirus vaccines, which were converted to enteric-coated tablets.

Vaccine Stability

The currently licensed Teva vaccines are stored between 2°C and 8°C with a shelf life of 30 months.

IMMUNOGENICITY OF VACCINE

Humoral Responses

Oral adenovirus vaccine recipients routinely shed virus fecally, which can potentially continue for up to 6 weeks.^{292,320} Wyeth vaccine recipients developed neutralizing humoral antibody (immunoglobulin [Ig] G, IgM, and IgA).²⁹² In soldiers who were free of preexisting antibody, an average of 80% to 95% developed a neutralizing antibody level of greater than 1:8, while fewer than 50% demonstrated complement-fixing antibody.^{289,321} Neutralizing antibody responses were detected 2 to 3 weeks after vaccination. In general, antibody titers were less than titers achieved after natural infection.^{8,282} Local secretory IgA antibody was not induced by the oral vaccine, and reinfection of the respiratory tract was possible, but usually mild or asymptomatic.³²² Since viremia and viruria could occur in patients with febrile disease, invasiveness beyond mucosal surfaces could be important in the pathogenesis of the disease and infection.³²³ The serum-neutralizing antibody produced as a result of vaccination may have prevented the typical febrile disease associated with natural infection.³²¹ Local IgA antibody could be produced experimentally by the intranasal inoculation of an Ad4 vaccine in liquid.³²² A Phase I/II study of the Teva oral Ad4/7 vaccine showed 73% and 63% of immunonaïve subjects seroconverted to Ad4 and Ad7 at 28 days postvaccination.³¹³ A Phase III study of the same vaccines documented seroconversion at 94.5% (Ad4) and 95% (Ad7).³²⁴

Cellular Responses

The cellular response to Wyeth Ad4 and Ad7 vaccines has not been well studied. However, the cellular immune responses to some adenovirus-vectored vaccines have been excellent.³²⁵

Correlates of Protection

A SN assay value of 1:4 against a specific virus is considered as protective.³¹⁵ A fourfold rise in titer is considered as evidence of infection.

Special Groups (e.g., Immunosuppressed)

Although some early Ad1, Ad2, Ad4, and Ad5 vaccine trials were conducted among civilians,^{320,326,327} the Teva vaccines have not been used in groups other than healthy military trainees.

EFFICACY AND EFFECTIVENESS OF VACCINE

Military recruits who received Ad4 vaccine exhibited increased resistance to respiratory disease caused by this virus.^{278,281,328} A number of major well-controlled studies of more than 42,000 soldiers were performed from 1963 to 1966.³²⁹ Use of the vaccine reduced ARD by 50% and adenovirus infection in recruits by more than 90%.

Field trials began in 1969 for the Ad7 vaccine. The trials indicated protection against disease in susceptible persons.^{288,289,330} In addition, it was found that the two vaccines (for Ad4 and Ad7) could be administered simultaneously without interference or loss of efficacy.^{288,289,331} As a result of using the vaccines at one large Air Force base during a 9-year period, ARD caused by adenovirus disappeared.³³² The adenovirus surveillance program demonstrated that the combination of the two adenovirus vaccines was highly effective in controlling epidemic ARD.¹⁴ From 1971 to 1999, the live enteric-coated Ad4 and Ad7 vaccines were administered to new military recruits and successfully controlled ARD caused by adenoviruses.

The Phase III double-blind, placebo-controlled efficacy trial of the Teva Ad4 and Ad7 vaccines conducted in 2006 found the vaccines to be 99.3% effective in reducing Ad4-caused ARD.³²⁴ Two years after their reintroduction, the Teva vaccines were credited with a 100-fold decline in military trainee adenovirus cases and with the prevention of 1100 to 2700 hospitalizations each year.³⁰³

Duration of Immunity and Protection, Including Description of Reinfection, If Any

The duration of immunity and persistence of circulating antibody following immunization has not been well studied. The adenovirus vaccines were developed to protect new members of the military (recruits) against Ad4 and Ad7 respiratory disease during the first months of military service. The adenovirus vaccines were very effective in accomplishing that objective, and immunity beyond 56 days has not been studied.

Safety

Common Adverse Events

Four study groups were followed up for inpatient and outpatient episodes of illness during vaccine safety and immunogenicity trials at Lackland Air Force Base, TX, in 1976. In addition to the placebo group, one group received three vaccines simultaneously (Ad4, Ad7, and Ad21), another group received two vaccines simultaneously (Ad4 and Ad7), and the last group received one vaccine only (Ad21).²⁹⁴ No appreciable differences were noted in the inpatient or outpatient experiences of the different groups.²⁹⁴

Studies of the various live adenovirus vaccines have documented fecal and pharyngeal shedding of vaccine virus and

transmission to nonvaccinated study participants.^{295,320,321,324,333} Although recent trials of the Teva vaccine demonstrated sparse transmission among military personnel,³²⁴ the possibility of transmission of vaccine virus to other susceptible personnel should be considered for up to 4 weeks after receipt of modern vaccines.

Rare Adverse Events

In 1973, a Navy trainee was hospitalized with fever, malaise, and dyspnea for 11 days after receiving the Ad4 and Ad7 vaccines.³³⁴ The patient died on the 10th hospital day with the diagnosis of Ad7 pneumonia, based on the isolation of the virus by cell culture. However, it was not possible at the time to determine whether his infection was caused by a wild Ad7 before the development of vaccine-induced immunity or to an Ad7 vaccine strain.³³⁴

Immunocompromised Recipients

As adenoviruses may cause severe disease among immunocompromised persons, when HIV was first discovered, concern grew among U.S. military public health officials that adenovirus vaccine might cause severe disease among HIV-infected military trainees. Subsequently, a study was conducted among HIV-infected and uninfected military trainees. Although significantly fewer HIV-infected subjects responded to the Ad4 vaccine than did soldiers without HIV infection, no clinically apparent adverse reactions were detected.³¹⁹ Response to the Ad7 vaccine was difficult to define because many vaccine recipients in both groups had high neutralizing antibody titers at the time of vaccination.³¹⁹ More severely immunocompromised persons, who might inadvertently be infected with adenovirus vaccine virus through fecal shedding from “normal” vaccine recipients, have not been studied.

Pregnancy

The live, oral adenovirus vaccines were never subjected to animal reproduction studies because of the lack of a suitable animal model. Therefore, it is not known if these vaccines had the potential to cause fetal damage if given to a pregnant woman or to affect reproductive capacity. Female military trainees are screened with pregnancy tests before they received the adenovirus vaccines. During the recent Teva vaccine trials four of the subjects were found to be pregnant after they had received the Ad4 and Ad7 vaccines.³¹⁸ All four subjects delivered healthy babies at 36 to 40 weeks’ gestation. Unusual morbidity among pregnant vaccine recipients has not been reported.

Spread to Contacts

Because the virus may be shed fecally for up to 6 weeks, the virus may be spread to family members or very close contacts.^{320,326} Hence, vaccine recipients are instructed to maintain good hygiene and to avoid contact with children younger than 7 years of age, the immunocompromised, and pregnant women for 28 days following vaccination.³¹⁸ The spread of vaccine virus has not been well studied. In the recent Phase II trial of the Teva vaccines, participants were followed shedding on days 0, 7, 14, 21, 28, and 56. Periodic fecal shedding was noted 7 to 21 days following vaccination.³¹³

INDICATIONS FOR VACCINE—WHO AND WHY

Adenovirus vaccines have been indicated for the prevention and control of specific adenovirus-associated ARD in populations with a high risk of exposure, a high level of susceptibility,

and a high risk of subsequent infection and disease. Adenovirus vaccines were recommended for use in military populations at risk of developing ARD from adenoviruses. Use of these vaccines was not recommended for other populations. Current and future studies of virus transmission and disease occurrence in pediatric populations, chronic care, and other civilian institutional settings; high-level healthcare organizations; and colleges may identify other potential indications for adenovirus vaccines.

In 1971, the U.S. military began routinely administering both vaccines to males reporting to recruit training centers, but only during the winter months.¹⁵ The incoming recruits were given the vaccines within hours of arrival at a training center to obtain protection as early as possible in the training program. Initially, women were not given the vaccines because ARD outbreaks caused by adenoviruses had never been documented among military women, and there was concern about the possibility of administering the vaccines to women who were pregnant.¹⁹

The program of administering the vaccines only during the high-risk winter months was directed at control, rather than eradication of ARD caused by adenoviruses.¹⁵ With this schedule, late spring and early fall outbreaks occurred. These outbreaks prompted the U.S. Army and the U.S. Navy to adopt policies of year-round administration of both adenovirus vaccines in 1983.¹⁹ Taking a different course, the U.S. Air Force stopped the administration of adenovirus vaccines at its only recruit training center in Texas in the mid-1980s and adopted a program of surveillance with use of the vaccines only as indicated.¹⁹

When the military started immunizations to protect against ARD caused by adenoviruses, training programs were segregated by gender. These separate training programs have since been combined, leading to concern that the risk for ARD caused by adenoviruses would be the same for men and women. The U.S. military regulation for immunizations required that, based on risk, Ad4 and Ad7 vaccines be administered simultaneously and only once to Army, Navy, and Marine Corps recruits.³³⁵ In the Air Force and Coast Guard, the vaccines were administered when directed by the appropriate authority.³³⁵ The same regulation described precautions to be taken to avoid unintentional administration of the vaccines during pregnancy and counseling instructions regarding the possibility of pregnancy within 3 months following immunization.³³⁵

In response to the cessation of adenovirus vaccine production in 1996, the Army, Navy, and Marine Corps modified their policies of year-round vaccine administration to conserve the remaining vaccine for use in higher-risk months only. The modified policies directed that the vaccines be given to arriving military recruits only during the period of September 1 through March 31 until all vaccine stocks were exhausted in 1999.

CONTRAINDICATIONS AND PRECAUTIONS

Teva uses no antibiotics in the virus propagation steps. The vaccine is contraindicated for pregnant women, immunocompromised patients, persons outside the target age range of 17 to 50 years, persons who cannot swallow the tablets whole, or persons who are allergic to the vaccines.

PUBLIC HEALTH CONSIDERATIONS

Epidemiologic Effects of Vaccination, Including Herd and Contact Immunity

In the prevaccine era, ARD from adenoviruses caused significant morbidity at military training centers. These outbreaks

were extremely costly in terms of medical care requirements and time lost from training.¹⁹ The routine administration of the Ad4 and Ad7 vaccines, which began in 1971, resulted in an extremely efficacious, cost-effective, and safe immunization program.^{19,336–338} Initially, the vaccines were administered only during the high-risk winter months. The occurrence of spring and fall outbreaks prompted a modification of the program to administration of the vaccines year-round.¹⁹ From 1971 to 1999, except for a period in the 1970s when ineffective vaccines thought to be contaminated with solvent were used,¹⁵ there has never been a reported outbreak of ARD caused by Ad4 or Ad7 in U.S. military units that received the vaccines. Outbreaks of ARD caused by Ad14 and Ad21 have occurred, but these have been sporadic.^{19,294,339}

A problem in the administrative system used to procure vaccines for the U.S. military resulted in an adenovirus vaccine production delay that began in the spring of 1994 and lasted until late February 1995.¹⁹ There was only one outbreak of ARD caused by adenoviruses reported in conjunction with that production delay.²¹ When Wyeth Laboratories announced in 1994 that it would no longer produce the adenovirus vaccines, many in the U.S. Department of Defense and the different military services asked if the vaccines were still needed. Some thought that improved military barracks with modern heating, ventilation, and air-conditioning systems may have significantly reduced the risk of transmission of the adenoviruses and subsequent disease in susceptible people. The assessment of current risk was hampered by the lack of modern seroprevalence data for Ad4 and Ad7 among new military entrants and the lack of studies of adenovirus carriage, transmission, infection, and disease because the vaccines had been so successful. The U.S. Department of Defense also found it had lost the capability to perform serologic studies for adenovirus infections. Unfortunately, with the loss of the vaccines, adenovirus-associated ARD increased in all uniformed services, including the U.S. Air Force.

Disease Control Strategies

Ad4 and Ad7 vaccines have been the primary means for controlling adenovirus-induced ARD among U.S. military trainees. Secondary control measures have included reducing crowding, aligning trainees' beds head-to-foot, making frequent hand washing a requirement, increasing air exchanges, and keeping contact between different cohorts of recruits to a minimum (cohorting).^{340–342} Although interventions like frequent hand washing³⁴⁰ caused a 45% reduction in upper respiratory tract infections among trainees, the numerous attempted interventions fell far short of the effectiveness of administering adenovirus vaccines in reducing ARD.

Cost-to-Benefit Information

The excess morbidity caused by ARD and associated costs are of significant concern to the U.S. military. A number of cost-to-benefit analyses have been performed each showing significant benefit in using the vaccines.^{336–338} Using an estimated cost for receipt of both vaccines at \$150 and using the vaccines year-round in approximately 200,000 trainees per year, Radin and colleagues,³⁰³ estimated the U.S. Department of Defense would prevent approximately 6000 to 13,000 clinical adenovirus cases and 1100 to 2700 hospitalizations per year, saving the U.S. government approximately \$20 million per year in training and medical costs.

FUTURE VACCINES

Subunit Vaccine

In 1963, soluble viral subunit antigens were found to be highly immunogenic on parenteral administration in animals.^{321,343} Crystalline hexon and fiber antigens from Ad5 have been shown to induce neutralizing antibody and protection in human volunteers who were challenged.³⁴⁴ There has been no additional development of these antigens as potential vaccines. However, adenovirus vaccines consisting of soluble viral subunit antigens would be free of DNA and could alleviate fear of the oncogenic potential of adenoviruses.³²¹

Recombinant Vaccines

Advances in molecular biology have permitted the *in vivo* and *in vitro* gene transfer into mammalian cells using various adenovirus constructs. Genetically engineered adenoviruses have been altered to carry genes from a wide array of other pathogens including influenza A virus,³⁴⁵ Ebola virus,³⁴⁶ and Middle East respiratory syndrome–coronavirus,³⁴⁷ and to introduce genes as a therapy against various chronic diseases. Adenovirus vaccine construct research is quite dynamic. A detailed description is beyond the scope of this textbook. However, excellent reviews have been published discussing the advantages and drawbacks of recombinant adenovirus vaccine strategies.^{348–355}

Acknowledgments

We thank John Shaw of Teva Pharmaceuticals, Inc., and Andrew Towle, PhD, AVP LLC, for their critical review of this work.

References for this chapter are available at ExpertConsult.com.



REFERENCES

- Rowe W, Huebner R, Gilmore L, Parrott R, Ward T. Isolation of a cytopathogenic agent from human adenoids undergoing spontaneous degeneration in tissue culture. *Proc Soc Exp Biol Med.* 1953;84:570-573.
- Hilleman M, Werner J. Recovery of new agent from patients with acute respiratory illness. *Proc Soc Exp Biol Med.* 1954;85:183-188.
- Enders J, Bell J, Dingle J, et al. Adenoviruses: group name proposed for new respiratory tract viruses. *Science.* 1956;124:119-120.
- Dingle JH, Langmuir AD. Epidemiology of acute, respiratory disease in military recruits. *Am Rev Respir Dis.* 1968;97(6):suppl:1-65.
- Hilleman M, Gauld R, Butler R, et al. Appraisal of occurrence of adenovirus-caused respiratory illness in military populations. *Am J Hyg.* 1957;66:29-41.
- Chany C, Lepine P, Lelong M, et al. Severe and fatal pneumonia in infants and young children associated with adenovirus infections. *Am J Hyg.* 1958;67(3):367-378.
- Johansson ME, Brown M, Hierholzer JC, et al. Genome analysis of adenovirus type 31 strains from immunocompromised and immunocompetent patients. *J Infect Dis.* 1991;163(2):293-299.
- Bellanti JA, Artenstein MS, Brandt BL, Klutinis BS, Buescher EL. Immunoglobulin responses in serum and nasal secretions after natural adenovirus infections. *J Immunol.* 1969;103(5):891-898.
- Hierholzer JC. Adenoviruses in the immunocompromised host. *Clin Microbiol Rev.* 1992;5(3):262-274.
- Pinto A, Beck R, Jadavji T. Fatal neonatal pneumonia caused by adenovirus type 35. Report of one case and review of the literature. *Arch Pathol Lab Med.* 1992;116(1):95-99.
- Stalder H, Hierholzer JC, Oxman MN. New human adenovirus (candidate adenovirus type 35) causing fatal disseminated infection in a renal transplant recipient. *J Clin Microbiol.* 1977;6(3):257-265.
- Hierholzer JC, Wigand R, Anderson LJ, Adrian T, Gold JW. Adenoviruses from patients with AIDS: a plethora of serotypes and a description of five new serotypes of subgenus D (types 43-47). *J Infect Dis.* 1988;158(4):804-813.
- Ambinder RF, Burns W, Forman M, et al. Hemorrhagic cystitis associated with adenovirus infection in bone marrow transplantation. *Arch Intern Med.* 1986;146(7):1400-1401.
- Dudding BA, Top FH, Winter PE, et al. Acute respiratory disease in military trainees: the adenovirus surveillance program, 1966-1971. *Am J Epidemiol.* 1973;97:187-198.
- Top F. Control of adenovirus acute respiratory disease in U.S. Army trainees. *Yale J Biol Med.* 1975;48:185-195.
- Chang C, Lepine P, Lelong M, Le-Tan-Vinn S. Severe and fatal pneumonia in infants associated with adenovirus infections. *Am J Hyg.* 1958;67:367-378.
- Pingleton SK, Pingleton WW, Hill RH, et al. Type 3 adenoviral pneumonia occurring in a respiratory intensive care unit. *Chest.* 1978;73(4):554-555.
- Albert MJ. Enteric adenoviruses. Brief review. *Arch Virol.* 1986;88(1-2):1-17.
- Gaydos CA, Gaydos JC. Adenovirus vaccines in the U.S. military. *Mil Med.* 1995;160(6):300-304.
- Ryan MA, Gray GC, Smith B, et al. Large epidemic of respiratory illness due to adenovirus types 7 and 3 in healthy young adults. *Clin Infect Dis.* 2002;34(5):577-582.
- Barraza EM, Ludwig SL, Gaydos JC, Brundage JF. Reemergence of adenovirus type 4 acute respiratory disease in military trainees: report of an outbreak during a lapse in vaccination. *J Infect Dis.* 1999;179(6):1531-1533.
- Hendrix RM, Lindner JL, Benton FR, et al. Large, persistent epidemic of adenovirus type 4-associated acute respiratory disease in U.S. army trainees. *Emerg Infect Dis.* 1999;5(6):798-801.
- Russell KL, Broderick MP, Franklin SE, et al. Transmission dynamics and prospective environmental sampling of adenovirus in a military recruit setting. *J Infect Dis.* 2006;194(7):877-885.
- Erdman DD, Xu W, Gerber SI, et al. Molecular epidemiology of adenovirus type 7 in the United States, 1966-2000. *Emerg Infect Dis.* 2002;8(3):269-277.
- Hashido M, Mukouyama A, Sakae K, et al. Molecular and serological characterization of adenovirus genome type 7h isolated in Japan. *Epidemiol Infect.* 1999;122(2):281-286.
- Lebeck MG, McCarthy TA, Capuano AW, et al. Emergent US adenovirus 3 strains associated with an epidemic and serious disease. *J Clin Virol.* 2009;46(4):331-336.
- Landry ML, Lebeck MG, Capuano AW, McCarthy T, Gray GC. Adenovirus type 3 outbreak in Connecticut associated with a novel variant. *J Med Virol.* 2009;81(8):1380-1384.
- Kajon AE, Lu X, Erdman DD, et al. Molecular epidemiology and brief history of emerging adenovirus 14-associated respiratory disease in the United States. *J Infect Dis.* 2010;202(1):93-103.
- Rebello-de-Andrade H, Pereira C, Giria M, et al. Outbreak of acute respiratory infection among infants in Lisbon, Portugal, caused by human adenovirus serotype 3 and a new 7/3 recombinant strain. *J Clin Microbiol.* 2010;48(4):1391-1396.
- International Committee on Taxonomy of Viruses. <https://talk.ictvonline.org/files/ictv_official_taxonomy_updates_since_the_8th_report/m/vertebrate-official/4853>.
- Seto D, Chodosh J, Brister JR, MS J. Using the whole-genome sequence to characterize and name human adenoviruses. *J Virol.* 2011;85:5701-5702.
- Ghebremedhin B. Human adenovirus: viral pathogen with increasing importance. *Eur J Microbiol Immunol (Bp).* 2014;4(1):26-33.
- Robinson CM, Singh G, Lee JY, et al. Molecular evolution of human adenoviruses. *Sci Rep.* 2013;3:1812.
- Seto D, Jones MS, Dyer DW, Chodosh J. Characterizing, typing, and naming human adenovirus 55 n the era of whole genome data. *J Clin Virol.* 2013;58:741-742.
- Kajon AE, Echavarría M, de Jong JC. Designation of human adenovirus types based on sequence data: an unfinished debate. *J Clin Virol.* 2013;58(4):743-744.
- Aoki K, Benko M, Davison AJ, et al. Toward an integrated human adenovirus designation system that utilizes molecular and serological data and serves both clinical and fundamental virology. *J Virol.* 2011;85(11):5703-5704.
- Lion T. Adenovirus infections in immunocompetent and immunocompromised patients. *Clin Microbiol Rev.* 2014;27(3):441-462.
- Hierholzer JC, Pumarola A. Antigenic characterization of intermediate adenovirus 14-11 strains associated with upper respiratory illness in a military camp. *Infect Immun.* 1976;13(2):354-359.
- Foy HM. Adenoviruses. In: Evans AS, ed. *Viral Infections of Humans.* 3rd ed. New York, NY: Plenum Medical; 1989:77-94.
- Hierholzer J. Adenoviruses. In: Murray P, Baron E, Pfaller M, Tenover F, Tenover R, eds. *Manual of Clinical Microbiology.* Washington, DC: ASM; 1997:947-953.
- Brandt CD, Kim HW, Jeffries BC, et al. Infections in 18,000 infants and children in a controlled study of respiratory tract disease. II. Variation in adenovirus infections by year and season. *Am J Epidemiol.* 1972;95(3):218-227.
- Fox JP, Brandt CD, Wassermann FE, et al. The virus watch program: a continuing surveillance of viral infections in metropolitan New York families. VI. Observations of adenovirus infections: virus excretion patterns, antibody response, efficiency of surveillance, patterns of infections, and relation to illness. *Am J Epidemiol.* 1969;89(1):25-50.
- Foy HM, Cooney MK, McMahan R, Grayston JT. Viral and mycoplasmal pneumonia in a prepaid medical care group during an eight-year period. *Am J Epidemiol.* 1973;97(2):93-102.
- Brandt CD, Kim HW, Vargosko AJ, et al. Infections in 18,000 infants and children in a controlled study of respiratory tract disease. I. Adenovirus pathogenicity in relation to serologic type and illness syndrome. *Am J Epidemiol.* 1969;90(6):484-500.
- Schmitz H, Wigand R, Heinrich W. Worldwide epidemiology of human adenovirus infections. *Am J Epidemiol.* 1983;117(4):455-466.
- Pacini DL, Collier AM, Henderson FW. Adenovirus infections and respiratory illnesses in children in group day care. *J Infect Dis.* 1987;156(6):920-927.
- Cole R, Mastrota F, Floyd T, Chanock R. Illness and microbial experiences of nursery children at Junior Village. *Am J Hyg.* 1961;74:267-292.

48. Porter JD, Teter M, Traister V, et al. Outbreak of adenoviral infections in a long-term paediatric facility, New Jersey, 1986/87. *J Hosp Infect.* 1991;18(3):201-210.
49. Gerber SI, Erdman DD, Pur SL, et al. Outbreak of adenovirus genome type 7d2 infection in a pediatric chronic-care facility and tertiary-care hospital. *Clin Infect Dis.* 2001;32(5):694-700.
50. Mitchell LS, Taylor B, Reimels W, Barrett FF, Devincenzo JP. Adenovirus 7a: a community-acquired outbreak in a children's hospital. *Pediatr Infect Dis J.* 2000;19(10):996-1000.
51. Zhao S, Wan C, Ke C, et al. Re-emergent human adenovirus genome type 7d caused an acute respiratory disease outbreak in Southern China after a twenty-one year absence. *Sci Rep.* 2014;4:7365.
52. Commission on Acute Respiratory Disease. Experimental transmission of minor respiratory illness to human volunteers by filter-passing agents. Demonstration of two types of illness characterized by long and short incubation periods and different clinical features. *J Clin Invest.* 1947;26:957-973.
53. Baum S. Adenoviruses. In: Gorbach S, Bartlett J, Blacklow N, eds. *Infectious Diseases.* Philadelphia, PA: W.B. Saunders; 1992:1663-1667.
54. Benyesh-Melnick M, Rosenberg HS. The isolation of adenovirus type 7 from a fatal case of pneumonia and disseminated disease. *J Pediatr.* 1964;64:83-87.
55. Simila S, Ylikorkala O, Wasz-Hockert O. Type 7 adenovirus pneumonia. *J Pediatr.* 1971;79(4):605-611.
56. Straube RC, Thompson MA, Van Dyke RB, et al. Adenovirus type 7b in a children's hospital. *J Infect Dis.* 1983;147(5):814-819.
57. Liu J, Nian QG, Zhang Y, et al. In vitro characterization of human adenovirus type 55 in comparison with its parental adenoviruses, types 11 and 14. *PLoS ONE.* 2014;9(6):e100665.
58. Rubin B. Clinical picture and epidemiology of adenovirus infections. *Acta Microbiol Hung.* 1993;40:303-323.
59. Baum S. Adenovirus. In: Mandell G, Bennett J, Dolin R, eds. *Principles and Practice of Infectious Diseases.* New York, NY: Churchill Livingstone; 1995:1382-1387.
60. Dudding B, Wagner S, Zeller J. Fatal pneumonia associated with adenovirus type 7 in three military trainees. *N Engl J Med.* 1972;286:1289-1292.
61. Couch RB, Cate TR, Douglas RG Jr, Gerone PJ, Knight V. Effect of route of inoculation on experimental respiratory viral disease in volunteers and evidence for airborne transmission. *Bacteriol Rev.* 1966;30(3):517-529.
62. Bell JA, Rowe WP, Engler JL, Parrott RH, Huebner RJ. Pharyngoconjunctival fever; epidemiological studies of a recently recognized disease entity. *JAMA.* 1955;157(13):1083-1092.
63. Huebner RJ, Rowe WP, Ward TG, Parrott RH, Bell JA. Adenoidal-pharyngeal-conjunctival agents: a newly recognized group of common viruses of the respiratory system. *N Engl J Med.* 1954;251(27):1077-1086.
64. Bahn C. Swimming bath conjunctivitis. *N Orleans Med Sci J.* 1927;79:586-590.
65. D'Angelo LJ, Hierholzer JC, Keenlyside RA, Anderson LJ, Martone WJ. Pharyngoconjunctival fever caused by adenovirus type 4: report of a swimming pool-related outbreak with recovery of virus from pool water. *J Infect Dis.* 1979;140(1):42-47.
66. Foy HM, Cooney MK, Hatlen JB. Adenovirus type 3 epidemic associated with intermittent chlorination of a swimming pool. *Arch Environ Health.* 1968;17(5):795-802.
67. Parrott RH, Kim HW, Arrobio JO, et al. Epidemiology of respiratory syncytial virus infection in Washington, D.C. II. Infection and disease with respect to age, immunologic status, race and sex. *Am J Epidemiol.* 1973;98(4):289-300.
68. Jawetz E. The story of shipyard eye. *Br Med J.* 1959;15(5126):873-876.
69. Hogan M, Crawford J. Epidemic keratoconjunctivitis superfluous punctate keratitis, keratitis sub-epithelialis, keratitis maculosa, keratitis nummularis. *Am J Ophthalmol.* 1942;25:1057-1078.
70. D'Angelo LJ, Hierholzer JC, Holman RC, Smith JD. Epidemic keratoconjunctivitis caused by adenovirus type 8: epidemiologic and laboratory aspects of a large outbreak. *Am J Epidemiol.* 1981;113(1):44-49.
71. Hilleman M. Epidemiology of adenovirus respiratory infections in military recruit populations. *Ann N Y Acad Sci.* 1957;76:262-272.
72. Darougar S, Grey RH, Thaker U, McSwiggan DA. Clinical and epidemiological features of adenovirus keratoconjunctivitis in London. *Br J Ophthalmol.* 1983;67(1):1-7.
73. Guyer B, O'Day DM, Hierholzer JC, Schaffner W. Epidemic keratoconjunctivitis: a community outbreak of mixed adenovirus type 8 and type 19 infection. *J Infect Dis.* 1975;132(2):142-150.
74. Kemp MC, Hierholzer JC, Cabradilla CP, Obijeski JF. The changing etiology of epidemic keratoconjunctivitis: antigenic and restriction enzyme analyses of adenovirus types 19 and 37 isolated over a 10-year period. *J Infect Dis.* 1983;148(1):24-33.
75. de Jong JC, Wigand R, Wadell G, et al. Adenovirus 37: identification and characterization of a medically important new adenovirus type of subgroup D. *J Med Virol.* 1981;7(2):105-118.
76. Tabrizi SN, Ling AE, Bradshaw CS, Fairley CK, Garland SM. Human adenoviruses types associated with non-gonococcal urethritis. *Sex Health.* 2007;4(1):41-44.
77. Sprague JB, Hierholzer JC, Currier RW 2nd, Hattwick MA, Smith MD. Epidemic keratoconjunctivitis. A severe industrial outbreak due to adenovirus type 8. *N Engl J Med.* 1973;289(25):1341-1346.
78. Keenlyside RA, Hierholzer JC, D'Angelo LJ. Keratoconjunctivitis associated with adenovirus type 37: an extended outbreak in an ophthalmologist's office. *J Infect Dis.* 1983;147(2):191-198.
79. Mufson MA, Belshe RB. A review of adenoviruses in the etiology of acute hemorrhagic cystitis. *J Urol.* 1976;115(2):191-194.
80. Numazaki Y, Kumasaka T, Yano N, et al. Further study on acute hemorrhagic cystitis due to adenovirus type 11. *N Engl J Med.* 1973;289(7):344-347.
81. Koga S, Shindo K, Matsuya F, et al. Acute hemorrhagic cystitis caused by adenovirus following renal transplantation: review of the literature. *J Urol.* 1993;149(4):838-839.
82. Londergan TA, Walzak MP. Hemorrhagic cystitis due to adenovirus infection following bone marrow transplantation. *J Urol.* 1994;151(4):1013-1014.
83. Echavarría MS, Ray SC, Ambinder R, Dumler JS, Charache P. PCR detection of adenovirus in a bone marrow transplant recipient: hemorrhagic cystitis as a presenting manifestation of disseminated disease. *J Clin Microbiol.* 1999;37(3):686-689.
84. Horwitz M. Adenoviridae and their replication. In: Fields BN, Chanock RM, Hirsch MS, et al., eds. *Virology.* New York, NY: Raven Press; 1990:1679-1721.
85. Cromeans TL, Lu X, Erdman DD, Humphrey CD, Hill VR. Development of plaque assays for adenoviruses 40 and 41. *J Virol Methods.* 2008;151(1):140-145.
86. Chiba S, Nakata S, Nakamura I, et al. Outbreak of infantile gastroenteritis due to type 40 adenovirus. *Lancet.* 1983;2(8356):954-957.
87. Rodriguez WJ, Kim HW, Brandt CD, et al. Fecal adenoviruses from a longitudinal study of families in metropolitan Washington, D.C.: laboratory, clinical, and epidemiologic observations. *J Pediatr.* 1985;107(4):514-520.
88. Uhnou I, Wadell G, Svensson L, Johansson ME. Importance of enteric adenoviruses 40 and 41 in acute gastroenteritis in infants and young children. *J Clin Microbiol.* 1984;20(3):365-372.
89. Baschat AA, Towbin J, Bowles NE, Harman CR, Weiner CP. Is adenovirus a fetal pathogen? *Am J Obstet Gynecol.* 2003;189(3):758-763.
90. Petrikovsky BM, Lipson SM, Kaplan MH. Viral studies on amniotic fluid from fetuses with and without abnormalities detected by prenatal sonography. *J Reprod Med.* 2003;48(4):230-232.
91. Bajanowski T, Wiegand P, Cecchi R, et al. Detection and significance of adenoviruses in cases of sudden infant death. *Virchows Arch.* 1996;428(2):113-118.
92. Simila S, Jouppila R, Salmi A, Pohjonen R. Encephalomyelitis in children associated with an adenovirus type 7 epidemic. *Acta Paediatr Scand.* 1970;59(3):310-316.
93. Kelsey DS. Adenovirus meningoencephalitis. *Pediatrics.* 1978;61(2):291-293.
94. Chatterjee NK, Samsonoff WA, Balasubramaniam N, et al. Isolation and characterization of adenovirus 5 from the brain of an infant with fatal cerebral edema. *Clin Infect Dis.* 2000;31(3):830-833.
95. Ooi MH, Wong SC, Clear D, et al. Adenovirus type 21-associated acute flaccid paralysis during an outbreak of

- hand-foot-and-mouth disease in Sarawak, Malaysia. *Clin Infect Dis*. 2003;36(5):550-559.
96. Sturdy PM, Court SD, Gardner PS. Viruses and whooping-cough. *Lancet*. 1971;2(7731):978-979.
 97. Nelson KE, Gavitt F, Batt MD, et al. The role of adenoviruses in the pertussis syndrome. *J Pediatr*. 1975;86(3):335-341.
 98. Melon S, Mendez S, Iglesias B, et al. Involvement of adenovirus in clinical mononucleosis-like syndromes in young children. *Eur J Clin Microbiol Infect Dis*. 2005;24(5):314-318.
 99. Abzug MJ, Levin MJ. Neonatal adenovirus infection: four patients and review of the literature. *Pediatrics*. 1991;87(6):890-896.
 100. Price NO, Hacker JK, Silvers JH, et al. Adenovirus Type 3 Viremia in an Adult with Toxic Shock-Like Syndrome. *Clin Infect Dis*. 2001;33(2):260-262.
 101. Ruskanen O, Meurman O, Kusjavi A. Adenoviruses. In: Richman D, Whitley R, Hayden F, eds. *Clinical Virology*. New York, NY: Churchill Livingstone; 1997:525-547.
 102. Robinson C, Echavarría M. Adenoviruses. In: Murray P, Baron E, Jorgensen J, Landry M, Pfaller M, eds. *Manual of Clinical Microbiology*. 9th ed. Washington, DC: ASM Press; 2007:1589-1600.
 103. de Jong PJ, Valderrama G, Spigland I, Horwitz MS. Adenovirus isolates from urine of patients with acquired immunodeficiency syndrome. *Lancet*. 1983;1(8337):1293-1296.
 104. Fife KH, Ashley R, Shields AF, et al. Comparison of neutralization and DNA restriction enzyme methods for typing clinical isolates of human adenovirus. *J Clin Microbiol*. 1985;22(1):95-100.
 105. Carmichael GP Jr, Zahradnik JM, Moyer GH, Porter DD. Adenovirus hepatitis in an immunosuppressed adult patient. *Am J Clin Pathol*. 1979;71(3):352-355.
 106. Ohori NP, Michaels MG, Jaffe R, Williams P, Yousem SA. Adenovirus pneumonia in lung transplant recipients. *Hum Pathol*. 1995;26(10):1073-1079.
 107. Gelfand MS, Cleveland KO, Lancaster D, Corbett CE, Florendo NT. Adenovirus parotitis in patients with AIDS. *Clin Infect Dis*. 1994;19(6):1045-1048.
 108. Flomenberg PR, Chen M, Munk G, Horwitz MS. Molecular epidemiology of adenovirus type 35 infections in immunocompromised hosts. *J Infect Dis*. 1987;155(6):1127-1134.
 109. Schnurr D, Dondero ME. Two new candidate adenovirus serotypes. *Intervirology*. 1993;36(2):79-83.
 110. Flomenberg P, Babbitt J, Drobyski WR, et al. Increasing incidence of adenovirus disease in bone marrow transplant recipients. *J Infect Dis*. 1994;169(4):775-781.
 111. Dolin R. Viral pneumonia. In: Gorbach SLBJ, Blacklow NB, eds. *Infectious Diseases*. Philadelphia, PA: W.B. Saunders; 1992:485-490.
 112. Echavarría M, Forman M, van Tol MJ, et al. Prediction of severe disseminated adenovirus infection by serum PCR. *Lancet*. 2001;358(9279):384-385.
 113. Ito M, Hirabayashi N, Uno Y, Nakayama A, Asai J. Necrotizing tubulointerstitial nephritis associated with adenovirus infection. *Hum Pathol*. 1991;22(12):1225-1231.
 114. Bruno B, Zager RA, Boeckh MJ, et al. Adenovirus nephritis in hematopoietic stem-cell transplantation. *Transplantation*. 2004;77(7):1049-1057.
 115. Howard DS, Phillips IG, Reece DE, et al. Adenovirus infections in hematopoietic stem cell transplant recipients. *Clin Infect Dis*. 1999;29(6):1494-1501.
 116. Courouclé XI, Welty SE, Ramsay PL, et al. Detection of microorganisms in the tracheal aspirates of preterm infants by polymerase chain reaction: association of adenovirus infection with bronchopulmonary dysplasia. *Pediatr Res*. 2000;47(2):225-232.
 117. Bowles NE, Ni J, Kearney DL, et al. Detection of viruses in myocardial tissues by polymerase chain reaction. evidence of adenovirus as a common cause of myocarditis in children and adults. *J Am Coll Cardiol*. 2003;42(3):466-472.
 118. Gabbert C, Donohue M, Arnold J, Schwimmer JB. Adenovirus 36 and obesity in children and adolescents. *Pediatrics*. 2010;126(4):721-726.
 119. Broderick MP, Hansen CJ, Irvine M, et al. Adenovirus 36 seropositivity is strongly associated with race and gender, but not obesity, among US military personnel. *Int J Obes (Lond)*. 2010;34(2):302-308.
 120. van Ginneken V, Sitnyakowsky L, Jeffery JE. Infectobesity": viral infections (especially with human adenovirus-36: Ad-36) may be a cause of obesity. *Med Hypotheses*. 2009;72(4):383-388.
 121. Esposito S, Preti V, Consolo S, Nazzari E, Principi N. Adenovirus 36 infection and obesity. *J Clin Virol*. 2012;55(2):95-100.
 122. International Committee on Taxonomy of Viruses, Fauquet C, International Union of Microbiological Societies. Virology Division. *Virus taxonomy: classification and nomenclature of viruses: eighth report of the International Committee on Taxonomy of Viruses*. Boston, MA: Elsevier/Academic Press; 2005.
 123. Hierholzer JC, Stone YO, Broderon JR. Antigenic relationships among the 47 human adenoviruses determined in reference horse antisera. *Arch Virol*. 1991;121(1-4):179-197.
 124. Jones MS 2nd, Harrach B, Ganac RD, et al. New adenovirus species found in a patient presenting with gastroenteritis. *J Virol*. 2007;81(11):5978-5984.
 125. Walsh MP, Chintakuntlawar A, Robinson CM, et al. Evidence of molecular evolution driven by recombination events influencing tropism in a novel human adenovirus that causes epidemic keratoconjunctivitis. *PLoS ONE*. 2009;4(6):e5635.
 126. Walsh MP, Seto J, Jones MS, et al. Computational analysis identifies human adenovirus type 55 as a re-emergent acute respiratory disease pathogen. *J Clin Microbiol*. 2010;48(3):991-993.
 127. Robinson CM, Singh G, Henquell C, et al. Computational analysis and identification of an emergent human adenovirus pathogen implicated in a respiratory fatality. *Virology*. 2011;409(2):141-147.
 128. Torres S, Chodosh J, Seto D, Jones MS. The revolution in viral genomics as exemplified by the bioinformatic analysis of human adenoviruses. *Viruses*. 2010;2:1367-1381.
 129. Liu H, Jin L, Koh SB, et al. Atomic structure of human adenovirus by cryo-EM reveals interactions among protein networks. *Science*. 2010;329(5995):1038-1043.
 130. Philipson J, Peterson U, Lindberg U. Molecular biology of adenoviruses. *Viral Monogr*. 1975;14:1-115.
 131. Ginsberg HS, Pereira HG, Valentine RC, Wilcox WC. A proposed terminology for the adenovirus antigens and virion morphological subunits. *Virology*. 1966;28(4):782-783.
 132. Horne R, Bonner S, Waterson A. The icosahedral form of an adenovirus. *J Mol Biol*. 1956;1:84-86.
 133. Maizel JV Jr, White DO, Scharff MD. The polypeptides of adenovirus. I. Evidence for multiple protein components in the virion and a comparison of types 2, 7A, and 12. *Virology*. 1968;36(1):115-125.
 134. van Oostrum J, Burnett RM. Molecular composition of the adenovirus type 2 virion. *J Virol*. 1985;56(2):439-448.
 135. International Committee on Taxonomy of Viruses, Van Regenmortel MHV, International Union of Microbiological Societies. Virology Division. *Virus taxonomy: classification and nomenclature of viruses: seventh report of the International Committee on Taxonomy of Viruses*. San Diego, CA: Academic Press; 2000.
 136. Wadell G. Adenoviridae: the adenoviruses. In: Lennett E, Halonen P, Murphy F, eds. *Laboratory Diagnosis of Infectious Disease: Principles and Practice*. New York, NY: Springer Verlag; 1988:284-300.
 137. Chen EC, Yagi S, Kelly KR, et al. Cross-species transmission of a novel adenovirus associated with a fulminant pneumonia outbreak in a new world monkey colony. *PLoS Pathog*. 2011;7(7):e1002155.
 138. Chen EC, Yagi S, Kelly KR, et al. Cross-species transmission of a novel adenovirus associated with a fulminant pneumonia outbreak in a New World monkey colony. *PLoS Pathog*. 2011;7(7):e1002155.
 139. Wevers D, Metzger S, Babweteera F, et al. Novel adenoviruses in wild primates: a high level of genetic diversity and evidence of zoonotic transmissions. *J Virol*. 2011;85(20):10774-10784.
 140. Rice AD, Adams MM, Wallace G, et al. Efficacy of CBX001 as a post exposure antiviral in New Zealand White rabbits infected with rabbitpox virus, a model for orthopoxvirus infections of humans. *Viruses*. 2011;3(1):47-62.
 141. Madisch I, Harste G, Pommer H, Heim A. Phylogenetic analysis of the main neutralization and hemagglutination determinants of all human adenovirus prototypes as a basis for molecular classification and taxonomy. *J Virol*. 2005;79(24):15265-15276.

142. Dehghan S, Seto J, Liu EB, et al. Computational analysis of four human adenovirus type 4 genomes reveals molecular evolution through two interspecies recombination events. *Virology*. 2013;443(2):197-207.
143. Rice AD, Adams MM, Lampert B, et al. Efficacy of CMX001 as a prophylactic and presymptomatic antiviral agent in New Zealand white rabbits infected with rabbitpox virus, a model for orthopoxvirus infections of humans. *Viruses*. 2011;3(2):63-82.
144. Centers for Disease Control and Prevention (CDC). Outbreak of adenovirus 14 respiratory illness—Prince of Wales Island, Alaska, 2008. *MMWR Morb Mortal Wkly Rep*. 2010;59(1):6-10.
145. Brummitt CF, Cherrington JM, Katzenstein DA, et al. Nosocomial adenovirus infections: molecular epidemiology of an outbreak due to adenovirus 3a. *J Infect Dis*. 1988;158(2):423-432.
146. Tate JE, Bunning ML, Lott L, et al. Outbreak of severe respiratory disease associated with emergent human adenovirus serotype 14 at a US air force training facility in 2007. *J Infect Dis*. 2009;199(10):1419-1426.
147. Centers for Disease Control and Prevention (CDC). Acute respiratory disease associated with adenovirus serotype 14—four states, 2006-2007. *MMWR Morb Mortal Wkly Rep*. 2007;56(45):1181-1184.
148. Lu QB, Tong YG, Wo Y, et al. Epidemiology of human adenovirus and molecular characterization of human adenovirus 55 in China, 2009-2012. *Influenza Other Respir Viruses*. 2014;8(3):302-308.
149. Evans AS. Latent adenovirus infections of the human respiratory tract. *Am J Hyg*. 1958;67(3):256-266.
150. Garnett CT, Talekar G, Mahr JA, et al. Latent species C adenoviruses in human tonsil tissues. *J Virol*. 2009;83(6):2417-2428.
151. Andiman WA, Miller G. Persistent infection with adenovirus types 5 and 6 in lymphoid cells from humans and woolly monkeys. *J Infect Dis*. 1982;145(1):83-88.
152. Zahradnik JM, Spencer MJ, Porter DD. Adenovirus infection in the immunocompromised patient. *Am J Med*. 1980;68(5):725-732.
153. Yolken RH, Bishop CA, Townsend TR, et al. Infectious gastroenteritis in bone-marrow-transplant recipients. *N Engl J Med*. 1982;306(17):1010-1012.
154. Lion T, Kosulin K, Landlinger C, et al. Monitoring of adenovirus load in stool by real-time PCR permits early detection of impending invasive infection in patients after allogeneic stem cell transplantation. *Leukemia*. 2010;24(4):706-714.
155. Reid JA, Breckon D, Hunter PR. Infection of staff during an outbreak of viral gastroenteritis in an elderly persons' home. *J Hosp Infect*. 1990;16(1):81-85.
156. Baum SG. Persistent adenovirus infections of nonpermissive monkey cells. *J Virol*. 1977;23(2):412-420.
157. Wold WS. Adenovirus genes that modulate the sensitivity of virus-infected cells to lysis by TNE. *J Cell Biochem*. 1993;53(4):329-335.
158. Horvath J, Palkonyay L, Weber J. Group C adenovirus DNA sequences in human lymphoid cells. *J Virol*. 1986;59(1):189-192.
159. Huebner RJ, Rowe WP, Lane WT. Oncogenic effects in hamsters of human adenovirus types 12 and 18. *Proc Natl Acad Sci USA*. 1962;48:2051-2058.
160. Ginsberg HS, Lundholm-Beauchamp U, Horswood RL, et al. Role of early region 3 (E3) in pathogenesis of adenovirus disease. *Proc Natl Acad Sci USA*. 1989;86(10):3823-3827.
161. Ginsberg HS, Prince GA. The molecular basis of adenovirus pathogenesis. *Infect Agents Dis*. 1994;3(1):1-8.
162. Elliott WM, Hayashi S, Hogg JC. Immunodetection of adenoviral E1A proteins in human lung tissue. *Am J Respir Cell Mol Biol*. 1995;12(6):642-648.
163. Gold DR, Tager IB, Weiss ST, Tosteson TD, Speizer FE. Acute lower respiratory illness in childhood as a predictor of lung function and chronic respiratory symptoms. *Am Rev Respir Dis*. 1989;140(4):877-884.
164. Liu F, Green MR. Promoter targeting by adenovirus E1a through interaction with different cellular DNA-binding domains. *Nature*. 1994;368(6471):520-525.
165. Shenk T, Flint J. Transcriptional and transforming activities of the adenovirus E1A proteins. *Adv Cancer Res*. 1991;57:47-85.
166. Matsuse T, Hayashi S, Kuwano K, et al. Latent adenoviral infection in the pathogenesis of chronic airways obstruction. *Am Rev Respir Dis*. 1992;146(1):177-184.
167. Bergelson JM, Cunningham JA, Droguett G, et al. Isolation of a common receptor for Coxsackie B viruses and adenoviruses 2 and 5. *Science*. 1997;275(5304):1320-1323.
168. Wang H, Li ZY, Liu Y, et al. Desmoglein 2 is a receptor for adenovirus serotypes 3, 7, 11 and 14. *Nat Med*. 2011;17(1):96-104.
169. Robinson C, Echavarría M. Adenoviruses. In: Versalovic J, Carroll K, Funke G, Jorgensen J, Landry M, eds. *Manual of Clinical Microbiology*. 10th ed. Washington, DC: ASM Press; 2011.
170. Smith CD, Craft DW, Shiromoto RS, Yan PO. Alternative cell line for virus isolation. *J Clin Microbiol*. 1986;24(2):265-268.
171. Woods GL, Young A. Use of A-549 cells in a clinical virology laboratory. *J Clin Microbiol*. 1988;26(5):1026-1028.
172. Rabalais GP, Stout GG, Ladd KL, Cost KM. Rapid diagnosis of respiratory viral infections by using a shell vial assay and monoclonal antibody pool. *J Clin Microbiol*. 1992;30(6):1505-1508.
173. Goldsmith CS, Miller SE. Modern uses of electron microscopy for detection of viruses. *Clin Microbiol Rev*. 2009;22(4):552-563.
174. Madeley CR. The emerging role of adenoviruses as inducers of gastroenteritis. *Pediatr Infect Dis*. 1986;5(1 suppl):S63-S74.
175. Brown M, Petric M, Middleton PJ. Diagnosis of fastidious enteric adenoviruses 40 and 41 in stool specimens. *J Clin Microbiol*. 1984;20(3):334-338.
176. Schmidt W, Schneider T, Heise W, et al. Stool viruses, coinfections, and diarrhea in HIV-infected patients. Berlin Diarrhea/Wasting Syndrome Study Group. *J Acquir Immune Defic Syndr Hum Retrovirol*. 1996;13(1):33-38.
177. Gray GC, McCarthy T, Lebeck MG, et al. Genotype prevalence and risk factors for severe clinical adenovirus infection, United States 2004-2006. *Clin Infect Dis*. 2007;45(9):1120-1131.
178. Thomas EE, Roscoe DL, Book L, et al. The utility of latex agglutination assays in the diagnosis of pediatric viral gastroenteritis. *Am J Clin Pathol*. 1994;101(6):742-746.
179. Kok T, Mickan LD, Burrell CJ. Routine diagnosis of seven respiratory viruses and *Mycoplasma pneumoniae* by enzyme immunoassay. *J Virol Methods*. 1994;50(1-3):87-100.
180. Wood SR, Sharp IR, de Jong JC, Verweij-Uijterwaal MW. Development and preliminary evaluation of an enzyme immunosorbent assay for the detection of adenovirus type 8. *J Med Virol*. 1994;44(4):348-352.
181. Hara M, Takao S, Fukuda S, Shimazu Y, Sasaki Y. Rapid diagnosis of adenovirus respiratory tract infections using the chromatographic immunoassay test in the pediatric outpatient setting. *Pediatr Infect Dis J*. 2010;29(3):267-269.
182. Simsir A, Grenebaum E, Nuovo G, Schulman LL. Late fatal adenovirus pneumonitis in a lung transplant recipient. *Transplantation*. 1998;65(4):592-594.
183. Goto E. Performance evaluation of detecting adenovirus by using rapid diagnostic kits among Japanese people. *J Infect Chemother*. 2012;18(3):361-369.
184. Arbizu RA, Aljomah G, Kozielski R, Baker SS, Baker RD. Intussusception associated with adenovirus. *J Pediatr Gastroenterol Nutr*. 2014;59(5):e41.
185. Wong S, Pabbaraju K, Pang XL, Lee BE, Fox JD. Detection of a broad range of human adenoviruses in respiratory tract samples using a sensitive multiplex real-time PCR assay. *J Med Virol*. 2008;80(5):856-865.
186. Heim A, Ebnet C, Harste G, Pring-Akerblom P. Rapid and quantitative detection of human adenovirus DNA by real-time PCR. *J Med Virol*. 2003;70(2):228-239.
187. Freymuth F, Vabret A, Cuvillon-Nimal D, et al. Comparison of multiplex PCR assays and conventional techniques for the diagnostic of respiratory virus infections in children admitted to hospital with an acute respiratory illness. *J Med Virol*. 2006;78(11):1498-1504.
188. Popowitch EB, O'Neill SS, Miller MB. Comparison of the Biofire FilmArray RP, Genmark eSensor RVP, Luminex xTAG RVPv1, and Luminex xTAG RVP fast multiplex assays for detection of respiratory viruses. *J Clin Microbiol*. 2013;51(5):1528-1533.
189. Echavarría M. Adenoviruses in immunocompromised hosts. *Clin Microbiol Rev*. 2008;21(4):704-715.

190. Morfin F, Dupuis-Girod S, Mundweiler S, et al. In vitro susceptibility of adenovirus to antiviral drugs is species-dependent. *Antivir Ther.* 2005;10(2):225-229.
191. Chiu CY, Yagi S, Lu X, et al. A novel adenovirus species associated with an acute respiratory outbreak in a baboon colony and evidence of coincident human infection. *MBio.* 2013;4(2):e00084.
192. Xu W, McDonough MC, Erdman DD. Species-specific identification of human adenoviruses by a multiplex PCR assay. *J Clin Microbiol.* 2000;38(11):4114-4120.
193. Pring-Akerblom P, Adrian T. Type- and group-specific polymerase chain reaction for adenovirus detection. *Res Virol.* 1994;145(1):25-35.
194. Adhikary AK, Inada T, Banik U, Numaga J, Okabe N. Identification of subgenus C adenoviruses by fiber-based multiplex PCR. *J Clin Microbiol.* 2004;42(2):670-673.
195. Lu X, Trujillo-Lopez E, Lott L, Erdman DD. Quantitative real-time PCR assay panel for detection and type-specific identification of epidemic respiratory human adenoviruses. *J Clin Microbiol.* 2013;51(4):1089-1093.
196. Buckwalter SP, Teo R, Espy MJ, et al. Real-time qualitative PCR for 57 human adenovirus types from multiple specimen sources. *J Clin Microbiol.* 2012;50(3):766-771.
197. Elnifro EM, Cooper RJ, Klapper PE, Bailey AS. PCR and restriction endonuclease analysis for rapid identification of human adenovirus subgenera. *J Clin Microbiol.* 2000;38(6):2055-2061.
198. Allard A, Albinsson B, Wadell G. Rapid typing of human adenoviruses by a general PCR combined with restriction endonuclease analysis. *J Clin Microbiol.* 2001;39(2):498-505.
199. Lin B, Vora GJ, Thach D, et al. Use of oligonucleotide microarrays for rapid detection and serotyping of acute respiratory disease-associated adenoviruses. *J Clin Microbiol.* 2004;42(7):3232-3239.
200. Ecker DJ, Sampath R, Blyn LB, et al. Rapid identification and strain-typing of respiratory pathogens for epidemic surveillance. *Proc Natl Acad Sci USA.* 2005;102(22):8012-8017.
201. Sarantis H, Johnson G, Brown M, Petric M, Tellier R. Comprehensive detection and serotyping of human adenoviruses by PCR and sequencing. *J Clin Microbiol.* 2004;42(9):3963-3969.
202. Ebner K, Pinsker W, Lion T. Comparative sequence analysis of the hexon gene in the entire spectrum of human adenovirus serotypes: phylogenetic, taxonomic, and clinical implications. *J Virol.* 2005;79(20):12635-12642.
203. Adhikary AK, Banik U, Numaga J, et al. Heterogeneity of the fibre sequence in subgenus C adenoviruses. *J Clin Pathol.* 2004;57(6):612-617.
204. Lu X, Erdman DD. Molecular typing of human adenoviruses by PCR and sequencing of a partial region of the hexon gene. *Arch Virol.* 2006;151(8):1587-1602.
205. Xu W, Erdman DD. Type-specific identification of human adenovirus 3, 7, and 21 by a multiplex PCR assay. *J Med Virol.* 2001;64(4):537-542.
206. Crawford-Miksza L, Schnurr DP. Analysis of 15 adenovirus hexon proteins reveals the location and structure of seven hypervariable regions containing serotype-specific residues. *J Virol.* 1996;70(3):1836-1844.
207. Biere B, Schweiger B. Human adenoviruses in respiratory infections: sequencing of the hexon hypervariable region reveals high sequence variability. *J Clin Virol.* 2010;47(4):366-371.
208. Madisch I, Wolfel R, Harste G, Pommer H, Heim A. Molecular identification of adenovirus sequences: a rapid scheme for early typing of human adenoviruses in diagnostic samples of immunocompetent and immunodeficient patients. *J Med Virol.* 2006;78(9):1210-1217.
209. Vabret A, Gouarin S, Joannes M, et al. Development of a PCR-and hybridization-based assay (PCR Adenovirus Consensus) for the detection and the species identification of adenoviruses in respiratory specimens. *J Clin Virol.* 2004;31(2):116-122.
210. Pehler-Harrington K, Khanna M, Waters CR, Henrickson KJ. Rapid detection and identification of human adenovirus species by adenoplex, a multiplex PCR-enzyme hybridization assay. *J Clin Microbiol.* 2004;42(9):4072-4076.
211. Pierce VM, Hodinka RL. Comparison of the GenMark Diagnostics eSensor respiratory viral panel to real-time PCR for detection of respiratory viruses in children. *J Clin Microbiol.* 2012;50(11):3458-3465.
212. McCarthy T, Lebeck MG, Capuano AW, Schnurr DP, Gray GC. Molecular typing of clinical adenovirus specimens by an algorithm which permits detection of adenovirus coinfections and intermediate adenovirus strains. *J Clin Virol.* 2009;46(1):80-84.
213. Wadell G. Molecular epidemiology of human adenoviruses. *Curr Top Microbiol Immunol.* 1984;110:191-220.
214. Adrian T, Becker M, Hierholzer JC, Wigand R. Molecular epidemiology and restriction site mapping of adenovirus 7 genome types. *Arch Virol.* 1989;106(1-2):73-84.
215. Li QG, Wadell G. Analysis of 15 different genome types of adenovirus type 7 isolated on five continents. *J Virol.* 1986;60(1):331-335.
216. Adrian T. Genome type analysis of adenovirus type 4. *Intervirology.* 1992;34(4):180-183.
217. Cooper RJ, Bailey AS, Killough R, Richmond SJ. Genome analysis of adenovirus 4 isolated over a six year period. *J Med Virol.* 1993;39(1):62-66.
218. Ariga T, Shimada Y, Ohgami K, et al. New genome type of adenovirus serotype 4 caused nosocomial infections associated with epidemic conjunctivitis in Japan. *J Clin Microbiol.* 2004;42(8):3644-3648.
219. McCarthy AJ, Bergin M, De Silva LM, Stevens M. Intravenous ribavirin therapy for disseminated adenovirus infection. *Pediatr Infect Dis J.* 1995;14(11):1003-1004.
220. Ljungman P. Treatment of adenovirus infections in the immunocompromised host. *Eur J Clin Microbiol Infect Dis.* 2004;23(8):583-588.
221. Chin J, Stallones RA, Lennette EH. The effect of gamma globulin on acute respiratory illness in military recruits. *Am J Epidemiol.* 1967;86(1):193-198.
222. Maisch B, Hufnagel G, Kolsch S, et al. Treatment of inflammatory dilated cardiomyopathy and (peri)myocarditis with immunosuppression and i.v. immunoglobulins. *Herz.* 2004;29(6):624-636.
223. Ramano A, Ladizinski E, Guarari-Rotman D, Revel M. Clinical effect of human fibroblast derived interferon in treatment of adenovirus epidemic keratoconjunctivitis and its complications. *Tex Rep Biol Med.* 1981;41:559-565.
224. Pauschinger M, Bowles NE, Fuentes-Garcia FJ, et al. Detection of adenoviral genome in the myocardium of adult patients with idiopathic left ventricular dysfunction. *Circulation.* 1999;99(10):1348-1354.
225. Sandkovsky U, Vargas L, Florescu DF. Adenovirus: current epidemiology and emerging approaches to prevention and treatment. *Curr Infect Dis Rep.* 2014;16(8):416.
226. Trousdale MD, Goldschmidt PL, Nobrega R. Activity of ganciclovir against human adenovirus type-5 infection in cell culture and cotton rat eyes. *Cornea.* 1994;13(5):435-439.
227. Buchdahl RM, Taylor P, Warner JD. Nebulised ribavirin for adenovirus pneumonia. *Lancet.* 1985;2(8463):1070-1071.
228. Cassano WF. Intravenous ribavirin therapy for adenovirus cystitis after allogeneic bone marrow transplantation. *Bone Marrow Transplant.* 1991;7(3):247-248.
229. Murphy GF, Wood DP Jr, McRoberts JW, Henslee-Downey PJ. Adenovirus-associated hemorrhagic cystitis treated with intravenous ribavirin. *J Urol.* 1993;149(3):565-566.
230. Kapelushnik J, Or R, Delukina M, et al. Intravenous ribavirin therapy for adenovirus gastroenteritis after bone marrow transplantation. *J Pediatr Gastroenterol Nutr.* 1995;21(1):110-112.
231. Sabroe I, McHale J, Tait DR, et al. Treatment of adenoviral pneumonitis with intravenous ribavirin and immunoglobulin. *Thorax.* 1995;50(11):1219-1220.
232. La Rosa AM, Champlin RE, Mirza N, et al. Adenovirus infections in adult recipients of blood and marrow transplants. *Clin Infect Dis.* 2001;32(6):871-876.
233. Ljungman P, Ribaud P, Eyrich M, et al. Cidofovir for adenovirus infections after allogeneic hematopoietic stem cell transplantation: a survey by the Infectious Diseases Working Party of the European Group for Blood and Marrow Transplantation. *Bone Marrow Transplant.* 2003;31(6):481-486.
234. Florescu DF, Pergam SA, Neely MN, et al. Safety and efficacy of CMX001 as salvage therapy for severe adenovirus infections in immunocompromised patients. *Biol Blood Marrow Transplant.* 2012;18(5):731-738.

235. Florescu DF, Keck MA. Development of CMX001 (Brincidofovir) for the treatment of serious diseases or conditions caused by dsDNA viruses. *Expert Rev Anti Infect Ther.* 2014;12(10):1171-1178.
236. Cooney MK, Hall CE, Fox JP. The Seattle virus watch. 3. Evaluation of isolation methods and summary of infections detected by virus isolations. *Am J Epidemiol.* 1972;96(4):286-305.
237. Hall CE, Brandt CD, Frothingham TE, et al. The virus watch program: a continuing surveillance of viral infections in metropolitan New York families. IX. A comparison of infections with several respiratory pathogens in New York and New Orleans families. *Am J Epidemiol.* 1971;94(4):367-385.
238. Crawford-Miksza LK, Schnurr DP. Adenovirus serotype evolution is driven by illegitimate recombination in the hyper-variable regions of the hexon protein. *Virology.* 1996;224(2):357-367.
239. Kajon AE, Mistchenko AS, Videla C, et al. Molecular epidemiology of adenovirus acute lower respiratory infections of children in the south cone of South America (1991-1994). *J Med Virol.* 1996;48(2):151-156.
240. Kajon AE, Wadell G. Sequence analysis of the E3 region and fiber gene of human adenovirus genome type 7h. *Virology.* 1996;215(2):190-196.
241. Larranaga C, Kajon A, Villagra E, Avendano LF. Adenovirus surveillance on children hospitalized for acute lower respiratory infections in Chile (1988-1996). *J Med Virol.* 2000;60(3):342-346.
242. Palomino MA, Larranaga C, Avendano LF. Hospital-acquired adenovirus 7h infantile respiratory infection in Chile. *Pediatr Infect Dis J.* 2000;19(6):527-531.
243. Gray GC, Setterquist SF, Jirsa SJ, Desjardin LE, Erdman DD. Emergent strain of human adenovirus endemic in Iowa. *Emerg Infect Dis.* 2005;11(1):127-128.
244. Esposito DH, Gardner TJ, Schneider E, et al. Outbreak of pneumonia associated with emergent human adenovirus serotype 14—Southeast Alaska, 2008. *J Infect Dis.* 2010;202(2):214-222.
245. Lewis PF, Schmidt MA, Lu X, et al. A community-based outbreak of severe respiratory illness caused by human adenovirus serotype 14. *J Infect Dis.* 2009;199(10):1427-1434.
246. Brosch L, Tchandja J, Marconi V, et al. Adenovirus serotype 14 pneumonia at a basic military training site in the United States, spring 2007: a case series. *Mil Med.* 2009;174(12):1295-1299.
247. Yun HC, Prakash V. Transmission of adenovirus serotype 14 in the health care setting. *Clin Infect Dis.* 2008;46(12):1935-1936.
248. Binn LN, Sanchez JL, Gaydos JC. Emergence of adenovirus type 14 in US military recruits—a new challenge. *J Infect Dis.* 2007;196(10):1436-1437.
249. Huang G, Yu D, Zhu Z, et al. Outbreak of febrile respiratory illness associated with human adenovirus type 14p1 in Gansu Province, China. *Influenza Other Respir Viruses.* 2013;7(6):1048-1054.
250. Anderson BD, Barr KL, Heil GL, Friary JA, Gray GC. A comparison of viral fitness and virulence between emergent adenovirus 14p1 and prototype adenovirus 14p strains. *J Clin Virol.* 2012;54(3):265-268.
251. Hough HS, Gong H, Kajon AE, et al. Genome sequences of human adenovirus 14 isolates from mild respiratory cases and a fatal pneumonia, isolated during 2006-2007 epidemics in North America. *Respir Res.* 2010;11:116.
- 251a. Lam E, Ramke M, Warnecke G, et al. Effective apical infection of differentiated human bronchial epithelial cells and induction of proinflammatory chemokines by the highly pneumotropic human adenovirus type 14p1. *PLoS ONE.* 2015;10(7):e0131201.
252. Calder JA, Erdman DD, Ackelsberg J, et al. Adenovirus type 7 genomic-type variant, New York City, 1999. *Emerg Infect Dis.* 2004;10(1):149-152.
253. van der Veen J, Oei KG, Abarbanel MF. Patterns of infections with adenovirus types 4, 7 and 21 in military recruits during a 9-year survey. *J Hyg (Lond).* 1969;67(2):255-268.
254. Raty R, Kleemola M, Melen K, Stenvik M, Julkunen I. Efficacy of PCR and other diagnostic methods for the detection of respiratory adenoviral infections. *J Med Virol.* 1999;59(1):66-72.
255. Mantjarvi R. Adenovirus infections in servicemen in Finland. *Ann Med Exp Biol Fenn.* 1966;44(suppl):1-43.
256. Chmielewicz B, Benzler J, Pauli G, et al. Respiratory disease caused by a species B2 adenovirus in a military camp in Turkey. *J Med Virol.* 2005;77(2):232-237.
257. Hwang SM, Park DE, Yang YI, et al. Outbreak of febrile respiratory illness caused by adenovirus at a South Korean military training facility: clinical and radiological characteristics of adenovirus pneumonia. *Jpn J Infect Dis.* 2013;66(5):359-365.
258. Yu P, Ma C, Nawaz M, et al. Outbreak of acute respiratory disease caused by human adenovirus type 7 in a military training camp in Shaanxi, China. *Microbiol Immunol.* 2013;57(8):553-560.
259. Fraser PK, Hatch LA. Outbreak of adenovirus infection in the Portsmouth Naval Command, 1958. *Br Med J.* 1959;1(5120):470-473.
260. Kajon AE, Dickson LM, Metzgar D, et al. Outbreak of febrile respiratory illness associated with adenovirus 11a infection in a Singapore military training cAMP. *J Clin Microbiol.* 2010;48(4):1438-1441.
261. Nishio O, Matsui K, Goh KT, Matsunaga Y, Inouye S. Prevalence of adenovirus types 3 and 7 antibodies in Singapore. *Jpn J Infect Dis.* 2001;54(3):128-129.
262. To KK, Tse H, Chan WM, et al. A novel psittacine adenovirus identified during an outbreak of avian chlamydiosis and human psittacosis: zoonosis associated with virus-bacterium coinfection in birds. *PLoS Negl Trop Dis.* 2014;8(12):e3318.
263. Kayali G, Ortiz EJ, Chorazy ML, Gray GC. Lack of evidence of avian adenovirus infection among turkey workers. *J Agromedicine.* 2009;14(3):299-305.
264. Prado V, O'Ryan ML. Acute gastroenteritis in Latin America. *Infect Dis Clin North Am.* 1994;8(1):77-106.
265. Ankers HE, Klapper PE, Cleator GM, Bailey AS, Tullo AB. The role of a rapid diagnostic test (adenovirus immune dot-blot) in the control of an outbreak of adenovirus type 8 keratoconjunctivitis. *Eye (Lond).* 1993;7(suppl Pt 3):15-17.
266. Klapper PE, Cleator GM. Adenovirus cross-infection: a continuing problem. *J Hosp Infect.* 1995;30(suppl):262-267.
267. Richmond S, Burman R, Crosdale E, et al. A large outbreak of keratoconjunctivitis due to adenovirus type 8. *J Hyg (Lond).* 1984;93(2):285-291.
268. Holladay RC, Campbell GD Jr. Nosocomial viral pneumonia in the intensive care unit. *Clin Chest Med.* 1995;16(1):121-133.
269. Larsen RA, Jacobson JT, Jacobson JA, Strikas RA, Hierholzer JC. Hospital-associated epidemic of pharyngitis and conjunctivitis caused by adenovirus (21/H21 + 35). *J Infect Dis.* 1986;154(4):706-709.
270. Blanshard C, Francis N, Gazzard BG. Investigation of chronic diarrhoea in acquired immunodeficiency syndrome. A prospective study of 155 patients. *Gut.* 1996;39(96):824-832.
271. Stallones R, Hilleman M, Gauld R, Warfield M, Anderson S. Adenovirus (R1-APC-ARD) vaccine for prevention of acute respiratory illness: II. field evaluation. *JAMA.* 1957;163:9-15.
272. Hilleman M, Greenberg J, Warfield M, Anderson S, Glabere R. Second field evaluation of bivalent types 4 and 7 adenovirus vaccines. *Arch Intern Med.* 1958;102:428-436.
273. Culver JO, Lennette EH, Flintjer JD. Adenovirus vaccine; a field evaluation of protective capacity against respiratory disease. *Am J Hyg.* 1959;69(2):120-126.
274. Sherwood RW, Buescher EL, Nitz RE, Cooch JW. Effects of adenovirus vaccine on acute respiratory disease in U. S. Army recruits. *JAMA.* 1961;178:1125-1127.
275. Rollison DE, Page WF, Crawford H, et al. Case-control study of cancer among US Army veterans exposed to simian virus 40-contaminated adenovirus vaccine. *Am J Epidemiol.* 2004;160(4):317-324.
276. Hitchcock G, Tyrrell DA, Bynoe ML. Vaccination of man with attenuated live adenovirus. *J Hyg (Lond).* 1960;58:277-282.
277. Couch RB, Chanock RM, Cate TR, et al. Immunization with types 4 and 7 adenovirus by selective infection of the intestinal tract. *Am Rev Respir Dis.* 1963;88(suppl):394-403.
278. Chanock RM, Ludwig W, Heubner RJ, Cate TR, Chu LW. Immunization by selective infection with type 4 adenovirus grown in human diploid tissue cultures. I. Safety and lack of oncogenicity and tests for potency in volunteers. *JAMA.* 1966;195(6):445-452.

279. Trentin JJ, Van Hoosier GL Jr, Samper L. The oncogenicity of human adenoviruses in hamsters. *Proc Soc Exp Biol Med.* 1968;127(3):683-689.
280. McBride WD, Wiener A. In vitro transformation of hamster kidney cells by human adenovirus type 12. *Proc Soc Exp Biol Med.* 1964;115:870-874.
281. Edmondson WP, Purcell RH, Gundelfinger BF, et al. Immunization by selective infection with type 4 adenovirus grown in human diploid tissue culture. II. Specific protective effect against epidemic disease. *JAMA.* 1966;195(6):453-459.
282. Rosenbaum MJ, De Berry P, Sullivan EJ, et al. Characteristics of vaccine-induced and natural infection with adenovirus type 4 in naval recruits. *Am J Epidemiol.* 1968;88(1):45-54.
283. Huebner R. *The problem of oncogenicity of adenoviruses* The First International Conference on Vaccines Against Viral and Rickettsial Diseases of Man: PanAmerican Health Org; Nov 7-11, 1966.
284. Gilden RV, Kern J, Lee YK, et al. Serologic surveys of human cancer patients for antibody to adenovirus T antigens. *Am J Epidemiol.* 1970;91(5):500-509.
285. McAllister RM, Gilden RV, Green M. Adenoviruses in human cancer. *Lancet.* 1972;1(7755):831-833.
286. Division of Cancer Cause and Prevention. *The Virus Cancer Program.* Bethesda, MD: National Cancer Institute; 1974.
287. Top F, Grossman R, Bartelloni P, et al. Immunization with live types 7 and 4 adenovirus vaccines. I. Safety, infectivity, antigenicity, and potency of adenovirus type 7 vaccine in humans. *J Infect Dis.* 1971;124(2):148-154.
288. Top F, Dudding B, Russell P, Buescher E. Control of respiratory disease in recruits with types 4 and 7 adenovirus vaccines. *Am J Epidemiol.* 1971;94(2):142-146.
289. Top F, Buescher E, Bancroft W, Russell P. Immunization with live types 7 and 4 adenovirus vaccines. II. Antibody response and protective effect against acute respiratory disease due to adenovirus type 7. *J Infect Dis.* 1971;124(2):155-160.
290. Van der Veen J, Dijkman J. Association of type 21 adenovirus with acute respiratory illness in military recruits. *Am J Hyg.* 1962;76:149-159.
291. Dudding BA, Bartelloni PJ, Scott RM, et al. Enteric immunization with live adenovirus type 21 vaccine. I. Tests for safety, infectivity, immunogenicity, and potency in volunteers. *Infect Immun.* 1972;5(3):295-299.
292. Scott RM, Dudding BA, Romano SV, Russell PK. Enteric immunization with live adenovirus type 21 vaccine. II. Systemic and local immune responses following immunization. *Infect Immun.* 1972;5(3):300-304.
293. Top F, Brandt W, Russell P. Adenovirus ARD in basic combat trainees. *Research in Biological and Medical Sciences: Annual Progress Report.* 1976;462-465.
294. Takafuji ET, Gaydos JC, Allen RG, Top FH Jr. Simultaneous administration of live, enteric-coated adenovirus types 4, 7 and 21 vaccines: safety and immunogenicity. *J Infect Dis.* 1979;140(1):48-53.
295. Schwartz AR, Togo Y, Hornick RB. Clinical evaluation of live, oral types 1, 2, and 5 adenovirus vaccines. *Am Rev Respir Dis.* 1974;109(2):233-239.
296. Lee T, Miller R. *Remanufacture of adenovirus vaccines type 4 and 7.* Depart of Defense Recruit and Trainee Healthcare Symposium 2002;2002 April 15-18; Towson, MD; 2002.
297. Dewitt C, Laural V, Walter E, et al. *Adenovirus type 4 outbreak in military basic trainees: reemergence of disease due to lack of vaccine availability.* International Conference on Emerging Infectious Diseases. Atlanta, GA; July 16-19, 2000.
298. McNeill K, Gaydos J, Watts B, Goldenbaum M, Poropatich R. *Clinical presentations of otherwise healthy young soldiers with reemerging adenovirus type 4-associated acute respiratory disease* International Conference on Emerging Infectious Diseases. Atlanta, GA; July 16-19, 2000.
299. McNeill KM, Ridgely Benton F, Monteith SC, Tuchscherer MA, Gaydos JC. Epidemic spread of adenovirus type 4-associated acute respiratory disease between U.S. Army installations. *Emerg Infect Dis.* 2000;6(4):415-419.
300. Ludwig SL, Brundage JF, Kelley PW, et al. Prevalence of antibodies to adenovirus serotypes 4 and 7 among unimmunized US Army trainees: results of a retrospective nationwide seroprevalence survey. *J Infect Dis.* 1998;178(6):1776-1778.
301. Sanchez JL, Binn LN, Innis BL, et al. Epidemic of adenovirus-induced respiratory illness among US military recruits: Epidemiologic and immunologic risk factors in healthy, young adults. *J Med Virol.* 2001;65(4):710-718.
302. Gray GC, Goswami PR, Malasig MD, et al. Adult adenovirus infections: loss of orphaned vaccines precipitates military respiratory disease epidemics. For the Adenovirus Surveillance Group. *Clin Infect Dis.* 2000;31(3):663-670.
303. Radin JM, Hawksworth AW, Blair PJ, et al. Dramatic decline of respiratory illness among US military recruits after the renewed use of adenovirus vaccines. *Clin Infect Dis.* 2014;59(7):962-968.
304. Russell KL, Hawksworth AW, Ryan MA, et al. Vaccine-preventable adenoviral respiratory illness in US military recruits, 1999-2004. *Vaccine.* 2006;24(15):2835-2842.
305. Gray GC, Callahan JD, Hawksworth AW, Fisher CA, Gaydos JC. Respiratory diseases among U.S. military personnel: countering emerging threats. *Emerg Infect Dis.* 1999;5(3):379-385.
306. Institute of Medicine. *Urgent attention needed to restore lapsed adenovirus vaccine availability.* Washington, DC: Institute of Medicine Report; 2000.
307. Katz SL. A tale of two vaccines. *Clin Infect Dis.* 2000;31(3):671-672.
308. Russell P. Adenovirus infection is not trivial [letter]. *US Med.* 1998 November.
309. Lueck S. *Outbreak. A common virus, a military recruit—and a mysterious death.* *The Wall Street Journal.* 2001 July 13;Sect. 1.
310. Centers for Disease Control and Prevention (CDC). Two fatal cases of adenovirus-related illness in previously healthy young adults—Illinois, 2000. *MMWR Morb Mortal Wkly Rep.* 2001;50(26):553-555.
311. Department of Defense Armed Forces Epidemiological Board. *Recommendation for the use of adenovirus vaccine.* Falls Church, VA: Department of Defense Armed Forces Epidemiological Board; 1998.
312. Purkayastha A, Su J, McGraw J, et al. Genomic and bioinformatics analyses of HAdV-4vac and HAdV-7vac, two human adenovirus (HAdV) strains that constituted original prophylaxis against HAdV-related acute respiratory disease, a reemerging epidemic disease. *J Clin Microbiol.* 2005;43(7):3083-3094.
313. Lyons A, Longfield J, Kuschner R, et al. A double-blind, placebo-controlled study of the safety and immunogenicity of live, oral type 4 and type 7 adenovirus vaccines in adults. *Vaccine.* 2008;26(23):2890-2898.
314. Potter RN, Cantrell JA, Mallak CT, Gaydos JC. Adenovirus-associated deaths in US military during postvaccination period, 1999-2010. *Emerg Infect Dis.* 2012;18(3):507-509.
315. Hoke CH Jr, Snyder CE Jr. History of the restoration of adenovirus type 4 and type 7 vaccine, live oral (Adenovirus Vaccine) in the context of the Department of Defense acquisition system. *Vaccine.* 2013;31(12):1623-1632.
316. Towle A, Azad A. *Restoring adenovirus vaccine.* *Contract Pharma.* March 4, 2015.
317. Tint H, Stone JL, Minecci LC, Rubin BA. Type 4 adenovirus vaccine, live, prepared in human diploid cell system for oral administration. *Prog Immunobiol Stand.* 1969;3:113-122.
318. Barr Labs, Inc. *Package insert-adenovirus type 4 and adenovirus type 7 vaccine, live, oral.* March 2011, 2011. <<http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/ucm247508.htm>>.
319. Rhoads JL, Bix DL, Wright DC, et al. Safety and immunogenicity of multiple conventional immunizations administered during early HIV infection. *J Acquir Immune Defic Syndr.* 1991;4(7):724-731.
320. Stanley ED, Jackson GG. Spread of enteric live adenovirus type 4 vaccine in married couples. *J Infect Dis.* 1969;119(1):51-59.
321. Lee S, Hung P. Vaccines for control of respiratory disease caused by adenoviruses. *Rev Med Virol.* 1993;3:209-216.
322. Smith TJ, Buescher EL, Top FH Jr, Altemeier WA, McCown JM. Experimental respiratory infection with type 4 adenovirus vaccine in volunteers: clinical and immunological responses. *J Infect Dis.* 1970;122(4):239-248.

323. Gutekunst RR, Heggie AD. Viremia and viruria in adenovirus infections. Detection in patients with rubella or rubelliform illness. *N Engl J Med.* 1961;264:374-378.
324. Kuschner RA, Russell KL, Abuja M, et al. A phase 3, randomized, double-blind, placebo-controlled study of the safety and efficacy of the live, oral adenovirus type 4 and type 7 vaccine, in U.S. military recruits. *Vaccine.* 2013;31(28):2963-2971.
325. Barratt-Boyes SM, Soloff AC, Gao W, et al. Broad cellular immunity with robust memory responses to simian immunodeficiency virus following serial vaccination with adenovirus 5- and 35-based vectors. *J Gen Virol.* 2006;87(Pt 1):139-149.
326. Mueller RE, Muldoon RL, Jackson GG. Communicability of enteric live adenovirus type 4 vaccine in families. *J Infect Dis.* 1969;119(1):60-66.
327. Schwartz AR, Togo Y, Hornick RB. Clinical evaluation of live, oral types 1, 2, and 5 adenovirus vaccines. *Am Rev Respir Dis.* 1974;109(2):233-239.
328. Van der Veen J, Abarbanel N, Oki K. Vaccination with live type 4 adenoviruses: evaluation of antibody responses and protective efficacy. *J Hyg (Lond).* 1968;66:499-511.
329. Peckinpaugh RO, Pierce WE, Rosenbaum MJ, Edwards EA, Jackson GG. Mass enteric live adenovirus vaccination during epidemic ARD. *JAMA.* 1968;205(1):75-80.
330. Rosenbaum M, Edwards E, Hoefler D, et al. Recent experiences with live adenovirus vaccines in Navy recruits. *Mil Med.* 1975;140:251-257.
331. Gooch W, Mogabgab W. Simultaneous oral administration of live adenovirus types 4 and 7 vaccines. *Am J Epidemiol.* 1972;25:388-394.
332. Meiklejohn G. Viral respiratory disease at Lowry Air Force Base in Denver, 1952-1982. *J Infect Dis.* 1983;148(5):775-784.
333. Mueller RE, Muldoon RL, Jackson GG. Communicability of enteric live adenovirus type 4 vaccine in families. *J Infect Dis.* 1969;119(1):60-66.
334. Loker EF Jr, Hodges GR, Kelly DJ. Fatal adenovirus pneumonia in a young adult associated with ADV-7 vaccine administered 15 days earlier. *Chest.* 1974;66(2):197-199.
335. Departments of the Army (AR40-562) TNB61 TNB, The Air Force (Joint Instruction 48-110), and Transportation (CG COMDTINST M623.4E). *Immunizations and Chemoprophylaxis.* Washington, DC: US Government Printing Office; 1995 p. 1996-404-611(November 1):20059.
336. Collis PB, Dudding BA, Winter PE, Russell PK, Buescher EL. Adenovirus vaccines in military recruit populations: a cost-benefit analysis. *J Infect Dis.* 1973;128(6):745-752.
337. Howell M, Nang R, Gaydos C, Gaydos J. Prevention of adenoviral acute respiratory disease in Army recruits: cost-effectiveness of a military vaccination policy. *Am J Prev Med.* 1998;14:168-175.
338. Hyer RN, Howell MR, Ryan MA, Gaydos JC. Cost-effectiveness analysis of reacquiring and using adenovirus types 4 and 7 vaccines in naval recruits. *Am J Trop Med Hyg.* 2000;62(5):613-618.
339. Mayo MA, Haenni AL. Report from the 36th and the 37th meetings of the Executive Committee of the International Committee on Taxonomy of Viruses. *Arch Virol.* 2006;151(5):1031-1037.
340. Ryan MA, Christian RS, Wohlrahe J. Handwashing and respiratory illness among young adults in military training. *Am J Prev Med.* 2001;21(2):79-83.
341. Gaydos JC. Returning to the past: respiratory illness, vaccines, and handwashing. *Am J Prev Med.* 2001;21(2):150-151.
342. Lee T, Jordan NN, Sanchez JL, Gaydos JC. Selected nonvaccine interventions to prevent infectious acute respiratory disease. *Am J Prev Med.* 2005;28(3):305-316.
343. Wilcox WC, Ginsberg HS. Production of specific neutralizing antibody with soluble antigens of type 5 adenovirus. *Proc Soc Exp Biol Med.* 1963;114:37-42.
344. Couch RB, Kasel JA, Perreira HG, Haase AT, Knight V. Induction of immunity in man by crystalline adenovirus type 5 capsid antigens. *Proc Soc Exp Biol Med.* 1973;143(4):905-910.
345. Weaver EA. Vaccines within vaccines: the use of adenovirus types 4 and 7 as influenza vaccine vectors. *Hum Vaccin Immunother.* 2014;10(3):544-556.
346. Wong G, Richardson JS, Pillet S, et al. Adenovirus-vectored vaccine provides postexposure protection to Ebola virus-infected nonhuman primates. *J Infect Dis.* 2015;212(suppl 2):S379-S383.
347. Guo X, Deng Y, Chen H, et al. Systemic and mucosal immunity in mice elicited by a single immunization with human adenovirus type 5 or 41 vector-based vaccines carrying the spike protein of Middle East respiratory syndrome coronavirus. *Immunology.* 2015;145(4):476-484.
348. Xiang K, Ying G, Yan Z, et al. Progress on adenovirus-vectored universal influenza vaccines. *Hum Vaccin Immunother.* 2015;11(5):1209-1222.
349. Majhen D, Calderon H, Chandra N, et al. Adenovirus-based vaccines for fighting infectious diseases and cancer: progress in the field. *Hum Gene Ther.* 2014;25(4):301-317.
350. Deal C, Pekosz A, Ketner G. Prospects for oral replicating adenovirus-vectored vaccines. *Vaccine.* 2013;31(32):3236-3243.
351. Tucker SN, Tingley DW, Scallan CD. Oral adenoviral-based vaccines: historical perspective and future opportunity. *Expert Rev Vaccines.* 2008;7(1):25-31.
352. Zhang J. Advances and future challenges in recombinant adenoviral vectored H5N1 influenza vaccines. *Viruses.* 2012;4(11):2711-2735.
353. Baden LR, Walsh SR, Seaman MS, et al. First-in-human evaluation of the safety and immunogenicity of a recombinant adenovirus serotype 26 HIV-1 Env vaccine (IPCAVD 001). *J Infect Dis.* 2013;207(2):240-247.
354. Majhen D, Calderon H, Chandra N, et al. Adenovirus-based vaccines for fighting infectious diseases and cancer: progress in the field. *Hum Gene Ther.* 2014;25:301-317.
355. Wold WS, Toth K. Adenovirus vectors for gene therapy, vaccination and cancer gene therapy. *Curr Gene Ther.* 2013;13(6):421-433.