

The epidemiologic factors associated with breast density: A review

Dong-Man Ye, Tao Yu

Department of Medical Imaging, Cancer Hospital of China Medical University, Liaoning Cancer Hospital and Institute, Shenyang, Liaoning Province, China

In recent years, some studies have evaluated the epidemiologic factors associated with breast density. However, the variant and inconsistent results exist. In addition, breast density has been proved to be a significant risk factor associated with breast cancer. Our review summarized the published studies and emphasized the crucial factors including epidemiological factors associated with breast density. In addition, we also discussed the potential reasons for the discrepant results with risk factors. To decrease the incidence and mortality rates for breast cancer, in clinical practice, breast density should be included for clinical risk models in addition to epidemiological factors, and physicians should get more concentrate on those women with risk factors and provide risk-based breast cancer screening regimens.

Key words: Breast cancer, breast density, epidemiological factors

How to cite this article: Ye DM, Yu T. The epidemiologic factors associated with breast density: A review. *J Res Med Sci* 2022;27:53.

INTRODUCTION

Breast cancer has been a worldwide burden of women's health, as 2.3 million newly cases were diagnosed in 2020.^[1] Breast density has been proved to be a significant risk factor associated with breast cancer.^[2,3] To evaluate the factors associated with breast density, it has been regarded as a more important understanding of the etiology and potential pathways of breast cancer. Several epidemiologic studies have found that women with high breast density were more likely to develop breast cancer than those with low breast density.^[3-5] In recent years, some studies have evaluated the epidemiologic factors associated with high breast density, such as age, body mass index (BMI), reproductive factors, and life style.^[6-10] However, efforts are still warranted, as studies from different countries and ethnicities may contribute to inconsistent results.

The breasts were composed of fat, connective, epithelial tissues that reflected on the mammography according

to their X-ray attenuation coefficient.^[11] Fat was represented as light areas on mammography, while fibroglandular was dark. To assess breast density for consistent standard by radiologists, some measurement methods were proposed, such as Wolfe grades,^[12] Tabar grades,^[13] Breast Imaging Reporting and Data System (BIRADS),^[14] and quantitative measurement depending on the proportion of area of dense breast tissue through planimetry, and computer assisted methods.^[15] BIRADS was widely used in clinical practice and the 5th edition was proposed in 2013.^[16] The 5th edition BIRADS system has four categories: (a) almost entirely fatty; (b) scattered areas of fibro glandular density; (c) heterogeneously dense; (d) extremely dense. High breast density including categories c and d may obscure detection of small masses and lower the sensitivity. Studies have reported that the sensitivity of detecting breast cancer by radiologists was 30%–64% in high density while 80%–98% in low density.^[17,18] Higher breast density had the possibility to mask the tumor and lower the sensitivity for detecting cancer.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

Access this article online

Quick Response Code:



Website:

www.jmsjournal.net

DOI:

10.4103/jrms.jrms_962_21

Address for correspondence: Dr. Tao Yu, Department of Medical Imaging, Cancer Hospital of China Medical University, Liaoning Cancer Hospital and Institute, No. 44 Xiaohuyan Road, Dadong District, Shenyang, Liaoning Province 110042, China.
E-mail: yutao@cancerhosp-in-cmu.com

Submitted: 31-Oct-2021; **Revised:** 14-Jan-2022; **Accepted:** 26-Jan-2022; **Published:** 29-Jul-2022

Previous studies have elucidated that higher breast density might be associated with more aggressive characteristics, such as histopathological grade, tumor size, molecular subtypes.^[19-27]

Breast density was modifiable and highly heritable. It was not clear whether the epidemiologic risk factors related to breast cancer were mediated through breast density. It is necessary to review previous epidemiologic studies and clarify the association between these epidemiologic factors and breast density. In our review, epidemiologic factors such as age, race, BMI, reproductive factors, hormone exposures, diet and lifestyle, influenced on breast density were addressed.

EPIDEMIOLOGIC FACTORS

Age and race

The association between age and breast density has been investigated in many studies. It was suggested that breast density decreased with aging.^[28-34] The International Consortium on Mammographic Density investigated over 11,000 women from 22 countries worldwide and found that the breast density of both premenopausal women and postmenopausal women declined with increasing age.^[35] The meta-analyses and pooled analyses showed that the decrease of percent breast density was 0.24 cm among premenopausal women and 0.38 cm among postmenopausal women for the 10-year increase of age.^[35] Some studies found that the association between decreased breast density and increasing age was nonlinear,^[28,29,31,32] and breast density was at plateau at the age of 65 years.^[28,32] As age was regarded as one of the most important factors reflecting breast density, it was used to adjust for multivariable models in many studies. However, there were some outliers. Kang *et al.* reported that there was not significant association between breast density and age in women in the United Arab Emirates.^[36] Another study based on the New York University Langone Medical Center in 2008 found that a small proportion (19%) of young women <40 years old had low breast density, while 44% of women at the age of 60s and 36% of women at the age of 70s had high breast density.^[37] In several studies limited the sample size, different race and geographic area, the association between age and breast density was not clear, however, generally, the age was inversely associated with breast density.

The studies based on ethnic differences in breast density have been investigated. A number of studies have been investigated to find the association between breast density and races. Carmen *et al.*^[38] compared the breast density among white, African Americans, and Asian women and did not find any differences across ethnic groups. Ursin *et al.*^[39] reported that the breast density in Asians

was higher than African Americans. After adjusting for age, BMI, and other factors, Chen *et al.*^[40] investigated that the percent breast density in Asian women was lower than African women but was not varied from Caucasians. Another study reported that the percent breast density of Chinese women was significantly higher than Malays and Indians.^[41] McCarthy *et al.* found that black women had higher breast density than white women after adjusting some risk factors.^[42] Generally speaking, the breast density varied across different ethnic groups, and Asian women had higher breast density than western women.^[43,44] Thus, the relationship of race to breast density may explain variation in breast cancer risk among different ethnic groups.

Body mass index

Most previous studies have investigated the inverse association between breast density and BMI.^[45-52] Especially, BMI was negatively related to breast density in both premenopausal and postmenopausal women.^[45] However, not all of the studies reported the consistent results, as Jeon *et al.* found that there was no association between BMI and breast density in premenopausal women.^[53] The inconsistent results remained unexplained but largely caused by differences among the population. Compared to fibrous and glandular tissue, the amount of fat increased with the increasement of BMI, thus the absolute measurement of density was considered as a more important method.^[54]

A number of studies have also been conducted to find whether the adiposity at early age influences the breast density. Some studies have reported the inverse association of adiposity at the early age with breast density.^[7,55-59] Alimujiang *et al.*^[55] found that a 1 kg/m² increase in BMI at age 10 was related to a 6.4% decrease in volumetric percent density and a 6.9% increase in nondense volume. The author also found that higher BMI at age 18 was associated with lower breast density in premenopausal women.^[60] However, in another study, Rice^[61] have found that there was not significant association between body fatness during childhood and breast density among Mexican women. The possible mechanism was adiposity in earlier life could modulate hormone exposure and growth factor levels, therefore, the breast cellular proliferation and breast density tissue development were influenced.^[62-65]

Reproductive factors

Reproductive factors included menopause status, age at menarche, age at menopause, parity, breastfeeding, age at first birth, number of children, and so on. A large number of studies have investigated the reproductive factors associated with breast density.

Menopause status was one of the most important factors influencing the breast density. As lots of studies have been

reported, postmenopausal women usually had lower breast density and premenopausal women had higher density.^[53,66-72] A longitudinal study included postmenopausal women and age-matched group of premenopausal women has investigated that menopause led to a reduction in the area of dense tissue, and an increase of nondense area.^[71] Previous studies have reported that the average annual decline in percent density was 0.5%–2% and the maximum reduction was occurred during the menopausal transition.^[30-32,71] According to Pike's hypothesis, the decline of breast density over the menopause was caused by the decreased rate of breast tissue aging.^[73] Many studies had used subgroups analysis by stratifying the population based on menopause status to correct for confounding factors.

Age at menarche and menopause represented the time of a women exposure to endogenous and exogenous hormones that could influence breast tissue proliferation. The association between age at menarche and breast density was inconsistent reported by previous studies.^[10,51,66,74-82] Some studies investigated the positive relationship,^[51,80-82] while others reported no association between age at menarche and breast density.^[32,74,77-79] Tehranifar *et al.* had found the inverse association between age at menarche and breast density.^[66] Some studies have reported that late age at menopause was inversely associated with breast density,^[83,84] while others investigated no significant association.^[33,85,86] The evidence was far from conclusive, further researches were still warranted for verification in large sample size.

The association between other reproductive factors, such as parity, age at first pregnancy, and breastfeeding, and breast density has been far from conclusive among people from different races. Several studies have investigated that nulliparity or lower parity was negatively associated with breast density,^[33,53,68,80,87] while others found no significant association.^[67,77] Some studies showed that the age at first birth was positively associated with breast density,^[33,69,74,88] however, other studies investigated the inverse association.^[80,89] No significant association had been found between age at first birth and breast density.^[67,77] Based on the Pikes model, the study had reported that later age at first birth could increase the cumulative exposure to breast density,^[73] however, in different races, exceptions could still exist. Breastfeeding has been regarded as a protective factor for breast cancer in many studies, as during lactation, the level of prolactin increased and promotes cell proliferation.^[90] Sung *et al.* have reported the positive association between breastfeeding and breast density,^[53] while Yaghjian *et al.* investigated that there was positive association between breastfeeding and absolute dense and nondense areas among premenopausal women.^[72] Further researches were needed to clarify the association between breastfeeding and breast density.

Hormone exposures

Breast density could be influenced by hormonal environment, such as hormone replacement therapy (HRT), oral contraception that were used widely. Some studies have reported that HRT may increase the breast density among a substantial proportion of women.^[91-94] The influence of different HRTs on the breast density needs more evidence to clarify. Tibolone was commonly used as a tissue-specific compound for treatment in postmenopausal women. It could rapidly convert into 2 estrogenic metabolites and the $\Delta 4$ isomer, which has progestogenic and androgenic activities.^[95] *In vitro* and *in vivo* data have suggested that the effect of tibolone on breast density was different from conventional HRT.^[96] Lundström *et al.* reported that women receiving continuous combined HRT had higher breast density than those receiving tibolone and placebo treatment.^[92] However, another study illustrated that tibolone treatment was associated with lower breast density than continuous combination HRT among postmenopausal women.^[97] Junkermann *et al.*^[98] conducted a prospective, randomized study to evaluate the difference regimens between continuous combined HRT and sequential combined HRT, and the results showed that no significant difference between two treatment groups. Another study investigated four different regimens, including Group I which received conjugated equine estrogens (CEE), Group II which received CEE and continuous medroxyprogesterone acetate (MPA), Group III which received CEE and cyclic MPA, Group IV which received tibolone treatment, the results showed that there was significant difference between Group II and Group IV which indicated that continuous combination HRT was more likely to be associated with increased breast density than other forms of HRT regimens.^[99] Interestingly, Harvey *et al.*^[100] compared the transdermal and oral HRT associated with breast density and found that oral HRT had a higher incidence of increased breast density than transdermal HRT. Some studies also investigated whether the suspension of HRT influences breast density. Colacurci *et al.*^[101] found that the suspension of HRT for 3 weeks might decrease the increased breast density before, while Weaver *et al.*^[102] had an inverse conclusion that the cessation of HRT did not change breast density. However, it is worth to point that the change of breast density was dynamic during HRT, as it is increased with initiation and decreased with discontinuation.^[103]

Some studies investigated the effect of different hormones, such as progestogen, oestrogen, corticoid, androgens, and so on. Gabrielson^[104] emphasized that baseline breast density was influenced by hormones from the progestogen, estrogen, and corticoid pathways, however, the main driver was androgens. Another cross-sectional study reported that progesterone was significant associated with percentage

of breast density than estrogen among premenopausal women.^[105]

White *et al.*^[106] found that breast tissue is more radiographically dense in the luteal phase than in the follicular phase among premenopausal women aged 40–49, as some pathologic studies illustrated that epithelial cell proliferation, lobule size, and stromal edema are all greater in the luteal phase for breast tissues.^[107-109] Colacurci *et al.* found that suspension of HRT for about 3 weeks could increase breast density, however, the included samples were limited.^[101]

Diet and lifestyle

Previous epidemiologic studies have reported inconsistent findings about dietary, as the food was consumed by some combinations and nutrients intakes were highly related by interactive and synergistic effect. A study had found a weak and negative association between vegetables intake and breast density among Japanese women.^[110] Another study found that decreased total or saturated fat intake or cholesterol intake was positively related to dense area of breast density.^[111]

Some studies were conducted among Mediterranean population.^[112-116] A study compared three patterns, such as a fruit–vegetable–cereal pattern, a salad–sauce–pasta/grain pattern, and a meat–starch pattern, and found that the fruit–vegetable–cereal and salad–sauce–pasta/grain patterns were negatively associated with breast density among smokers.^[112] Another study found that high breast density was inversely associated with increased consumption of vegetables, olive oil, frequent intakes of cheese, high intakes of b-carotene, Vitamin C, calcium, and potassium.^[113] Castelló *et al.* examined the difference between two different patterns (Western and Mediterranean) associated with breast density and found that women with western pattern were more likely to have higher breast density, while no significant association was found between Mediterranean pattern and breast density.^[115]

Some studies not only concentrated on the dietary patterns but also the energy intake. A study investigated the positive association between energy intake and percent breast density during adulthood, and the effect of high fat and sugar dietary pattern greatly attenuated considering the total energy.^[117] Another study dietary energy density was positively associated with breast density, as each 1 kcal/g unit increased in food-only energy density was associated with a 25.9% increase in percent dense breast volume.^[118]

Vitamin D and calcium have been found to modulate cellular proliferation and differentiation in several tissues, including breast tissue.^[119,120] The studies were conducted

to find the association between Vitamin D and calcium and breast density. Be'rube *et al.* investigated that Vitamin D and calcium were inversely associated with breast density and suggested that they might regard as protective factors of breast cancer.^[121] The results about Vitamin D were highly similar to other studies.^[122,123] Vitamin D from sunlight and dietary intake was hydroxylated in the liver into 25-hydroxy Vitamin D (25[OH]D), and 25(OH) D was hydroxylated to the biologically active metabolite 1,25-dihydroxy vitamin D (1,25[OH] 2D) that probably in breast tissue.^[124,125] Green *et al.*^[126] found that there was not significant association between breast density and plasma 25(OH) D levels. However, another study found that high levels of 25(OH) D and Vitamin D substitution were negatively associated with breast density among premenopausal women.^[127]

Soy effect was proposed to be a factor on breast cell proliferation. Ursin investigated the inverse association between soy and breast density, as the breast density was significantly different between the highest and lowest quartiles of soy intake.^[88] A meta-analysis reported that isoflavones from soy and red clover would not change the breast density among postmenopausal women, however, it might cause a small increase among premenopausal women.^[128] Another study investigated the association between milk intake and breast density in premenopausal women; the results showed that the low/reduced-fat milk intake was negatively related to volumetric percent density.^[129]

Alcohol has been regarded as an established risk factor associated with breast cancer.^[130-132] The evidence of the association between alcohol consumption and breast density was inconsistent. Most studies reported the positive association between alcohol consumption and breast density.^[79,114,133-137] Jacobsen *et al.*^[137] evaluated the types of different alcohol and found that there were not significant difference among beer, wine, or fortified wine. Alcohol could change the levels of estrogen, that might contribute to the increase of breast density.^[138,139] However, another study found that alcohol consumption was not related to breast density among premenopausal women during adolescence and adulthood in the Nurses' Health Study II cohort.^[140] In contrast, the inverse association was found between breast density and alcohol consumption in two US birth cohorts.^[141] The evidence was not consistent, as the population from different geographic areas, the types of alcohol, detail timing of exposures, etc.

Previous studies have found inconsistent association between smoking and breast density. Some studies illustrated that smoking was inversely associated with breast density.^[142-145] Cigarette decreasing the levels of circulation estrogen levels might influence the breast

density.^[142] Another study reported the positive association among premenopausal women.^[77] Other three studies found no significant association.^[45,146,147]

Physical activity was supposed to reduce breast cancer through different mechanisms, such as weight loss, reduced hormone exposures and levels of insulin and insulin-like growth factor exposure, and so on.^[148-150] The association between physical activity and breast density has also been evaluated in recent years.^[67,146,151-156] Some studies have found that physical activity was not a determinant of breast density.^[67,151,153,156] Other studies found that physical activity actually impacts the breast density.^[152,154,155] Pilar *et al.*^[152] found that those who reported at least 3.5 h of physical inactivity per day had higher breast density compared to those who reported 1 h or less. Another longitudinal study was carried out among Mediterranean population and reported that the increasing levels of leisure time for physical activity were negatively associated with higher breast density.^[154] The biological mechanisms of the association between physical activity and breast density had not been established, however, several interesting possibilities might exist. Physical activity could reduce the hormone exposures that played important part in breast cell proliferation.^[157] In addition, the levels of insulin-like growth factor-I (IGF-I) modulated by physical activity could influence breast epithelial cell mitogens.^[158] The results have not been applied to different ethnic groups, further researches are needed to clarify the association between physical activity and breast density.

Others

Some studies evaluated the association between the breast density and other factors, such as residential traffic noise, education level, occupation, metabolic syndrome, socioeconomic status, and diabetes.^[68,159-168] Roswall *et al.*^[159] found no association between breast density and traffic noise exposure, either residential road or railway noise in a Danish cohort. Another study found that the socioeconomic status played a role in determining the amount of fat in breast.^[160] However, another study found early life socioeconomic status, including education and income was not significantly associated with breast density in adulthood.^[167] Lope *et al.*^[161] found that the occupational exposures were associated with breast density, the women that exposure to perchloroethylene, ionizing radiation, mold spores, or aliphatic/alcyclic hydrocarbon had higher breast density. Another study was conducted to evaluate the influence of occupations and occupational exposure to physical, chemical, and microbiological agents on MD, the results found that premenopausal women in Spanish occupationally exposed to fungicides, herbicides, and insecticides were more likely to have lower breast density.^[162] In addition, some diseases such as diabetes

and the metabolic syndrome were reported to be related to breast density.^[163,168]

Breast density has been regarded as having a polygenic basis.^[169] Genome-wide association studies have suggested that multiple low and moderate penetrance breast cancer susceptibility loci in women, such as single nucleotide polymorphisms (SNPs).^[169-177] However, it's not clear whether the etiological pathways increase the breast cancer risk due to the dense tissue. As Sartor *et al.*^[171] had found that compared to major homozygotes, minor homozygotes of SNPs rs9383589, CCDC170 and rs6557161, ESR1 were related to high breast density and poorer breast cancer survival in a large independent breast cancer cohort. Stone *et al.*^[174] also found associations between rs6001930 (MKL1) and both adjusted absolute dense and nondense areas and between rs17356907 (NTN4) and adjusted absolute nondense area. Vachon *et al.*^[169] found that LSP1-rs3817198 and RAD51 L1-rs10483813 were associated with breast density and in the same direction as the breast cancer association, while another study found no evidence for the association between majority of SNPs and breast density.^[173] Further studies based on the understanding the underlying mechanisms could help identify etiological pathways implicated in how dense density increased breast cancer risk.

CONCLUSION

Our review summarized the epidemiological factors associated with breast density. Although evidence has suggested that age, race, BMI, reproductive factors, hormone exposures, diet, and lifestyle have an influence on breast density, inconsistency results were observed due to a different size of populations and different race/ethnicity. Future studies with sufficient sample size and the same race/ethnicity and methodology are warranted. Given breast density was the most prevalent risk factor for developing breast cancer, in clinical practice, breast density should be included for clinical risk models and should be taken into account by the physicians. The consideration of breast density and other risk factors could provide comprehensive and individualized plans for breast screening, ultimately decrease the incidence and death rates for breast cancer.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, *et al.* Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in

- 185 countries. *CA Cancer J Clin* 2021;71:209-49.
2. Boyd NF, Guo H, Martin LJ, Sun L, Stone J, Fishell E, *et al.* Mammographic density and the risk and detection of breast cancer. *N Engl J Med* 2007;356:227-36.
3. McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: A meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:1159-69.
4. Boyd NF, Rommens JM, Vogt K, Lee V, Hopper JL, Yaffe MJ, *et al.* Mammographic breast density as an intermediate phenotype for breast cancer. *Lancet Oncol* 2005;6:798-808.
5. Boyd NF, Martin LJ, Bronskill M, Yaffe MJ, Duric N, Minkin S. Breast tissue composition and susceptibility to breast cancer. *J Natl Cancer Inst* 2010;102:1224-37.
6. Yaghjian L, Colditz G, Rosner B, Rich S, Egan K, Tamimi RM. Adolescent caffeine consumption and mammographic breast density in premenopausal women. *Eur J Nutr* 2020;59:1633-9.
7. Han Y, Berkey CS, Herman CR, Appleton CM, Alimujiang A, Colditz GA, *et al.* Adiposity change over the life course and mammographic breast density in postmenopausal women. *Cancer Prev Res (Phila)* 2020;13:475-82.
8. Alexeeff SE, Odo NU, McBride R, McGuire V, Achacoso N, Rothstein JH, *et al.* Reproductive factors and mammographic density: Associations among 24,840 women and comparison of studies using digitized film-screen mammography and full-field digital mammography. *Am J Epidemiol* 2019;188:1144-54.
9. Yaghjian L, Colditz G, Rosner B, Gasparova A, Tamimi RM. Associations of coffee consumption and caffeine intake with mammographic breast density. *Breast Cancer Res Treat* 2018;169:115-23.
10. McDonald JA, Tehranifar P, Flom JD, Terry MB, James-Todd T. Hair product use, age at menarche and mammographic breast density in multiethnic urban women. *Environ Health* 2018;17:1.
11. AL Mousa DS, Ryan EA, Mello-Thoms C, Brennan PC. What effect does mammographic breast density have on lesion detection in digital mammography? *Clin Radiol* 2014;69:333-41.
12. Wolfe JN. Breast patterns as an index of risk for developing breast cancer. *AJR Am J Roentgenol* 1976;126:1130-7.
13. Gram IT, Funkhouser E, Tabár L. The Tabár classification of mammographic parenchymal patterns. *Eur J Radiol* 1997;24:131-6.
14. Radiology ACO. American College of Radiology Breast Imaging Reporting and Data System (BI-RADS). Reston, VA: Radiology ACO; 1993.
15. Boyd NF, Jensen HM, Cooke G, Han HL. Relationship between mammographic and histological risk factors for breast cancer. *J Natl Cancer Inst* 1992;84:1170-9.
16. American College of Radiology, ACR BI-RADS atlas: breast imaging reporting and data system (5th ed.), Reston, Virginia (2013).
17. Mandelson MT, Oestreicher N, Porter PL, White D, Finder CA, Taplin SH, *et al.* Breast density as a predictor of mammographic detection: Comparison of interval- and screen-detected cancers. *J Natl Cancer Inst* 2000;92:1081-7.
18. Carney PA, Miglioretti DL, Yankaskas BC, Kerlikowske K, Rosenberg R, Rutter CM, *et al.* Individual and combined effects of age, breast density, and hormone replacement therapy use on the accuracy of screening mammography. *Ann Intern Med* 2003;138:168-75.
19. Yang WT, Dryden M, Broglio K, Gilcrease M, Dawood S, Dempsey PJ, *et al.* Mammographic features of triple receptor-negative primary breast cancers in young premenopausal women. *Breast Cancer Res Treat* 2008;111:405-10.
20. Eriksson L, Hall P, Czene K, Dos Santos Silva I, McCormack V, Bergh J, *et al.* Mammographic density and molecular subtypes of breast cancer. *Br J Cancer* 2012;107:18-23.
21. Yaghjian L, Colditz GA, Collins LC, Schnitt SJ, Rosner B, Vachon C, *et al.* Mammographic breast density and subsequent risk of breast cancer in postmenopausal women according to tumor characteristics. *J Natl Cancer Inst* 2011;103:1179-89.
22. Shaikh AJ, Mullooly M, Sayed S, Ndumia R, Abayo I, Orwa J, *et al.* Mammographic breast density and breast cancer molecular subtypes: The Kenyan-African aspect. *Biomed Res Int* 2018;2018:6026315.
23. Eriksson L, Czene K, Rosenberg L, Humphreys K, Hall P. The influence of mammographic density on breast tumor characteristics. *Breast Cancer Res Treat* 2012;134:859-66.
24. Chen JH, Hsu FT, Shih HN, Hsu CC, Chang D, Nie K, *et al.* Does breast density show difference in patients with estrogen receptor-positive and estrogen receptor-negative breast cancer measured on MRI? *Ann Oncol* 2009;20:1447-9.
25. Phipps AI, Buist DS, Malone KE, Barlow WE, Porter PL, Kerlikowske K, *et al.* Breast density, body mass index, and risk of tumor marker-defined subtypes of breast cancer. *Ann Epidemiol* 2012;22:340-8.
26. Li E, Guida JL, Tian Y, Sung H, Koka H, Li M, *et al.* Associations between mammographic density and tumor characteristics in Chinese women with breast cancer. *Breast Cancer Res Treat* 2019;177:527-36.
27. Razzaghi H, Troester MA, Gierach GL, Olshan AF, Yankaskas BC, Millikan RC. Association between mammographic density and basal-like and luminal A breast cancer subtypes. *Breast Cancer Res* 2013;15:R76.
28. McCormack VA, Perry NM, Vinnicombe SJ, Dos Santos Silva I. Changes and tracking of mammographic density in relation to Pike's model of breast tissue aging: A UK longitudinal study. *Int J Cancer* 2010;127:452-61.
29. Kelemen LE, Pankratz VS, Sellers TA, Brandt KR, Wang A, Janney C, *et al.* Age-specific trends in mammographic density: The Minnesota Breast Cancer Family study. *Am J Epidemiol* 2008;167:1027-36.
30. Busana MC, De Stavola BL, Sovio U, Li J, Moss S, Humphreys K, *et al.* Assessing within-woman changes in mammographic density: A comparison of fully versus semi-automated area-based approaches. *Cancer Causes Control* 2016;27:481-91.
31. Lokate M, Stellato RK, Veldhuis WB, Peeters PH, van Gils CH. Age-related changes in mammographic density and breast cancer risk. *Am J Epidemiol* 2013;178:101-9.
32. Maskarinec G, Pagano I, Lurie G, Kolonel LN. A longitudinal investigation of mammographic density: The multiethnic cohort. *Cancer Epidemiol Biomarkers Prev* 2006;15:732-9.
33. Riza E, dos Santos Silva I, De Stavola B, Perry N, Karadedou-Zafiriadou E, Linos D, *et al.* Correlates of high-density mammographic parenchymal patterns by menopausal status in a rural population in Northern Greece. *Eur J Cancer* 2005;41:590-600.
34. Dung Yun Trieu P, Mello-Thoms C, Peat JK, Doan Do T, Brennan PC. Associations of breast density with demographic, reproductive, and lifestyle factors in a developing Southeast Asian population. *Asia Pac J Public Health* 2017;29:377-87.
35. Burton A, Maskarinec G, Perez-Gomez B, Vachon C, Miao H, Lajous M, *et al.* Mammographic density and ageing: A collaborative pooled analysis of cross-sectional data from 22 countries worldwide. *PLoS Med* 2017;14:e1002335.
36. Kang YJ, Ahn SK, Kim SJ, Oh H, Han J, Ko E. Relationship between mammographic density and age in the United Arab Emirates population. *J Oncol* 2019;2019:7351350.
37. Checka CM, Chun JE, Schnabel FR, Lee J, Toth H. The relationship of mammographic density and age: Implications for breast cancer screening. *AJR Am J Roentgenol* 2012;198:W292-5.
38. del Carmen MG, Halpern EF, Kopans DB, Moy B, Moore RH, Goss PE, *et al.* Mammographic breast density and race. *AJR Am J*

- Roentgenol 2007;188:1147-50.
39. Ursin G, Ma H, Wu AH, Bernstein L, Salane M, Parisky YR, *et al.* Mammographic density and breast cancer in three ethnic groups. *Cancer Epidemiol Biomarkers Prev* 2003;12:332-8.
 40. Chen Z, Wu AH, Gauderman WJ, Bernstein L, Ma H, Pike MC, *et al.* Does mammographic density reflect ethnic differences in breast cancer incidence rates? *Am J Epidemiol* 2004;159:140-7.
 41. Mariapun S, Li J, Yip CH, Taib NA, Teo SH. Ethnic differences in mammographic densities: An Asian cross-sectional study. *PLoS One* 2015;10:e0117568.
 42. McCarthy AM, Keller BM, Pantalone LM, Hsieh MK, Synnstedt M, Conant EF, *et al.* Racial differences in quantitative measures of area and volumetric breast density. *J Natl Cancer Inst* 2016;108:djw104.
 43. McTiernan A, Martin CF, Peck JD, Aragaki AK, Chlebowski RT, Pisano ED, *et al.* Estrogen-plus-progestin use and mammographic density in postmenopausal women: Women's Health Initiative randomized trial. *J Natl Cancer Inst* 2005;97:1366-76.
 44. Maskarinec G, Pagano I, Lurie G, Wilkens LR, Kolonel LN. Mammographic density and breast cancer risk: The multiethnic cohort study. *Am J Epidemiol* 2005;162:743-52.
 45. Ishihara S, Taira N, Kawasaki K, Ishibe Y, Mizoo T, Nishiyama K, *et al.* Association between mammographic breast density and lifestyle in Japanese women. *Acta Med Okayama* 2013;67:145-51.
 46. Akinyemiju TF, Tehranifar P, Flom JD, Liao Y, Wei Y, Terry MB. Early life growth, socioeconomic status, and mammographic breast density in an urban US birth cohort. *Ann Epidemiol* 2016;26:540-5.e2.
 47. Yang Y, Liu J, Gu R, Hu Y, Liu F, Yun M, *et al.* Influence of factors on mammographic density in premenopausal Chinese women. *Eur J Cancer Prev* 2016;25:306-11.
 48. Perry N, Moss S, Dixon S, Milner S, Mokbel K, Lemech C, *et al.* Mammographic breast density and urbanization: Interactions with BMI, environmental, lifestyle, and other patient factors. *Diagnostics (Basel)* 2020;10:418.
 49. Reeves KW, Stone RA, Modugno F, Ness RB, Vogel VG, Weissfeld JL, *et al.* Longitudinal association of anthropometry with mammographic breast density in the Study of Women's Health Across the Nation. *Int J Cancer* 2009;124:1169-77.
 50. Gram IT, Funkhouser E, Tabar L. Anthropometric indices in relation to mammographic patterns among peri-menopausal women. *Int J Cancer* 1997;73:323-6.
 51. Haars G, van Noord PA, van Gils CH, Grobbee DE, Peeters PH. Measurements of breast density: No ratio for a ratio. *Cancer Epidemiol Biomarkers Prev* 2005;14:2634-40.
 52. Maskarinec G, Meng L, Ursin G. Ethnic differences in mammographic densities. *Int J Epidemiol* 2001;30:959-65.
 53. Jeon JH, Kang JH, Kim Y, Lee HY, Choi KS, Jun JK, *et al.* Reproductive and hormonal factors associated with fatty or dense breast patterns among Korean women. *Cancer Res Treat* 2011;43:42-8.
 54. Pinsky RW, Helvie MA. Mammographic breast density: Effect on imaging and breast cancer risk. *J Natl Compr Canc Netw* 2010;8:1157-64.
 55. Alimujiang A, Imm KR, Appleton CM, Colditz GA, Berkey CS, Toriola AT. Adiposity at age 10 and mammographic density among premenopausal women. *Cancer Prev Res (Phila)* 2018;11:287-94.
 56. Alexeeff SE, Odo NU, Lipson JA, Achacoso N, Rothstein JH, Yaffe MJ, *et al.* Age at menarche and late adolescent adiposity associated with mammographic density on processed digital mammograms in 24,840 women. *Cancer Epidemiol Biomarkers Prev* 2017;26:1450-8.
 57. Dorgan JF, Klifa C, Shepherd JA, Egleston BL, Kwiterovich PO, Himes JH, *et al.* Height, adiposity and body fat distribution and breast density in young women. *Breast Cancer Res* 2012;14:R107.
 58. Andersen ZJ, Baker JL, Bihmann K, Vejborg I, Sørensen TI, Lyng E. Birth weight, childhood body mass index, and height in relation to mammographic density and breast cancer: A register-based cohort study. *Breast Cancer Res* 2014;16:R4.
 59. Hopper JL, Nguyen TL, Stone J, Aujard K, Matheson MC, Abramson MJ, *et al.* Childhood body mass index and adult mammographic density measures that predict breast cancer risk. *Breast Cancer Res Treat* 2016;156:163-70.
 60. Alimujiang A, Appleton C, Colditz GA, Toriola AT. Adiposity during early adulthood, changes in adiposity during adulthood, attained adiposity, and mammographic density among premenopausal women. *Breast Cancer Res Treat* 2017;166:197-206.
 61. Rice MS, Bertrand KA, Lajous M, Tamimi RM, Torres-Mejia G, Biessy C, *et al.* Body size throughout the life course and mammographic density in Mexican women. *Breast Cancer Res Treat* 2013;138:601-10.
 62. Guo YP, Martin LJ, Hanna W, Banerjee D, Miller N, Fishell E, *et al.* Growth factors and stromal matrix proteins associated with mammographic densities. *Cancer Epidemiol Biomarkers Prev* 2001;10:243-8.
 63. Martin LJ, Boyd NF. Mammographic density. Potential mechanisms of breast cancer risk associated with mammographic density: Hypotheses based on epidemiological evidence. *Breast Cancer Res* 2008;10:201.
 64. Houghton LC, Sisti JS, Hankinson SE, Xie J, Xu X, Hoover RN, *et al.* Estrogen metabolism in premenopausal women is related to early life body fatness. *Cancer Epidemiol Biomarkers Prev* 2018;27:585-93.
 65. Yochum L, Tamimi RM, Hankinson SE. Birthweight, early life body size and adult mammographic density: A review of epidemiologic studies. *Cancer Causes Control* 2014;25:1247-59.
 66. Tehranifar P, Reynolds D, Flom J, Fulton L, Liao Y, Kudadjie-Gyamfi E, *et al.* Reproductive and menstrual factors and mammographic density in African American, Caribbean, and white women. *Cancer Causes Control* 2011;22:599-610.
 67. Rice MS, Bertrand KA, Lajous M, Tamimi RM, Torres G, López-Ridaura R, *et al.* Reproductive and lifestyle risk factors and mammographic density in Mexican women. *Ann Epidemiol* 2015;25:868-73.
 68. Lee E, Doanvo N, Lee M, Soe Z, Lee AW, Van Doan C, *et al.* Immigration history, lifestyle characteristics, and breast density in the Vietnamese American Women's Health Study: A cross-sectional analysis. *Cancer Causes Control* 2020;31:127-38.
 69. Dai H, Yan Y, Wang P, Liu P, Cao Y, Xiong L, *et al.* Distribution of mammographic density and its influential factors among Chinese women. *Int J Epidemiol* 2014;43:1240-51.
 70. Engmann NJ, Scott C, Jensen MR, Winham SJ, Ma L, Brandt KR, *et al.* Longitudinal changes in volumetric breast density in healthy women across the menopausal transition. *Cancer Epidemiol Biomarkers Prev* 2019;28:1324-30.
 71. Boyd N, Martin L, Stone J, Little L, Minkin S, Yaffe M. A longitudinal study of the effects of menopause on mammographic features. *Cancer Epidemiol Biomarkers Prev* 2002;11:1048-53.
 72. Yaghjian L, Colditz GA, Rosner B, Bertrand KA, Tamimi RM. Reproductive factors related to childbearing and mammographic breast density. *Breast Cancer Res Treat* 2016;158:351-9.
 73. Pike MC, Krailo MD, Henderson BE, Casagrande JT, Hoel DG. 'Hormonal' risk factors, 'breast tissue age' and the age-incidence of breast cancer. *Nature* 1983;303:767-70.
 74. Heng D, Gao F, Jong R, Fishell E, Yaffe M, Martin L, *et al.* Risk factors for breast cancer associated with mammographic features in Singaporean Chinese women. *Cancer Epidemiol Biomarkers Prev* 2004;13:1751-8.
 75. Wong CS, Lim GH, Gao F, Jakes RW, Offman J, Chia KS, *et al.*

- Mammographic density and its interaction with other breast cancer risk factors in an Asian population. *Br J Cancer* 2011;104:871-4.
76. Nguyen TL, Schmidt DF, Makalic E, Dite GS, Stone J, Apicella C, *et al.* Explaining variance in the cumulus mammographic measures that predict breast cancer risk: A twins and sisters study. *Cancer Epidemiol Biomarkers Prev* 2013;22:2395-403.
 77. Gapstur SM, López P, Colangelo LA, Wolfman J, Van Horn L, Hendrick RE, *et al.* Associations of breast cancer risk factors with breast density in Hispanic women. *Cancer Epidemiol Biomarkers Prev* 2003;12:1074-80.
 78. McCormack VA, Perry N, Vinnicombe SJ, Silva Idos S. Ethnic variations in mammographic density: A British multiethnic longitudinal study. *Am J Epidemiol* 2008;168:412-21.
 79. Vachon CM, Kuni CC, Anderson K, Anderson VE, Sellers TA. Association of mammographically defined percent breast density with epidemiologic risk factors for breast cancer (United States). *Cancer Causes Control* 2000;11:653-62.
 80. Butler LM, Gold EB, Greendale GA, Crandall CJ, Modugno F, Oestreicher N, *et al.* Menstrual and reproductive factors in relation to mammographic density: The Study of Women's Health Across the Nation (SWAN). *Breast Cancer Res Treat* 2008;112:165-74.
 81. Dite GS, Gurrin LC, Byrnes GB, Stone J, Gunasekara A, McCredie MR, *et al.* Predictors of mammographic density: Insights gained from a novel regression analysis of a twin study. *Cancer Epidemiol Biomarkers Prev* 2008;17:3474-81.
 82. Warren R, Skinner J, Sala E, Denton E, Folklerd E, Healey CS, *et al.* Associations among mammographic density, circulating sex hormones, and polymorphisms in sex hormone metabolism genes in postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2006;15:1502-8.
 83. Gram IT, Funkhouser E, Tabar L. Reproductive and menstrual factors in relation to mammographic parenchymal patterns among perimenopausal women. *Br J Cancer* 1995;71:647-50.
 84. Carlile T, Kopecky KJ, Thompson DJ, Whitehead JR, Gilbert FI Jr., Present AJ, *et al.* Breast cancer prediction and the Wolfe classification of mammograms. *JAMA* 1985;254:1050-3.
 85. Lim SE, Ahn H, Lee ES, Kong SY, Jung SY, Lee S, *et al.* Interaction effect between breast density and reproductive factors on breast cancer risk in Korean population. *J Cancer Prev* 2019;24:26-32.
 86. Modugno F, Ngo DL, Allen GO, Kuller LH, Ness RB, Vogel VG, *et al.* Breast cancer risk factors and mammographic breast density in women over age 70. *Breast Cancer Res Treat* 2006;97:157-66.
 87. Dorgan JF, Klifa C, Deshmukh S, Egleston BL, Shepherd JA, Kwiterovich PO Jr., *et al.* Menstrual and reproductive characteristics and breast density in young women. *Cancer Causes Control* 2013;24:1973-83.
 88. Ursin G, Sun CL, Koh WP, Khoo KS, Gao F, Wu AH, *et al.* Associations between soy, diet, reproductive factors, and mammographic density in Singapore Chinese women. *Nutr Cancer* 2006;56:128-35.
 89. Sung J, Song YM, Stone J, Lee K, Lee D. Reproductive factors associated with mammographic density: A Korean co-twin control study. *Breast Cancer Res Treat* 2011;128:567-72.
 90. Das R, Vonderhaar BK. Prolactin as a mitogen in mammary cells. *J Mammary Gland Biol Neoplasia* 1997;2:29-39.
 91. Persson I, Thurfjell E, Holmberg L. Effect of estrogen and estrogen-progestin replacement regimens on mammographic breast parenchymal density. *J Clin Oncol* 1997;15:3201-7.
 92. Lundström E, Christow A, Kersemaekers W, Svane G, Azavedo E, Söderqvist G, *et al.* Effects of tibolone and continuous combined hormone replacement therapy on mammographic breast density. *Am J Obstet Gynecol* 2002;186:717-22.
 93. Stomper PC, Van Voorhis BJ, Ravnikaar VA, Meyer JE. Mammographic changes associated with postmenopausal hormone replacement therapy: A longitudinal study. *Radiology* 1990;174:487-90.
 94. Berkowitz JE, Gatewood OM, Goldblum LE, Gayler BW. Hormonal replacement therapy: Mammographic manifestations. *Radiology* 1990;174:199-201.
 95. Rymer JM. The effects of tibolone. *Gynecol Endocrinol* 1998;12:213-20.
 96. Palacios S. Tibolone: What does tissue specific activity mean? *Maturitas* 2001;37:159-65.
 97. Marchesoni D, Driul L, Ianni A, Fabiani G, Della Martina M, Zuiani C, *et al.* Postmenopausal hormone therapy and mammographic breast density. *Maturitas* 2006;53:59-64.
 98. Junkermann H, von Holst T, Lang E, Rakov V. Influence of different HRT regimens on mammographic density. *Maturitas* 2005;50:105-10.
 99. Erel CT, Esen G, Seyisoglu H, Elter K, Uras C, Ertungealp E, *et al.* Mammographic density increase in women receiving different hormone replacement regimens. *Maturitas* 2001;40:151-7.
 100. Harvey J, Scheurer C, Kawakami FT, Quebe-Fehling E, de Palacios PI, Ragavan VV. Hormone replacement therapy and breast density changes. *Climacteric* 2005;8:185-92.
 101. Colacurci N, Fornaro F, De Francis P, Mele D, Palermo M, del Vecchio W. Effects of a short-term suspension of hormone replacement therapy on mammographic density. *Fertil Steril* 2001;76:451-5.
 102. Weaver K, Kataoka M, Murray J, Muir B, Anderson E, Warren R, *et al.* Does a short cessation of HRT decrease mammographic density? *Maturitas* 2008;59:315-22.
 103. Rutter CM, Mandelson MT, Laya MB, Seger DJ, Taplin S. Changes in breast density associated with initiation, discontinuation, and continuing use of hormone replacement therapy. *JAMA* 2001;285:171-6.
 104. Gabrielson M, Azam S, Hardell E, Holm M, Ubhayasekera KA, Eriksson M, *et al.* Hormonal determinants of mammographic density and density change. *Breast Cancer Res* 2020;22:95.
 105. Noh JJ, Maskarinec G, Pagano I, Cheung LW, Stanczyk FZ. Mammographic densities and circulating hormones: A cross-sectional study in premenopausal women. *Breast* 2006;15:20-8.
 106. White E, Velentgas P, Mandelson MT, Lehman CD, Elmore JG, Porter P, *et al.* Variation in mammographic breast density by time in menstrual cycle among women aged 40-49 years. *J Natl Cancer Inst* 1998;90:906-10.
 107. Vogel PM, Georgiade NG, Fetter BF, Vogel FS, McCarty KS Jr. The correlation of histologic changes in the human breast with the menstrual cycle. *Am J Pathol* 1981;104:23-34.
 108. Longacre TA, Bartow SA. A correlative morphologic study of human breast and endometrium in the menstrual cycle. *Am J Surg Pathol* 1986;10:382-93.
 109. Pike MC, Spicer DV, Dahmouch L, Press MF. Estrogens, progestogens, normal breast cell proliferation, and breast cancer risk. *Epidemiol Rev* 1993;15:17-35.
 110. Takata Y, Maskarinec G, Park SY, Murphy SP, Wilkens LR, Kolonel LN. Mammographic density and dietary patterns: The multiethnic cohort. *Eur J Cancer Prev* 2007;16:409-14.
 111. Knight JA, Martin LJ, Greenberg CV, Lockwood GA, Byng JW, Yaffe MJ, *et al.* Macronutrient intake and change in mammographic density at menopause: Results from a randomized trial. *Cancer Epidemiol Biomarkers Prev* 1999;8:123-8.
 112. Tseng M, Vierkant RA, Kushi LH, Sellers TA, Vachon CM. Dietary patterns and breast density in the Minnesota Breast Cancer Family study. *Cancer Causes Control* 2008;19:481-9.
 113. Masala G, Ambrogetti D, Assedi M, Giorgi D, Del Turco MR, Palli D. Dietary and lifestyle determinants of mammographic

- breast density. A longitudinal study in a Mediterranean population. *Int J Cancer* 2006;118:1782-9.
114. Voevodina O, Billich C, Arand B, Nagel G. Association of Mediterranean diet, dietary supplements and alcohol consumption with breast density among women in south Germany: A cross-sectional study. *BMC Public Health* 2013;13:203.
 115. Castelló A, Ascunce N, Salas-Trejo D, Vidal C, Sanchez-Contador C, Santamariña C, *et al.* Association between Western and Mediterranean dietary patterns and mammographic density. *Obstet Gynecol* 2016;128:574-81.
 116. Tseng M, Sellers TA, Vierkant RA, Kushi LH, Vachon CM. Mediterranean diet and breast density in the Minnesota Breast Cancer Family Study. *Nutr Cancer* 2008;60:703-9.
 117. Mishra GD, dos Santos Silva I, McNaughton SA, Stephen A, Kuh D. Energy intake and dietary patterns in childhood and throughout adulthood and mammographic density: Results from a British prospective cohort. *Cancer Causes Control* 2011;22:227-35.
 118. Jones JA, Hartman TJ, Klifa CS, Coffman DL, Mitchell DC, Vernarelli JA, *et al.* Dietary energy density is positively associated with breast density among young women. *J Acad Nutr Diet* 2015;115:353-9.
 119. Lin R, White JH. The pleiotropic actions of vitamin D. *Bioessays* 2004;26:21-8.
 120. Lipkin M, Newmark HL. Vitamin D, calcium and prevention of breast cancer: A review. *J Am Coll Nutr* 1999;18:392S-7S.
 121. Bérubé S, Diorio C, Mâsse B, Hébert-Croteau N, Byrne C, Côté G, *et al.* Vitamin D and calcium intakes from food or supplements and mammographic breast density. *Cancer Epidemiol Biomarkers Prev* 2005;14:1653-9.
 122. Fair AM, Lewis TJ, Sanderson M, Dupont WD, Fletcher S, Egan KM, *et al.* Increased vitamin D and calcium intake associated with reduced mammographic breast density among premenopausal women. *Nutr Res* 2015;35:851-7.
 123. Tseng M, Byrne C, Evers KA, Daly MB. Dietary intake and breast density in high-risk women: A cross-sectional study. *Breast Cancer Res* 2007;9:R72.
 124. Welsh J. Vitamin D and breast cancer: Insights from animal models. *Am J Clin Nutr* 2004;80:1721S-4S.
 125. Welsh J, Wietzke JA, Zinser GM, Byrne B, Smith K, Narvaez CJ. Vitamin D-3 receptor as a target for breast cancer prevention. *J Nutr* 2003;133:2425S-33S.
 126. Green AK, Hankinson SE, Bertone-Johnson ER, Tamimi RM. Mammographic density, plasma vitamin D levels and risk of breast cancer in postmenopausal women. *Int J Cancer* 2010;127:667-74.
 127. Straub L, Riedel J, Luppá PB, Wissing J, Artmann A, Kiechle M, *et al.* Mammographic density and vitamin D levels – A cross-sectional study. *Geburtshilfe Frauenheilkd* 2017;77:257-67.
 128. Hooper L, Madhavan G, Tice JA, Leinster SJ, Cassidy A. Effects of isoflavones on breast density in pre- and post-menopausal women: A systematic review and meta-analysis of randomized controlled trials. *Hum Reprod Update* 2010;16:745-60.
 129. Han Y, Zong X, Li Y, Colditz GA, Toriola AT. Milk intake and mammographic density in premenopausal women. *Breast Cancer Res Treat* 2019;174:249-55.
 130. Jayasekara H, MacInnis RJ, Room R, English DR. Long-term alcohol consumption and breast, upper aero-digestive tract and colorectal cancer risk: A systematic review and meta-analysis. *Alcohol Alcohol* 2016;51:315-30.
 131. Singletary KW, Gapstur SM. Alcohol and breast cancer: Review of epidemiologic and experimental evidence and potential mechanisms. *JAMA* 2001;286:2143-51.
 132. Romieu I, Scoccianti C, Chajès V, de Batlle J, Biessy C, Dossus L, *et al.* Alcohol intake and breast cancer in the European prospective investigation into cancer and nutrition. *Int J Cancer* 2015;137:1921-30.
 133. Okamoto T, Ito A. Association between alcohol consumption and mammographic density: A hospital-based cross-sectional study. *Breast Cancer* 2019;26:478-84.
 134. Maskarinec G, Takata Y, Pagano I, Lurie G, Wilkens LR, Kolonel LN. Alcohol consumption and mammographic density in a multiethnic population. *Int J Cancer* 2006;118:2579-83.
 135. McBride RB, Fei K, Rothstein JH, Alexeeff SE, Song X, Sakoda LC, *et al.* Alcohol and tobacco use in relation to mammographic density in 23,456 women. *Cancer Epidemiol Biomarkers Prev* 2020;29:1039-48.
 136. Vachon CM, Kushi LH, Cerhan JR, Kuni CC, Sellers TA. Association of diet and mammographic breast density in the Minnesota breast cancer family cohort. *Cancer Epidemiol Biomarkers Prev* 2000;9:151-60.
 137. Jacobsen KK, Lynge E, Tjønneland A, Vejborg I, von Euler-Chelpin M, Andersen ZJ. Alcohol consumption and mammographic density in the Danish Diet, Cancer and Health cohort. *Cancer Causes Control* 2017;28:1429-39.
 138. Seitz HK, Pelucchi C, Bagnardi V, La Vecchia C. Epidemiology and pathophysiology of alcohol and breast cancer: Update 2012. *Alcohol Alcohol* 2012;47:204-12.
 139. Dumitrescu RG, Shields PG. The etiology of alcohol-induced breast cancer. *Alcohol* 2005;35:213-25.
 140. Liu Y, Tamimi RM, Colditz GA, Bertrand KA. Alcohol consumption across the life course and mammographic density in premenopausal women. *Breast Cancer Res Treat* 2018;167:529-35.
 141. McDonald JA, Michels KB, Cohn BA, Flom JD, Tehranifar P, Terry MB. Alcohol intake from early adulthood to midlife and mammographic density. *Cancer Causes Control* 2016;27:493-502.
 142. Jacobsen KK, Lynge E, Vejborg I, Tjønneland A, von Euler-Chelpin M, Andersen ZJ. Cigarette smoking and mammographic density in the Danish Diet, Cancer and Health cohort. *Cancer Causes Control* 2016;27:271-80.
 143. Butler LM, Gold EB, Conroy SM, Crandall CJ, Greendale GA, Oestreicher N, *et al.* Active, but not passive cigarette smoking was inversely associated with mammographic density. *Cancer Causes Control* 2010;21:301-11.
 144. Cabanes A, Pastor-Barriuso R, García-López M, Pedraz-Pingarrón C, Sánchez-Contador C, Vázquez Carrete JA, *et al.* Alcohol, tobacco, and mammographic density: A population-based study. *Breast Cancer Res Treat* 2011;129:135-47.
 145. Jeffreys M, Warren R, Gunnell D, McCarron P, Smith GD. Life course breast cancer risk factors and adult breast density (United Kingdom). *Cancer Causes Control* 2004;15:947-55.
 146. Brand JS, Czene K, Eriksson L, Trinh T, Bhoo-Pathy N, Hall P, *et al.* Influence of lifestyle factors on mammographic density in postmenopausal women. *PLoS One* 2013;8:e81876.
 147. Roubidoux MA, Kaur JS, Griffith KA, Stillwater B, Novotny P, Sloan J. Relationship of mammographic parenchymal patterns to breast cancer risk factors and smoking in Alaska Native women. *Cancer Epidemiol Biomarkers Prev* 2003;12:1081-6.
 148. Carmichael AR, Daley AJ, Rea DW, Bowden SJ. Physical activity and breast cancer outcome: A brief review of evidence, current practice and future direction. *Eur J Surg Oncol* 2010;36:1139-48.
 149. Lahmann PH, Friedenreich C, Schuit AJ, Salvini S, Allen NE, Key TJ, *et al.* Physical activity and breast cancer risk: The European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev* 2007;16:36-42.
 150. Irwin ML. Randomized controlled trials of physical activity and breast cancer prevention. *Exerc Sport Sci Rev* 2006;34:182-93.
 151. Azam S, Kemp Jacobsen K, Aro AR, von Euler-Chelpin M, Tjønneland A, Vejborg I, *et al.* Regular physical activity and

- mammographic density: A cohort study. *Cancer Causes Control* 2018;29:1015-25.
152. López P, Van Horn L, Colangelo LA, Wolfman JA, Hendrick RE, Gapstur SM. Physical inactivity and percent breast density among Hispanic women. *Int J Cancer* 2003;107:1012-6.
 153. Yaghjian L, Colditz GA, Wolin K. Physical activity and mammographic breast density: A systematic review. *Breast Cancer Res Treat* 2012;135:367-80.
 154. Masala G, Assedi M, Ambrogetti D, Sera F, Salvini S, Bendinelli B, *et al.* Physical activity and mammographic breast density in a Mediterranean population: The EPIC Florence longitudinal study. *Int J Cancer* 2009;124:1654-61.
 155. Masala G, Assedi M, Sera F, Ermini I, Occhini D, Castaldo M, *et al.* Can dietary and physical activity modifications reduce breast density in postmenopausal women? The DAMA study, a randomized intervention trial in Italy. *Cancer Epidemiol Biomarkers Prev* 2019;28:41-50.
 156. Pettee Gabriel K, Klifa C, Pérez A, Kriska AM, Snetselaar L, Dorgan JF, *et al.* Adolescent and young adult exposure to physical activity and breast density. *Med Sci Sports Exerc* 2013;45:1515-23.
 157. Bernstein L, Ross RK. Endogenous hormones and breast cancer risk. *Epidemiol Rev* 1993;15:48-65.
 158. Sachdev D, Yee D. The IGF system and breast cancer. *Endocr Relat Cancer* 2001;8:197-209.
 159. Roswall N, Andersen ZJ, von Euler-Chelpin M, Vejborg I, Lynge E, Jensen SS, *et al.* Residential traffic noise and mammographic breast density in the Diet, Cancer, and Health cohort. *Cancer Causes Control* 2018;29:399-404.
 160. Sung J, Song YM. The relationship between education level and mammographic density. *Eur J Cancer Prev* 2015;24:491-6.
 161. Lope V, García-Pérez J, Pérez-Gómez B, Pedraza-Flechas AM, Alguacil J, González-Galarzo MC, *et al.* Occupational exposures and mammographic density in Spanish women. *Occup Environ Med* 2018;75:124-31.
 162. Jiménez T, García-Pérez J, van der Haar R, Alba MÁ, Lucas P, Sierra MÁ, *et al.* Occupation, occupational exposures and mammographic density in Spanish women. *Environ Res* 2021;195:110816.
 163. Tehranifar P, Protacio A, Schmitt KM, Desperito E, Oskar S, Potter AJ, *et al.* The metabolic syndrome and mammographic breast density in a racially diverse and predominantly immigrant sample of women. *Cancer Causes Control* 2015;26:1393-403.
 164. Aitken Z, Walker K, Stegeman BH, Wark PA, Moss SM, McCormack VA, *et al.* Mammographic density and markers of socioeconomic status: A cross-sectional study. *BMC Cancer* 2010;10:35.
 165. Marini M, Bendinelli B, Assedi M, Occhini D, Castaldo M, Fabiano J, *et al.* Low back pain in healthy postmenopausal women and the effect of physical activity: A secondary analysis in a randomized trial. *PLoS One* 2017;12:e0177370.
 166. Hassinger TE, Mehaffey JH, Knisely AT, Contrella BN, Brenin DR, Schroen AT, *et al.* The impact of bariatric surgery on qualitative and quantitative breast density. *Breast J* 2019;25:1198-205.
 167. Tehranifar P, Cohn BA, Flom JD, Protacio A, Cirillo P, Lumey LH, *et al.* Early life socioeconomic environment and mammographic breast density. *BMC Cancer* 2017;17:41.
 168. Buschard K, Thomassen K, Lynge E, Vejborg I, Tjønneland A, von Euler-Chelpin M, *et al.* Diabetes, diabetes treatment, and mammographic density in Danish Diet, Cancer, and Health cohort. *Cancer Causes Control* 2017;28:13-21.
 169. Vachon CM, Scott CG, Fasching PA, Hall P, Tamimi RM, Li J, *et al.* Common breast cancer susceptibility variants in LSP1 and RAD51L1 are associated with mammographic density measures that predict breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 2012;21:1156-66.
 170. Burton H, Chowdhury S, Dent T, Hall A, Pashayan N, Pharoah P. Public health implications from COGS and potential for risk stratification and screening. *Nat Genet* 2013;45:349-51.
 171. Sartor H, Brandt J, Grassmann F, Eriksson M, Czene K, Melander O, *et al.* The association of Single Nucleotide Polymorphisms (SNPs) with breast density and breast cancer survival: The Malmö diet and cancer study. *Acta Radiol* 2020;61:1326-34.
 172. Vachon CM, Scott CG, Tamimi RM, Thompson DJ, Fasching PA, Stone J, *et al.* Joint association of mammographic density adjusted for age and body mass index and polygenic risk score with breast cancer risk. *Breast Cancer Res* 2019;21:68.
 173. Keller BM, McCarthy AM, Chen J, Armstrong K, Conant EF, Domchek SM, *et al.* Associations between breast density and a panel of single nucleotide polymorphisms linked to breast cancer risk: A cohort study with digital mammography. *BMC Cancer* 2015;15:143.
 174. Stone J, Thompson DJ, Dos Santos Silva I, Scott C, Tamimi RM, Lindstrom S, *et al.* Novel associations between common breast cancer susceptibility variants and risk-predicting mammographic density measures. *Cancer Res* 2015;75:2457-67.
 175. Lindström S, Thompson DJ, Paterson AD, Li J, Gierach GL, Scott C, *et al.* Genome-wide association study identifies multiple loci associated with both mammographic density and breast cancer risk. *Nat Commun* 2014;5:5303.
 176. Li H, Giger ML, Sun C, Ponsukcharoen U, Huo D, Lan L, *et al.* Pilot study demonstrating potential association between breast cancer image-based risk phenotypes and genomic biomarkers. *Med Phys* 2014;41:031917.
 177. Varghese JS, Thompson DJ, Michailidou K, Lindström S, Turnbull C, Brown J, *et al.* Mammographic breast density and breast cancer: Evidence of a shared genetic basis. *Cancer Res* 2012;72:1478-84.