



Editorial

Air Pollution and Health: The Need for a Medical Reading of Environmental Monitoring Data

Marcello Iriti ^{1,2} , Prisco Piscitelli ^{1,3,*} , Eduardo Missoni ^{4,5} and Alessandro Miani ^{1,6}

¹ Italian Society of Environmental Medicine (SIMA), 20133 Milan, Italy; marcello.iriti@unimi.it (M.I.); alessandro.miani@gmail.com (A.M.)

² Department of Agricultural and Environmental Sciences, University of Milan, 20133 Milan, Italy

³ Euro Mediterranean Scientific Biomedical Institute (ISBEM), Rue de Bellard 20, 1040 Bruxelles, Belgium

⁴ Department of Sociology and Social Research, Bicocca University, 20133 Milan, Italy; eduardomissoni@gmail.com

⁵ Centre for Research on Health and Social Care Management (CERGAS), SDA Bocconi School of Management, 20133 Milan, Italy

⁶ Department of Environmental Science and Policy, University of Milan, 20133 Milan, Italy

* Correspondence: priscofreedom@hotmail.com; Tel./Fax: +32-(02)-4302408

Received: 12 March 2020; Accepted: 16 March 2020; Published: 25 March 2020



Abstract: Air pollution is a recent public health issue. In 2006, the World Health Organization (WHO) published updated air quality guidelines for a number of air pollutants (including PM₁₀ and PM_{2.5}), which recommended for particulate matter annual average concentration levels at half or less the limit values set by European legislation. In the European Union, around 80% of the European urban population is exposed to air pollution above the levels recommended by the WHO guidelines. Only in 2015 the WHO addressed for the first time the topic of the health impacts of air pollution in its general assembly, which adopted a resolution clearly defining air pollution as the world's largest single environmental health risk factor. Nowadays, the WHO considers air pollution as a major public health threat, causing a 7% increase in overall mortality for each increase of 10 µg/m³ in annual average of PM_{2.5}. This result has been achieved thanks to the outstanding efforts of the director of the WHO's Environment and Public Health Department, Dr. Maria Neira, who has devoted her full commitment to highlighting the consequences that air pollution has on people's health. More recently, at European level, the Air Quality Directive has been subject to a fitness check, published in 2019; the European Green Deal has since announced its aim to align EU air quality standards more closely with the WHO recommendations. Every year, the European Environment Agency (EEA) publishes its "Air Quality in Europe" Report to assess the figures on air pollution across Europe and related health impacts. However, environmental data provided by official regional or national agencies—used by decision makers to adopt preventive measures such as limitations on urban traffic or domestic heating—refer to legal thresholds established by the law (usually on the basis of values set at European level, at least for the EU). These legal thresholds, however, are not adequate to fully protect population against all impacts from air pollution as recommended by WHO and scientific evidence. Therefore, we point out the need for a medical reading of environmental monitoring data that should be performed both at national and regional or local level by health authorities, to foster population health protection against air pollution and guarantee the application of the precautionary principle. A stronger cooperation between environmental agencies and health authorities is needed to address the new challenges to human and planetary health arising from air pollution and climate change. Health authorities should integrate their medical staff with new professionals and researchers with adequate training in environmental sciences to foster population health protection against air pollution. For this purposes, multi-disciplinary research units or teams should be established by local health authorities on environmental health topics, working together with medical staff and environmental agencies for a mutual integration of competencies.

Keywords: air pollution; health protection; particulate matter; air quality standards

1. Air Pollution and Health: A Recent Issue

In 2006, the World Health Organization (WHO) published updated Air Quality Guidelines for a number of air pollutants, which recommend for particulate matter annual average concentration levels at half or less the limit values set by European legislation [1]. According to these air quality guidelines, annual average concentrations of PM₁₀ should not exceed 20 µg/m³ (compared to the current limit value set by the EU of 40 µg/m³) and PM_{2.5} should not exceed 10 µg/m³ (compared to the current EU limit value, set by EU legislation at 25 µg/m³) [1]. A number of countries have already adopted air quality standards for annual averages of PM₁₀ and PM_{2.5} that are closer to the 2005 WHO guidelines (including Switzerland, Canada, Norway, Japan, and the United States of America). In the European Union, around 80% of the European urban population is exposed to air pollution above the levels recommended by the WHO guidelines [2]. The World Health Organization has furthermore noted that the adverse effects on health of particulate matter are well documented, and that there is no evidence of a safe level of exposure or a threshold below which no adverse health effects occur, including exacerbations of respiratory diseases (e.g., asthma, especially in children).

In 2013, the International Agency for Research on Cancer (IARC) had already classified outdoor air pollution in general, and particulate matter in particular, as *carcinogenic to humans* (IARC Class 1), pointing out that fine dusts are known to produce severe health impacts even at very low concentrations [3]. Surprisingly, only on 26 May 2015 the World Health Organization addressed for the first time the topic of the health impacts of air pollution, and its general assembly adopted a resolution clearly defining air pollution as the world's largest single environmental health risk factor. Nowadays, WHO considers air pollution a major public health threat, causing a 7% increase in overall mortality for each increase of 10 µg/m³ in annual average of PM_{2.5} [4]. This result has been achieved thanks to the outstanding efforts of the director of the WHO's Environment & Public Health Department, Dr. Maria Neira, who has devoted her full commitment to highlighting the consequences that air pollution has on people's health. More recently, at European level, the Air Quality Directive has been subject to a fitness check, published in 2019; the European Green Deal has since announced its aim to align EU air quality standards more closely with WHO recommendations, but the cut to the limits on PM₁₀ and PM_{2.5} has not yet been accomplished.

2. Air Pollution: An under-Perceived Threat For public Health and Society

The European Environment Agency (EEA) publishes, every year, a specific Air Quality Report concerning 41 European countries (including the EU Member States). The most recent report was issued in autumn 2019, and again recognizes air pollution as a significant threat to human health in European cities, resulting in considerable impacts in terms of premature mortality, medical costs, and loss of productivity [2]. According to the EEA's "Air Quality in Europe—2019 Report", the most harmful pollutants for human health are particulate matter (PM), nitrogen dioxide (NO₂) and ground-level ozone (O₃), resulting in about 538,000 estimated premature deaths across Europe and almost 506,000 in the 28 countries belonging to the European Union, including the UK (Table 1). As a proxy for the economic impact of air pollution, we can refer to the indicator "years of life lost" (YLL) attributable to air pollution via PM_{2.5} and NO₂ and O₃ exposure, which has been estimated by the EEA at about 4,150,000 for Europe as whole and 4,466,000 for the 28 EU member States (Table 1). If we consider that, as Chiabai, Spadaro and Neuman have recently assessed [5], each year of life lost for European people corresponds to an annual value of 100,000 euros, it comes up more clearly how huge is the impact of air pollution both in terms of population health and economic consequences: more than four billion euros per year only for this indicator. A recent fitness check of the EU's air quality legislation also confirmed the substantial economic impact of air pollution.

Table 1. Premature deaths and years of life lost (YYL) attributable to PM_{2.5}, NO₂ and O₃ exposure in 41 European countries and the EU-28. Data concerning year 2018 (modified from EEA 2019 Air Quality Report)².

Country	Population (1000)	Premature Deaths Due to PM _{2.5}	YYL Due to PM _{2.5}	Premature Deaths Due to NO ₂	YYL Due to NO ₂	Premature Deaths Due to O ₃	YYL Due to O ₃
Austria	8700	5300	60,200	1000	12,200	270	4000
Belgium	11,311	7600	77,600	1600	16,200	180	2400
Bulgaria	7154	13,100	142,000	1100	6400	280	3700
Croatia	4191	5300	46,900	260	4500	190	2500
Cyprus	1184	580	7400	240	300	30	410
Czechia	10,554	9600	105,500	240	5100	350	5000
Denmark	5707	2700	30,100	80	860	90	980
Estonia	1316	500	6300	<1	40	20	230
Finland	5487	1500	16,000	<1	470	60	570
France	64,977	33,200	414,700	7500	112,400	1400	21,600
Germany	82,176	59,600	638,500	11,900	134,200	2400	31,800
Greece	10,784	12,900	120,700	2900	23,100	640	6400
Hungary	9830	12,100	139,300	770	14,300	380	6000
Ireland	4726	1100	12,000	50	310	30	230
Italy	60,666	58,600	593,700	14,600	200,700	3000	32,100
Latvia	1969	1700	17,600	60	1400	60	600
Lithuania	2889	2600	27,400	20	760	70	940
Luxembourg	576	230	2700	50	510	10	110
Malta	450	210	2700	<1	180	20	180
Netherlands	16,979	9200	103,800	1500	19,900	270	3300
Poland	37,967	43,100	533,800	1500	20,400	1100	3300
Portugal	9809	4900	56,300	610	9100	320	16,600
Romania	19,761	23,400	271,600	2600	14,100	490	3200
Slovakia	5426	4800	59,900	20	2700	160	2600
Slovenia	2064	1700	20,000	70	1800	70	1100
Spain	44,145	24,100	290,500	7700	92,400	1500	19,100
Sweden	9851	2900	28,300	30	1000	120	1400
UK	65,379	31,800	324,900	11,800	99,700	530	6400
Albania	2876	5100	14,500	70	1300	180	6890
Andorra	73	40	540	<1	40	<5	40
Bosnia and Herzegovina	3516	5400	41,700	20	1700	120	2000
Iceland	333	60	670	<1	30	<5	5
Kosovo	1772	3800	36,300	20	650	100	1300
Liechtenstein	38	20	210	<1	30	<5	20
Monaco	38	30	290	10	270	<5	20
Montenegro	622	630	7300	<1	260	20	410
North Macedonia	2071	3400	30,400	110	1200	70	1100
Norway	5211	1300	12,900	130	2000	50	550
San Marino	33	30	280	<1	10	<5	20
Serbia	7076	13,700	127,800	1500	1796	280	4300
Switzerland	8327	3700	42,800	620	10,500	240	3300
EU-28	506,028	374,000	4,150,000	68,000	795,000	14,000	180,000
Total	538,014	412,000	4,446,000	71,000	821,000	15,100	193,800

This evidence demonstrates that individual health risks deriving from the inhalation of air pollutants are likely under-perceived, as well as the related societal and economic burden due to premature deaths and years of life lost. It took decades to define the risks related to cigarette smoking, based on scientific evidence, but the perception of individual risk arising from smoking is still not clear to smokers. One might argue that the same is happening with the issue of health consequences of air pollution. The fact that risk perception in the public opinion concerning this dramatic topic is largely inadequate is confirmed by figures provided by the EEA, which show that WHO guideline values for fine particulate matter (PM_{2.5}) are exceeded at 69% of all reporting monitoring stations; as a result, 77% of the urban population across the EU was exposed to air pollution at levels indicated as harmful by the World Health Organization [1,2]. Indeed, the recent fitness check of the air quality legislation of the EU highlights that current EU air quality standards are not as ambitious as established scientific advice suggests for several air pollutants, especially fine particulate matter (PM_{2.5}). The European Green Deal

has announced that the Commission will strive to steer Europe towards a zero-pollution ambition and, in that context, draw on the lessons learnt from the evaluation of the current air quality legislation, with a view to revise EU air quality standards and align them more closely with the WHO recommendations.

3. The Need for a Medical Reading of Environmental Monitoring Data

A huge number of reliable epidemiological studies have associated air pollution with increased mortality due to all causes in the general population, especially cardio-cerebral-vascular accidents and respiratory diseases or lung cancers [6–21]. Moreover, an increase in the number of hospitalizations has been proved within 48–72 hours from peak concentrations of fine particulate matter [22–29]. Long term negative effects of air pollution have also been documented for neurodegenerative diseases as well as in terms of potentiality to cause type-2 diabetes and leukemia [30–32]. Scientific evidence indicates that all the health effects due to PM 2.5 and PM 10 are already displayed at the current limits of particulate concentrations allowed in EU Member States, both in terms of all-cause mortality and cause-specific mortality (i.e., cardiopulmonary diseases and lung cancer) [21]. However, environmental data provided by official regional or national agencies—used by decision makers to adopt preventive measures such as limitations on urban traffic or domestic heating—are still referred to air quality standards established by the law (usually on the basis of standards set at European level, at least for the EU). These legal thresholds, however, are not fully in line with the levels recommended by the WHO to protect the population against all impacts from air pollution, particularly children or more vulnerable people [24]. A stronger cooperation between environmental agencies and health authorities is needed to address the new challenges to human and planetary health arising from air pollution and climate change. Finally, it is important to provide, in a systematic way, a medical reading of environmental monitoring data, and guarantee the application of the precautionary principle enshrined in Article 191 of the Treaty on the Functioning of the European Union and in the communication of 22 February 2000 of the European Commission. Health authorities, both at national and regional or local level, should integrate their medical staff with new professionals and researchers adequately trained in environmental sciences to foster population health protection against air pollution. For this purposes, multi-disciplinary research units or teams should be established by local health authorities on environmental health topics, working together with medical staff and environmental agencies for a mutual integration of competencies.

Author Contributions: M.I., P.P., E.M. and A.M. equally contributed to conceive, write and revise the article. All authors have read and agreed to the published version of the manuscript.

Funding: No external funding source.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization; Occupational and Environmental Health Team. *WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide: Global Update 2005: Summary of Risk Assessment*; World Health Organization: Geneva, Switzerland, 2006; Available online: http://apps.who.int/iris/bitstream/10665/69477/1/WHO_SDE_PHE_OEH_06.02_eng.pdf (accessed on 2 March 2020).
2. EEA. *Air Quality in Europe—2019 Report*; No 10/2019; European Environment Agency: Copenhagen, Denmark, 2019.
3. IARC. *Outdoor Air Pollution*; IARC MonogrEvalCarcinog Risks Hum; World Health Organization: Geneva, Switzerland, 2013; Volume 109.
4. World Health Organization Media Centre. Available online: <https://www.who.int/mediacentre/news/releases/2014/air-pollution/en/> (accessed on 3 April 2019).
5. Chiabai, A.; Spadaro, J.V.; Neumann, M.B. Valuing deaths or years of life lost? Economic benefits of avoided mortality from early heat warning systems. *Mitig. Adapt. Strateg. Glob. Chang.* **2018**, *23*, 1159–1176. [CrossRef] [PubMed]

6. Shi, L.; Zanobetti, A.; Kloog, I.; Coull, B.A.; Koutrakis, P.; Melly, S.J.; Schwartz, J.D. Low-concentration PM_{2.5} and mortality: Estimating acute and chronic effects in a population-based study. *Environ. Health Perspect.* **2016**, *124*, 46–52. [CrossRef] [PubMed]
7. Chan, Y.L.; Wang, B.; Chen, H.; Ho, K.F.; Cao, J.; Hai, G.; Jalaludin, B.; Herbert, C.; Thomas, P.S.; Saad, S.; et al. Pulmonary inflammation induced by low dose particulate matter exposure in mice. *Am. J. Physiol. Lung Cell.Mol. Physiol.* **2019**. [CrossRef] [PubMed]
8. Badaloni, C.; Cesaroni, G.; Cerza, F.; Davoli, M.; Brunekreef, B.; Forastiere, F. Effects of long-term exposure to particulate matter and metal components on mortality in the Rome longitudinal study. *Environ. Int.* **2017**, *109*, 146–154. [CrossRef]
9. Samoli, E.; Stafoggia, M.; Rodopoulou, S.; Ostro, B.; Declercq, C.; Alessandrini, E.; Díaz, J.; Karanasiou, A.; Kelessis, A.G.; Le Tertre, A.; et al. MED-PARTICLES Study Group. Associations between fine and coarse particles and mortality in Mediterranean cities: Results from the MED-PARTICLES project. *Environ. Health Perspect.* **2013**, *121*, 932–938. [CrossRef]
10. Dominici, F.; McDermott, A.; Daniels, M.; Zeger, S.L.; Samet, J.M. Revised analyses of the National Morbidity, Mortality, and Air Pollution Study: Mortality among residents of 90 cities. *J. Toxicol. Environ. Health A* **2005**, *68*, 1071–1092. [CrossRef]
11. Ostro, B.; Feng, W.Y.; Broadwin, R.; Green, S.; Lipsett, M. The effects of components of fine particulate air pollution on mortality in California: Results from CALFINE. *Environ. Health Perspect.* **2007**, *115*, 13–19. [CrossRef]
12. Gryparis, A.; Forsberg, B.; Katsouyanni, K.; Analitis, A.; Touloumi, G.; Schwartz, J.; Samoli, E.; Medina, S.; Anderson, H.R.; Niciu, E.M.; et al. Acute effects of ozone on mortality from the “air pollution and health: A European approach” project. *Am. J. Respir. Crit. Care Med.* **2004**, *170*, 1080–1087. [CrossRef]
13. Forastiere, F.; Stafoggia, M.; Berti, G.; Bisanti, L.; Cernigliaro, A.; Chiusolo, M.; Mallone, S.; Miglio, R.; Pandolfi, P.; Rognoni, M.; et al. Particulate matter and daily mortality: A case-crossover analysis of individual effect modifiers. *Epidemiology* **2008**, *19*, 571–580. [CrossRef]
14. Samet, J.M.; Dominici, F.; Curriero, F.C.; Coursac, I.; Zeger, S.L. Fine Particulate Air Pollution and Mortality in 20 U.S. Cities, 1987–1994. *N. Engl. J. Med.* **2000**, *343*, 1742–1749. [CrossRef]
15. Gehring, U.; Heinrich, J.; Krämer, U.; Grote, V.; Hochadel, M.; Sugiri, D.; Kraft, M.; Rauchfuss, K.; Eberwein, H.; Wichmann, H. Long-Term Exposure to Ambient Air Pollution and Cardiopulmonary Mortality in Women. *Epidemiology* **2006**, *17*, 545–551. [CrossRef] [PubMed]
16. Brunekreef, B.; Beelen, R.; Hoek, G.; Schouten, L.; Bausch-Goldbohm, S.; Fischer, P.; Armstrong, B.; Hughes, E.; Jerrett, M.; Van den Brandt, P. Effects of long-term exposure to traffic-related air pollution on respiratory and cardiovascular mortality in the Netherlands: The NLCS-AIR study. *Res. Rep. Health Eff. Inst.* **2009**, *139*, 5–89.
17. Raaschou-Nielsen, O.; Andersen, Z.J.; Beelen, R.; Samoli, E.; Stafoggia, M.; Weinmayr, G.; Homann, B.; Fischer, P.; Nieuwenhuijsen, M.J.; Brunekreef, B.; et al. Air pollution and lung cancer incidence in 17 European cohorts: Prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol.* **2013**, *14*, 813–822. [CrossRef]
18. Faustini, A.; Stafoggia, M.; Berti, G.; Bisanti, L.; Chiusolo, M.; Cernigliaro, A.; Mallone, S.; Primerano, R.; Scarnato, C.; Simonato, L.; et al. The relationship between ambient particulate matter and respiratory mortality: A multi-city study in Italy. *Eur. Respir. J.* **2011**, *38*, 538–547. [CrossRef]
19. Vigotti, M.A.; Chiaverini, F.; Biagiola, P.; Rossi, G. Urban air pollution and emergency visits for respiratory complaints in Pisa, Italy. *J. Toxicol. Environ. Health A* **2007**, *70*, 266–269. [CrossRef]
20. Pope, C.A., III; Burnett, R.T.; Thun, M.J.; Calle, E.E.; Krewski, D.; Ito, K.; Thurston, G.D. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* **2002**, *287*, 1132–1141. [CrossRef]
21. European Commission Working Document (2019) 427. Fitness Check of the Ambient Air Quality Directives.
22. European Commission, COM(2019) 640 Final. The European Green Deal. Available online: https://ec.europa.eu/environment/air/pdf/SWD_2019_427_F1_AAQ%20Fitness%20Check.pdf (accessed on 13 March 20).
23. WHO. *Review of Evidence on Health Aspects of Air Pollution—REVIHAAP Project: Final Technical Report*; WHO: Geneva, Switzerland, 2013; Available online: http://www.euro.who.int/__data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report-final-version.pdf?ua=1 (accessed on 13 March 2020).

24. Beelen, R.; Hoek, G.; Van den Brandt, P.A.; Goldbohm, R.A.; Fischer, P.; Schouten, L.J.; Armstrong, B.; Brunekreef, B. Long-term exposure to traffic-related air pollution and lung cancer risk. *Epidemiology* **2008**, *5*, 702–710. [[CrossRef](#)]
25. Vineis, P.; Hoek, G.; Krzyzanowski, M.; Vigna-Taglianti, F.; Veglia, F.; Airoidi, L.; Autrup, H.; Dunning, A.; Garte, S.; Hainaut, P.; et al. Air pollution and risk of lung cancer in a prospective study in Europe. *Int. J. Cancer* **2006**, *119*, 169–174. [[CrossRef](#)]
26. Colais, P.; Serinelli, M.; Faustini, A.; Stafoggia, M.; Randi, G.; Tessari, R.; Chiusolo, M.; Pacelli, B.; Mallone, S.; Vigotti, M.A.; et al. Air pollution and urgent hospital admissions in nine Italian cities. Results of the EpiAir Project. *Epidemiol. Prev.* **2009**, *33* (Suppl. 1), 77–94.
27. Biggeri, A.; Bellini, P.; Terracini, B. Meta-analysis of the Italian studies on short-term effects of air pollution—MISA 1996–2002. *Epidemiol. Prev.* **2004**, *28*, 4–100.
28. Anderson, H.R.; Spix, C.; Medina, S.; Schouten, J.P.; Castellsague, J.; Rossi, G.; Zmirou, D.; Touloumi, G.; Wojtyniak, B.; Ponka, A.; et al. Air pollution and daily admissions for chronic obstructive pulmonary disease in 6 European cities: Results from the APHEA project. *Eur. Respir. J.* **1997**, *10*, 1064–1071. [[CrossRef](#)] [[PubMed](#)]
29. Magnani, C.; Mattioli, S.; Miligi, L.; Ranucci, A.; Rondelli, R.; Salvan, A.; Bisanti, L.; Masera, G.; Rizzari, C.; Zambon, P.; et al. SETIL: Italian multicentric epidemiological case-control study on risk factors for childhood leukaemia, non-hodgkin lymphoma and neuroblastoma: Study population and prevalence of risk factors in Italy. *Ital. J. Pediatr.* **2014**, *40*, 103. [[CrossRef](#)] [[PubMed](#)]
30. Power, M.C.; Adar, S.D.; Yanosky, J.D.; Weuve, J. Exposure to air pollution as a potential contributor to cognitive function, cognitive decline, brain imaging, and dementia: A systematic review of epidemiologic research. *Neurotoxicology* **2016**, *56*, 235–253. [[CrossRef](#)] [[PubMed](#)]
31. Renzi, M.; Cerza, F.; Gariazzo, C.; Agabiti, N.; Cascini, S.; Di Domenicantonio, R.; Davoli, M.; Forastiere, F.; Cesaroni, G. Air pollution and occurrence of type 2 diabetes in a large cohort study. *Environ. Int.* **2018**, *112*, 68–76. [[CrossRef](#)]
32. EUR-Lex. Access to European Union Law. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=LEGISSUM:I32042> (accessed on 2 March 2020).



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).