

## R E V I E W

## Sudden unexplained death due to cardiac arrest

*Vincenza Precone<sup>1</sup>, Giulia Guerri<sup>2</sup>, Geraldo Krasi<sup>3</sup>, Laura Lupi<sup>4</sup>, Ilaria Papa<sup>4</sup>, Tommaso Beccari<sup>5</sup>, Paolo Enrico Maltese<sup>2</sup>, Elena Manara<sup>1</sup>, Matteo Bertelli<sup>1</sup>*

<sup>1</sup> MAGI Euregio, Bolzano, Italy; <sup>2</sup> MAGI's Lab, Rovereto (TN), Italy; <sup>3</sup> MAGI Balkans, Tirana, Albania; <sup>4</sup> Section of Cardiovascular Diseases, Department of Clinical and Surgical Specialities, Radiological Sciences and Public Health, University and Spedali Civili Hospital of Brescia, Italy; <sup>5</sup> Department of Pharmaceutical Sciences, University of Perugia, Perugia, Italy

**Summary.** Sudden unexplained death due to cardiac arrest refers to a group of heterogeneous heart disorders characterized by sudden cessation of cardiac activity followed by hemodynamic collapse. It may be associated with structural heart disease or may occur in the absence of structural abnormalities. These inherited conditions increase the risk of sudden unexplained death in living relatives when there is a family history of sudden death. It is recommended to screen other family members of sudden unexplained death victims, as studies have revealed affected individuals in 40% of families. ([www.actabiomedica.it](http://www.actabiomedica.it))

**Key words:** sudden unexplained death, cardiac arrest, arrhythmia

Sudden unexplained death (SUD) due to cardiac arrest refers to a group of heterogeneous heart disorders characterized by sudden cessation of cardiac activity followed by hemodynamic collapse. In the elderly, it is a relatively common cause of death, mostly due to structural heart disease. In the non-elderly it is less frequent and can be called sudden unexplained death syndrome, which includes all sudden autopsy-negative deaths occurring after the first year of life and involving previously healthy children, adolescents and young adults (1). The world incidence of SUD due to cardiac arrest is 4-5 million cases per year (4).

Sudden unexplained death may be associated with structural heart disease, such as hypertrophic cardiomyopathy, dilated cardiomyopathy, arrhythmogenic cardiomyopathy and arrhythmogenic syndromes, or may occur in the absence of structural abnormalities, as in the case of long or short QT syndrome, Brugada syndrome, catecholaminergic polymorphic ventricular tachycardia, early repolarization syndrome, idiopathic ventricular fibrillation, primary arrhythmogenic disorders, Wolff-Parkinson-White syndrome or SUD in

epilepsy (2,3). Approximately 80% of cases are ascribed to coronary artery disease, 10-15% are associated with non-ischemic cardiomyopathy and 5% are caused by arrhythmic disorders (4). These inherited conditions increase the risk of SUD in living relatives when there is a family history of sudden death. It is recommended to screen other family members of SUD victims, as studies have revealed affected individuals in 40% of families (5). First and second-degree relatives of the deceased should undergo comprehensive cardiovascular evaluation, physical examination, 12-lead electrocardiogram, treadmill stress test, 24-h Holter monitoring and echocardiogram, and full personal and family medical histories should be recorded (6).

Genetic testing is useful for confirming diagnosis of SUD and for differential diagnosis, recurrence risk evaluation and prenatal diagnosis in families with a known mutation. Pathogenic variants may be sequence missense, nonsense, splicing and small indels, as well as large deletions/duplications (Supplementary Table 1). A multi-gene NGS panel is used by MAGI to detect nucleotide variations in coding exons and flank-

ing introns of the above genes. Tests comprehensive of all genes associated with disorders that may lead to sudden unexplained death are not currently listed in Orphanet, but are offered by 16 accredited medical genetic laboratories in the US, listed in the GTR database. The guidelines for clinical use of genetic testing are described by the American Heart Association (7).

## Conclusions

We created a NGS panel to detect nucleotide variations in coding exons and flanking regions of all the genes associated with cardiac disorders. When a family with a previous case of sudden unexplained death comes to our attention we perform the analysis of all the associated genes.

In order to have a high diagnostic yield, we developed a NGS test that reaches an analytical sensitivity (proportion of true positives) and an analytical specificity (proportion of true negatives) of  $\geq 99\%$  (coverage depth  $\geq 10x$ ).

**Conflict of interest:** Each author declares that he or she has no commercial associations (e.g. consultancies, stock ownership, equity interest, patent/licensing arrangement etc.) that might pose a conflict of interest in connection with the submitted article

## References

1. Ellsworth EG, Ackerman MJ. The changing face of sudden cardiac death in the young. *Heart Rhythm* 2005; 2: 1283-5.
2. Noseworthy PA, Newton-Cheh C. Genetic determinants of sudden cardiac death. *Circulation* 2008; 118: 1854-63.
3. Barsheshet A, Brenyo A, Moss AJ, Goldenberg I. Genetics of sudden cardiac death. *Curr Cardiol Rep* 2011; 13: 364-76.
4. Chugh SS, Reinier K, Teodorescu C, et al. Epidemiology of sudden cardiac death: clinical and research implications. *Prog Cardiovasc Dis* 2008; 51: 213-28.
5. Tan HL, Hofman N, van Langen IM, van der Wal AC, Wilde AA. Sudden unexplained death: heritability and diagnostic yield of cardiological and genetic examination in surviving relatives. *Circulation* 2005; 112: 207-13.
6. Tester DJ, Ackerman MJ. Cardiomyopathic and channelopathic causes of sudden unexplained death in infants and children. *Annu Rev Med* 2009; 60: 69-84.
7. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death – executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm* 2018; 15: e190-252.

---

Received: 5 August 2019

Accepted: 5 September 2019

Correspondence:

Stefano Paolacci

Via delle Maioliche 57/D

38068 Rovereto (TN), Italy

E-mail: stefano.paolacci@assomagi.org