



HHS Public Access

Author manuscript

Nat Commun. Author manuscript; available in PMC 2015 March 26.

Published in final edited form as:

Nat Commun. ; 5: 5015. doi:10.1038/ncomms6015.

A Non-inactivating High-voltage-activated Two-Pore Na⁺ Channel that Supports Ultra-long Action Potentials and Membrane Bistability

Chunlei Cang^{1,*}, Kimberly Aranda¹, and Dejian Ren^{1,*}

¹Department of Biology, University of Pennsylvania, 415 S. University Ave., Philadelphia, Pennsylvania 19104, USA

Abstract

Action potentials (APs) are fundamental cellular electrical signals. The genesis of short APs lasting milliseconds is well understood. Ultra-long APs (ulAPs) lasting seconds to minutes also occur in eukaryotic organisms, but their biological functions and mechanisms of generation are largely unknown. Here, we identify TPC3, a previously uncharacterized member of the two-pore channel protein family, as a new voltage-gated Na⁺ channel (Na_v) that generates ulAPs, and that establishes membrane potential bistability. Unlike the rapidly inactivating Na_vs that generate short APs in neurons, TPC3 has a high activation threshold, activates slowly, and does not inactivate—three properties that help generate long-lasting APs and guard the membrane against unintended perturbation. In amphibian oocytes, TPC3 forms a channel similar to channels induced by depolarization and sperm entry into eggs. TPC3 homologs are present in plants and animals, and they may be important for cellular processes and behaviors associated with prolonged membrane depolarization.

Action potentials (APs) are the most fundamental electrical means that neurons use to communicate with each other. In mammals, neuronal APs generally last only a few milliseconds. This short period allows for energy conservation and a high capacity for information coding¹. Beginning with the Hodgkin-Huxley studies of squid giant axon APs², the functions and mechanisms underlying the generation of short APs have been extensively studied. All of the major ion channel proteins and genes responsible for initiating APs have been discovered, and many of them have been studied to atomic resolution³. There are nine plasma membrane voltage-gated Na⁺-selective channel (Na_v) pore-forming subunits (Na_v1.1–Na_v1.9) in mammals. These proteins have four homologous repeats of six transmembrane-spanning domains (4×6TM)⁴. The pores of voltage-gated K⁺ channels (K_vs) are made of four subunits of 1×6TM proteins⁵. The rapid inactivation of Na_vs, within

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Correspondence should be addressed to DR (dren@sas.upenn.edu) or CC (ccang@sas.upenn.edu).

Author Contributions

C.C. contributed all the patch-clamp recording data. K.A. contributed the imaging data. C.C. and D.R. designed the experiments. K.A. and D.R. developed reagents. D.R. supervised the research. C.C. and D.R. wrote the manuscript.

Competing Financial Interests

The authors declare that they have no competing financial interests.

milliseconds, and the activation of K_v s are responsible for the falling phase of APs and their short duration.

Ultra-long AP-like depolarizations (ulAPs) that can last seconds to tens of minutes have also been recorded in many eukaryote organisms, including unicellular amoebae^{6, 7} and multicellular organisms, like algae⁸, *Arabidopsis thaliana*^{9, 10}, marine worms^{11, 12}, tunicates¹³, and amphibians^{14, 15}. The functions of ulAPs are not well established, but they are thought to mediate ion homeostasis, cytoplasmic streaming, tissue repair, intercellular communication, response to environmental stimuli, and polyspermy block¹⁶. The mechanisms underlying the generation of ulAPs are also poorly understood. In cells in low ionic environments, such as pond water, Cl^- efflux may mediate depolarization^{8, 17}. Many Cl^- channels have been identified, but they have not been shown to generate long APs when introduced into heterologous expression systems¹⁸. Some ulAPs depend on a novel Na^+ channel on the plasma membrane^{11, 14, 15}. The biophysical properties of this channel are markedly different from the rapidly inactivating Na_v s. However, the molecular identity of the channel has not been verified.

Here, we identify TPC3 as such a Na^+ -selective ion channel. TPC3 has a 2×6 TM structure, half the size of the 4×6 TM rapidly inactivating Na_v s. TPC3 homologs are present in many different organisms that have been reported to have ulAPs. Unlike the 4×6 TM Na_v s, TPC3 has a high threshold voltage (~ 0 mV) for activation, and the channel does not inactivate. In fibroblasts and neurons transfected with TPC3, a brief depolarization stimulus elicits APs that last several seconds to minutes. When expressed in *Xenopus* oocytes, TPC3 forms an inducible voltage-gated Na^+ channel that does not inactivate and has an activity profile similar to that of the native conductance reported for a previously uncharacterized Na_v .

RESULTS

TPC3 is a plasma membrane-localized TPC family member

To uncover the molecular mechanisms underlying the generation of ulAPs, we searched candidate proteins that can form non-inactivating Na_v s. Previous studies have shown that plasma membrane Na_v s can be formed by tetramers of 1×6 TM proteins from the Na_v Bac protein family in bacteria¹⁹ and by monomers of 4×6 TM proteins from the Na_v family ($Na_v1.1$ – $Na_v1.9$) in animals^{20, 21}. However, depolarizing holding potentials inactivate all members of these protein families. The third family of ion channels that has sequence similarity to the Na_v Bac and Na_v families is the two-pore channel (TPC) family of 2×6 TM proteins, which has three vertebrate members (TPC1–3). TPCs are thought to function in intracellular organelles^{22, 23}. Whole organelle patch-clamp experiments have shown that both native and heterologously expressed mammalian TPC1 and TPC2 create functional ion channels on endosomes and lysosomes^{24, 25, 26, 27, 28, 29, 30, 31}.

TPC3 is the only one of the 25 proteins found in animals from the three ion channel families (1×6 TM, 2×6 TM, and 4×6 TM) with sequence similarities to Na_v s and Ca_v s that has not been electrophysiologically characterized. Humans and mice do not have a functional *tpc3* gene^{32, 33}. Therefore, we isolated full-length TPC3 cDNA from zebrafish, *Danio rerio* (drTPC3). The sequence of drTPC3 is similar to human TPC1 (47% similarity and 23%

identity), Na_vs (46% similarity and 26% identity to hNa_v1.1), and Ca_vs (43% similarity and 23% identity to hCa_v1.1). Interestingly, the S4 segments in the two repeats, IS4 and IIS4, have three charged residues (R/K) displaced in an every-third-residue fashion in their N-terminal regions, which is characteristic of voltage-gated channels, such as the *Shaker* K_v channel (Fig. 1a), Na_vs, and Ca_vs.

In contrast to the intracellular localization of TPC1 and TPC2²⁴, GFP-tagged drTPC3 was detected on the plasma membrane (Fig. 1b). As expected for many overexpressed membrane proteins, a weak GFP signal was observed on endolysosome-like organelles. Similar to mammalian TPC1s and TPC2s, Zebrafish TPC2 (drTPC2) was primarily localized to intracellular organelles (Fig. 1c).

TPC3 forms a high-voltage-activated Na⁺ channel

As the electrophysiological properties of all other TPC members have been recorded from endolysosomes^{25, 26, 27, 28, 29, 30}, we first characterized drTPC3 expressed on endolysosomes using whole-endolysosomal recordings with symmetrical Na⁺ concentration (145 mM) in the bath and pipette. Organelles expressing drTPC3 protein were identified by GFP fused to drTPC3. No obvious current was detected at voltages of -100 to +70 mV. When the voltage applied was increased above +70 mV, currents were detectable in a subset of GFP-positive endolysosomes (18/42, >100 pA at +150 mV; Supplementary Fig. 1). These results were in contrast to hTPC1- or hTPC2-expressing endolysosomes, which give robust currents under similar conditions^{25, 26, 27}. In endolysosomes where a TPC3 current (I_{TPC3}) was detected, the current's activation threshold of +70 mV was much higher than that of hTPC1 (~0 mV) under identical recording conditions (pH_{lumen} 4.6)²⁶ and is unlikely to be reached by lysosomes under physiological conditions.

To test whether drTPC3 formed a functional plasma membrane ion channel, we recorded whole-cell currents from HEK293T cells transfected with drTPC3. Little voltage-activated current was observed in HEK293T cells that were not transfected (Fig. 2a) or were transfected with the lysosome-localized drTPC2 (Fig. 2b). Remarkably, transfection with drTPC3, whether GFP-tagged (Fig. 2c) or untagged (Fig. 2d), generated robust currents elicited by a depolarizing membrane potential (14/15, >500 pA at +100 mV).

The drTPC3 was activated by depolarization (Fig. 2d, e) with slow kinetics (Fig. 2c; $\tau = 224 \pm 34$ ms at +50 mV). Under more physiological asymmetric Na⁺ conditions (150 mM Na⁺ in bath and 8 mM in pipette), drTPC3 generated an inward current at voltages from approximately 0 to 60 mV (Fig. 2f). The current-voltage (I - V) relationship was similar to that of other Na_vs but was right-shifted toward higher voltages. The channel's activation threshold of approximately 0 mV was much higher than the activation threshold for 1×6TM and 4×6TM Na_vs (approximately -50 to -40 mV^{19, 34}).

TPC3 does not inactivate

The I_{TPC3} was not obviously inactivated at test voltages lasting as long as 20 s (Fig. 3a-c). In addition, pre-pulses of -100 to +150 mV, which have been shown to inactivate other plasma membrane Na_vs at high voltages, did not cause steady-state inactivation of drTPC3

(Fig. 3d, e). The lack of obvious inactivation observed for TPC3 is unlike any of the other plasma membrane Na_V s. Therefore, we conclude that drTPC3 forms a novel voltage-gated Na^+ channel with a high threshold for activation that does not inactivate.

TPC3 is Na^+ -selective

The sequence of drTPC3 is similar to those of the $4\times 6\text{TM Ca}_V$ s and Na_V s, but does not have an obvious sequence equivalent to the “DEKA” or “EEEE” ion selectivity-determining motif found in the pore-filter regions of the $4\times 6\text{TM Na}_V$ s or Ca_V s. Under conditions where the major cations were supplied by 150 mM Na^+ in the bath and 150 mM *N*-methyl-D-glucamine (NMDG) in the pipette, an inward I_{TPC3} could be elicited with depolarizing membrane potentials (Fig. 4a). However, replacing the Na^+ in the bath with 150 mM K^+ , 100 mM Ca^{2+} , or 150 mM NMDG^+ abolished the inward current, suggesting that drTPC3 was primarily permeable to Na^+ but not to K^+ , Ca^{2+} , Cl^- , or NMDG^+ (Fig. 4a, b). We used the reversal potentials measured under bi-ionic conditions to estimate the relative permeability of drTPC3 (Fig. 4c, d). The $P_{\text{Na}}/P_{\text{Cs}}$ ratio was 88.1 ± 14.5 ($n = 6$) and $P_{\text{Na}}/P_{\text{K}}$ was 117.3 ± 15.1 ($n = 11$). The selectivity that TPC3 displayed for Na^+ was comparable to that of the $4\times 6\text{TM Na}_V$ s.

I_{TPC3} was not inhibited by TTX (Fig. 5a, b), a toxin that blocks several $4\times 6\text{TM Na}_V$ s at concentrations in the sub- μM range. In contrast, drTPC3 was highly sensitive to inhibition by high-voltage-gated L-type Ca_V inhibitors (Cd^{2+} IC_{50} : $151 \pm 28 \mu\text{M}$, Fig. 5c, d; verapamil: $0.60 \pm 0.07 \mu\text{M}$; Fig. 5e, f; nifedipine: $3.35 \pm 0.26 \mu\text{M}$, Fig. 5g, h).

TPC3 is insensitive to $\text{PI}(3,5)\text{P}_2$ and $\text{PI}(4,5)\text{P}_2$

$\text{PI}(3,5)\text{P}_2$, a phosphatidylinositol bisphosphate enriched on endosomal and lysosomal membranes, potentiates several mammalian ion channels, including TRPM4, TRPMLs, TPC1 and TPC2^{26, 27, 35, 36, 37}. We measured I_{TPC3} from endolysosomes to test the sensitivity of TPC3 to $\text{PI}(3,5)\text{P}_2$. Unlike hTPC1 and hTPC2^{26, 27}, drTPC3 was not potentiated or inhibited by exogenous $\text{PI}(3,5)\text{P}_2$ (Fig. 6a, b).

In addition to requiring $\text{PI}(3,5)\text{P}_2$ for optimal activation, the endolysosome channel TRPML1 is inhibited by $\text{PI}(4,5)\text{P}_2$ ³⁸, a plasma membrane-enriched lipid required for the normal functions of many ion channels^{39, 40}. TRPML1 activity is thought to be restricted to intracellular organelles, in part due to the opposing regulatory functions of $\text{PI}(3,5)\text{P}_2$ and $\text{PI}(4,5)\text{P}_2$ ³⁸. In contrast to TRPML1, drTPC3 was not blocked by $\text{PI}(4,5)\text{P}_2$, even at a concentration of 20 μM (Fig. 6c, d). As a control, we also tested the effects of $\text{PI}(3,5)\text{P}_2$ and $\text{PI}(4,5)\text{P}_2$ on the lysosome channel drTPC2. The drTPC2 was potentiated by $\text{PI}(3,5)\text{P}_2$ (Supplementary Fig. 2a) and inhibited by $\text{PI}(4,5)\text{P}_2$ (Supplementary Fig. 2b). The contrasting sensitivity to lipid regulation between TPC3 and other TPCs separates TPC3 from the endolysosome-confined channels.

TPC3 supports ultra-long action potentials and bistability

We used current-clamp recordings to test whether TPC3 supports the generation of ulAPs when transfected into HEK293T fibroblast cells. Without transfection, HEK293T fibroblasts are non-excitable and do not generate regenerative responses (Fig. 7a, $n = 7$). In the drTPC3-

transfected cells, depolarizing stimuli, upon reaching a threshold of approximately 0 mV, elicited AP-like depolarizations that lasted seconds to many minutes (9/12, Fig. 7b–d). Further increasing the stimulus strength sometimes “drove” the membrane potential of the cell to a steady state of approximately +50 to +80 mV (Fig. 7b), close to the equilibrium potential of Na⁺ (+91 mV). Interestingly, the cells remained at the newly established depolarized membrane potential until it was terminated by a hyperpolarizing pulse (Fig. 7c), a characteristic of cell membrane bistability. Thus, transfecting drTPC3 alone was sufficient to convert non-excitable cells into excitable ones.

We also tested the ability of TPC3 to support ulAPs and membrane bistability in mammalian neurons, which express high levels of K_vs. In cultured mouse hippocampal neurons transfected with drTPC3, but not untransfected control neurons, a brief stimulus induced a long-depolarization (4/18, Supplementary Fig. 3) and sometimes induced membrane bistability.

TPC3 forms the Na⁺ channel found in amphibian oocytes

We next searched for a model system that is known to have a native Na⁺ current similar to *I*_{TPC3}. Among the commonly studied cellular model systems, frog oocytes and eggs from *Xenopus laevis* and *Rana pipiens* generate ulAPs and are known to have a non-inactivating Na⁺ channel activated at high voltages whose molecular identity is unknown^{14, 15}. A peculiar but little-understood property of the native current in these models is its “induction” by depolarization. A small current can be recorded from non-induced *R. pipiens* oocytes¹⁵, but robust currents in *X. laevis* oocytes can only be recorded after induction by a pre-pulse depolarization¹⁴ (Fig. 8a–c). The native *Xenopus* oocyte channel had a high activation threshold, activated slowly, and did not inactivate (Fig. 8b, c), similar to what has been previously reported¹⁴. The channel was inhibited by verapamil (IC₅₀, 0.8 μM) and nifedipine (IC₅₀, 7.19 μM) (Supplementary Fig. 4). Thus, the biophysical and pharmacological properties of the unknown *Xenopus* channel are similar to those of drTPC3 expressed in HEK293T cells.

To test whether the *Xenopus* oocyte Na⁺ channel is encoded by TPC3, we cloned the full-length TPC3 homolog (xITPC3) from *X. laevis* oocytes. xITPC3 is highly similar to drTPC3 (57% identity, 75% similarity). When expressed in HEK293T cells, xITPC3 formed a high-threshold voltage-activated Na⁺ channel on the plasma (Supplementary Fig. 5A, B; n = 5) and endolysosomal (Supplementary Fig. 5c, d; n = 10) membranes. Similar to drTPC3, xITPC3 was blocked by verapamil and Cd²⁺ (Supplementary Fig. 6).

We then expressed xITPC3 in its native environment by injecting xITPC3 cRNA into *Xenopus* oocytes, and compared the properties of the resulting channel to those of the native one. In oocytes injected with xITPC3 cRNA, but not in oocytes injected with H₂O, a depolarizing pre-pulse induced a non-inactivating current that, compared to the native one, was increased by 3-fold 2 days after injection and 11-fold 7 days after (Fig. 8d–f). These data suggest that TPC3 forms the native *Xenopus* oocyte channel.

DISCUSSION

We have identified the first high-voltage-activated, non-inactivating, Na⁺-selective plasma membrane ion channel and provided a molecular mechanism for the generation of ulAPs. These results and earlier extensive studies on short APs^{2, 3, 41} make it clear that both short and ultra-long APs can be generated from voltage-gated Na⁺ channels. The short APs are generated by rapidly inactivating Na_Vs, whereas the ulAPs are generated by non-inactivating Na_Vs. Cl⁻ channels and slowly inactivating Ca_Vs may also generate APs lasting several seconds under certain conditions, which require further study.

With the addition of TPC3 as a new family member, Na_Vs are now known to be formed by 1×6TM, 2×6TM, and 4×6TM proteins, the three most likely combinations of 6TM channel pore-forming building blocks⁴². Like Ca_Vs, Na_Vs can also be divided according to their activation thresholds into low-voltage- and high-voltage-activated families. TPC3 is the only known high-voltage-activated Na_V with an activation threshold of ~ 0 mV. The electrophysiological properties of all three members of the vertebrate TPC family have now been described. They all form Na⁺ channels, with TPC1 and TPC3 being depolarization-activated^{25, 26, 27, 28}. In contrast to what was previously believed, the TPC family also forms plasma membrane ion channels. Thus, future studies addressing the physiology of TPC proteins should also include plasma membrane functions.

In addition to their vastly different durations, there are other clear distinctions between short and ultra-long APs. For example, short APs can only be elicited from relatively negative resting membrane potentials because the Na_Vs are subject to steady-state inactivation at more depolarized voltages, even if still below action potential threshold. In contrast, ulAPs can be generated from much more depolarized membrane potentials (Fig. 7d) because TPC3 lacks steady-state inactivation. Short APs in mammalian neurons can also be induced by short weak depolarization stimuli, conditions achievable by synaptic inputs. The ulAPs, however, generally require much stronger stimuli with a longer duration of approximately 100 milliseconds. In frog oocytes, for instance, ulAPs are induced by a large depolarization of >0 mV or by sequential repetitive sub-threshold stimuli^{14, 43}. The egg ulAPs can also be activated by pricking the membrane or by sperm entry, mechanical or chemical stimuli that presumably lead to the membrane breaking and depolarization to approximately 0 mV for many milliseconds. A Ca_V-mediated depolarization that lasts many milliseconds also precedes Na_V opening in the “sperm-activated” ulAPs in marine worm eggs^{11, 44}.

If these ulAPs are indeed generated by TPC3, its slow activation kinetics and activation threshold close to 0 mV presumably explain the high stimulation threshold in terms of duration and amplitude for triggering the ulAPs. Because TPC3 activation can lead to a long-lasting large depolarization that sometimes does not repolarize, the high threshold for TPC3 activation may provide the cells with a low-pass filter mechanism that guards against unintended large variations in membrane potential caused by small changes in the environment. In practical terms, TPC3 and its derivatives with lower activation thresholds could serve as bistable switches for neuronal control when used in conjunction with channelrhodopsin^{45, 46}.

Unlike short APs, the *in vivo* functions of ulAPs are largely unknown. The most well-studied ulAPs are observed in oocytes and eggs from several invertebrates and vertebrates^{11, 16, 43, 47}, which do not appear to have short APs. The long depolarization in some invertebrate eggs is proposed to mediate a fast electrical polyspermy block (but see also⁴⁸). In frog oocytes and eggs under artificial conditions in a bath containing approximately 100 mM Na⁺, the ulAPs observed after electrical or mechanical induction appear to be dependent on a TPC3-like conductance^{14, 49}. However, it is unlikely that the TPC3-like conductance is used to generate an electrical polyspermy block triggered by sperm entry. Under natural conditions, the eggs are fertilized in pond water that contains low [Na⁺]. The equilibrium potential for Na⁺ is likely too low for TPC3 to generate a large depolarization. Studies have also implicated ulAPs in the oocyte maturation process when the cells are exposed to bodily fluids containing >100 mM Na⁺⁵⁰. However, we did not observe an obvious inhibition of progesterone-induced maturation of *X. laevis* oocytes treated with 5 μM verapamil, which effectively blocked xTPC3. Depolarizations similar to the TPC3-mediated one may have function besides in reproduction. For example, prolonged depolarizations induced by cell damage have been suggested to aid in wound repair and tissue regeneration, which have been characterized in non-mammals that have a higher regenerative capability than mammals⁵¹. In addition, an EST (expressed sequence tag) database search suggests that the dog TPC3 (accession # XM_005630075) is also expressed in brain (#CO590579), omentum smooth muscle (accession #HS091858) and aorta (#CO671199). The discovery of TPC3 in systems that are amenable to genetic manipulation (e.g., zebrafish) allows these hypotheses to be further tested.

High-amplitude ulAPs do not appear to be present in all animals. For example, in oocytes and eggs from several mammalian species, such as the mouse, a depolarizing current only elicits APs that last <1 s, presumably resulting from Ca_vs. These cells also lack TPC3-like currents^{52, 53, 54}. The *tpc3* gene is found in sea urchins, fish, reptiles, amphibians, birds, cats, and dogs, but it is not present in other mammals, such as humans and mice^{32, 33}. The absence of functional *tpc3* could explain the apparent lack of ulAPs in some mammals. Future comparative physiological studies with organisms from different kingdoms/phyla will determine whether differences in the presence of the *tpc3* gene in the genome are related to differences in behavior (e.g., animal posture) and cellular responses (e.g., tissue regeneration and stress responses).

METHODS

Cell culture

Animal use protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Pennsylvania. Mammalian cells were cultured in a humidified CO₂ (5%) incubator at 37°C. HEK293T cells (from ATCC) were maintained in DMEM (Gibco) supplemented with 10% FBS (Lonza) and 1× Glutamax (Gibco). Mouse hippocampal neurons were cultured as previously described⁵⁵. Hippocampi were dissected out from P0 pups, chopped into small pieces in ice-cold Ca²⁺-free HBSS, and digested in Ca²⁺-free HBSS containing 20 units/ml papain and 1 mM L-cysteine hydrochloride for 30 min at 37 °C. Neurons were dissociated by pipetting up and down with a 1 ml tip and plated

onto poly-lysine-coated cover slips. In the 24 hours after plating, neurons were cultured in DMEM (Biowhittaker) supplemented with 10% Ham's F-12 (Biowhittaker), 10% bovine calf serum (iron supplemented, Hyclone) and 0.5× Pen/strep (Gibco). Thereafter the medium was changed to Neurobasal A (Gibco) supplemented with 1× B-27 (Gibco), 1× Glutamax and 0.5× Pen/strep. Transfections were done using Lipofectamine 2000 or Lipofectamine LTX (Invitrogen).

Xenopus laevis oocytes (defolliculated with collagenase treatment) were maintained in ND96 solution⁵⁶ containing (in mM): 96 NaCl, 2 KCl, 1 MgCl₂, 1.8 CaCl₂, 5 HEPES, pH 7.4 (adjusted with NaOH), and 50 µg/ml gentamycin, at 19 ± 1 °C. Stage V and VI oocytes were selected for experiments. Each oocyte was injected with 50 nl cRNA (10 ng) or H₂O using a Nanoject II microinjector (Drummond), and was recorded two to seven days after injection.

Cloning of zebrafish and *Xenopus* TPCs

Full-length Zebrafish, *Danio rerio*, TPCs (drTPC2, drTPC3) and *Xenopus laevis* TPC3 (xlTPC3) cDNAs were amplified with PCR using first-strand cDNA synthesized from whole zebrafish ovary or *Xenopus* oocytes. The forward and reverse PCR primers containing cloning sites had the following sequences: drTPC2 (forward, TAGTCGACATGGAAGAAGAACCGCTGCTGGCTGGC; reverse, TACCCGGGTAATTATGTATTAAAGGCCGAGGTGAGGG), drTPC3 (forward, TAGTCGACATGTCTGAGGGGAAGACTGAGAAAAGTAGC; reverse, TACCCGGGTTATACGGAATCAAAAACAGGGTTTGAGAAGTTGG), xlTPC3 (forward, ATGGTACCATGTCAGAAGCTGAAGTCACAAGTAATG; reverse, TGGATCCACTGTAGTTCAACACACCGCCCAATG). All the clones were completely sequenced to ensure no artificially introduced mutation in the open reading frames (ORFs). N-terminally GFP-tagged TPCs were cloned into the Sall/XmaI sites (drTPC3) or the KpnI/BamHI sites (xlTPCs) of pEGFP-C1. Non-tagged drTPC3 were cloned into the HindIII/EcoRV sites of a pTracer-CMV2-based vector.

For cRNA synthesis used for *Xenopus* oocyte injection, ORFs without tag were transferred into pGEM-HE vector, which contains UTR sequences from *Xenopus* β-globin to enhance RNA stability and expression in the oocytes⁵⁷. cRNAs were synthesized from PCR-generated templates using an mMESAGE mMACHINE T7 kit (Ambion), precipitated with LiCl and resuspended in RNase-free H₂O.

Imaging

Transfected cells were plated onto glass coverslips, fixed with 2% paraformaldehyde and mounted with Fluoroshield™ with DAPI (Sigma). Images were taken with a laser-scanning confocal microscope (Leica TMRE TCS SL) using a 63× oil objective.

Electrophysiology

Recordings were carried out at room temperature using a Multiclamp 700B (Molecular Device) for HEK293T cells and neurons, or a GeneClamp 500B amplifier (Axon) for *Xenopus* oocytes. A Digidata 1440A (Molecular Device) under the control of pClamp

software (Molecular Device) was used for data acquisition. Clampfit (Molecular Device) software was used for data analysis. Liquid junction potentials (LJPs) were corrected for all patch-clamp recordings.

Unless otherwise stated, bath solution for whole-cell voltage-clamp recordings with HEK293T cells contained (in mM): 145 Na-methanesulfonate, 5 NaCl, 10 HEPES, 10 glucose, pH7.4. Pipette solution contained (in mM): 130 Na- methanesulfonate, 2.5 MgCl₂, 10 HEPES, 10 EGTA, 20 NaOH, pH 7.2 (LJP = 0.9 mV).

For current-clamp recordings with HEK293T cells (Fig. 7), bath solution contained (in mM): 140 NaCl, 5 KCl, 2 CaCl₂, 1 MgCl₂, 10 glucose, 20 HEPES, pH 7.4 by NaOH. Pipette solution contained (in mM): 140 K-gluconate, 4 NaCl, 2 MgCl₂, 0.39 CaCl₂, 1 EGTA, 10 HEPES, pH 7.2 by KOH (LJP = 15.3 mV). For current-clamp recordings with neurons (Supplementary Fig. 3), bath solution contained (in mM): 150 NaCl, 3.5 KCl, 1 MgCl₂, 1.2 CaCl₂, 10 HEPES, 20 glucose, pH 7.4 by NaOH. Pipette solution contained (in mM): 135 K-aspartate, 5 KCl, 5 NaCl, 1 MgCl₂, 1 EGTA, 4 Mg-ATP, 0.3 Na-GTP, 14 phosphocreatine (di-tris), 10 HEPES, pH 7.2 by KOH (LJP = 16.0 mV).

Whole-endolysosome recordings were performed with enlarged endolysosomes from HEK293T cells treated with vacuolin-1 (1 μM, overnight) as previously described²⁵. Bath solution used for endolysosomal recordings contained (in mM): 140 Na-gluconate (or K-gluconate for Supplementary Fig. 2), 4 KCl (or NaCl for Supplementary Fig. 2), 2 MgCl₂, 0.39 CaCl₂, 1 EGTA, 10 HEPES, pH 7.2 by NaOH (or KOH for Supplementary Fig. 2). Pipette solution contained (in mM): 145 NaCl, 5 KCl, 1 MgCl₂, 2 CaCl₂, 10 glucose, 10 HEPES, 10 MES, 10 glucose, pH 4.6 (adjusted with HCl; LJP = -15.9 mV for Supplementary Fig.2; LJP = -12.5 mV for the others.).

The voltage-dependence of channel activation was studied with voltage-step protocols (illustrated in Fig. 2a) using bath and pipette solutions containing 150 mM Na⁺ as the major cation. The activation curve (Fig. 2e) was fitted with Boltzmann equation:

$$G/G_{\max} = 1 / (1 + \exp[(V_{1/2} - V) / \kappa])$$

where G is the conductance, $V_{1/2}$ is the half maximum activation voltage, V is the testing voltage, and κ is the slope factor.

The pipette solution used to determine the relative permeability $P_{\text{Na}}/P_{\text{Cs}}$ consisted of (in mM) 150 CsOH, 2.5 MgCl₂, 10 HEPES, 10 EGTA, pH 7.2 by methanesulfonic acid. The bath solution contained (in mM) 50 Na-methanesulfonate, 100 NMDG, 5 HCl, 10 HEPES, 10 glucose, pH 7.4 by methanesulfonic acid (LJP = 11.4 mV). The pipette solution used to determine the relative permeability of $P_{\text{Na}}/P_{\text{K}}$ consisted of (in mM) 140 K-gluconate, 10 KOH, 2.5 MgCl₂, 10 TEA, 10 EGTA, 10 HEPES, pH 7.2 by methanesulfonic acid. The bath solution contained (in mM) 50 Na-methanesulfonate, 90 NMDG, 10 TEA, 5 HCl, 10 HEPES, 10 glucose, pH 7.4 by methanesulfonic acid (LJP = 17.3 mV). The relative permeabilities of TPC3 (Fig. 4) were calculated using the following equations:

$$P_{\text{Na}}/P_{\text{Cs}} = [C]_i \exp(E_{\text{rev}}F/RT)/[Na]_o \quad P_{\text{Na}}/P_{\text{K}} = [K]_i \exp(E_{\text{rev}}F/RT)/[Na]_o$$

where E_{rev} is the reverse potential, F is Faraday's constant, R is the gas constant, and T is the absolute temperature.

We recorded no appreciable current when Ca^{2+} was used as the bath cation, indicating that Ca^{2+} is impermeable (Fig. 4a). We did not calculate $P_{\text{Ca}}/P_{\text{Cs}}$ because the current was too small to obtain a reversal potential that was not significantly contaminated by the endogenous HEK293 cell currents. We also used Ca^{2+} as the major cation in the pipette and Na^+ as in the bath in an attempt to obtain $P_{\text{Na}}/P_{\text{Ca}}$; however, for reasons to be determined, high $[\text{Ca}^{2+}]$ in the pipette abolished the depolarization-activated inward Na^+ current, making it difficult to determine the accurate reversal potential for permeability calculation.

The inhibition curves presented in Fig. 5, Supplementary Figs. 4 and 6 were fitted with the Hill equation:

$$I/I_0 = 1/[1+(X/IC_{50})^h]$$

where I and I_0 are the current amplitudes obtained in the presence and absence of channel blockers, respectively, X is the blocker concentration, IC_{50} is the concentration required for half-maximal inhibition, and h is the Hill coefficient.

For two-electrode voltage-clamp (TEVC) recordings with *Xenopus* oocytes, ND96 was used as the bath solution. Pipettes were filled with 3 M KCl. In an attempt to knockdown xITPC3 in *Xenopus* oocytes, we injected an antisense morpholino oligo against xITPC3, but did not find obvious reduction of the native current, perhaps because of the current's already small size (~0.3 μA whole cell), the presence of two copies of xITPC3 genes with differing sequences (not shown) from the tetraploid genome, and/or a long lifetime of xITPC3's mRNA or protein.

Data analysis

Data were analyzed using Clampfit (Molecular Device), Excel (Microsoft) and OriginPro (Origin lab) and plotted using OriginPro. Numeric data were shown as mean \pm s.e.m.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Drs. H. Zhang, M. Mullins, Y. Ramu, Z. Lu, K. Foskett and Z. Ma for kindly providing us with zebrafish cDNAs and *Xenopus* oocytes. We thank useful discussions with Drs. I. Raman, B. Bean, L. Jaffe and C. Deutsch. This work was funded, in part, by NIH (grants 2R01NS055293 and 5R01NS074257 to D.R.), and the University of Pennsylvania Research Foundation.

References

1. Alle H, Roth A, Geiger JR. Energy-efficient action potentials in hippocampal mossy fibers. *Science*. 2009; 325:1405–1408. [PubMed: 19745156]
2. Hodgkin AL, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol*. 1952; 117:500–544. [PubMed: 12991237]
3. Hille B. *Ion Channels of Excitable Membranes* (ed. 3 edn). 2001
4. Catterall WA. Voltage-gated sodium channels at 60: structure, function and pathophysiology. *J Physiol*. 2012; 590:2577–2589. [PubMed: 22473783]
5. Jan YN, Jan LY. Voltage-gated potassium channels and the diversity of electrical signaling. *J Physiol*. 2012; 590:2592–2599.
6. Tasaki I, Kamiya N. A Study on Electrophysiological Properties of Carnivorous Amoebae. *Journal of cellular physiology*. 1964; 63:365–380. [PubMed: 14187282]
7. Tasaki I, Kamiya N. Electrical response of a slime mold to mechanical and electrical stimuli. *Protoplasma*. 1950; 39:333–343.
8. Findlay G, Hope AB. Ionic relations of cells of *Chara Australis* VII. The separate electrical characteristics of the plasmalemma and tonoplast. *Aust J Biol Sci*. 1964; 17:62–77.
9. Favre P, Agosti RD. Voltage-dependent action potentials in *Arabidopsis thaliana*. *Physiologia plantarum*. 2007; 131:263–272. [PubMed: 18251897]
10. Fromm J, Lautner S. Electrical signals and their physiological significance in plants. *Plant, cell & environment*. 2007; 30:249–257.
11. Jaffe LA, Kado RT, Kline D. A calcium-activated sodium conductance produces a long-duration action potential in the egg of a nemertean worm. *J Physiol*. 1986; 381:263–278. [PubMed: 2442351]
12. Kline D, Jaffe LA, Kado RT. A calcium-activated sodium conductance contributes to the fertilization potential in the egg of the nemertean worm *Cerebratulus lacteus*. *Dev Biol*. 1986; 117:184–193. [PubMed: 3091428]
13. Miyazaki S, Takahashi K, Tsuda K. Calcium and sodium contributions to regenerative responses in the embryonic excitable cell membrane. *Science*. 1972; 176:1441–1443. [PubMed: 5033653]
14. Baud C, Kado RT, Marcher K. Sodium channels induced by depolarization of the *Xenopus laevis* oocyte. *Proc Natl Acad Sci U S A*. 1982; 79:3188–3192. [PubMed: 6285341]
15. Schlichter LC. Ionic currents underlying the action potential of *Rana pipiens* oocytes. *Dev Biol*. 1989; 134:59–71. [PubMed: 2471665]
16. Jaffe LA, Cross NL. Electrical regulation of sperm-egg fusion. *Annu Rev Physiol*. 1986; 48:191–200. [PubMed: 2423022]
17. Jaffe LA, Schlichter LC. Fertilization-induced ionic conductances in eggs of the frog, *Rana pipiens*. *J Physiol*. 1985; 358:299–319. [PubMed: 2580083]
18. Duran C, Thompson CH, Xiao Q, Hartzell HC. Chloride channels: often enigmatic, rarely predictable. *Annu Rev Physiol*. 2010; 72:95–121. [PubMed: 19827947]
19. Ren D, Navarro B, Xu H, Yue L, Shi Q, Clapham DE. A prokaryotic voltage-gated sodium channel. *Science*. 2001; 294:2372–2375. [PubMed: 11743207]
20. Noda M, et al. Primary structure of *Electrophorus electricus* sodium channel deduced from cDNA sequence. *Nature*. 1984; 312:121–126. [PubMed: 6209577]
21. Catterall WA. From ionic currents to molecular mechanisms: the structure and function of voltage-gated sodium channels. *Neuron*. 2000; 26:13–25. [PubMed: 10798388]
22. Zhu MX, Ma J, Parrington J, Galione A, Evans AM. TPCs: Endolysosomal channels for Ca^{2+} mobilization from acidic organelles triggered by NAADP. *FEBS Lett*. 2010; 584:1966–1974. [PubMed: 20159015]
23. Galione A. NAADP receptors. *Cold Spring Harb Perspect Biol*. 2011; 3:a004036. [PubMed: 21047915]
24. Calcraft PJ, et al. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature*. 2009; 459:596–600. [PubMed: 19387438]

25. Cang C, et al. mTOR regulates lysosomal ATP-sensitive two-pore Na⁺ channels to adapt to metabolic state. *Cell*. 2013; 152:778–790. [PubMed: 23394946]
26. Cang C, Bekele B, Ren D. The voltage-gated sodium channel TPC1 confers endolysosomal excitability. *Nat Chem Biol*. 2014; 10:463–469. [PubMed: 24776928]
27. Wang X, et al. TPC proteins are phosphoinositide-activated sodium-selective ion channels in endosomes and lysosomes. *Cell*. 2012; 151:372–383. [PubMed: 23063126]
28. Jha A, Ahuja M, Patel S, Brailoiu E, Muallem S. Convergent regulation of the lysosomal two-pore channel-2 by Mg²⁺, NAADP, PI(3,5)P2 and multiple protein kinases. *EMBO J*. 2014; 33:501–511. [PubMed: 24502975]
29. Schieder M, Rotzer K, Bruggemann A, Biel M, Wahl-Schott CA. Characterization of two-pore channel 2 (TPCN2)-mediated Ca²⁺ currents in isolated lysosomes. *J Biol Chem*. 2010; 285:21219–21222. [PubMed: 20495006]
30. Pitt SJ, et al. TPC2 is a novel NAADP-sensitive Ca²⁺ release channel, operating as a dual sensor of luminal pH and Ca²⁺. *J Biol Chem*. 2010; 285:35039–35046. [PubMed: 20720007]
31. Brailoiu E, et al. An NAADP-gated two-pore channel targeted to the plasma membrane uncouples triggering from amplifying Ca²⁺ signals. *J Biol Chem*. 2010; 285:38511–38516. [PubMed: 20880839]
32. Cai X, Patel S. Degeneration of an intracellular ion channel in the primate lineage by relaxation of selective constraints. *Mol Biol Evol*. 2010; 27:2352–2359. [PubMed: 20463046]
33. Zhu MX, Ma J, Parrington J, Calcraft PJ, Galione A, Evans AM. Calcium signaling via two-pore channels: local or global, that is the question. *Am J Physiol Cell Physiol*. 2010; 298:C430–C441. [PubMed: 20018950]
34. Raman IM, Sprunger LK, Meisler MH, Bean BP. Altered subthreshold sodium currents and disrupted firing patterns in Purkinje neurons of Scn8a mutant mice. *Neuron*. 1997; 19:881–891. [PubMed: 9354334]
35. Zhang Z, Okawa H, Wang Y, Liman ER. Phosphatidylinositol 4,5-bisphosphate rescues TRPM4 channels from desensitization. *J Biol Chem*. 2005; 280:39185–39192. [PubMed: 16186107]
36. Nilius B, et al. The Ca²⁺-activated cation channel TRPM4 is regulated by phosphatidylinositol 4,5-bisphosphate. *EMBO J*. 2006; 25:467–478. [PubMed: 16424899]
37. Dong XP, et al. PI(3,5)P2 controls membrane trafficking by direct activation of mucolipin Ca²⁺ release channels in the endolysosome. *Nat Commun*. 2010; 1:38. [PubMed: 20802798]
38. Zhang X, Li X, Xu H. Phosphoinositide isoforms determine compartment-specific ion channel activity. *Proc Natl Acad Sci U S A*. 2012; 109:11384–11389. [PubMed: 22733759]
39. Hilgemann DW, Feng S, Nasuhoglu C. The complex and intriguing lives of PIP2 with ion channels and transporters. *Sci STKE*. 2001; 11:re19. [PubMed: 11734659]
40. Suh BC, Hille B. PIP2 is a necessary cofactor for ion channel function: how and why? *Annu Rev Biophys*. 2008; 37:175–195. [PubMed: 18573078]
41. Catterall WA, Raman IM, Robinson HP, Sejnowski TJ, Paulsen O. The Hodgkin-Huxley heritage: from channels to circuits. *J Neurosci*. 2012; 32:14064–14073. [PubMed: 23055474]
42. Ren D. Sodium leak channels in neuronal excitability and rhythmic behaviors. *Neuron*. 2011; 72:899–911. [PubMed: 22196327]
43. Hagiwara S, Jaffe LA. Electrical properties of egg cell membranes. *Annual review of biophysics and bioengineering*. 1979; 8:385–416.
44. Jaffe LA, Gould-Somero M, Holland L. Ionic mechanism of the fertilization potential of the marine worm, *Urechis caupo* (Echiura). *J Gen Physiol*. 1979; 73:469–492. [PubMed: 571895]
45. Schoenberger P, Gerosa D, Oertner TG. Temporal control of immediate early gene induction by light. *PLoS One*. 2009; 4:e8185. [PubMed: 19997631]
46. Berndt A, Yizhar O, Gunaydin LA, Hegemann P, Deisseroth K. Bi-stable neural state switches. *Nat Neurosci*. 2009; 12:229–234. [PubMed: 19079251]
47. Steinhardt RA, Lundin L, Mazia D. Bioelectric responses of the echinoderm egg to fertilization. *Proc Natl Acad Sci U S A*. 1971; 68:2426–2430. [PubMed: 5289876]
48. DeFelice LJ, Dale B. Voltage response to fertilization and polyspermy in sea urchin eggs and oocytes. *Dev Biol*. 1979; 72:327–341. [PubMed: 510790]

49. Schlichter LC. A role for action potentials in maturing *Rana pipiens* oocytes. *Dev Biol.* 1983; 98:60–69. [PubMed: 6305748]
50. Schlichter LC. Spontaneous action potentials produced by Na and Cl channels in maturing *Rana pipiens* oocytes. *Dev Biol.* 1983; 98:47–59. [PubMed: 6305747]
51. Beane WS, Morokuma J, Lemire JM, Levin M. Bioelectric signaling regulates head and organ size during planarian regeneration. *Development.* 2013; 140:313–322. [PubMed: 23250205]
52. Murnane JM, DeFelice LJ. Electrical maturation of the murine oocyte: an increase in calcium current coincides with acquisition of meiotic competence. *Zygote.* 1993; 1:49–60. [PubMed: 8081801]
53. Georgiou P, Bountra C, Bland KP, House CR. Calcium-evoked opening of potassium channels in hamster eggs. *Quarterly journal of experimental physiology.* 1983; 68:687–700. [PubMed: 6316399]
54. Okamoto H, Takahashi K, Yamashita N. Ionic currents through the membrane of the mammalian oocyte and their comparison with those in the tunicate and sea urchin. *J Physiol.* 1977; 267:465–495. [PubMed: 559759]
55. Lu B, Su Y, Das S, Liu J, Xia J, Ren D. The neuronal NALCN channel contributes resting sodium permeability and is required for normal respiratory rhythm. *Cell.* 2007; 129:371–383. [PubMed: 17448995]
56. Leonard JP, Nargeot J, Snutch TP, Davidson N, Lester HA. Ca channels induced in *Xenopus* oocytes by rat brain mRNA. *J Neurosci.* 1987; 7:875–881. [PubMed: 2435866]
57. Liman ER, Tytgat J, Hess P. Subunit stoichiometry of a mammalian K⁺ channel determined by construction of multimeric cDNAs. *Neuron.* 1992; 9:861–871. [PubMed: 1419000]




Figure 1. TPC3 is expressed on the plasma membrane

(a) *Upper*: predicted membrane topology of TPC3s. *Lower*: Sequences of IS4 and IIS4 of TPC3 proteins from zebrafish (*Danio rerio*, *dr*), frog (*Xenopus laevis*, *xl*), chicken (*Gallus gallus*, *gg*), sea urchin (*Strongylocentrotus purpuratus*, *sp*), tunicate (*Ciona intestinalis*, *ci*) and marine acorn worm (*Saccoglossus kowalevskii*, *sk*), aligned against the S4 region of the Shaker K_v channel. (b, c) Localization of N-terminally GFP-tagged drTPC3 (b) and drTPC2 (c) transfected into HEK293T cells. Scale bars, 5 μ m.




Figure 2. TPC3 forms a high voltage-activated plasma membrane Na⁺ channel

(a–d) Whole-cell currents recorded with symmetrical [Na⁺] (150 mM) in the bath and pipette from nontransfected HEK293T cells (a), or cells transfected with GFP-tagged drTPC2 (GFP-drTPC2) (b), GFP-drTPC3 (c) or non-tagged drTPC3 (d). (e) Normalized conductance (G/G_{\max}) – voltage (V) relationships reconstructed from d. (f) Similar to panel c but recorded under asymmetrical [Na⁺] conditions (150 mM in the bath and 8 mM in the pipette). Left panels show representative traces obtained with a step voltage protocol (shown in a for a–d, and in f). Current (I)–voltage (V) relationships constructed using the current sizes at the end of the voltage steps are in right panels. Data are represented as mean \pm s.e.m.




Figure 3. TPC3 channels do not inactivate

(a) Representative whole-cell currents activated by 20-s-long step pulses to +50 and +100 mV ($V_h = -70$ mV). (b) Average amplitudes of the current peak and those at the end of pulses (20 s). (c) Current amplitudes obtained at the end of pulses normalized to those at the beginning (2 s). (d) Steady-state inactivation of TPC3. Currents were recorded with a test pulse to +100 mV following a pre-pulse of 2 s in duration, at voltages ranging from -100 to +150 mV (10-mV step). (e) Current amplitudes at the end of the test pulses (arrow in d) were plotted against pre-pulse voltages. Whole-cell currents were recorded from HEK293T cells transfected with drTPC3 (non-tagged). GFP-tagged drTPC3 generated similar non-inactivating currents (not shown; see also Fig. 2). Numbers of cells are indicated in parentheses. Data are represented as mean \pm s.e.m.




Figure 4. TPC3 channels are Na⁺-selective

Whole-cell currents were recorded from GFP-drTPC3-transfected HEK293T cells.

Representative traces (**a**) and average amplitudes (**b**) of currents recorded with a pipette solution containing 150 mM NMDG⁺ and bath containing 150 mM Na⁺, 150 mM K⁺, 100 mM Ca²⁺, or 150 mM NMDG⁺ as indicated. (**c**) Representative currents elicited by voltage pulses (illustrated) under bi-ionic condition (150 mM K⁺ in pipette and 50 mM Na⁺ in bath).

(**d**) Current sizes obtained from **c** plotted against test pulse voltages to obtain reversal potential (E_{rev}) for the calculation of relative permeability. Data are represented as mean \pm s.e.m.




Figure 5. TPC3 is insensitive to the Na^+ blocker TTX but is inhibited by Ca^{2+} blockers
 Whole-cell currents were elicited with a test pulse to +100 mV ($V_h = -70$ mV) and recorded under the symmetric $[\text{Na}^+]$ (150 mM in the bath and pipette) condition from GFP-drTPC3-transfected HEK293T cells. Inhibitors were added as indicated. Representative currents were recorded in the presence and absence of each drug. **(a)** Representative currents and **(b)** average current amplitudes in the presence or absence of TTX. **(c)** Representative currents and **(d)** dose response curve for Cd^{2+} . **(e)** Representative currents and **(f)** dose response curve for verapamil. **(g)** Representative currents and **(h)** dose response curve for nifedipine. Current amplitudes normalized to that obtained without drug were fitted with the Hill equations. Data are represented as mean \pm s.e.m.




Figure 6. TPC3 is insensitive to PI(3,5)P₂ and PI(4,5)P₂

Endolysosomal currents were recorded from GFP-drTPC3-transfected HEK293T cells. Concentrations of PI(3,5)P₂ and PI(4,5)P₂ are indicated in the figure. (a) Representative current traces and (b) average current amplitudes in the absence (basal) or presence of PI(3,5)P₂. (c) Representative current traces and (d) average current amplitudes in the absence or presence of PI(4,5)P₂. Data are represented as mean \pm s.e.m.




Figure 7. TPC3 supports ultra-long action potentials and membrane bistability

Representative whole-cell current clamp recordings were performed in HEK293T cells without (**a**) or with (**b–d**) GFP-drTPC3 transfection. Depolarizing current injection stimuli (illustrated below each recording with the sizes indicated) were applied. ulAPs were elicited from resting membrane potentials of -50 mV in **b** and **c**, or -20 mV in **d**.




Figure 8. TPC3-like current in *Xenopus* oocytes

Two-electrode voltage-clamp recordings were performed with *Xenopus* oocytes without (a–c) or with (d–f) xITPC3 RNA injection. (a, b, d, e) Representative currents recorded with a voltage step protocol (–50 to +80 mV with a 10-mV step, $V_h = -60$ mV, shown in a) before (a, d) and after (b, e) the oocyte was “induced” with a 60-s depolarization of +50 mV. Capacitive transient currents are masked for clarity. (c, f) Average current-voltage relationships. Data are represented as mean \pm s.e.m.