

NATURE NOTES

Spider traps amphibian in northeastern Madagascar

Thio Rosin Fulgence^{1,2,3}  | Dominic Andreas Martin³  | Holger Kreft^{3,4}  |
Fanomezana Mihaja Ratsoavina²  | Aristide Andrianarimisa² ¹Natural and Environmental Sciences, Regional University Centre of the SAVA Region (CURSA), Antalaha, Madagascar²Zoology and Animal Biodiversity, Faculty of Sciences, University of Antananarivo, Antananarivo, Madagascar³Biodiversity, Macroecology and Biogeography, University of Goettingen, Goettingen, Germany⁴Centre for Biodiversity and Sustainable Land Use (CBL), University of Goettingen, Goettingen, Germany**Correspondence**Thio Rosin Fulgence, Natural and Environmental Sciences, Regional University Centre of the SAVA Region (CURSA), Antalaha, Madagascar and Zoology and Animal Biodiversity, Faculty of Sciences, University of Antananarivo, Antananarivo, Madagascar and Biodiversity, Macroecology and Biogeography, University of Goettingen, Goettingen, Germany.
Email: thiorosinf@yahoo.fr**Funding information**

German Academic Exchange Service (DAAD), Grant/Award Number: 57449386; Diversity Turn in Land Use Science, Grant/Award Number: 11-76251-99-35/13 (ZN3119)

Abstract

Predation can take unexpected turns. For instance, various invertebrate species—most commonly spiders—may prey on vertebrates. Here, we report one observation of a spider (Sparassidae, *Damastes* sp.) feeding on an amphibian (Hyperoliidae, *Heterixalus andrakata*) inside a retreat in northeastern Madagascar. To our knowledge, this is the second report of vertebrate predation by spiders in Madagascar. Three additional observations of retreats built by the same spider species show that the spiders built similar retreats and were hiding at the rear end of the retreat. The retreats were built by weaving two green leaves together which were still attached to the tree. We speculate from the observations, that the retreat serves as a targeted trap that deceives frogs seeking shelter during daytime.

KEYWORDSamphibians, behavior, *Heterixalus andrakata*, Madagascar, predation, spider**1 | INTRODUCTION**

Finding food is an important component of animal behavior, encompassing on average more than 50% of a lifetime activity budget (Fennessy, 2004). Predation is an important technique to acquire food (Bertram, 1979; Kie, 1999) and occurs between many different taxa, such as vertebrates preying on other vertebrates, for example, a bird preying on a gecko (Koski & Merçon, 2015; Lopes et al., 2005), snakes feeding on lizards (Raselimanana, 2018), and amphibians eating amphibians (Glaw & Vences, 2007; Ndriantsoa et al., 2014; Rasolonjatovo et al., 2018), or between vertebrates and invertebrates, for example, a bird eating a

butterfly (Bowers et al., 1985; Collins & Watson, 1983; Olofsson et al., 2010; Pinheiro & Cintra, 2017; Stefanescu, 2000; Su et al., 2015). However, invertebrates can also prey on vertebrates, thereby turning the “expected order” around. Reported cases are geographically widespread and highly diverse: for example, crabs preying on frogs (Pyke et al., 2013; Rosa et al., 2014), dragonfly larvae (Barej et al., 2009) and water scorpions eating tadpoles (von May et al., 2019), water bugs preying on fish (von May et al., 2019), praying mantis feeding on lizards (Jehle et al., 1996), and carabid beetles feeding on amphibians (Wizen & Gasith, 2011). These case and further examples are collated in a recently published review (Valdez, 2020).

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd

The review also reveals that spiders are among those invertebrate predators which have been reported to prey on vertebrates and different types of prey by spider have been listed in several papers such as Jackson (1987); Pekár et al. (2012); Michalko and Pekár (2016). Among those vertebrate, preying by spider are mammals (von May et al., 2019), reptiles (Folt & Lapinski, 2017; von May et al., 2019; Shine & Tamayo, 2016), and amphibians (Amaral et al., 2015; Costapereira et al., 2010; Folly et al., 2017; Gaiarsa et al., 2012; Glos, 2003; Kirchmeyer et al., 2017; von May et al., 2019; Menin et al., 2005; Pedrozo et al., 2017). Generally, amphibians seem to be the most common vertebrate prey of spiders (Valdez, 2020), probably due to their soft skin (Valdez, 2020), but also due to their small to moderate size (Duellman & Trueb, 1986).

Most reports have documented spiders to catch their prey by active hunting (Kirchmeyer et al., 2017; Maffei et al., 2010) or by using orb webs to catch flying or jumping vertebrates such as bats, birds, and amphibians in midair (Folt & Lapinski, 2017; Kirchmeyer et al., 2017; Muscat et al., 2014; Nyffeler & Kno, 2013; Toledo, 2005).

Here, we report on a predation by a spider of the genus *Damastes* sp. catching a small frog (*Heterixalus andrakata*, Glaw and Vences, 1991, Least Concern) in northeastern Madagascar. As we understand, this is the second report of spider predation on amphibians in Madagascar.

2 | MATERIALS AND METHODS

2.1 | Study area

We conducted field observations around Ambodiala (commune Farahalana, Sambava District) and Antsikory (commune Ampanefena, Vohemar District) in northeastern Madagascar (Figure 1). The climate

in this part of Madagascar is tropical-humid. The landscape was formerly covered with humid evergreen forests (Du Puy & Moat, 1996), but forests are nowadays fragmented (Vieilledent et al., 2018) and the landscape is dominated by smallholder agriculture.

2.2 | Incidental observations


We made four incidental observations during ecological surveys in the study area. DAM made the first incidental observation, at which the spider was feeding on the amphibian, in the morning on 25th October 2017 in a woody fallow in Ambodiala (14°24'47" S, 50°5'17" E) after a bird point count. The woody fallow is a former slash-and-burn (*tavy*) field on which rice was last cultivated in 2001. The shrubs and trees inside the woody fallow were around two to three meters high.

All other observations, at which only the retreat was observed, were made by TRF inside vanilla plantations during additional ecological surveys in the area. The second observation was on 20th August 2018 at 7:40 p.m. in Antsikory (13°55'35.8" S, 50°02'40.1"E). The third observation was on the same date at 9:00 p.m. in the same village but in a different vanilla plantation (13°55'49.0" S, 50°02'26.3"E). The fourth observation was on 3rd October 2018 at 6:34 p.m. in Ambodiala (14°24'28" S, 50°5'8"E). Vanilla plantations in the study region represent agroforestry systems characterized by vanilla vines growing on small-statured support trees, while tall trees provide shade.

2.3 | Specimens

Two spider individuals were collected, euthanized, and fixed in 90% ethanol. We labeled voucher specimens with field numbers

FIGURE 1 Study area. (a) Location of the SAVA region in northeastern Madagascar, (b) General study region where the observations were conducted and (c) Location of Ambodiala and Antsikory, the villages where the observations were conducted



THC140 (first observation) and THC293 (fourth observation). We measured the specimen THC140 on millimeter paper (Figure 2a) to record prosoma and opisthosoma length. While we have not collected the frog specimen observed during the predation, we have collected one individual from the same locality of the same species, which was recorded with the field number THC144. It has been euthanized, fixed with 90% alcohol, conserved in 70% alcohol, and stored at the University Center of SAVA Region (CURSA). Tissue biopsies of frogs and spider specimens, preserved in 90% alcohol, were also deposited at the Evolutionary Biology laboratory at the University of Braunschweig, Germany. We verified the frog identification based on DNA sequences of the 16S rRNA gene of the Mitochondrial DNA. We identified the spiders to genus level with the help of an expert in arachnology, Dr. Peter Jäger, from the Senckenberg Research Institute and the Natural History Museum in Frankfurt, Germany.

3 | RESULTS

We found four different spider retreats from *Damastes* sp. (Sparassidae) that were built using leaves in three different species of trees. In all cases, the retreats consisted of leaves woven and pulled together with a spider silk. Thereby, the leaves became close to each other, closing roughly two thirds of the leaf edges. The leaves were still green and attached to the stem. The leaves were woven on apex and edge (Figure 2d), only on edges (Figure 2b) and in middle of the segmented of each other (case of *Phyllarthron madagascariensis*). The retreats were open at the leave's base and the spiders were well-hidden at the rear end of the trap (i.e., the apex of the leaves) and not visible from the entrance.

3.1 | First incidental observation

After a bird point count in the morning (6:45 a.m.), we saw how a spider (*Damastes* sp.) was feeding on a frog (*Heterixalus andrakata*, Hyperoliidae) near the woven leaves of *Tambourissa* sp. The spider held on the head of the amphibian with the fangs. The amphibian posterior legs were above the back of the spider, while the head was down. The amphibian did not move anymore, so it seemed already killed (Figure 2c). When we approached the scene, the spider with prey went hiding into the retreat (Figure 2b). We took photographs and left the predation event. The tree leaves measured circa 26 cm in length and circa 9 cm in width at the widest point. The height of the leaves from the ground was around 120 cm.

In the afternoon (4:15 p.m. of the same day), we came back to the same place and the spider was still at the same place (hiding between the leaves). We collected the specimen (Figure 2a) but could not find the frog prey anymore. Around the tree, within a 2-m-radius, we found four other living individuals of *Heterixalus andrakata*. During the second through fourth incidental observations, we found the same spider species hiding between leaves of different tree species but we could not observe any predation events.

3.2 | Second incidental observation

We found the spider during a nocturnal amphibian and reptile survey in a vanilla plantation hiding in the retreat built in two leaves of *Phyllarthron madagascariensis*. The tree leaves measured circa 29 cm in length and circa 8 cm in maximum width. The height of the woven leaves where the spider was hiding was around 180 cm from the ground.




FIGURE 2 Retreat and predation event near retreat of *Damastes* sp. (a) Spider specimen of *Damastes* sp. (THC140, adult female), the prosoma and opisthosoma are approximately 1.5 cm in length (smallest square = 0.1 cm)—Observation 1; (b) *Damastes* sp. feeding on *Heterixalus andrakata* (frog) inside of the retreat, built of leaves of *Tambourissa* sp.—Observation 1; (c) Predation event where *Damastes* sp. captured *Heterixalus andrakata* near the retreat—Observation 1; (d) *Damastes* sp. hiding in the retreat, built of leaves of *Cedrela odorata*—Observation 4

3.3 | Third incidental observation

The third observation resembled the second, but occurred in a different vanilla plantation within the same village, circa 300 m away from the second observation. The height of the woven leaves of *Phyllarthron madagascariensis* where the spider was hiding was around 170 cm above the ground.

3.4 | Fourth incidental observation

We found the spider hiding between leaves of *Cedrela odorata* (Figure 2d). Before we took the picture, we found the spider at the far end of the retreat. When we took the picture, the spider was flushed out from the retreat. The length of the leaf was circa 8 cm with a width of circa 3 cm. We found the woven leaves around 50 cm from the ground.

4 | DISCUSSION

4.1 | Predation event and retreat

We report one predation event of spider *Damastes* sp. eating a small frog *Heterixalus andrakata*. Additionally, we observed three individuals of the same species of spider sitting in retreats made by green leaves attached to the stem of the tree. The retreats generally show the same shape. Two leaves were woven by spider silk in the apex, edges and open in the base of the leaves seems enabling prey climbing up the stem of the tree to enter. The spiders do not seem to have a preference for a single tree species and the height from the ground also seems variable from those observations.

Spiders are the most cited invertebrate group preying on vertebrates (Barej et al., 2009). However, the majority of reports of amphibian predation by invertebrates stems from the Neotropics. Few predation events on Afrotropical anurans by invertebrates have been published (Babangenge et al., 2019). Reports from Africa are from Tanzania and Uganda, where fishing spiders prey on tadpoles (Vonesh, 2005), from South Africa, where crabs predate on amphibians, and from Cameroon, where wandering spiders prey on tree frogs (Barej et al., 2009). Whether this geographic bias concerning amphibian predation by invertebrates is indeed reflecting a difference in the frequency of such behavior or whether the bias is due to more research being conducted in the Neotropics (Martin et al., 2012; Meyer et al., 2015) remains, however, unclear. Our observation is, to our knowledge, the second report of spider predation on vertebrates from Madagascar after Glos (2003), which reports on spiders (Pisauridae) feeding on frogs (*Heterixalus tricolor*) on the reed grass stems within a pond in Kirindy dry forest.

4.2 | Characteristics of *Damastes* sp. spiders and *Heterixalus* sp. frogs

The genus *Damastes* is included in the Sparassidae family, a group that is called “huntsman spiders” (Rayor, 2018) and occurs around the world

(Jäger, 2012). Most huntsman spiders do not build webs to capture their prey (Rayor, 2018). Instead, some of them are known for their running and hunting habit. But, the genus *Damastes* is an exception since members of the genus typically use a sit-and-wait approach (Soutinho et al., 2018). However, some Sparassidae have been found in their own silk nest which is fastened with debris, leaving leaves or stems that are completely surrounded by silk (Jackson, 1987). Furthermore, most species in the family Sparassidae are nocturnal (Rayor, 2018).

The genus *Heterixalus* is predominantly arboreal and typically occurs in open areas within the human-dominated landscape (Blommers-Schlösser, 1982; Raharivololoniaina et al., 2003). The species *H. andrakata* is distributed in northern and northeastern Madagascar (Glaw & Vences, 2007). During our ecological survey, we found *H. andrakata* to be mostly active at night, but recorded some daytime activity in agroforests. However, the species is typically hiding away during daytime between leaves, possibly to avoid dehydration.

4.3 | Speculation on possible systematic trapping behavior

Previous reports of spiders preying on amphibians point to an opportunistic behavior and provide no evidence of specialization. Based on our report, we speculate that the spiders use targeted traps to prey on amphibians. We base this speculation on several strings of evidence. First, some reports describe spider retreats used by spiders as a protection from predators (Henschel & Jocqué, 1994; Nentwig & Heimer, 1987; Stradling, 1994; Thirunavukarasu et al., 1996). However, these retreats may also be modified to serve as a trap (Nentwig & Heimer, 1987). Second, the behavior was observed independently in four spider individuals at four different sites suggesting that the retreat building is frequently performed by *Damastes* sp. in northeastern Madagascar. Third, a key factor facilitating the trapping behavior of *Damastes* sp. may be that *Heterixalus andrakata* and possibly also other arboreal frogs try to hide from sunlight during the day in order to avoid dehydration (Rodel & Braun, 1999). When temperatures rise, the frogs look for shade and cover away from the ground, which the spiders provide in form of their retreat. The frogs might favor the seemingly protected traps in an attempt to hide from other predators such as birds that scan the vegetation for prey. Based on these strings of evidence, we speculate that amphibians may not only be an opportunistic, indiscriminate, or accidental prey, but rather a targeted systematically exploited food source of *Damastes* sp. spiders.

Nonetheless, further research is required to confirm this, especially so, as we only report a single observation of the spider feeding on the frog. Additionally, large prey such as the frog is more likely to catch the attention of an observer, thereby posing the risk of being overinterpreted.

ACKNOWLEDGMENTS

We would like to thank the landowners in Ambodiala and Antsikory who let us survey their lands. We also thank our field assistants

Romual Randrianantenaina and Erosiniot Befidimanana for their great support during fieldwork and James Herrera for his feedback that improved the quality of the first draft of this manuscript. We also thank Miguel Vences from Evolutionary Biology laboratory for genetic identification of the frog species. This study was financially supported by the Volkswagen Stiftung within the project "Diversity Turn in Land Use Science" (Grant number 11-76251-99-35/13 (ZN3119)) which is a collaboration between the University of Goettingen, the University of Antananarivo, Madagascar National Parks and the University of CURSA (Regional University Centre of SAVA Region). TRF was supported by the German Academic Exchange Service (DAAD) within the "Partnerships for Supporting Biodiversity in Developing Countries" initiative (Project Nr.57449386). We collected data under the research permit N°254/18/MEEF/SG/DGF/DSAP/SCB. Regranted by the Ministry for Water, Ecology and Forest (MEEF), Antananarivo.

CONFLICT OF INTEREST

The corresponding author declares on behalf of authors that there is no conflict of interest to disclose.

AUTHOR CONTRIBUTIONS

Thio Rosin Fulgence: Conceptualization (equal); methodology (equal); validation (equal); writing-original draft (lead). **Dominic Andreas Martin:** Conceptualization (equal); methodology (equal); validation (equal); writing-original draft (equal). **Holger Kreft:** Conceptualization (equal); methodology (equal); supervision (equal); validation (equal); writing-original draft (supporting). **Fanomezana Mihaja Ratsoavina:** Conceptualization (equal); methodology (equal); supervision (equal); validation (equal); writing-original draft (supporting). **Aristide Andrianarimisa:** Conceptualization (equal); methodology (equal); supervision (lead); validation (equal); writing-original draft (supporting).

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Thio Rosin Fulgence  <https://orcid.org/0000-0001-7205-7282>

Dominic Andreas Martin  <https://orcid.org/0000-0001-7197-2278>

Holger Kreft  <https://orcid.org/0000-0003-4471-8236>

Fanomezana Mihaja Ratsoavina  <https://orcid.org/0000-0003-1661-1669>

[org/0000-0003-1661-1669](https://orcid.org/0000-0003-1661-1669)

Aristide Andrianarimisa  <https://orcid.org/0000-0002-4839-7890>

REFERENCES

- Amaral, L. C., Castanheira, P. D. S., & De Carvalho-e-silva, S. P. (2015). Predation on the tropical bullfrog *Adenomera marmorata* (Anura: Leptodactylidae) by the wandering spider *Ctenus ornatus* (Araneae: Ctenidae) in southeastern Brazil. *Herpetology Notes*, 8, 329–330.
- Babangenge, G. B., Jocqué, R., Masudi, F. M., Rödel, M. O., Burger, M., Gvoždík, V., & Pauwels, O. S. G. (2019). Frog-eating spiders in the Afrotropics: An analysis of published and new cases. *Bulletin of the Chicago Herpetological Society*, 54, 57–63.
- Barej, M. F., Wurstner, J. A. M., & Böhme, W. (2009). Predation on the treefrog *Leptopelis brevirostris* (Anura: Arthroleptidae) by a wandering spider (Araneae: Ctenidae) in Cameroon. *Herpetology Notes*, 2, 137–139.
- Bertram, B. C. R. (1979). In J. J. R. Grimsdell (Ed.), *Studying predators* (50 pp.). 2nd No. 3. African Wildlife Foundation, P.O. Box 48177 Nairobi, Kenya.
- Blommers-Schlösser, R. M. A. (1982). Observations on the Malagasy frog *Heterixalus Laurent, 1944* (Hyperoliidae). *Beaufortia*, 32, 1–11.
- Bowers, M. D., Brown, I. L., & Wheye, D. (1985). Bird predation as a selective agent in a butterfly population. *Evolution (NY)*, 39, 93–103.
- Collins, C. T., & Watson, A. (1983). Field observations of bird predation on neotropical moths. *Biotropica*, 15, 53. <https://doi.org/10.2307/2387999>
- Costa-pereira, R., Martins, F. I., Sczesny-moraes, E. A., Brescovit, A., Grande, C., Grande, C., & Grande, C. (2010). Predation on young treefrog (*Osteocephalus taurinus*) by arthropods (Insecta, Mantodea and Arachnida, Araneae) in Central Brazil. *Biota Neotropica*, 10, 469–472. <https://doi.org/10.1590/S1676-06032010000300042>
- Du Puy, D. J., & Moat, J. (1996). A refined classification of the primary a refined classification of the primary vegetation of Madagascar based on the underlying geology: Using GIS to map its distribution and to assess its conservation status. In W. R. Lourenço (Ed.), *Proceedings of the international symposium on the "Biogeography de Madagascar", Paris, September 1995* (pp. 205–218).
- Duellman, W. E., & Trueb, L. (1986). *Biology of amphibians* (1st ed., 549 pp.).
- Fennessy, J. (2004). Behavioural ecology. In *Ecology of desert-dwelling giraffe Giraffa camelopardalis angolensis in northwestern Namibia. Sydney, Australia* (pp. 126–159).
- Folly, H., De Arruda, L. F., Gomes, V. F., Neves, M. O., & Feio, R. N. (2017). Predation on *Oolygon carnevallii* (Caramaschi and Kisteumacher, 1989) (Anura, Hylidae) by *Phoneutria nigriventer* (Keyserling, 1891) (Araneae, Ctenidae). *Herpetology Notes*, 10, 365–367.
- Folt, B., & Lapinski, W. (2017). New observations of frog and lizard prédation by wandering and orb-weaver spiders in Costa Rica. *Phyllomedusa*, 16, 269–277. <https://doi.org/10.11606/issn.2316-9079.v16i2p269-277>
- Gaiarsa, M. P., Rodrigues, L., De Alencar, V., Dias, C. J., & Martins, M. (2012). Predator or prey? Predatory interactions between the frog *Cycloramphus boraceiensis* and the spider *Trechaleoides biocellata* in the Atlantic Forest of Southeastern Brazil. *Herpetology Notes*, 5, 67–68.
- Glaw, F., & Vences, M. (2007). In M. Vences, & V. Glaw (Eds.), *Ny Toro-hay momba ny Amphibia sy ny Reptilia an'i Madagasikara* (3rd ed., 529 pp). Cologne.
- Glos, J. (2003). The amphibian fauna of the Kirindy dry forest in western Madagascar. *Salamandra*, 39, 75–90.
- Henschel, J. R., & Jocqué, R. (1994). Bauble spiders: A new species of acaearanea (Araneae: Theridiidae) with ingenious spiral retreats. *Journal of Natural History*, 28, 1287–1295. <https://doi.org/10.1080/00222939400770651>
- Jackson, R. R. (1987). The biology of *Olios* spp., huntsman spiders (Araneae, Sparassidae) from Queensland and Sri Lanka: Predatory behaviour and cohabitation with social spiders. *Bulletin of the British Arachnological Society*, 7, 133–136.
- Jäger, P. (2012). New species of the spider genus *Olios* Walckenaer, 1837 (Araneae: Sparassidae: Sparassinae) from Laos. *Zootaxa*, 68, 61–68. <https://doi.org/10.11646/zootaxa.3228.1.3>
- Jehle, R., Franz, A., Kapfer, M., Schramm, H., & Tunner, H. G. (1996). Lizards as prey of arthropods: Praying Mantis *Mantis religiosa* (Linnaeus, 1758) feeds on juvenile Sand Lizard *Lacerta agilis* Linnaeus, 1758 (Squamata: Sauria: Lacertidae). *Herpetozoa*, 9, 157–159.
- Kie, J. G. (1999). Optimal foraging and risk of predation: Effects on behavior and social structure in ungulates. *Journal of Mammalogy*, 80(4), 1114–1129. <https://doi.org/10.2307/1383163>

- Kirchmeyer, J., Amaral, L. C., Magaldi, A., Cerqueira, R. L., & De Carvalho-silva, S. P. (2017). Predation on the treefrog *Scinax similis* (Anura: Hylidae) by the orb-weaver spider *Eriophora fuliginea* (Araneae: Araneidae) in southeastern Brazil. *Phyllomedusa*, 16, 113–116.
- Koski, D. A., & Merçon, L. (2015). Predation on *Tropidurus torquatus* (Squamata: Tropiduridae) by the Guira Cuckoo *Guira guira* (Aves: Cuculiformes) in the state of Espírito Santo, Southeastern Brazil. *Herpetology Notes*, 8, 35–37.
- Lopes, L. E., Fernandes, A. M., & Marini, M. A. (2005). Predation on vertebrates by Neotropical passerine birds. *Lundiana*, 6, 57–66.
- Maffei, F., Ubaid, F. K., & Jim, J. (2010). Predation of herps by spiders (Araneae) in the Brazilian Cerrado. *Herpetology Notes*, 3, 167–170.
- Martin, L. J., Blossey, B., & Ellis, E. (2012). Mapping where ecologists work: Biases in the global distribution of terrestrial ecological observations. *Ecological Society of America*, 10(4), 195–201. <https://doi.org/10.1890/110154>
- Menin, M., Rodrigues, D. D. J., & De Azevedo, C. S. (2005). Predation on amphibians by spiders (Arachnida, Araneae) in the Neotropical region. *Phyllomedusa*, 4, 39–47. <https://doi.org/10.11606/issn.2316-9079.v4i1p39-47>
- Meyer, C., Kreft, H., Guralnick, R., & Jetz, W. (2015). Global priorities for an effective information basis of biodiversity distributions. *Nature Communications*, 6, 1–8. <https://doi.org/10.1038/ncomm59221>
- Michalko, R., & Pekár, S. (2016). Different hunting strategies of generalist predators result in functional differences. *Oecologia*, 181, 1187–1197. <https://doi.org/10.1007/s00442-016-3631-4>
- Muscat, E., Rotenberg, E. L., & Chagas, C. A. (2014). Predation of *Scinax littoralis* (Anura: Hylidae) by *Eriophora fuliginea* (Araneae: Araneidae) in Southeastern Brazil. *Herpetology Notes*, 7, 169–170.
- Ndriantsoa, S. H., Rakotonanahary, T., Dawson, J., & Edmonds, D. (2014). Predation of the critically endangered *Boophis williamsi* by *Boophis goudotii* at Ankaratra Massif, Madagascar. *Herpetology Notes*, 7, 343–345.
- Nentwig, W., & Heimer, S. (1987). Ecological aspects of spider webs. In W. Nentwig (Ed.), *Ecophysiology Spiders* (pp. 211–225). Berlin, Heidelberg: Springer. https://doi.org/10.1007/978-3-642-71552-5_15
- Nyffeler, M., & Kno, M. (2013). Bat predation by spiders. *PLoS One*, 8(3), e58120. <https://doi.org/10.1371/journal.pone.0058120>
- Olofsson, M., Vallin, A., Jakobsson, S., & Wiklund, C. (2010). Marginal eyespots on butterfly wings deflect bird attacks under low light intensities with UV wavelengths. *PLoS One*, 5(5), e10798. <https://doi.org/10.1371/journal.pone.0010798>
- Pedrozo, M., Almeida, L. D. S., Moroti, M. D. T., & Santana, D. J. (2017). Predation on *Physalaemus olfersii* (Anura: Leptodactylidae) by *Phoneutria nigriventer* (Araneae: Ctenidae) in Atlantic Forest, South-east of Brazil. *Herpetology Notes*, 10, 369–371.
- Pekár, S., Coddington, J. A., & Blackledge, T. A. (2012). Evolution of stenophagy in spiders (Araneae): Evidence based on the comparative analysis of spider diets. *Evolution* (N. Y.), 66, 776–806. <https://doi.org/10.5061/dryad.1d8761h1>
- Pinheiro, C. E. G., & Cintra, R. (2017). Butterfly predators in the neotropics: Which birds are involved? *Journal of the Lepidopterists' Society*, 71, 109–114. <https://doi.org/10.18473/lepi.71i2.a5>
- Pyke, G. H., Ah Yong, S. T., Fuessel, A., & Callaghan, S. (2013). Marine crabs eating freshwater frogs: Why are such observations so rare? *Herpetology Notes*, 6, 195–199.
- Raharivoloniaina, L., Vieites, D. R., Glaw, F., & Vences, M. (2003). Larval stages, habitat and distribution of the hyperoliid frog *Heterixalus rutenbergi* (Boettger, 1881). *Alytes*, 21, 59–65.
- Raselimanana, A. P. (2018). Field observations of predation events in Malagasy amphibians and reptiles. *Herpetology Notes*, 11, 659–662.
- Rasolonjatovo, S. M., Scherz, M. D., Raselimanana, A. P., & Vences, M. (2018). Tadpole predation by *Mantidactylus bellyi* Mocquard, 1895 with brief description of the site and morphological measurements of the specimen. *Herpetology Notes*, 11, 747–750.
- Rayor, L. S. (2018). *Huntsman spider biology: Life-history, reproduction & husbandry* (pp. 1–7).
- Rodel, M.-O., & Braun, U. (1999). Associations between Anurans and Ants in a West African Savanna (Anura: Microhylidae, Hyperoliidae, and Hymenoptera: Formicidae). *Biotropica*, 31, 178. <https://doi.org/10.2307/2663971>
- Rosa, G. M., Noël, J., Sabino-Pinto, J., & Andreone, F. (2014). Predation on the treefrog *Boophis rufioculius* (Anura, Mantellidae) by the freshwater crab *Hydrothelphusa* sp. (Decapoda, Potamonautidae) in Madagascar. *Crustaceana*, 87, 890–894. <https://doi.org/10.1163/15685403-00003331>
- Shine, R., & Tamayo, B. (2016). When predators become prey: The lizard-eating spiders of suburbia. *Zoologist*, 38, 212–213. <https://doi.org/10.7882/AZ.2016.021>
- Soutinho, J. G., Couto, H., Andreone, F., Crottini, A., & Rosa, G. M. (2018). When camouflage fails: Predation of a huntsman spider *Damastes* sp. (Araneae: Sparassidae) on a stick insect *Antongilia* sp. (Phasmatoidea: Bacillidae: Antongiliinae) from Madagascar. *Acta Arachnologica*, 67, 31–33. <https://doi.org/10.2476/asjaa.67.31>
- Stefanescu, C. (2000). Bird predation on cryptic larvae and pupae of a swallowtail butterfly. *Butlletí del Grup Català D'anellament*, 17, 39–49.
- Stradling, D. J. (1994). Distribution and behavioral ecology of an arboreal 'Tarantula' spider in Trinidad. *Biotropica*, 26, 84–97. <https://doi.org/10.2307/2389113>
- Su, S., Lim, M., & Kunte, K. (2015). Prey from the eyes of predators: Color discriminability of aposematic and mimetic butterflies from an avian visual perspective. *Evolution* (N. Y.), 69(11), 2985–2994. <https://doi.org/10.1111/evo.12800>
- Thirunavukarasu, P., Nicolson, M., & Elgar, M. A. (1996). Leaf selection by the leaf-curling spider *Phonognatha graeffei* (Keyserling) (Araneidae: Araneae). *Arachnology*, 10, 187–189.
- Toledo, L. F. (2005). Predation of juvenile and adult anurans by invertebrates: Current knowledge and perspectives. *Herpetological Review*, 36, 395–400.
- Valdez, J. W. (2020). Arthropods as vertebrate predators: A review of global patterns. *Global Ecology and Biogeography*, 29(10), 1691–1703. <https://doi.org/10.1111/geb.13157>
- Vieilledent, G., Grinand, C., Rakotomalala, F. A., Ranaivosoa, R., Rakotoarijaona, J. R., Allnutt, T. F., & Achard, F. (2018). Combining global tree cover loss data with historical national forest cover maps to look at six decades of deforestation and forest fragmentation in Madagascar. *Biological Conservation*, 222, 189–197. <https://doi.org/10.1016/j.biocon.2018.04.008>
- von May, R., Biggi, E., Cárdenas, H., & Rabosky, D. L. (2019). Ecological interactions between arthropods and small vertebrates in a lowland Amazon rainforest. *Amphibian & Reptile Conservation*, 13, 65–77.
- Vonesh, J. R. (2005). Sequential predator effects across three life stages of the African tree frog, *Hyperolius spinularis*. *Oecologia*, 143, 280–290. <https://doi.org/10.1007/s00442-004-1806-x>
- Wizen, G., & Gasith, A. (2011). Predation of amphibians by carabid beetles of the genus *Epomis* found in the central coastal plain of Israel. *ZooKeys*, 100, 181–191. <https://doi.org/10.3897/zookeys.100.1526>

How to cite this article: Fulgence TR, Martin DA, Kreft H, Ratsoavina FM, Andrianarimisa A. Spider traps amphibian in northeastern Madagascar. *Ecol Evol.* 2021;11:682–687. <https://doi.org/10.1002/ece3.7102>