Correction

## Correction: Evaluation of tenascin-C by tenatumomab in T-cell non-Hodgkin lymphomas identifies a new target for radioimmunotherapy


Giuseppe Gritti<sup>1</sup>, Andrea Gianatti<sup>2</sup>, Fiorella Petronzelli<sup>3</sup>, Rita De Santis<sup>3</sup>, Chiara Pavoni<sup>1</sup>, Riccardo Lorenzo Rossi<sup>4</sup>, Laura Cattaneo<sup>2</sup>, Luigi Giusto Spagnoli<sup>5</sup>, Silvia Ferrari<sup>1</sup>, Andrea Rossi<sup>1</sup>, Anna Maria Barbui<sup>1</sup> and Alessandro Rambaldi<sup>1,6</sup>


Published: March 30, 2018

C

**Copyright**: Gritti et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

This article has been corrected: The online version of figure 3 has been corrected:





Reference	<b>GEO</b> series	probeID	adj.P.Val	P.Value	logFC
Iqbal et al., 2010	GSE19069	201645_at	3.3E-19	6.4E-22	6.69
		216005 at	6.4E-12	7.0E-14	3.73
Piccaluga et al., 2006	GSE6338	201645_at	4.6E-15	6.0E-17	3.59
		216005 at	3.7E-07	3.5E-08	1.21

**Figure 3: (A)** Gene expression values from Iqbal et al. [25]: TNC average expression for resting cells and five type of lymphomas. **(B)** Gene expression values from Piccaluga et al [26]: TNC average expression for resting and activated normal CD4+ and CD8+ T cells and three types of lymphomas. All expression values are reported in relative fluorescence unit, as from original microarray datasets extracted with GEO2R functionality of the repository. **(C)** Differential expression values of TNC gene expression between normal tissues (resting or activated non lymphoma cells) and lymphomas from the two considered datasets relative to the TNC probes on the microarray: adjusted *p* values (BH correction), raw *p* Value and log fold change are shown for two TNC probes in both studies.

Original article: Oncotarget. 2018; 9:9766-9775. https://doi.org/10.18632/oncotarget.23919

<sup>&</sup>lt;sup>1</sup> Hematology and Bone Marrow Transplant Units, Ospedale Papa Giovanni XXIII, Bergamo, Italy

<sup>&</sup>lt;sup>2</sup> Pathology Unit, Ospedale Papa Giovanni XXII, Bergamo, Italy

<sup>&</sup>lt;sup>3</sup> Sigma Tau S.p.A. Biotech Products R and D, Pomezia, Italy

<sup>&</sup>lt;sup>4</sup> Bioinformatics, Istituto Nazionale Genetica Molecolare, Milan, Italy

<sup>&</sup>lt;sup>5</sup> Department of Biomedicine and Prevention, Università di Roma Tor Vergata, Rome, Italy

<sup>&</sup>lt;sup>6</sup> Department of Oncology and Oncohematology, Università degli Studi di Milano, Milan, Italy