



Group 13/15 Compounds Hot Paper

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 3806–3811

International Edition: doi.org/10.1002/anie.202013849

German Edition: doi.org/10.1002/ange.202013849

NHC-stabilized Parent Arsanylalanes and -gallanes

Michael A. K. Weinhart, Michael Seidl, Alexey Y. Timoshkin, and Manfred Scheer*

Dedicated to Professor Wolfgang Kaim on the occasion of his 70th birthday

Abstract: The synthesis and characterization of the unprecedented compounds $IDipp \cdot E'H_2AsH_2$ ($E' = Al, Ga$; $IDipp = 1,3$ -bis(2,6-diisopropylphenyl)imidazolin-2-ylidene) are reported, the first monomeric, parent representatives of an arsanylalane and arsanylgallane, respectively, stabilized only by a LB (LB = Lewis Base). They are prepared by a salt metathesis reaction of $KAsH_2$ with $IDipp \cdot E'H_2Cl$ ($E' = Al, Ga$). The H_2 -elimination pathway through the reaction of AsH_3 with $IDipp \cdot E'H_3$ ($E' = Al, Ga$) was found to be a possible synthetic route with some disadvantages compared to the salt metathesis reaction. The corresponding organo-substituted compounds $IDipp \cdot GaH_2AsPh_2$ (**1**) and $IDipp \cdot AlH_2AsPh_2$ (**2**) were obtained by the reaction of $KAsPh_2$ with $IDipp \cdot E'H_2Cl$ ($E' = Al, Ga$). The novel branched parent compounds $IDipp \cdot E'H(EH_2)_2$ ($E' = Al, Ga$; $E = P, As$) were synthesized by salt metathesis reactions starting from $IDipp \cdot E'HCl_2$ ($E' = Al, Ga$). Supporting DFT computations give insight into the different synthetic pathways and the stability of the products.

Introduction

The chemistry of group 13/15 compounds is an active research field and has influenced many areas of chemistry. For instance, unsaturated compounds of the type $H_2E'EH_2$ ($E' =$ Group 13 element, $E =$ Group 15 element) are isoelectronic to alkenes. They are of interest as starting materials for semiconducting applications^[1] or as precursor for composite 13/15 materials.^[2] In comparison to aminoboranes $LB \cdot BR_2NR_2 \cdot LA$ (LB = Lewis base, LA = Lewis acid) the

chemistry of the heavier group 13/15 element analogs is rarely investigated. The few known compounds of arsanylalanes and -gallanes $LB \cdot [E'R_2AsR_2]_n \cdot LA$ ($E' = Al, Ga$) exist as dimers (**A**, $n = 2$),^[3] trimers ($n = 3$),^[4] or LB/LA-stabilized monomers depending on the steric demands of the organic substituents^[5] (**B**, Figure 1) as well as the LA/LB. Since these compounds are precursors for the synthesis of binary GaAs or AlAs materials via MOCVD processes (metalorganic chemical vapor deposition),^[6] the parent compounds of these precursors are of particular interest for improving the current MOCVD process which involves the reaction of trimethylgallium with the toxic gas AsH_3 at elevated temperatures. In contrast to the phosphorus analog $E'H_2PH_2$ ($E' = Al, Ga$), for which we recently succeeded in the synthesis of the first only LB-stabilized parent compounds $IDipp \cdot E'H_2PH_2$ ($E' = Al, Ga$; $IDipp = 1,3$ -bis(2,6-diisopropylphenyl)imidazolin-2-ylidene),^[7] the heavier arsenic analogs exhibit a higher lability of the Ga–As/Al–As bond, which is why they have so far only been studied by theoretical methods.^[8] In fact, because of their toxicity, light sensitivity, and tendency to decompose, as well as the unsuitable NMR activity of the As nucleus, the handling and characterization of such compounds are hampered by numerous difficulties. Moreover, only a few examples of stable primary arsines, such as $(2,6-Tipp_2C_6H_3)AsH_2$ ($Tipp = 2,4,6$ -*i*-Pr₃C₆H₂), $TriptAsH_2$ ($Tript =$ tribenzobarrelene),^[9] or $NMe_3 \cdot BH_2AsH_2$ ^[10a] containing bulky or special substituents have so far been reported. Therefore, the question arises whether compounds containing AsH_2 bound to alanes and gallanes can be synthesized. In any case, a stabilization via a LB and a LA or at least via a LB alone would be needed if organic substitution at the As and the Al and Ga atoms, respectively, was to be avoided. Even from this perspective, it is astonishing that only parent arsanylboranes exist as LA/LB-^[10b] or LB-stabilized^[10a] molecules. No LA/LB-stabilized arsanylalanes or -gallanes have been reported yet, only their phosphanyl analogs,^[10c] which reflects the

[*] M. A. K. Weinhart, M. Seidl, Prof. Dr. M. Scheer
Institut für Anorganische Chemie
Universität Regensburg
93040 Regensburg (Germany)
E-mail: manfred.scheer@ur.de
Homepage: <http://www.uni-regensburg.de/chemie-pharmazie/anorganische-chemie-scheer>

Prof. Dr. A. Y. Timoshkin
Institute of Chemistry
Saint Petersburg State University
Universitetskaya emb. 7/9, 199034 St. Petersburg (Russia)

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202013849>.

© 2020 The Authors. *Angewandte Chemie International Edition* published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.





Figure 1. Examples of dimeric (**A**) and monomeric arsanyltrielanes (**B** and **C**).

specific lability of the corresponding E'–As bonds (E' = Al, Ga).

Herein, we report the synthesis and characterization of the first monomeric parent compound of an arsanylgallane, IDipp·GaH₂AsH₂ (**3**), and an arsanylalane, IDipp·AlH₂AsH₂ (**4**), as well as their organo-substituted analogs IDipp·E'H₂AsPh₂ (**1**: E' = Ga, **2**: E' = Al; **C**), only stabilized by a LB. The initially formed unprecedented side products IDipp·E'H·(EH₂)₂ (E' = Al, Ga; E = As, P; **5–8**) could be synthesized and characterized on a selective route.

Results and Discussion

The organo-substituted compounds IDipp·GaH₂AsPh₂ (**1**) and IDipp·AlH₂AsPh₂ (**2**) can be synthesized by the reaction of IDipp·E'H₂Cl (E' = Ga, Al)^[11] with KAsPh₂·dioxane in Et₂O at –80 °C [Eq. (1)]. Compound **1** was isolated at –30 °C as colorless crystals in a yield of 63 % and **2** as pale yellow blocks in a yield of 52 %.



In the solid state, **1** and **2** can be stored at ambient temperatures in an inert atmosphere for more than two months without decomposition. The molecular ion peak of **1** is detected at *m/z* 688.2142 in the mass spectrum (LIFDI-MS). The LIFDI-MS spectrum of **2** shows a fragment peak of IDipp⁺ due to decomposition of **2** during the ionization process. The ¹H NMR spectra of **1** and **2** show a broad singlet at δ = 4.28 ppm for the GaH₂ moiety in **1** and a broad singlet at δ = 3.95 ppm for the AlH₂ moiety in **2**, respectively. The ²⁷Al NMR spectrum of **2** reveals a broad singlet at δ = 126.5 ppm, which partially overlays with the signal of the NMR sample head and the NMR tube material.

The structures of **1** and **2**, determined by single-crystal X-ray analysis, are depicted in Figure 2 and Figure S35 (cf. SI), respectively. The Al–As bond in **2** shows a length of 2.4929(4) Å and is therefore slightly longer than the Al–As bond (2.485(2) Å) in tmp₂AlAsPh₂^[12] (tmp = 2,2,6,6-tetramethylpiperidine). Compound **1** reveals a Ga–As bond length of 2.4659(5) Å, which is in good agreement with the sum of the covalent radii (2.46 Å) of Ga and As.^[13] Compared to the few other known examples of monomeric arsanylgallanes, the Ga–As bond in **1** is slightly longer than in (C₅Me₅)₂GaAs(SiMe₃)₂ (2.433 Å)^[5a] and similar to (Me₅As)₃Ga (2.433–2.508 Å)^[14] and (*t*-Bu)₂GaAs(*t*-Bu)₂ (2.466 Å).^[5b] In contrast, dimeric structures of the type [R₂GaAsR'₂]₂ feature larger Ga–As bond lengths of 2.558, 2.550, and 2.524 Å in [*n*-




Figure 2. Molecular structure of **1** in the solid state; thermal ellipsoids at 50% probability.^[19] Selected bond lengths [Å] and angles [°]: Ga–As 2.4659(5), Ga–C1 2.068(3), C1–Ga–As 109.33(8), H1–Ga–As–C4 134.4(1).

Bu₂GaAs(*t*-Bu)₂]₂,^[15] [Me₂GaAs(*t*-Bu)₂]₂,^[15] and [Ph₂GaAs(CH₂SiMe₃)₂]₂^[4] respectively. These larger Ga–As distances are not the result of the tetracoordination of the Ga atom or the ring formation, since the trimer [Br₂GaAs(CH₂SiMe₂)₂]₃ exhibits shorter Ga–As bond lengths of 2.432(2)–2.464(1) Å. A more plausible explanation is the steric repulsion and the ring strain due to endocyclic bond angles of 83–96° in the dimers in contrast to 103–121° in the trimer [Br₂GaAs(CH₂SiMe₂)₂]₃.

Compounds **1** and **2** reveal an eclipsed conformation with a torsion angle of H1–Ga–As–C4 = 134.4° and H1–Al–As–C4 = 138.1°, respectively. The E'–C1 bond lengths in **1** (2.068(3) Å, E' = Ga) and **2** (2.0634(12) Å, E' = Al) are in the range of usual E'–C single bonds and are similar to the Ga–C1 bond length in IDipp·GaH₂Pcy₂ (2.090(2) Å,^[7] Cy = cyclohexyl) and to the Al–C1 (2.056(2) Å) bond length in IDipp·AlH₂PH₂,^[7] respectively. The C1–Ga–As angle of **1** (109.33(8)°) is in good agreement with the C1–Al–As angle in **2** (109.53(3)°).

For the synthesis of the parent compounds IDipp·GaH₂AsH₂ (**3**) and IDipp·AlH₂AsH₂ (**4**), two different routes were used [Eq. (2)]. Similarly to the substituted analogs, compounds **3** and **4** are accessible by a salt metathesis reaction between IDipp·E'H₂Cl (E' = Al, Ga) and KAsH₂ at –80 °C in THF (route 1). Furthermore, **3** and **4** can be synthesized by H₂-elimination reactions of IDipp·E'H₃ (E' = Al, Ga) and AsH₃ (route 2). For this purpose, an excess of AsH₃ is condensed onto a solution of IDipp·E'H₃ in toluene at –70 °C and stirred for 3 days at this temperature. Unfortunately, **3** and **4** were formed only in minor amounts via route 2 according to ¹H NMR spectroscopic monitoring (Figure S1 and S2). The low yield of these H₂-elimination reactions is obviously caused by the applied temperature of –70 °C, which significantly slows down the exergonic reaction between IDipp·E'H₃ and AsH₃ but was needed throughout the reaction to keep AsH₃ condensed (see below, Table 1, process 1). Compound **3** can be isolated at –30 °C in a crystalline yield of 39 % via route 1. In the mass spectrum

Table 1: Thermodynamic characteristics of studied reactions (gas phase compounds if not noted otherwise).^[a]

Entry	Process	E' = Al			E' = Ga		
		ΔH°_{298}	ΔS°_{298}	ΔG°_{298}	ΔH°_{298}	ΔS°_{298}	ΔG°_{298}
1	IDipp-E'H ₃ + AsH ₃ = H ₂ + IDipp-E'H ₂ AsH ₂	-27.6	-26.3	-19.7	-29.2	-26.3	-21.4
2	IDipp-E'H ₃ + AsHPh ₂ = H ₂ + IDipp-E'H ₂ AsPh ₂	-11.2	-61.8	7.2	-15.7	-60.6	2.3
3	IDipp-E'H ₂ Cl + KAsH ₂ = KCl _(s) + IDipp-E'H ₂ AsH ₂	-227.7	-179.8	-174.1	-261.9	-182.8	-207.4
4	IDipp-E'H ₂ Cl + KAsPh ₂ -dioxane = KCl _(s) + dioxane + IDipp-E'H ₂ AsPh ₂	-97.2	98.6	-126.6	-134.2	96.7	-163.1
5	IDipp-E'H ₂ AsH ₂ = 1/3(E'H ₂ AsH ₂) ₃ + IDipp	65.4	76.5	42.6	52.9	75.4	30.4
6	IDipp-E'H ₂ AsPh ₂ = 1/3(E'Ph ₂ AsH ₂) ₃ + IDipp	44.5	70.3	23.6	33.8	75.1	11.4
7	IDipp-E'H ₂ AsH ₂ + AsH ₃ = H ₂ + IDipp-E'H(AsH ₂) ₂	-23.0	-43.3	-10.1	-25.4	-39.0	-13.8
8	IDipp-E'H ₂ AsH ₂ + PH ₃ = H ₂ + IDipp-E'H(PH ₂) ₂	-13.0	-38.5	-1.6	-11.9	-40.6	0.2
9	IDipp-E'HCl ₂ + 2NaPH ₂ = 2NaCl _(s) + IDipp-E'(PH ₂) ₂	-468.6	-354.8	-362.8	-536.0	-343.8	-433.5
10	IDipp-E'HCl ₂ + 2KAsH ₂ = 2KCl _(s) + IDipp-E'(AsH ₂) ₂	-461.8	-367.9	-352.1	-535.9	-352.9	-430.7

[a] Standard enthalpies ΔH°_{298} and standard Gibbs energies ΔG°_{298} in kJ mol⁻¹, standard entropies ΔS°_{298} in J mol⁻¹ K⁻¹. B3LYP/def2-TZVP level of theory.

(LIFDI-MS) the molecular ion peak of **3** is detected at m/z 535.1239 [$M-H$]⁺. The ¹H NMR spectrum of **3** in C₆D₆ shows a triplet at $\delta = -0.18$ ppm (³J_{H,H} = 3.68 Hz) for the AsH₂ moiety and a broad singlet at $\delta = 4.31$ ppm for the GaH₂ moiety. Compound **3** co-crystallizes with the starting material IDipp-GaH₂Cl (for more information see SI). The structure of **3** in solid state is shown in Figure 3. With a distance of 2.4503(12) Å the Ga-As bond length in **3** is between the Ga-As bond lengths in **1** (2.4659(5) Å), (C₅Me₅)₂GaAs(SiMe₃)₂ (2.433 Å),^[5a] and (*t*-Bu)₂GaAs(*t*-Bu)₂ (2.466 Å).^[5b] The Ga-C1 bond in **3** (2.0476(17) Å) is shorter compared to the Ga-C1 distance in **1** (2.068(3) Å) which reveals the repulsion between the NHC and the phenyl groups in **1**. Since the H substituents at the As atom had to be restrained, no statement about the conformation of **3** can be made. The C1-Ga-As angle in **3** (107.99(6)°) is slightly smaller compared to the substituted analog **1** (109.35(3)°) and to the phosphorus derivative IDipp-GaH₂PH₂ (109.19(5)°).^[7]

IDipp-AlH₂AsH₂ (**4**) can be isolated at -30 °C as colorless plates in a yield of 40% via route 1. The LIFDI-MS spectrum of **4** only shows the fragment ion peak of IDipp⁺ due to the decomposition of **4** during the ionization process. The ¹H NMR spectrum of **4** in C₆D₆ reveals a triplet at $\delta =$

-0.47 ppm (³J_{H,H} = 3.23 Hz) for the AsH₂ moiety and a broad singlet at $\delta = 4.1$ ppm for the AlH₂ moiety. In the ¹H NMR spectrum, besides **4** a side product IDipp-AlH(AsH₂)₂ (**5**) can be detected as two doublets of doublets at $\delta = -0.15$ ppm and $\delta = -0.04$ ppm, respectively, for the AsH₂ moieties (²J_{H,H} = 12.59 Hz, ³J_{H,H} = 2.80 Hz). The signals for these two AsH₂ moieties split in two separated signals because of the prochirality of the entities. The ²⁷Al NMR spectrum of **4** shows a broad signal at $\delta = 133.5$ ppm which is partly superimposed with the signal of the NMR sample head and the NMR tube material. Compound **4** (Figure 4) crystallizes in the monoclinic space group *I2/a* and co-crystallizes with IDipp-AlH(AsH₂)₂ (**5**) (for more information, see SI). The Al-As distance in **4** is in the range of 2.399(6)-2.473(4) Å. The Al-C1 bond length (2.060(2) Å) is very similar to the bond length in **1** (2.0634(12) Å) and IDipp-AlH₂PH₂ (2.056(2) Å).^[7] The C1-Al-As angle varies between 107.83(17)° and 114.3(2)° because of the disorder of the AsH₂ moiety.

The formation of IDipp-AlH(AsH₂)₂ (**5**) as a side product led us to the question if the selective synthesis of compounds of the type IDipp-E'H(AsH₂)₂ (E' = Al, Ga) was possible, and indeed we were able to synthesize **5** and IDipp-GaH(AsH₂)₂




Figure 3. Molecular structure of **3** in the solid state; thermal ellipsoids at 50% probability.^[19] Selected bond lengths [Å] and angles [°]: Ga-As 2.4503(12), Ga-C1 2.0476(17), C1-Ga-As 107.99(6).





Figure 4. Molecular structure of **4** in solid state (part 1); thermal ellipsoids at 50% probability.^[19] Selected bond lengths [Å] and angles [°]: Al-As1 2.399(6), C1-Al 2.060(2), C1-Al-As1 107.83(17)-114.3(2).

(6) via the corresponding salt metathesis route [Eq. (3)], which was supported by DFT computations (see Table 1, process 10). In fact, such branched alkane-like parent compounds are so far unknown and only additional donor stabilized compounds of the type $(\text{Dipp}_2\text{Nacnac})\text{E}'(\text{EH}_2)_2$ ($\text{Dipp}_2\text{Nacnac} = \text{HC}[\text{C}(\text{Me})\text{N}(\text{Ar})]_2$, $\text{Ar} = 2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3$) exist for $\text{E}' = \text{N}$,^[16a] P, As.^[16b]



Compounds **5** and **6** crystallize as colorless thin needles at -30°C in a yield of 42% and 36%, respectively. The LIFDI-MS spectrum of **5** shows a fragment ion peak of IDipp^+ due to decomposition of **5** during the ionization process. In the mass spectrum of **6** (LIFDI-MS) the molecular ion peak is detected at m/z 611.0607 [$M-\text{H}$]⁺. Solutions of **5** show a strong tendency towards decomposition. The ^1H NMR spectrum of **5** in $[\text{D}_8]\text{toluene}$ at -80°C reveals two doublets of doublets at $\delta = -0.09$ ppm and $\delta = 0.14$ ppm ($^2J_{\text{H,H}} = 12.40$ Hz, $^3J_{\text{H,H}} = 2.71$ Hz) for the two AsH_2 moieties, a broad singlet at $\delta = 4.82$ ppm for the AlH moiety, as well as the formation of IDippH_2 and free IDipp as decomposition products. In the ^1H NMR spectrum of **6** in C_6D_6 the signals for the AsH_2 moieties and the GaH moiety are shifted downfield to $\delta = 0.20, 0.38$ ($^2J_{\text{H,H}} = 12.77$ Hz, $^3J_{\text{H,H}} = 3.46$ Hz), and $\delta = 5.09$ ppm compared to **5**.

Compounds **5** and **6** crystallize from concentrated *n*-hexane solutions as very thin colorless plates. Because of the thinness of the crystals the single-crystal X-ray analysis of **6** was only possible to a theta range of 47° . Nevertheless, it was possible to solve the structure and prove the framework of the heavy atoms of **6** (see Figure S42). Compound **5** co-crystallizes with 6% of the starting material $\text{IDipp}\cdot\text{AlHCl}_2$ (see Figure S41). Compounds **5** and **6** crystallize in the monoclinic space group $I2/a$. The molecular structure of **5** in solid state is depicted in Figure 5. The $\text{E}'\text{-As}$ distances in **5** and **6** are in the range of 2.451(4)–2.511(6) Å (**5**) and 2.4412(19)–2.446(2) Å (**6**), respectively, and therefore similar to the Al-As bonds in $(\text{Dipp}_2\text{Nacnac})\text{Al}(\text{AsH}_2)_2$ ($\text{Dipp}_2\text{Nacnac} = \text{HC}[\text{C}(\text{Me})\text{N}(\text{Ar})]_2$, $\text{Ar} = 2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3$).^[15] The $\text{E}'\text{-C1}$ bond lengths ($\text{Al-C1} = 2.066(3)$ Å, $\text{Ga-C1} = 2.064(9)$ Å) are not heavily affected by the presence of a second AsH_2 moiety compared to **3** (2.0476(17) Å) and **4** (2.060(2) Å), respectively. The $\text{C1-E}'\text{-As}$ angles are $114.24(9)^\circ$ and $114.38(10)^\circ$ for **5** as well as $111.7(2)^\circ$ and $113.3(2)^\circ$ for **6**.^[17]

Interestingly, during the synthesis of the phosphorus analog $\text{IDipp}\cdot\text{E}'\text{H}_2\text{PH}_2$ ($\text{E}' = \text{Al}, \text{Ga}$) by the reaction of $\text{IDipp}\cdot\text{E}'\text{H}_2\text{Cl}$ with NaPH_2 we did not find any sign for the formation of $\text{IDipp}\cdot\text{E}'\text{H}(\text{PH}_2)_2$ ($\text{E}' = \text{Al}, \text{Ga}$) as a side product.^[7] A possible pathway for the formation of **5** as a side product in the arsenic case is the reaction of the formed product $\text{IDipp}\cdot\text{E}'\text{H}_2\text{AsH}_2$ with in situ formed AsH_3 in an H_2 -elimination reaction. Computations confirm that this route is




Figure 5. Molecular structure of **5** in solid state; thermal ellipsoids at 50% probability.^[19] Selected bond lengths [Å] and angles [°]: Al-As1 2.451(4), Al-As2 2.474(3), Al-C1 2.066(3), As1-Al-C1 114.38(10), As2-Al-C1 114.24(9).

possible in the arsenic case (Table 1, process 7) while it is more unlikely for phosphorus (Table 1, process 8), which agrees with our experimental observations.

Similar to **5** and **6**, we were able to synthesize the parent branched compounds $\text{IDipp}\cdot\text{GaH}(\text{PH}_2)_2$ (**7**) and $\text{IDipp}\cdot\text{AlH}(\text{PH}_2)_2$ (**8**) selectively by the salt metathesis reaction of $\text{IDipp}\cdot\text{E}'\text{HCl}_2$ and NaPH_2 in Et_2O (Table 1, process 9). Compounds **7** and **8** can be isolated at -30°C in a yield of 57% and 48%, respectively. The ^1H NMR spectrum of **7** in C_6D_6 shows a doublet which splits into multiplets at $\delta = 0.54$ ppm ($^1J_{\text{PH}} = 175$ Hz) for the PH_2 moieties and a broad singlet at $\delta = 4.81$ ppm for the GaH moiety. In the ^1H NMR spectrum of **8** in $[\text{D}_8]\text{toluene}$ at -80°C the PH_2 moieties can be detected at $\delta = 0.42$ ppm ($^1J_{\text{PH}} = 175.4$ Hz) as a doublet of multiplets. The AlH moiety can be detected as a broad singlet at $\delta = 4.56$ ppm. The ^{31}P NMR spectra of **7** and **8** show a triplet of multiplets at $\delta = -255.4$ ppm (**7**, $^1J_{\text{PH}} = 175$ Hz, $^2J_{\text{PH}} = 18.17$ Hz) and at $\delta = -270.8$ ppm (**8**, $^1J_{\text{PH}} = 175.4$ Hz, $^2J_{\text{PH}} = 15.48$ Hz), respectively. Due to the prochirality of the PH_2 groups in **7** and **8** the signals in the ^1H and ^{31}P NMR spectra reveal a fine splitting which could not be resolved. Like **5**, solutions of **8** show a strong tendency towards decomposition. Compounds **7** and **8** crystallize in the monoclinic space group $I2/a$. The molecular structures of **7** and **8** in solid state are shown in Figure 6 and Figure S44, respectively. The $\text{E}'\text{-P}$ bonds are shorter compared to the arsenic analogs with 2.3437(10)–2.3574(9) Å (**7**) and 2.3075(10)–2.3418(9) Å (**8**). The $\text{E}'\text{-C1}$ bond lengths are again not affected by the change from arsenic substituents to phosphorus substituents on the E' atom. The Ga-C1 bond length is 2.075(3) Å and the Al-C1 bond length is 2.066(2) Å. The $\text{C1-E}'\text{-P}$ angles ($112.38(7)^\circ$ and $113.68(7)^\circ$ for **7**; $112.04(6)^\circ$ and $113.91(6)^\circ$ for **8**) are comparable to the $\text{C1-E}'\text{-As}$ angles in the arsenic analogs **5** and **6**.

Computational studies indicate that the salt elimination route via solid potassium chloride formation is highly exothermic and exergonic both for the parent and the substituted compounds, which could be experimentally verified by the synthesis of **1–4**. (Table 1, process 3 and 4). The hydrogen elimination route via the reaction of $\text{IDipp}\cdot\text{E}'\text{H}_3$




Figure 6. Molecular structure of **7** in solid state; thermal ellipsoids at 50% probability.^[19] Selected bond lengths [Å] and angles [°]: Ga-P1 2.3574(9), Ga-P2 2.3437(10), Ga-C1 2.075(3), P1-Ga-C1 113.68(7), P2-Ga-C1 112.38(7).

with AsH₃ (Table 1, process 1) is exothermic and at 298 K exergonic by about 20 kJ mol⁻¹, but slightly endergonic (2–7 kJ mol⁻¹) for the reaction with diphenylarsine (Table 1, process 2), which reflects that compounds **1** and **2** could not be accessed via route 2. Compounds **1–4** are predicted to be stable with respect to IDipp dissociation with formation of (E'H₂AsH₂)_n polymers, which were modeled by the formation of the trimer^[18] (Table 1, process 5 and 6). The interaction of IDipp-E'H₂AsH₂ with an arsine formed in situ (Table 1, process 7) is also exergonic (Al: –10.1 kJ mol⁻¹, Ga: –13.8 kJ mol⁻¹) and may explain the formation of **5** as a side product during the synthesis of IDipp-AlH₂AsH₂ via route 1. In contrast, a similar reaction for the phosphorus analogs (Table 1, process 8) is energetically less favored and has Gibbs energies close to zero at 298 K. Nevertheless, computations show that route 1 is an even more exergonic reaction for the synthesis of branched pnictogenylalanes and -gallanes than for the synthesis of the linear compounds (Table 1, process 9 and 10). This is confirmed by the synthesis of the unique molecules IDipp-E'H(AsH₂)₂ (**5**: Al, **6**: Ga) and IDipp-E'H(PH₂)₂ (**7**: Ga, **8**: Al) via route 1.

Conclusion

The results show that, regardless of the rather low E'–As bond stability (E' = Al, Ga), we succeeded in the synthesis of the first monomeric parent arsanylalanes and -gallanes stabilized only by a LB. Besides the synthesis of the organo-substituted arsenic derivatives by salt metathesis, it was shown that the monomeric parent compounds can be obtained by salt metathesis and H₂-eliminations, respectively. However, the latter method is incomplete, so that the first one is preferred. Furthermore, in contrast to the synthesis of the corresponding phosphanylalanes and -gallanes, the As derivatives exhibit a different reactivity and form the branched side products IDipp-E'H(AsH₂)₂ (E' = Al, Ga), obviously by AsH₃-caused substitution reactions. This kind of alkane-like branched parent derivatives had been unknown before and subsequently the double substituted parent compounds

IDipp-E'H(EH₂)₂ (E' = Al, Ga; E = As, P) could be selectively synthesized by salt metathesis reactions. They may serve as chelating ligands in coordination chemistry, which is currently being investigated. The monomeric compounds IDipp-E'H₂AsH₂ (E' = Al, Ga) represent unprecedented parent arsanylalanes and -gallanes without any prior sterical stabilization by a substituent but by a LB. In further studies, the focus will be on their reaction behavior towards cationation and as precursor for CVD processes to obtain Group 13/15 materials.

Acknowledgements

The authors thank the German Research Council (DFG) for comprehensive support in the project Sche 384/35-1. A.Y.T. is grateful to the SPSU grant 12.65.44.2017. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Keywords: alanes · arsenic · gallanes · group 13/15 compounds · Lewis bases

- [1] a) J. D. Masuda, A. J. Hoshkin, T. W. Graham, C. Beddic, M. C. Fermin, N. Etkin, D. W. Stephan, *Chem. Eur. J.* **2006**, *12*, 8696–8707; b) R. A. Fischer, J. Weiß, *Angew. Chem. Int. Ed.* **1999**, *38*, 2830–2850; *Angew. Chem.* **1999**, *111*, 3002–3022; c) R. L. Wells, W. L. Gladfelter, *J. Cluster Sci.* **1997**, *8*, 217–238; d) A. C. Jones, P. O'Brien, in *CVD of Compound Semiconductors: Precursor Synthesis Development and Applications*, VCH, Weinheim, **1996**.
- [2] a) S. Schulz, *Coord. Chem. Rev.* **2001**, *215*, 1–37; b) S. Schulz, *Adv. Organomet. Chem.* **2003**, *49*, 225–317; c) B. Neumüller, E. Irvani, *Coord. Chem. Rev.* **2004**, *248*, 817–834; d) A. Y. Timoshkin, *Coord. Chem. Rev.* **2005**, *249*, 2094–2131; e) T. J. Clark, K. Lee, I. Manners, *Chem. Eur. J.* **2006**, *12*, 8634–8648; f) A. Staubitz, A. P. Soto, I. Manners, *Angew. Chem. Int. Ed.* **2008**, *47*, 6212–6215; *Angew. Chem.* **2008**, *120*, 6308–6311.
- [3] a) G. E. Coates, J. Graham, *J. Chem. Soc.* **1963**, 233–237; b) D. E. Heaton, R. A. Jones, K. B. Kidd, A. H. Cowley, C. M. Nunn, *Polyhedron* **1988**, *7*, 1901–1908; c) R. L. Wells, A. T. McPhail, T. M. Speer, *Organometallics* **1992**, *11*, 960–963.
- [4] O. T. Beachley, G. E. Coates, *J. Chem. Soc.* **1965**, 3241–3247.
- [5] a) E. K. Byrne, L. Parkanyi, K. H. Theopold, *Science* **1988**, *241*, 332–334; b) K. T. Higa, C. George, *Organometallics* **1990**, *9*, 275–277; c) D. A. Atwood, L. Contreras, A. H. Cowley, R. A. Jones, M. A. Mardones, *Organometallics* **1993**, *12*, 17–18.
- [6] a) M. R. Leys, *Chemtronics* **1987**, *2*, 155–164; b) G. B. Stringfellow, *Rep. Prog. Phys.* **1982**, *45*, 469–525.
- [7] M. A. K. Weinhart, A. S. Lisovenko, A. Y. Timoshkin, M. Scheer, *Angew. Chem. Int. Ed.* **2020**, *59*, 5541–5545; *Angew. Chem.* **2020**, *132*, 5586–5590.
- [8] A. Y. Timoshkin, *Phosphorus Sulfur Silicon Relat. Elem.* **2001**, *168*, 275–280.
- [9] M. Brynda, *Coord. Chem. Rev.* **2005**, *249*, 2013–2034.
- [10] a) C. Marquardt, A. Adolf, A. Stauber, M. Bodensteiner, A. V. Virovets, A. Y. Timoshkin, M. Scheer, *Chem. Eur. J.* **2013**, *19*, 11887–11891; b) U. Vogel, P. Hoemensch, K.-Ch. Schwan, A. Y. Timoshkin, M. Scheer, *Chem. Eur. J.* **2003**, *9*, 515–519; c) U.

- Vogel, A. Y. Timoshkin, M. Scheer, *Angew. Chem. Int. Ed.* **2001**, *40*, 4409–4412; *Angew. Chem.* **2001**, *113*, 4541–4544.
- [11] A. Hock, L. Werner, C. Luis, U. Radius, *Dalton Trans.* **2020**, *49*, 11108–11119.
- [12] K. Knabel, I. Krossing, H. Nöth, H. Schwenk-Kircher, M. Schmidt-Amelunxen, T. Seifert, *Eur. J. Inorg. Chem.* **1998**, 1095–1114.
- [13] P. Pyykkö, M. Atsumi, *Chem. Eur. J.* **2009**, *15*, 186–197.
- [14] C. G. Pitt, K. T. Higa, A. T. McPhail, R. L. Wells, *Inorg. Chem.* **1986**, *25*, 2483–2484.
- [15] A. M. Arif, B. L. Benac, A. H. Cowley, R. Geerts, R. A. Jones, K. B. Kidd, J. M. Power, S. T. Schwab, *J. Chem. Soc. Chem. Commun.* **1986**, 1543–1545.
- [16] a) V. Jancik, L. W. Pineda, J. Pinkas, H. W. Roesky, D. Neculai, A. M. Neculai, R. Herbst-Irmer, *Angew. Chem. Int. Ed.* **2004**, *43*, 2142–2145; *Angew. Chem.* **2004**, *116*, 2194–2197; b) B. Li, S. Bauer, M. Seidl, A. Y. Timoshkin, M. Scheer, *Chem. Eur. J.* **2019**, *25*, 13714–13718.
- [17] Due to the limited quality of the X-ray structure analysis the bond lengths of **6** might deviate a bit.
- [18] C. Marquardt, O. Hegen, A. Vogel, A. Stauber, M. Bodensteiner, A. Y. Timoshkin, M. Scheer, *Chem. Eur. J.* **2018**, *24*, 360–363.
- [19] Deposition Numbers 2035397 (for **1**), 2035398 (for **2**), 2035399 (for **3**), 2035340 (for **4**), 2035341 (for **5**), 2035342 (for **6**), 2035343 (for **7**), and 2035344 (for **8**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service www.ccdc.cam.ac.uk/structures.

Manuscript received: October 14, 2020

Revised manuscript received: November 16, 2020

Accepted manuscript online: November 16, 2020

Version of record online: December 15, 2020