

POSTER PRESENTATION

Open Access

INCF plasmids responsible by dissemination of *bla*_{KPC} gene among enterobacteriaceae

F Calisto^{1*}, L Lito², J Melo-Cristino², A Duarte¹

From International Conference on Prevention & Infection Control (ICPIC 2011)
Geneva, Switzerland. 29 June – 2 July 2011

Introduction / objectives

In recent years, carbapenem resistance has emerged among *Enterobacteriaceae* in many geographical locations due to *Klebsiella pneumoniae* producing KPC carbapenemase. The aim of this study was to characterize the genetic elements involved in *bla*_{KPC} gene mobilization and diffusion.

Methods

A total of 22 isolates, *K. pneumoniae* (n=20), *Escherichia coli* (n=1) and *Enterobacter aerogenes* (n=1), were collected from a university hospital, in Lisbon, Portugal. MIC were determined by Etest and interpreted according to Clinical and Laboratory Standards Institute guidelines. PCR was performed with primers designed to amplify *bla*_{SHV}, *bla*_{TEM} and *bla*_{KPC} genes. Molecular typing was performed by M13-PCR fingerprinting and in representative strains by Multilocus Sequence Typing (MLST). Replicon typing was used to define plasmid incompatibility groups (Inc).

Results

Twenty-two isolates had MIC ranging between 2 and 32 mg/L for imipenem and meropenem. All isolates encoding KPC-3. Associated to carbapenemase 7 isolates encoded for TEM-1 or SHV-1 and 6 isolates for both TEM-1 and SHV-type. Although M13-PCR fingerprint analysis showed four predominant profiles, but with MLST only one sequence type ST11 was found among *K. pneumoniae* isolates. The predominant plasmid was included in IncF (90.5%) and was found plasmids belong to different replicons FIC (52.4%), HI1, HI2, and I1-Iy (47.8%), T (42.8%), FIAs (33.3%), W and P (32.8%) and FIB (9.5%).

Conclusion

The *bla*_{KPC} gene, often associated with other β-lactamases, was transferred between *K. pneumoniae*, *E. aerogenes* and *E. coli* strains by IncF group plasmid. The clustering of resistance genes in plasmids and their organization with regard to cotransfer underlines the importance of these plasmids in the spread of antimicrobial multiresistance.

Disclosure of interest

None declared.

Author details

¹i-Med, Faculdade de Farmácia, CHLN, EPE Lisboa, Lisbon, Portugal.

²Laboratory of Microbiology, CHLN, EPE Lisboa, Lisbon, Portugal.

Published: 29 June 2011

doi:10.1186/1753-6561-5-S6-P132

Cite this article as: Calisto et al.: INCF plasmids responsible by dissemination of *bla*_{KPC} gene among enterobacteriaceae. *BMC Proceedings* 2011 **5**(Suppl 6):P132.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹i-Med, Faculdade de Farmácia, CHLN, EPE Lisboa, Lisbon, Portugal
Full list of author information is available at the end of the article