



# Child alcohol use disorder in Eastern Uganda: screening, diagnostics, risk factors and management of children drinking alcohol in Uganda (TREAT C-AUD): a mixed-methods research protocol

Vilde Skylstad <sup>1</sup>, Harriet Aber,<sup>2</sup> Victoria Bakken,<sup>3</sup> Jutta Dierkes,<sup>4,5</sup> Silje Akselberg Iversen,<sup>6</sup> Esther Kisaakye,<sup>2</sup> Melf-Jakob Kühl,<sup>1</sup> Joyce Sserunjogi Nalugya,<sup>7,8</sup> Divya Rayamajhi,<sup>6</sup> Deogratious Sebuwufu,<sup>2</sup> Ane-Marthe Solheim Skar,<sup>1,9</sup> Norbert Skokauskas,<sup>3</sup> Neda Valeckaite,<sup>1</sup> Henry Wamani,<sup>2</sup> Ingunn Marie S. Engebretsen <sup>1</sup>, Juliet Ndimwibo Babirye,<sup>2</sup> The TREAT C-AUD consortium

**To cite:** Skylstad V, Aber H, Bakken V, *et al.* Child alcohol use disorder in Eastern Uganda: screening, diagnostics, risk factors and management of children drinking alcohol in Uganda (TREAT C-AUD): a mixed-methods research protocol. *BMJ Paediatrics Open* 2021;**5**:e001214. doi:10.1136/bmjpo-2021-001214

VS and HA are joint first authors.  
IMSE and JNB are joint senior authors.

Received 1 July 2021  
Accepted 13 July 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**  
Dr Vilde Skylstad; Vilde.skylstad@uib.no

## ABSTRACT

**Background** Following a finding of alcohol use among children aged 5–8 years old in Mbale, Uganda, this project investigates the magnitude of alcohol and substance use among children aged 6–13 years old and related household, community, school, health system and clinical factors.

**Methods** The project includes four larger work packages (WPs). WP1 comprises management, WP2 and 3 include the scientific components and WP4 includes integration of results, dissemination, policy and implementation advice. This protocol presents the planned research work in WP 2 and 3. WP2 comprises the adaptation and validation of the alcohol use screening tool Car-Relax-Alone-Forget-Family and Friends-Trouble (CRAFT) to the age group and setting. WP3 comprises four substudies (SS). SS1 is a cross-sectional community household survey with an estimated sample size of 3500 children aged 6–13 years and their caregivers. We apply cluster sampling and systematic sampling within the clusters. Data collection includes a structured questionnaire for caregiver and child, measuring social and demographic factors, mental health status, alcohol and substance use, nutrition history and anthropometry. Urine samples from children will be collected to measure ethyl glucuronide (EtG), a biological marker of alcohol intake. Further, facilitators, barriers and response mechanisms in the health system (SS2) and the school system (SS3) is explored with surveys and qualitative assessments. SS4 includes qualitative interviews with children. Analysis will apply descriptive statistics for the primary outcome of establishing the magnitude of alcohol drinking and substance use, and associated factors will be assessed using appropriate regression models. The substudies will be analysed independently, as well as inform each other through mixed methods strategies at the stages of design, analysis, and dissemination.

## What is already known on this topic?

- ▶ Uganda has a relatively high consumption of alcohol, estimated to 12.2 (10.1–14.6) litres of pure alcohol, according to the 2019 Global Health Observatory data.
- ▶ Any level of alcohol consumption, regardless of the amount, leads to loss of healthy life and children are at greatest risk of harm.
- ▶ In Uganda, regional and demographic patterns vary, and for some alcohol production and consumption is widely practised part of their culture.

## What this study hopes to add?

- ▶ This research aims to describe the prevalence of alcohol consumption and substance use among children aged 6–13 years old in Mbale District, Eastern Uganda.
- ▶ Substudies investigate the health and school system's approaches to management of children that have challenges with substance use.
- ▶ The study explores risk and protective factors for childhood alcohol use to inform screening, policy formulation and implementation of prevention and intervention programmes.

**Ethics and dissemination** Data protection and ethical approvals have been obtained in Uganda and Norway, and referral procedures developed. Dissemination comprises peer-reviewed, open access research papers, policy recommendations and intersectoral dialogues. Trial registration number Clinicaltrials.gov 29.10.2020 (#NCT04743024).

## INTRODUCTION

Harmful use of alcohol is an important public health issue.<sup>1</sup> Limited attention has been given to preadolescence or childhood alcohol drinking,<sup>2</sup> and existing research mainly come from high income countries.<sup>2</sup> In sub-Saharan Africa, there has been limited research about child mental health in general, and alcohol and substance use in particular.<sup>3</sup> However, some reports on early substance use do exist. In 2018, a study with data from the Global School-Based Student Health Survey (GSHS) showed that among participants in African countries, 13.4%–15.5% reported alcohol intake before age 11.<sup>4</sup> The most recent Ugandan GSHS from 2003 reported that 15.2% of 13–15 years had been ‘really drunk’ and 8.5% had used other drugs.<sup>5</sup> These high rates of adolescent drinking suggest an early initiation of alcohol use, although more recent data are lacking.

Early-onset alcohol and substance use is an important predictor for developing alcohol and substance use disorders and mental health conditions later in life.<sup>6</sup> Neurological, psychological and social child development<sup>7</sup> may be disturbed by alcohol and substance use. It is also associated with a variety of negative mental and somatic health outcomes in addition to social risky behaviour including violence, crime and school drop-out.<sup>2,8</sup> If drinking starts before the age of 10 years, structures in the child’s developing brain involved in functions sentinel for social and academic achievements such as planning, learning, attention, appetite, reward and emotional and behavioural control may be affected.<sup>7</sup>

Uganda is a low-income country in East Africa with more than 42 million inhabitants, equivalent to almost 180 people per square km, ranking among the top ten most densely populated African countries. Almost half the population is younger than 15 years of age. It ranks 159/189 on the human development index report from 2020 and has an average of 6 years of formal schooling.<sup>9</sup> Like many other sub-Saharan countries, Uganda has a comparably high alcohol consumption per capita,<sup>10</sup> and traditions of home brewing are common.<sup>11</sup> In 2014, we identified a prevalence of 8.4% clinically defined harmful alcohol use or dependence in a sample of children aged 5–8 years (n=119) with a high mental health symptom load.<sup>12</sup> This prompted further qualitative investigation, which corroborated our finding.

The research project: TREAT Child Alcohol Use Disorder (C-AUD) in Eastern Uganda: screening, diagnostics, risk factors and management of children drinking alcohol, funded by the Norwegian Research Council (TREAT C-AUD) (#285489), is a 5-year project initiated in 2019 to answer questions regarding alcohol and other substance use in childhood. ‘Alcohol use disorder’ (AUD) is a diagnostic term from the Diagnostic and Statistical Manual of Mental Disorders (DSM–5), classified into mild, moderate and severe disorder. Our project has adapted this term to childhood and preadolescence and constructed the term C-AUD. In our study, we will investigate both the prevalence and associated factors of childhood substance and alcohol use (beyond sipping and tasting), as well as evaluate screening and diagnostic criteria for the term C-AUD. The project’s objectives are to investigate the magnitude of substance use, alcohol use and C-AUD in children, and related household factors, its management in the school and health sector and improve contextual understanding, and to engage in the public health dialogue and policy recommendations in Uganda.

## METHODS

### Site description

The study site is Mbale District in Eastern Uganda, 245 km from the capital Kampala. Situated close to the Kenyan border, Mbale City is the main urban trading area in the district, which is predominantly rural and dominated by peasant farming. The staple crops include millet and maize grain which are also commonly used for home brewing. The urban area has a regional referral hospital with a psychiatric department.

### Study design

The TREAT C-AUD project will use both qualitative and quantitative methods (table 1). The project includes four larger work packages (WPs) (figure 1). WP1 comprises management, WP2 and 3 include the scientific components and WP4 includes integration of results, dissemination, policy and implementation advice. This protocol presents the planned research work in WP 2 and 3. WP2 comprises the adaptation and validation of a screening tool for childhood alcohol and substance use. WP3 has

**Table 1** The work packages and substudies in TREAT C-AUD

Work package 2	Work package 3			
Adapt and validate the Car-Relax-Alone-Forget-Family and Friends-Trouble screening tool to the age group and setting	Substudy 1: Community household survey Cross sectional cluster survey with children and caregivers	Substudy 2: Health system study Survey and qualitative data collection on C-AUD and health system response	Substudy 3: School study Survey and qualitative data collection on school response to childhood alcohol and substance	Substudy 4: Contextual understanding Qualitative assessments of children to understand the context-specific factors to aid interpretation

C-AUD, Child Alcohol Use Disorder.

WP1: Coordination and management	
WP 2: Tool development - CRAFFT <ul style="list-style-type: none"> <li>Expert consultations for content and language</li> <li>Qualitative study with children</li> <li>Validation study</li> </ul>	WP 3: Data collection and analysis <ul style="list-style-type: none"> <li>SS 1: Community survey with 3500 households</li> <li>SS 2: Health system study with survey and qualitative assessments</li> <li>SS 3: School study with survey and qualitative assessments</li> <li>SS4: Child perspectives using qualitative assessments</li> </ul>
WP 4: Dissemination, implementation, and policy consultations	

**Figure 1** TREAT C-AUD Protocol: work packages (WP) and substudies (SS)\*. \*The mixed-methods diagram illustrates how we apply interlinked qualitative and quantitative research methods in the design and the analysis of the TREAT C-AUD research project. C-AUD, Child Alcohol Use Disorder; CRAFFT, Car-Relax-Along-Forget-Family and Friends-Trouble.


four substudies, and it includes quantitative and qualitative data collection methods on epidemiological and contextual data from the community households, health system, school system and children's perspectives.

Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the 'Patient and Public Involvement' section for further details.

Separate data collection from the substudies will be analysed independently, as well as inform each other through mixed-methods strategies at the stages of design, analysis and dissemination. There are many ways to mix methods and procedures at the design, analysis and interpretation stages.<sup>13</sup> Thus, according to Zhang and Creswell, 2013, our project qualifies as a mixed methods project having both concurrent and sequential elements of mixing qualitative and quantitative methods at the design, conduct and interpretation stages.

### WP2: screening tool adaption and validation

We will adapt the Car-Relax-Along-Forget-Family and Friends-Trouble (CRAFFT) screening tool to the age group and setting by multidisciplinary committee review, as recommended by WHO.<sup>14</sup> This includes forward and backward translations of tools by bilingual independent translators; rereview of translated versions and conciliation with authors; pilot testing with qualitative assessment and quantitative reliability and validity assessment. We will conduct psychiatric interviews using the MINI International Neuropsychiatric Interview for Children and Adolescents (MINI-KID)<sup>15</sup> diagnostic tool (adapted for DSM-5 based on the English V.7.0.2)<sup>16</sup> as the gold standard for validation. We will evaluate the use of the screening tool and diagnostic criteria for C-AUD in synergy with the WP3 health systems component, where we investigate the clinical practice related to substance and alcohol use, and possible C-AUD in children.



**Figure 2** Conceptual framework of the interplay between C-AUD, risk factors and outcomes. C-AUD, Child Alcohol Use Disorder.

### WP3: community household, health system, school system and children's perspectives

WP3 builds on a conceptual framework (figure 2) where the development of C-AUD and its effect is influenced by factors related to the child, caregiver and household. The WP includes four substudies: a community household survey, a health system study, a school study and qualitative studies of children's perspectives. The health system and school studies use mixed methodologies.

#### Community household survey

A two-stage stratified cluster survey as recommended by the 2018 WHO coverage survey reference manual will be used<sup>17</sup> since the district does not have a registration system for the household or household members.

A sample size of 3500 children is estimated with the following assumptions: prevalence of alcohol consumption of 4%, a precision of 3.6%, a 95% confidence level and a design effect (DEFF) of 2.<sup>18</sup> The number of households to visit is estimated at 2982, computed from the effective sample size multiplied by 'DEFF', the proportion of households with eligible children and the nonresponse rate (10%) as demonstrated in table 2. Non-response is considered if caregivers are absent from the household, refusal to provide consent to study participation, uninhabited households among other reasons.

A two-stage sampling procedure will be used. In the first stage, 99 clusters (villages) will be selected based on probability proportionate to population size. The second stage of selection will be based on the household listing, which shall be done by the research teams in the selected clusters. Serial numbers shall be allocated for unique identification for each household along with their respective GPS coordinates. It is from this stage that systematic sampling without replacement shall be done using Open Data Kit app (ODK) (<https://opendatakit>).

**Table 2** Sample size estimates for the target group (6–13 years), TREAT C-AUD study

Total no of households (NHH) with target population in the district, NHH to visit	Target no of households to be visited in the district	No of clusters needed per stratum (N clusters per stratum)	Total households to visit per cluster	Total no of clusters in the survey
5963	2982	99	30	199

C-AUD, Child Alcohol Use Disorder.

org/) predesigned template for the selection of the 30 households in each cluster. The starting point for each cluster will be randomly selected by the ODK. All selected households will then be visited irrespective of having an eligible child. Household visits will be attempted on three consecutive visits to the household on three separate days before the attempted interview is labelled void. The WHO reference manual for Cluster Surveys<sup>17</sup> recommends interviewing every eligible child in the household which we will adhere to. Survey data will be collected using the ODK platform.

The survey questionnaire includes several scales, as well as questions designed by the TREAT C-AUD research team. We address mental health, somatic health and social factors associated with the main objective: alcohol and substance use in children. A schematic list of the tools is provided in table 3. We shall assess caregiver mental health status using the WHO Self-Reporting Questionnaire-20,<sup>19</sup> which has been validated in Uganda.<sup>20</sup> Children's mental health status will be measured by child and caregiver response of the Paediatric Symptom Checklist-35,<sup>21</sup> which has been validated in Batswana.<sup>22</sup> We measure caregiver traumatic experiences with the Stressful Life Events Screening Questionnaire,<sup>23</sup> using the revised version developed to improve cross cultural validity.<sup>24</sup> Traumatic experiences in the children are assessed with the Child and Adolescent Trauma Screen, with good psychometric properties in international validation,<sup>25</sup> and previous use in Uganda.<sup>26</sup> For both child and caregiver, we assess post-traumatic stress symptoms using a 2-item abbreviated version of the PTSD Checklist-Civilian version.<sup>27</sup> The Alabama Parenting Questionnaire (APQ) has a parent and a child form, and measures parenting practices.<sup>28</sup> We will use a short version (APQ-15), which has been used in cross cultural settings and has been validated for the age group.<sup>29 30</sup> Both the child and caregiver will answer the Conflict Tactics Scale-Parent-Child version,<sup>31</sup> which has been validated for the age range 6–13 years<sup>32</sup> and has demonstrated good cross-cultural reliability and validity, including in East Africa.<sup>33</sup> We assess caregiver alcohol use with the WHO Alcohol Use Disorders Identification Test (AUDIT), which has been cross-culturally validated.<sup>34</sup> To assess child alcohol and substance use, we include the adapted CRAFFT screening tool. Further, we have included questions from the GSHS<sup>35</sup> to explore the extent and consequences of alcohol and substance use, and questions from the US National Survey of Parents and Youth to explore caregiver–child communication about substance use,<sup>36</sup> and the National Longitudinal

Study of Adolescent to Adult Health (Add Health study) to explore peer-use.<sup>37</sup> Disability will be assessed using the Washington Group/UNICEF Child Functioning Module.<sup>38</sup> Child nutrition status will be assessed by asking the caregiver about food diversity, frequency and food security.<sup>39</sup> We will take anthropometric measurements and collect urine from children to measure Ethylglucuronate, which is a metabolite of ethanol, reflecting recent alcohol consumption.<sup>40</sup>

Analysis will include standard descriptive statistics, using frequencies and percentages for categorical variables, and medians with ranges and means with 95% CIs for continuous variables. To establish the magnitude of alcohol drinking and substance use among all children in the survey, prevalence will be given as cluster-adjusted percentage. A positive screen on the CRAFFT tool is defined as a score  $\geq 1$  on the 6 CRAFFT items. Age group (6–9 and 10–13 years old) and sex stratified prevalence will be provided. Assessments of associated risk factors will include multiple dimensions of care and parenting, socio-economic status and education, and mental health symptoms of the child and carer.

#### Health system study

In the health system study, we adopt the WHO framework for health systems service delivery to explore the readiness of the health system in Mbale district to screen, diagnose and manage children with C-AUD. We apply a mixed-methods approach, including a review of patient and government records, as well as key informant interviews with health workers and caregivers of children aged 6–13 years. We will explore five main domains; service delivery, human resources, health financing, pharmaceuticals and other medical products, health management information system (HMIS). In broad terms, we seek to investigate the referral pathways, estimate the proportion of detected and treated C-AUD in the health system, assess the health system capacity and competence for managing C-AUD, explore the functionality of the Ugandan HMIS for C-AUD and evaluate existing health financing mechanisms for prevention and treatment of C-AUD. An item topic guide for each domain has been developed and is given in table 4. Data collection for these objectives is yet to commence.

#### School study

In the school study, we aim to understand how childhood alcohol and substance use is viewed and managed within the education sector, including the degree of awareness,



**Table 3** List of tools and measures in the community household survey in the TREAT C-AUD project

Theme	Questionnaire to caretaker	Questionnaire to child
<b>Sociodemography</b>	Contextualised questions	
<b>Mental health and psychometrics</b>		
Mental health (caregiver)	WHO self-report questionnaire-20	–
Exposure to potentially traumatising events	Stressful Life Events Screening Questionnaire	Child and Adolescent Trauma Screen
Post-traumatic stress symptoms	2-item PTSD Checklist-Civilian version (PCL-C)	PCL-C
Alcohol and substance use	AUDIT	CRAFFT
Further investigation of alcohol and substance use	Contextualised questions	<ul style="list-style-type: none"> <li>▶ Sections from the Global School Based Health Survey</li> <li>▶ Items from on parent discussion of substances from the U.S National Survey of Parents and Youth</li> <li>▶ Items on peer use from the Add Health-study</li> </ul>
Health system use and health seeking	Contextualised questions	–
Parenting	Alabama Parenting Questionnaire-15	Alabama Parenting Questionnaire-15
Violence	Conflict Tactics Scale-Parent-Child version	Conflict Tactic Scale-Parent-Child version
General mental health (child)	Paediatric Symptom Checklist 35-Parent Report	Paediatric Symptom Checklist 35-Youth Report
<b>Nutrition</b>	Food diversity (child)	Contextualised questions
	24-hour food frequency (child)	–
	Food security	–
<b>Somatic history</b>		
General (parent and child)	HIV, chronic diseases, recent symptoms and common illnesses	–
Developmental (child)	UNICEF- Module on Child Functioning and Disability	–
Injuries (child)	Contextualised questions	–
<b>Anthropometry (child)</b>		
		Height
		Weight
		Mid upper arm circumference
		Head circumference
<b>Urine sample (child)</b>		u-STIX
		Ethyl glucuronide (EtG)

AUDIT, Alcohol Use Disorders Identification Test ; CRAFFT, Car-Relax-Along-Forget-Family and Friends-Trouble.

formal and informal response mechanisms, and the schools' interaction with the health sector.

First, we will conduct a survey among teachers of 148 public and private primary schools, representative for Mbale district. We will investigate their awareness of childhood alcohol and substance and their use of formal and informal response mechanisms. Moreover, we aim to estimate to what degree teachers understand childhood substance and alcohol use, herein C-AUD, as a developmental challenge that requires their response. These

data will be analysed using both descriptive and analytical statistics.

Second, two qualitative components of the school study will explore the perceptions on childhood alcohol and substance among school heads and school health personnel from the health sector. This inquiry includes a section on current bottlenecks and untapped response potentials between the sectors. These components will be recoded, transcribed, and analysed using framework analysis. Further, we aim to triangulate these results

**Table 4** Item topic guide for the health system substudy

Service delivery:	What are the referral pathways for C-AUD? What C-AUD screening and treatment criteria is available? Is this screening and treatment for C-AUD according to guidelines? How can absence or interruptions in these services be avoided? How can adverse outcomes at screening and treatment be monitored?
Human resource for health:	What is the capacity in terms of human and material resource of selected health facilities to manage C-AUD? What are the skills of health workers to correctly diagnose and treat C-AUD? Can an education package be provided as on-job training locally to increase the capacity of the health workers?
Health financing:	How much does it cost to screen and treat C-AUD or other mental disorders? What is the source of funding and advocacy efforts to secure funding to meet finance gaps? What are the cost of equipment, training, space and human resources for the C-AUD management? How can additional and sustainable financing be invested in these areas? How do we measure the allocative and technical efficiency of these investments?
Pharmaceuticals and other medical products:	What is the level of access to drugs? Are there medical products for treatment of C-AUD? How can the supply-chain management for the drugs be improved?
Health Management Information System (HMIS):	What data needs to be collected in the HMIS for tracking C-AUD? How is data captured in the registers or other records? Examples of process indicators are: percentage of children 6–13 years meeting the screening criteria for C-AUD after examination; percentage of children 6–13 years meeting the treatment criteria who receive the appropriate intervention; and percentage of children 6–13 years needing treatment of C-AUD who are treated within 48 hours of the decision to treat.

C-AUD, Child Alcohol Use Disorder; HMIS, Health management information system.

against the quantitative results to obtain a more robust and integrated picture of their views and existing school-based response mechanisms. We also probe for suggestions on better management of childhood alcohol and substance in schools and between the sectors. The tools for data collection are not yet pretested and data collection is scheduled for 2022.

### Children's perspectives

We will conduct qualitative interviews with children to gain a more context specific and in depth understanding of childhood alcohol and substance. Children will be involved at various stages during the project period. They will be asked to express their thoughts during qualitative interviews on general questions related to what alcohol and substance use are, what they are taught about it and who they rely on if they need it. We aim to understand children's understanding and perceptions on the topic. Further, they will be interviewed to give their views on the various elements of the adapted CRAFFT tool. Both qualitative interviews and focus group discussions will be used.

### Analysis, interpretation and integration

Quantitative descriptive analysis will include proportions, prevalence estimates and 95% CIs. The cluster design will be adjusted for, and sampling weights considered. The cross-sectional study designs have an inbuilt dilemma with reverse causalities, and only associated factors can be given. However, we plan for the binary substance use outcome to assess in logistical analysis, which factors that have a higher odd of being associated with the practice.

Qualitative analysis will include content analysis. All substudies will be analysed and presented separately, and

integration of results will only happen after each substudy is individually analysed and interpreted. Mixing of results will happen at the level of synthesis, dissemination and policy advice.

### Patient and public involvement

Patient (child) and public (stakeholders and experts) involvement will be sought on multiple levels. The Ugandan Ministry of Health was consulted regarding the design of the protocol and will be invited for dissemination meetings. Children and experts will be involved in adapting the screening tool for alcohol use detection at the stage of development (translate, discuss and adjust) and testing (quality assurance and epidemiological study). Moreover, patients and public stakeholders will be involved in the reporting and dissemination plans of this research.

### Ethics and dissemination

Ethical approval was obtained from the Regional Committees for Medical Research Ethics-South East Norway 6 March 2020, reference number 50146 and Makerere University College of Health sciences, School of Public Health, Higher degrees, Research and Ethics Committee 7 February 2020, reference number 745. In Uganda, the research project was registered at the Uganda National Council for Science and Technology, reference number SS 5245. Dissemination comprises peer-reviewed, open access, English language research papers, and policy recommendations and intersectoral dialogues relevant for public health policy. Data will be stored and kept in Uganda, controlled by Makerere School of Public health, and transferred abroad based on data transfer and storage agreements. Only officially acknowledged secure

data management and analysis systems will be used in line with General Data Protection Regulations in both Norway and Uganda.

### Status of progress

The adaptation of the CRAFFT tool was done between September 2019 and March 2020. This included consultations with experts and insight from qualitative interviews with 56 children. Analysis and write up for this are ongoing. Validation of the CRAFFT tool was conducted between September 2020 and March 2021 with 470 primary school age children and data entry is on-going. Data collection for the community household survey commenced on the 12th of November 2021 and data cleaning is on-going. Preliminary numbers show that a total of 4648 households were visited in 99 enumeration areas. 2575 of the visited households had eligible children and 3887 children were eligible for study participation. A total of 3594 children were enrolled into the study. We collected urine samples and anthropometric measurements for each enrolled child. Data collection for the health system and school study has not yet commenced. Further qualitative data collection with children exploring explanatory factors and context are planned and has not yet commenced.

### Author affiliations

<sup>1</sup>Faculty of Medicine, Department of Global Public Health and Primary Care, Centre for International Health, University of Bergen, Bergen, Norway

<sup>2</sup>School of Public Health, Makerere University College of Health Sciences, Kampala, Uganda

<sup>3</sup>Department of Mental Health, Faculty of Medicine and Health Sciences, Regional Centre for Child and Youth Mental Health and Child Welfare, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

<sup>4</sup>Department of Clinical Medicine, Centre for Nutrition, Mohn Nutrition Research Laboratory, University of Bergen, Bergen, Norway

<sup>5</sup>Department of Medical Biochemistry and Pharmacology, Haukeland University Hospital, Bergen, Norway

<sup>6</sup>Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

<sup>7</sup>School of Public Health, School of Medicine, Makerere University College of Health Sciences, Kampala, Uganda

<sup>8</sup>Department of Psychiatry, Mulago National Referral Hospital, Kampala, Uganda

<sup>9</sup>Section for Implementation and Treatment Research, Norwegian Center for Violence and Traumatic Stress Studies, Oslo, Norway

**Twitter** Ingunn Marie S. Engebretsen @ingunnengebret1

**Collaborators** The TREAT C-AUD consortium. Principal investigators: Uganda: Juliet Ndimwibo Babirye, MD, MPH, PhD, Senior Lecturer, and Norway: Ingunn Marie S. Engebretsen, MD, PhD, Professor (made the first manuscript draft).

Participating countries, institutions, investigators (alphabetical) and roles: Uganda: Makerere University, College of Health Sciences, School of Public Health: Harriet Aber, BSc food Science and technology, MA Public Health Nutrition, Public health nutrition. Esther Kisaakye, BSc Nursing, MA Public Health, implementation. Henry Wamani, MD, PhD, Public health nutrition. Deogracious Sebuwufu, MA Statistics, MA Business Administration, Diploma in Law, Diploma in Systems Engineering, data management and statistics, Makerere University, College of Health Sciences, School of Public Health and School of Medicine, Department of Psychiatry, Mulago National Referral Hospital. Joyce Sserunjogi Nalugya, MD specialized in Child and Adolescent Psychiatry. Norway: University of Bergen, Faculty of Medicine, Department of Global Health and Primary Care, Centre for International Health. Melf-Jacob Kühl, MA Public Policy, MA Laws. Vilde Skylstad, MD, PhD candidate (corresponding author). Neda Valeckaitė, Medical Student. University of Bergen, Faculty of Medicine, Department Clinical Medicine, Centre for Nutrition, Mohn

Nutrition Research Laboratory: Jutta Dierkes, PhD. Professor, Nutrition. University of Bergen, Department of Global Health and Primary Care, Centre for International Health and Norwegian Centre for Violence and Traumatic Stress Studies (NKVTS). Ane-Marthe Solheim Skar, MA Psychology, PhD. Norwegian University of Science and Technology (NTNU), Department of Mental Health, Faculty of Medicine and Health Sciences, Regional Centre for Child and Youth Mental Health and Child Welfare: Norbert Skokauskas, MD, PhD Professor, Child and Adolescent Psychiatry. Victoria Bakken, MSc Health Psychology. Norwegian University of Science and Technology (NTNU), Faculty of Medicine and Health Sciences: Silje Akselberg Iversen, Medical Student. Divya Rayamajhi, Stud. MA Public Health.

**Contributors** JNB and IMSE are co-principal investigators. VS, HA, VB, JD, M-JK, JSN, DS, AMSS, NS, HW, IMSE and JNB contributed to the original protocol which formed the basis for this manuscript. HA, JN, EK, DS, JNB organised the data collection to date. VS, HA, IMSE and JNB led the writing of the manuscript for publication. VS and HA have shared first authorship and IMSE and JNB have shared last authorship. All authors contributed to the review of the manuscript for publication.

**Funding** This work was supported by The Norwegian Research Council, the GLOBVAC-programme with the project grant number 285 489.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.

**Data availability statement** Data sharing not applicable as no complete datasets are generated and/or analysed for this study. Data collection is not yet completed, and there are no available data to share from this study.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Vilde Skylstad <http://orcid.org/0000-0002-2811-6457>

Ingunn Marie S. Engebretsen <http://orcid.org/0000-0001-5852-3611>

### REFERENCES

- Griswold MG, Fullman N, Hawley C, *et al*. Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the global burden of disease study 2016. *The Lancet* 2018;392:1015–35.
- Donovan JE. The burden of alcohol use: focus on children and Preadolescents. *Alcohol Res* 2014;35:186–92.
- Cortina MA *et al*. Prevalence of child mental health problems in sub-Saharan Africa. *Arch Pediatr Adolesc Med* 2012;166:276.
- Chan GCK, Leung J, Kelly AB, *et al*. Familial alcohol supply, adolescent drinking and early alcohol onset in 45 low and middle income countries. *Addict Behav* 2018;84:178–85.
- Swahn MH, Ali B, Palmier J, *et al*. Early alcohol use and problem drinking among students in Zambia and Uganda. *J Public Health Afr* 2011;2:e20.
- Dawson DA, Goldstein RB, Chou SP, *et al*. Age at first drink and the first incidence of adult-onset DSM-IV alcohol use disorders. *Alcohol Clin Exp Res* 2008;32:2149–60.
- Zucker RA, Donovan JE, Masten AS, *et al*. Developmental processes and mechanisms: ages 0–10. *Alcohol Res Health* 2009;32:16–29.
- Kabiru CW, Beguy D, Crichton J, *et al*. Self-reported drunkenness among adolescents in four sub-Saharan African countries: associations with adverse childhood experiences. *Child Adolesc Psychiatry Ment Health* 2010;4:17.
- The United Nations Development Programme. *Human development report 2020—the next frontier: human development and the Anthropocene*. New York, NY, 2020.
- Global Health Observatory data repository, The World Health Organization [Internet]. Uganda, Alcohol, total per capita (15+) consumption (in litres of pure alcohol), 2019, Both sexes, 2019. Available: <https://apps.who.int/gho/data/node.main.A1029SDG3?lang=en>
- World Health Organization. *World Health organization, global status report on alcohol and health*, 2018.
- Engebretsen IMS, Nalugya JS, Skylstad V, *et al*. "I feel good when I drink"—detecting childhood-onset alcohol abuse and dependence in



- a Ugandan community trial cohort. *Child Adolesc Psychiatry Ment Health* 2020;14:42. doi:10.1186/s13034-020-00349-z
- 13 Zhang W, Creswell J. The use of "mixing" procedure of mixed methods in health services research. *Med Care* 2013;51:e51-7.
  - 14 Knight JR, Sherritt L, Harris SK, et al. Validity of brief alcohol screening tests among adolescents: a comparison of the audit, POSIT, cage, and CRAFFT. *Alcohol Clin Exp Res* 2003;27:67-73.
  - 15 Sheehan DV, Lecrubier Y, Sheehan KH, et al. The Mini-International neuropsychiatric interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry* 1998;59(Suppl 20):22-33. quiz 34-57.
  - 16 Lundin A, Hallgren M, Forsman M, et al. Comparison of DSM-5 classifications of alcohol use disorders with those of DSM-IV, DSM-III-R, and ICD-10 in a general population sample in Sweden. *J Stud Alcohol Drugs* 2015;76.
  - 17 World Health Organization. *World Health Organization vaccination coverage cluster surveys: reference manual*, 2018.
  - 18 Kaiser R, Woodruff BA, Bilukha O, et al. Using design effects from previous cluster surveys to guide sample size calculation in emergency settings. Vol. 30, economic outlook 2006.
  - 19 Beusenbergh M, Orley JH. *A User's guide to the self reporting questionnaire (SRQ / compiled by M. Beusenbergh and J. Orley [Internet]*. Geneva: World Health Organization. <https://apps.who.int/iris/handle/10665/61113>
  - 20 Nakimuli-Mpungu E, Mojtabai R, Alexandre PK, et al. Cross-cultural adaptation and validation of the self-reporting questionnaire among HIV+ individuals in a rural ART program in southern Uganda. *HIV/AIDS - Res Palliat Care* 2012;4.
  - 21 Jellinek MS, Murphy JM, Burns BJ. Brief psychosocial screening in outpatient pediatric practice. *J Pediatr* 1986;109:371-8.
  - 22 Lowenthal E, Lawler K, Harari N, et al. Validation of the pediatric symptom checklist in HIV-infected Batswana. *J Child Adolesc Ment Health* 2011;23:17-28.
  - 23 Goodman LA, Corcoran C, Turner K, et al. Assessing traumatic event exposure: general issues and preliminary findings for the stressful life events screening questionnaire. *J Trauma Stress* 1998;11:521-42.
  - 24 Green BL, Chung JY, Daroowalla A, et al. Evaluating the cultural validity of the stressful life events screening questionnaire. *Violence Against Women* 2006;12:1191-213.
  - 25 Sachser C, Berliner L, Holt T, et al. International development and psychometric properties of the child and adolescent trauma screen (cats). *J Affect Disord* 2017;210:189-195.
  - 26 Evarist A. Adolescent trauma and psychosocial wellbeing in Entebbe-Uganda. *Univers J Psychol* 2018;6.
  - 27 Lang AJ, Stein MB. An abbreviated PTSD checklist for use as a screening instrument in primary care. *Behav Res Ther* 2005;43:585-94.
  - 28 Shelton KK, Frick PJ, Wootton J. Assessment of parenting practices in families of elementary school-age children. *J Clin Child Psychol* 1996;25.
  - 29 Scott S, Briskman J, Dadds MR. Measuring parenting in community and public health research using brief child and parent reports. *J Child Fam Stud* 2011;20.
  - 30 Badahdah A, Le KT. Parenting young Arab children: psychometric properties of an adapted Arabic brief version of the Alabama parenting questionnaire. *Child Psychiatry Hum Dev* 2016;47:486-93.
  - 31 Straus MA. *Child-report, adult-recall, and sibling versions of the revised conflict tactics scale*. Durham, NC: Fam Res Lab, 1999.
  - 32 Kolko DJ, Kazdin AE, Day BT. Children's perspectives in the assessment of family violence: Psychometric characteristics and comparison to parent reports. *Child Maltreat* 1996;1.
  - 33 Nkuba M, Hermenau K, Hecker T. Violence and maltreatment in Tanzanian families—Findings from a nationally representative sample of secondary school students and their parents. *Child Abuse Negl* 2018;77.
  - 34 Babor TF. *AUDIT : the alcohol use disorders identification test : guidelines for use in primary health care*. 2nd edn. Geneva: World Health Organization, 2001.
  - 35 Global school-based student health survey (GSHS) questionnaire [Internet], 2013. Available: <https://www.who.int/ncds/surveillance/gshs/en/>
  - 36 David S, Hornik R, Maklan D. National Survey of Parents and Youth (NSPY), 1998-2004 - Restricted Use Files. Inter-university Consortium for Political and Social Research [distributor] 2011.
  - 37 Harris K, Udry J. National Longitudinal Study of Adolescent to Adult Health (Add Health), 1994-2008 [Public Use]. Carolina Population Center, University of North Carolina-Chapel Hill [distributor], Inter-university Consortium for Political and Social Research [distributor] 2018;2018.
  - 38 Loeb M, Cappa C, Cialesi R. *Measuring child functioning: the Unicef/ Washington group module*. 59. Salud Publica de Mexico, 2017.
  - 39 Coates J, Swindale A, Bilinsky P. *Household food insecurity access scale (HFIAS) for measurement of food access: indicator guide*. Washington, DC: Food Nutr Tech, 2007.
  - 40 Borucki K, Schreiner R, Dierkes J, et al. Detection of recent ethanol intake with new markers: comparison of fatty acid ethyl esters in serum and of ethyl glucuronide and the ratio of 5-hydroxytryptophol to 5-hydroxyindole acetic acid in urine. *Alcohol Clin Exp Res* 2005;29:781-7.