

Characterization of lipoprotein profiles in patients with hypertriglyceridemic Fredrickson-Levy and Lees dyslipidemia phenotypes: the Very Large Database of Lipids Studies 6 and 7

Renato Quispe^{1,2}, Aditya D. Hendrani^{1,3}, Behnoud Baradaran-Noveiry¹, Seth S. Martin^{1,4}, Emily Brown⁵, Krishnaji R. Kulkarni⁶, Maciej Banach⁷, Peter P. Toth^{1,8,9}, Eliot A. Brinton¹⁰, Steven R. Jones¹, Parag H. Joshi^{1,11}

¹Johns Hopkins Ciccarone Center for the Prevention of Heart Disease, Baltimore, MD, USA

²Department of Medicine, Albert Einstein College of Medicine, Jacobi Medical Center, Bronx, NY, USA

³Louisiana State University Health Science Center-Shreveport, LA, USA

⁴Welch Center for Prevention, Epidemiology, and Clinical Research, Baltimore, MD, USA

⁵Center for Inherited Heart Disease, Johns Hopkins Hospital, Baltimore, MD, USA

⁶VAP Diagnostics Laboratory, Birmingham, AL, USA

⁷Department of Hypertension, Medical University of Lodz, Lodz, Poland

⁸Department of Preventive Cardiology, CGH Medical Center, Sterling, IL, USA

⁹Department of Family and Community Medicine, University of Illinois College of Medicine, Peoria, IL, USA

¹⁰Utah Lipid Center, Salt Lake City, UT, USA

¹¹Department of Internal Medicine, Division of Cardiology, University of Texas Southwestern Medical Center, Dallas, TX, USA

Corresponding author:

Parag H. Joshi MD, MHS
UT Southwestern
Medical Center
5323 Harry Hines
Blvd, #E5-730F
Dallas, TX 75390-8830
USA
Phone: (214) 645-1268
E-mail: Parag.Joshi@
UTSouthwestern.edu

Submitted: 10 February 2019

Accepted: 13 February 2019

Arch Med Sci 2019; 15 (5): 1195–1202

DOI: <https://doi.org/10.5114/aoms.2019.87207>

Copyright © 2019 Termedia & Banach

Abstract

Introduction: The association between triglycerides (TG) and cardiovascular diseases is complex. The classification of hypertriglyceridemic (HTG) phenotypes proposed by Fredrickson, Levy and Lees (FLL) helps inform treatment strategies. We aimed to describe levels of several lipoprotein variables from individuals with HTG FLL phenotypes from the Very Large Database of Lipids.

Material and methods: We included fasting samples from 979,539 individuals from a contemporary large study population of US adults. Lipids were directly measured by density-gradient ultracentrifugation using the Vertical Auto Profile test while TG levels were measured in whole plasma using the Abbott ARCHITECT C-8000 system. Hyperchylomicronemic (Hyper-CM) and non-chylomicronemic (non-CM) phenotypes were defined using computationally derived models. Individuals with FLL type IIa phenotype were excluded. Distributions of lipid variables were compared using medians and Kruskal-Wallis test.

Results: A total of 11.9% ($n = 116,925$) of individuals met criteria for HTG FLL phenotypes. Those with hyper-CM phenotypes ($n = 5$, < 0.1% of population) had two-fold higher TG levels compared with non-chylomicronemic (non-CM) individuals (11.9% of population) ($p < 0.001$). Type IIb individuals had the highest non-HDL-C levels (median 242 mg/dl). Cholesterol in large VLDL₁₊₂ particles was higher than in small VLDL₃ particles in all phenotypes except FLL type III. Hyper-CM phenotypes had significantly lower HDL-C levels but greater HDL₂/HDL₃-C ratio compared to non-CM phenotypes. Cholesterol content of the lipoprotein (a) peak was significantly higher in the hyper-CM groups compared to non-CM phenotypes ($p < 0.0001$).

Conclusions: This observational hypothesis-generating study provides insight into the complexity of lipid metabolism in HTG phenotypes, including less traditional lipid measures such as LDL density, HDL subclasses and Lp(a)-C.

Key words: hypertriglyceridemia, dyslipidemia, Fredrickson-Levy phenotypes.

Introduction

The pathophysiologic role of triglycerides (TG) in the development of cardiovascular disease (CVD) is complex [1, 2]. TG-related CVD risk is likely driven by atherogenic properties of multiple TG-rich lipoproteins (TRLs) including large and small very low-density lipoproteins (VLDL), intermediate-density lipoproteins (IDL) and possibly chylomicrons (CM) and CM remnants [1].

Fredrickson, Levy and Lees (FLL) defined five primary lipoprotein phenotypes (type I to V) based on dyslipoproteinemias and characterized by the presence of hypercholesterolemia and/or hypertriglyceridemia [3, 4], and this classification system was formally adopted by the World Health Organization (WHO) in 1970 [5]. Since each FLL phenotype has elevations of one or more distinct lipoprotein classes (or subclasses), this classification may better inform treatment strategies based on the dyslipoproteinemia [3, 6, 7].

Significant clinical heterogeneity exists among hypertriglyceridemic (HTG) phenotypes. For instance, the presence of elevated chylomicrons (type I and V) confers remarkably distinct clinical manifestations, prognosis and management with more prominent risk of acute pancreatitis rather than CVD [8, 9]. Patients with other HTG phenotypes, such as type IIb [10] and III [11] are at markedly increase risk to develop premature coronary artery disease and early myocardial infarction. Emerging therapies, such as apolipoprotein C-III inhibition through antisense oligonucleotide [12] or monoclonal antibodies [13], may have unique and differing impact on the distinct FLL phenotypes.

To our knowledge, no large-scale population-based studies exist with detailed characterization of the lipid profile in individuals with hypertriglyceridemic FLL phenotypes. The aim of this study is to gain insight into the complex pathophysiology of hypertriglyceridemic FLL phenotypes by characterizing them in the Very Large Database of Lipids, which is an ongoing contemporary big-data project with lipoprotein data from more than 5 million unique patients from US-based clinics.

Material and methods

Study population

This is a cross-sectional analysis of lipid profile data from 979,539 unique US adults from

the Very Large Database of Lipids with fasting samples. The methods for the creation of this database have been described [14]. In summary, the database was constructed using de-identified data from the clinical operations of a single laboratory (VAP Diagnostics Lab, Birmingham, AL). The samples originated predominantly from primary care clinics throughout the US between 2009 and 2015.

Lipid measurements

Lipid profiles were measured with direct ultracentrifugation by the Vertical Auto Profile (VAP) test. The VAP test is a density gradient ultracentrifugation technique, validated in comparison to the beta quantification method [15], both of which separate lipoproteins according to their hydrated densities. The VAP directly measures cholesterol concentrations of all lipoprotein classes sequentially across the density spectrum (during gentle evacuation from the centrifuge tube). The resulting cholesterol peaks are defined as high-density lipoprotein cholesterol (HDL-C), lipoprotein (a) cholesterol (Lp(a)-C), “real” low-density lipoprotein cholesterol (LDL-C), intermediate-density lipoprotein cholesterol (IDL-C) and very low-density lipoprotein cholesterol (VLDL-C). Note that the total LDL-C level reported in this study is the sum of real LDL-C + Lp(a)-C + IDL-C, equivalent to non-HDL-C minus VLDL-C, consistent with the Friedewald equation of LDL-C. The standard lipid profile in clinical practice often presents an estimated version of this LDL-C value based on the Friedewald equation when TG < 400 mg/dl. Cholesterol concentrations in lipoprotein subclasses (HDL₂-C, HDL₃-C, LDL₁-C, LDL₂-C, LDL₃-C, LDL₄-C, VLDL₁-C, VLDL₂-C, and VLDL₃-C) were also measured by VAP [15]. TG were directly measured in whole plasma using the Abbott ARCHITECT C-8000 system (Abbott Park, IL) [14].

Non-HDL-C was calculated as total cholesterol (TC) minus HDL-C. Remnant lipoprotein particle cholesterol (RLP-C) was defined as the sum of the densest VLDL-C subfraction (VLDL₃-C) and IDL-C [16].

Further, to better characterize TG-related lipoprotein parameters we calculated several ratios of lipid factors: TG/TC, VLDL-C/TG, HDL₂-C/HDL₃-C and TG/HDL-C. We calculated average LDL particle density as the LDL Density Ratio (LLDR) by the following formula: LLDR = $\ln([\text{LDL}_3\text{-C} + \text{LDL}_4\text{-C}]/$

[LDL₁-C + LDL₂-C]), previously shown by us to correlate very strongly with ultracentrifugally-determined modal LDL density ($R^2 = 0.80$) [14].

Definition of FLL phenotypes

The definitions for FLL phenotype categorization and distributions in the study are shown in Table I. First, we categorized individuals by the presence or absence of hyperchylomicronemia (hyper-CM), defined as a total plasma TG/plasma TC > 10 or < 10 respectively [1]. We further stratified patients with hyper-CM into two categories: excess of chylomicrons only (FLL type I), and excess of both chylomicrons and VLDL (defined as VLDL-C above the 90th percentile for the general study population) (FLL type V).

Among patients without chylomicronemia (non-CM), we sub-classified individuals as 1) type IIb, based on the presence of both VLDL-C excess (> 90th percentile) and LDL-C excess (> 90th percentile) [17]; 2) type III, based on VLDL-C/TG > 0.3 and TG > 130 without LDL-C excess (< 90th percentile) [18]; and 3) type IV based on VLDL-C excess (> 90th percentile) without LDL-C excess (< 90th percentile) [17]. Individuals who met criteria for type IIa (LDL-C > 90th percentile and VLDL-C < 90th percentile and VLDL-C/TG < 0.3) were not included as regular subjects in this study, because they do not have HTG. Finally, the vast majority of individuals in the database population did not meet criteria for any FLL phenotype, and were classified into the “continuum” group.

Statistical analysis

Within each FLL phenotype and the continuum group, we examined TG levels and lipid parameters obtained from the VAP test. Distributions of lipid parameters are shown using medians, 25th and 75th percentile, and were compared using the Kruskal-Wallis test with a *p*-value < 0.001 arbitrarily required to be considered significant given our sample size.

Results

For the present study, lipid data from the first accessioned blood sample in the VLDL database from a subset of 979,539 unique individuals reported to be in the fasting state were included. A total of 768,161 (79.1%) individuals did not meet criteria for any FLL phenotype were classified in the continuum group, whereas 94,453 (9.7%) met criteria for type IIa hyperlipidemia, and 116,925 (11.9%) had one of several possible HTG phenotypes.

The frequency of HTG FLL phenotypes was as follows: 5 type I, 20,144 type IIb, 2,151 type III, 85,540 type IV and 246 type V, as shown in Table I.

Table I. The Very Large Database of Lipids: FLL phenotypes and continuum group

Classical dyslipidemia	Main features	Diagnostic criteria	N (%)	Age, median (25 th –75 th)	Male (%)
Type I	Chylomicron excess	TG/TC ratio > 10 and VLDL-C < 90 th percentile	5 (< 0.00004)	50.01 (35.0–52.8)	5 (100)
Type IIa	LDL-C excess	LDL-C > 90 th percentile, VLDL-C < 90 th percentile, and VLDL-C/TG < 0.3	94,453 (9.7)	55.21 (46.6–63.8)	39,437 (42.0)
Type IIb	LDL-C and VLDL-C excess	LDL-C, VLDL-C > 90 th percentile, and VLDL-C/TG < 0.3	20,144 (2.1)	54.99 (46.4–63.7)	9,583 (47.8)
Type III	Remnant lipoprotein cholesterol excess	VLDL-C/TG > 0.3, TG > 130, and LDL-C < 90 th percentile	2,151 (0.2)	57.45 (49.2–66.3)	1,073 (50.1)
Type IV	VLDL-C excess	VLDL-C > 90 th percentile, LDL-C < 90 th percentile, and VLDL-C/TG < 0.3	85,540 (8.8)	55.09 (46.0–65.04)	51,195 (60.1)
Type V	Chylomicron and VLDL-C excess	TG/TC ratio > 10 and VLDL-C > 90 th percentile	246 (0.03)	47.86 (40.2–54.6)	200 (81.3)
Continuum*	N/A	N/A	768,161 (79.1)	57.44 (46.6–67.7)	353,310 (46.2)

N/A – not applicable. *Continuum: individuals who did not meet criteria for any FLL phenotype.

Table II. Lipid profile of individuals with hypertriglyceridemic FLL phenotypes and continuum group

Lipid parameters	Hyper-chylomicronemic phenotypes			Non-chylomicronemic phenotypes			Continuum
	Type I	Type V	Type IIb	Type III	Type IV	Type IV	
TC [mg/dl]	102 (90-117)	222 (167-288)	289 (269-314)	234 (201-268)	214 (187-238)	186 (160-211)	
TG [mg/dl]	1858 (1026-2439)	3286.5 (2203-4823)	247 (208-302)	192 (157-252)	281 (228-367)	103 (75-140)	
TG/TC	17.61 (15.88-19.37)	13.66 (11.28-19.38)	0.86 (0.71-1.05)	0.88 (0.72-1.08)	1.36 (1.06-1.83)	0.56 (0.41-0.77)	
VLDL-C/TG	0.01 (0.01-0.02)	0.05 (0.03-0.06)	0.19 (0.17-0.22)	0.34 (0.31-0.38)	0.18 (0.15-0.21)	0.20 (0.18-0.23)	
VLDL-C [mg/dl]	29 (28-34)	140.5 (100-211)	46 (42-55)	67 (54-91)	48 (42-60)	20 (16-26)	
VLDL ₁₊₂ -C [mg/dl]	17.5 (11.2-20.2)	111.2 (75.1-174.4)	22.7 (19.8-27.6)	28.6 (21.9-41.7)	24.5 (20.8-32.2)	8.4 (6.5-11.3)	
VLDL ₃ -C [mg/dl]	12 (12-16)	28 (21-43)	24 (22-28)	38 (31-50)	24 (21-28)	12 (10-15)	
IDL-C [mg/dl]	5 (5-12)	14 (11-18)	30 (24-37)	39 (30-48)	22 (18-28)	12 (9-16)	
RLP-C [mg/dl]	17 (17-28)	43 (34-60)	54 (47-64)	77 (63-96)	47 (40-55)	24 (19-30)	
Total LDL-C [mg/dl]	41 (35-62)	45 (33-59)	192 (179-212)	109 (85-140)	120 (96-142)	109 (87-131)	
LDL-C [mg/dl]	25 (23-39)	19 (12-26)	155 (145-172)	57 (39-94)	91 (68-111)	89 (69-109)	
Lp(a)-C [mg/dl]	7 (4-11)	11 (7-16)	5 (3-7)	6 (4-9)	6 (4-9)	6 (4-10)	
LDL ₁ -C [mg/dl]	3.3 (2.4-9.9)	5.8 (3.4-8.6)	29.8 (25.6-35)	21.5 (16.7-26.6)	17 (12.8-21.4)	14 (9.6-19)	
LDL ₂ -C [mg/dl]	0.3 (0-4.4)	0.55 (0-2.4)	21.2 (11.6-34.7)	8.9 (4.9-16.5)	8.7 (3.9-16.8)	21.9 (12.2-33.7)	
LDL ₃ -C [mg/dl]	11.1 (1.8-12.6)	4.9 (2.4-8.1)	77.9 (69.5-88)	17.5 (10.3-38.1)	41.5 (29.1-53.4)	38.2 (27.4-50.4)	
LDL ₄ -C [mg/dl]	4.3 (1.5-4.7)	6 (4.2-7.9)	24.5 (13.3-37.1)	4.6 (2.6-8.6)	15.9 (8.5-24.9)	8.6 (4.8-13.6)	
LLDR	0.24 (-1.29-1.66)	0.50 (0.05-0.88)	0.75 (0.34-1.09)	-0.27 (-0.65-0.14)	0.81 (0.36-1.18)	0.29 (-0.20-0.81)	
HDL-C [mg/dl]	31 (23-33)	22 (16-27)	44 (38-52)	47 (40-54)	38 (33-45)	52 (43-64)	
HDL ₂ -C [mg/dl]	8 (6-10)	7 (4-10)	9 (7-12)	12 (9-15)	8 (6-11)	13 (9-18)	
HDL ₃ -C [mg/dl]	20 (15-23)	14 (12-18)	35 (31-40)	35 (30-40)	30 (26-35)	40 (33-47)	
HDL ₂ /HDL ₃ -C	0.5 (0.43-0.50)	0.44 (0.33-0.60)	0.27 (0.23-0.33)	0.33 (0.27-0.4)	0.27 (0.22-0.34)	0.32 (0.26-0.39)	
TG/HDL-C	80.61 (73.91-80.78)	149.20 (91.32-248.44)	5.59 (4.35-7.30)	4.18 (3.16-5.88)	7.34 (5.39-10.58)	1.95 (1.25-3.02)	
Non-HDL-C [mg/dl]	69 (55-94)	195 (144-259)	242 (225-266)	186 (154-220)	174 (150-196)	130 (107-154)	

LDL-C 90th percentile: 141 mg/dl and VLDL-C 90th percentile: 40 mg/dl. Numbers shown above are medians (25th-75th percentile). Medians were compared using the Kruskal-Wallis test. P-value was found to be < 0.001 for all the rows.

Triglyceride-rich lipoproteins: TG, RLP-C and VLDL subclasses

The hyper-CM groups had extremely high TG levels (> 1000 mg/dl), as shown in Table II. Type V individuals had the highest TG levels among FLL phenotypes; of note, more than 3/4 of type V individuals had TG levels above 2000 mg/dl (median: 3286.5; 2203–4823). Individuals categorized as type I had lower TG levels with a median of 1858 mg/dl (1026–2439 mg/dl; $p < 0.001$).

In contrast, TG levels in approximately 90% of non-CM individuals were less than 500 mg/dl. Individuals in the type IV phenotype had the highest average TG levels among the three non-CM phenotypes, while type III individuals had the lowest overall TG levels (Table II).

As specified by our definition, individuals with the type I phenotype had significantly lower VLDL-C levels than with type V; 29 mg/dl (28–34) vs. 140.5 mg/dl (100–211), respectively ($p < 0.001$). The proportion of cholesterol content in large VLDL particles (VLDL₁₊₂-C) relative to small VLDL particle (VLDL₃-C) was increased in all FLL phenotypes, except type III, in comparison to the continuum group. In addition, this proportion of VLDL cholesterol in larger particles was significantly higher in type V individuals than all other HTG phenotypes.

With the exception of type I, all individuals with a HTG FLL phenotype had higher RLP-C levels than the continuum group. RLP-C levels (IDL-C +

VLDL₃-C) in type III patients were 1.5-times higher than that of the other HTG groups.

Cholesterol lipoproteins: TC, HDL-C and LDL-C

Type IIb had higher TC levels compared to the other HTG FLL phenotypes and continuum group. Both total and “real” LDL-C levels were significantly lower in hyper-CM (FLL types I and V) compared to non-CM HTG and continuum groups (Table II). However, Lp(a)-C was higher in hyper-CM groups compared to non-CM phenotypes and continuum group.

Levels of LDL-C subclasses among HTG FLL phenotypes are shown in Table II. Finally, type IIb individuals had the highest non-HDL-C levels (median 242 mg/dl). Hyper-CM phenotypes had significantly lower HDL-C levels compared to non-CM phenotypes. The former also had significantly elevated HDL₂/HDL₃-C ratio compared to the latter (0.44–0.50 vs. 0.27–0.33, $p < 0.001$). Levels of triglycerides and cholesterol-based parameters are summarized in Figure 1 for both hyper-CM and non-CM phenotypes.

Discussion

The present study includes comprehensive directly measured lipoprotein data on individuals with HTG FLL phenotypes from the Very Large Database of Lipids, the largest lipid study to date. In our study population, we estimated the frequency




Figure 1. Levels of triglycerides and cholesterol measures in hyper-chylomicronemic and non-chylomicronemic phenotypes in the Very Large Database of Lipids study

of HTG FLL phenotypes, under-recognized entities and often labeled as rare disorders, in a contemporary cross-sectional population. Whereas hyper-CM phenotypes are truly rare diseases, non-CM HTG phenotypes, which carry significant risk of premature CVD through pathways other than LDL-C [1], were seen in ~10% of our study population.

Hyperchylomicronemic phenotypes

Our findings provide insight into lipid metabolism in the two hyper-CM phenotypes. As expected, individuals in the hyper-CM phenotypes had extremely high TG levels (over 1000 mg/dl), at which the risk of acute pancreatitis is strikingly elevated compared to the general population [19].

We observed that individuals with the type V phenotype had TG levels almost two times higher than type I perhaps due to elevation of both chylomicrons and VLDL particles, both of which carry TG.

Hypertriglyceridemia stimulates the activity of cholesteryl ester transfer protein (CETP), enhancing the equimolar neutral lipid exchange of triglycerides from TRL to HDL and LDL while CE is transferred in the opposite direction. This loading of HDL and LDL particles with TG renders these particles better substrates for lipolysis by hepatic lipase, which promotes catabolism and renal elimination of HDL and production of smaller, denser and more numerous LDL particles. As HDL-C levels decrease, TG/HDL-C ratio increases [20]. HDL-C levels are significantly lower in the hyper-CM phenotypes compared to the non-CM phenotypes. This could be explained by the higher relative TG concentration gradient from chylomicrons to HDL particles, compared to VLDL or RLP to HDL particles [17]. Among the hyper-CM phenotypes in which we found higher TG levels among type V individuals, there was a ~2 fold higher TG/HDL-C ratio among type V individuals compared with type I. Additionally, levels of HDL₂-C were similar between both hyper-CM groups, but HDL₃-C levels were significantly greater in type I. Overall, we found that the HDL₂/HDL₃-C ratio in both hyper-CM phenotypes was greater than all other individuals in the database. Endogenous HTG, more often driven by VLDL overproduction like in patients with diabetes mellitus, is known for provoking a shift toward smaller sizes of HDL, such as pre-beta1-HDL and HDL_{3a} [21]. As such, our findings could be the result of different pathophysiological mechanisms in hyper-CM phenotypes that are not seen in hypertriglyceridemia secondary to clinical conditions such as diabetes mellitus. Similarly, it could also be a manifestation of chylomicron particles being a less important source of triglyceride in the CETP pathway.

Levels of RLP-C in individuals with type V were almost two-fold higher than those with type I phe-

notype, likely reflecting the excess of VLDL particles since RLP-C contains VLDL₃-C as one of its components [16]. Another consequence of such VLDL excess is reflected in the greater levels of TC, non-HDL-C and LDL-C observed in type V as compared to type I, the result of the metabolism of VLDL particles [22]. Similarly, levels of IDL-C (the other component of RLP-C) were also ~2 times higher in type V compared to type I phenotype. Finally, VLDL-C and VLDL₁₊₂-C in type V dyslipidemia were almost 4 fold higher than other HTG phenotypes.

Surprisingly, logarithmic LDL density ratio (LLDR) was greater in type I (~ two-fold) compared to type V, despite the higher TG levels found in type V. This unexpected finding may be due to a greater presence of buoyant LDL which would naturally result from increased VLDL in type V, absent in type I. However, it should be noted that only 5 individuals were classified as type I, thus these results should be interpreted cautiously. Interestingly, Lp(a)-C was found to be higher among both hyper-CM compared to non-CM phenotypes, despite previous studies reporting a negative correlation between TG and Lp(a)-C levels. More studies are needed to confirm and further elucidate our novel finding of increased Lp(a) levels in patients with fasting chylomicronemia [23–25].

Non-chylomicronemic phenotypes

Among the non-CM phenotypes, individuals with type IV phenotype had the highest TG and the lowest HDL-C levels. Those with type IV phenotype had a lower HDL₂/HDL₃-C ratio compared to the type III and continuum groups, likely as a result of a given level of CETP activity leading to smaller denser HDL (HDL₃) in the setting of the higher TG levels in type IV [26]. Individuals with the type III phenotype were noted to have a higher HDL₂/HDL₃-C ratio, which appears to be related to higher HDL₂-C levels, and may indirectly reflect lower TG levels and lesser CETP activity. However, it is important to consider that individuals with other entities, such as hepatic lipase deficiency, can develop type III-like lipoprotein profile [27].

The type III (familial dysbetalipoproteinemia) FLL phenotype, also known as familial dysbetalipoproteinemia, is characterized by increased RLP-C and cholesterol enriched VLDL particles. This is believed to result from overproduction of VLDL combined with apolipoprotein-E (Apo E) structural variation reducing the clearance of RLP-C via the LDL receptor related protein [28]. Type III individuals had the highest levels of total VLDL-C among non-CM phenotypes, associated with significantly higher levels of remnant VLDL-C (VLDL₃-C), since VLDL₁₊₂-C levels were relatively similar among the three non-CM phenotypes. This is expected from

the definition of type III, an elevated VLDL-C/TG ratio. Also, as expected, IDL-C levels were highest in type III patients, an elevation that was even slightly more striking than for VLDL₃-C.

Interestingly, LLDR (LDL particle density) was lowest in type III dyslipidemia compared to other HTG phenotypes. The predominance of larger, buoyant LDL particle in type III might be related to the fact that median TG was lowest in type III among all HTG FLL phenotypes, and TG levels are known to be positively correlated with denser smaller LDL particles [29]. This is to our knowledge, however, a novel observation. Finally, our type III patients had a higher HDL₂/HDL₃-C ratio, related to higher HDL₂-C levels, which might reflect lower net CETP effect (assuming similar CETP activity) due to lower TG levels. These findings of lower average density of both LDL and HDL activity would predict lesser ASCVD risk, which nonetheless is strikingly elevated in type III patients, thus, likely due to the VLDL and IDL changes. To our knowledge, these observations of decreased LDL and HDL particle density are novel in the scientific literature.

There are limitations of this study. First, intra-individual temporal variation was not addressed since we included only one-time sample collection for lipid measurement. Second, demographic variables such as biophysical status, race/ethnicity, or dietary habits were not available. Third, we did not have clinical data, such as lipid-lowering treatment (including TG-lowering therapies) or history of CVD. Fourth, genetic information, well characterized in HTG phenotypes [30], could not be obtained and so we relied on phenotypic characterization of the FLL classifications. Lastly, the cross-sectional nature of the study does not warrant the drawing of any conclusions regarding cause-effect relationships among the various elements of our lipoprotein data, and the lack of any ASCVD data does not allow any inference regarding our lipoprotein data and ASCVD event risk.

In conclusion, our findings provide important information about the frequency of FLL phenotypes in a large contemporary population, as well as the comprehensive distribution of lipoprotein levels in the setting of HTG phenotypes that include TG levels in the chylomicronemic range. The present study, although observational and hypothesis-generating, might help gain insight into the complexity of dyslipidemia, particularly in hypertriglyceridemic settings.

Conflict of interest

Drs. Martin and Jones are listed as coinventors on a pending patent filed by Johns Hopkins University for LDL-C estimation. Dr Jones has served as an advisor to Sano/Regeneron. Dr Martin has

served as a consultant to Quest Diagnostics, Sano/Regeneron, Amgen, and the Pew Research Center. Dr. Joshi received non-compensated site PI for studies funded by AstraZeneca, Pfizer, Merck and Regeneron/Sanofi as well as modest consulting income from Regeneron. Dr. Toth has served as consultant and speaker for Amarin, Amgen, Kowa, Merck, Novo-Nordisk, Regeneron, and Sanofi; consultant to Gemphire. Dr. Brinton has received research support from Amarin, Kowa and the National Institutes of Health; honoraria as consultant for Akcea, Alexion, Amarin, Amgen, Janssen, Kastle, Kowa, Merck, Precision Biosciences, PTS Diagnostics, Regeneron, Sanofi-Aventis; honoraria as speaker for Akcea, Alexion, Amarin, Amgen, Boehringer-Ingelheim, Janssen, Kastle, Kowa, Merck, Novo-Nordisk, Regeneron, Sanofi-Aventis.

References

1. Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation* 2011; 123: 2292-333.
2. Soran H, Adam S, Mohammad JB, et al. Hypercholesterolaemia – practical information for non-specialists. *Arch Med Sci* 2018; 14: 1-21.
3. Fredrickson DS, Lees RS. A system for phenotyping hyperlipoproteinemia. *Circulation* 1965; 31: 321-7.
4. Brahm A, Hegele RA. Hypertriglyceridemia. *Nutrients* 2013; 5: 981-1001.
5. Beaumont JL, Carlson LA, Cooper GR, Fejfar Z, Fredrickson DS, Strasser T. Classification of hyperlipidaemias and hyperlipoproteinaemias. *Bull World Health Organ* 1970; 43: 891-915.
6. Sniderman A, Couture P, de Graaf J. Diagnosis and treatment of apolipoprotein B dyslipoproteinemias. *Nat Rev Endocrinol* 2010; 6: 335-46.
7. Stulc T, Lanska V, Snejdrlova M, Vrablik M, Prusikova M, Ceska R. A comprehensive guidelines-based approach reduces cardiovascular risk in everyday practice: the VARO study. *Arch Med Sci* 2017; 13: 705-10.
8. Goldberg JJ. Hypertriglyceridemia: impact and treatment. *Endocrinol Metab Clin North Am* 2009; 38: 137-49.
9. Banach M, Jankowski P, Jozwiak J, et al. PoLA/CFPI/P/PCS guidelines for the management of dyslipidaemias for family physicians 2016. *Arch Med Sci* 2017; 13: 1-45.
10. Genest JJ Jr, Martin-Munley SS, McNamara JR, et al. Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* 1992; 85: 2025-33.
11. Hopkins PN, Brinton EA, Nanjee MN. Hyperlipoproteinemia type 3: the forgotten phenotype. *Curr Atheroscler Rep* 2014; 16: 440.
12. Khetarpal SA, Qamar A, Millar JS, Rader DJ. Targeting apoC-III to reduce coronary disease risk. *Curr Atheroscler Rep* 2016; 18: 54.
13. Huynh K. Dyslipidaemia: monoclonal antibody targeting lipoprotein-bound human apoC-III. *Nat Rev Cardiol* 2017; 14: 632-3.
14. Ahmed HM, Elshazly MB, Martin SS, Blaha MJ, Kulkarni KR, Jones SR. Ratio of dense to buoyant LDL subclass is associated with LDL density phenotype (VLDL-5). *Open Chem Biomed Methods J* 2013; 6: 1-5.

15. Kulkarni KR. Cholesterol profile measurement by vertical auto profile method. *Clin Labor Med* 2006; 26: 787-802.
16. Jones SR, Martin SS, Brinton EA. Letter by Jones et al regarding article, "Elevated remnant cholesterol causes both low-grade inflammation and ischemic heart disease, whereas elevated low-density lipoprotein cholesterol causes ischemic heart disease without inflammation". *Circulation* 2014; 129: e655.
17. Quispe R, Al-Hijji M, Swiger KJ, et al. Lipid phenotypes at the extremes of high-density lipoprotein cholesterol: the very large database of lipids-9. *J Clin Lipidol* 2015; 9: 511-8 e1-5.
18. Wang T, Nakajima K, Leary ET, et al. Ratio of remnant-like particle-cholesterol to serum total triglycerides is an effective alternative to ultracentrifugal and electrophoretic methods in the diagnosis of familial type III hyperlipoproteinemia. *Clin Chem* 1999; 45: 1981-7.
19. Leaf DA. Chylomicronemia and the chylomicronemia syndrome: a practical approach to management. *Am J Med* 2008; 121: 10-2.
20. Foger B, Ritsch A, Doblinger A, Wessels H, Patsch JR. Relationship of plasma cholesteryl ester transfer protein to HDL cholesterol. Studies in normotriglyceridemia and moderate hypertriglyceridemia. *Arterioscler Thromb Vasc Biol* 1996; 16: 1430-6.
21. Gou L, Fu M, Xu Y, Tian Y, Yan B, Yang L. Alterations of high-density lipoprotein subclasses in endogenous hypertriglyceridemia. *Am Heart J* 2005; 150: 1039-45.
22. Sparks CE, Sparks JD. Lipid metabolism: insights into the complexity of VLDL metabolic pathways. *Curr Opin Lipidol* 2010; 21: 280-1.
23. McConathy WJ, Trieu VN, Koren E, Wang CS, Corder CC. Triglyceride-rich lipoprotein interactions with Lp(a). *Chem Phys Lipids* 1994; 67-68: 105-13.
24. Werba JP, Safa O, Gianfranceschi G, Michelagnoli S, Sirtori CR, Franceschini G. Plasma triglycerides and lipoprotein(a): inverse relationship in a hyperlipidemic Italian population. *Atherosclerosis* 1993; 101: 203-11.
25. Cohn JS, Lam CW, Sullivan DR, Hensley WJ. Plasma lipoprotein distribution of apolipoprotein(a) in the fed and fasted states. *Atherosclerosis* 1991; 90: 59-66.
26. Durrington PN. Cholesteryl ester transfer protein (CETP) inhibitors. *Br J Cardiol* 2012; 19: 126-33.
27. Jansen H, Verhoeven AJ, Sijbrands EJ. Hepatic lipase: a pro- or anti-atherogenic protein? *J Lipid Res* 2002; 43: 1352-62.
28. Mahley RW, Huang Y, Rall SC Jr. Pathogenesis of type III hyperlipoproteinemia (dysbetalipoproteinemia). Questions, quandaries, and paradoxes. *J Lipid Res* 1999; 40: 1933-49.
29. Cho Y, Lee SG, Jee SH, Kim JH. Hypertriglyceridemia is a major factor associated with elevated levels of small dense LDL cholesterol in patients with metabolic syndrome. *Ann Labor Med* 2015; 35: 586-94.
30. Johansen CT, Hegele RA. Genetic bases of hypertriglyceridemic phenotypes. *Curr Opin Lipidol* 2011; 22: 247-53.