



OPEN ACCESS

Educational gradients in all-cause mortality in two cohorts in the Czech Republic during the early stage of the postcommunist transition

Ceciel Pauls,¹ Andrea Dalecká ,² Wentian Lu,¹ Jaroslav Hubáček ,³ Hynek Pikhart,^{1,2} Martin Bobák^{1,2}

¹Department of Epidemiology and Public Health, University College London, London, UK
²RECETOX, Faculty of Science, Masaryk University, Brno, Czech Republic
³Center for Experimental Medicine, Institute of Clinical and Experimental Medicine, Prague, Czech Republic

Correspondence to

Dr Andrea Dalecká, Masaryk University RECETOX, Brno, Jihomoravský kraj, Czech Republic; andrea.dalecka@recetox.muni.cz

CP and AD are joint first authors.

Received 4 March 2022
Accepted 31 May 2022
Published Online First 21 June 2022

ABSTRACT

Objectives We investigated whether social gradient in all-cause mortality in the Czech Republic changed during the postcommunist transition by comparing two cohorts, recruited before and after the political changes in 1989.

Methods Participants (aged 25–64 years) in two population surveys (n=2530 in 1985, n=2294 in 1992) were followed up for mortality for 15 years (291 and 281 deaths, respectively). Education was classified into attainment categories and years of schooling (both continuous and in tertiles). Cox regression was used to estimate HR of death by educational indices in each cohort over a 15-year follow-up.

Results All three educational variables were significantly associated with reduced risk of death in both cohorts when men and women were combined; for example, the adjusted HRs of death in the highest versus lowest tertile of years of schooling were 0.65 (95% CI 0.47 to 0.89) in 1985 and 0.67 (95% CI 0.48 to 0.93) in 1992. Adjustment for covariates attenuated the gradients. In sex-specific analysis, the gradient was more pronounced and statistically significant in men. There were no significant interactions between cohort and educational indices.

Conclusions The educational gradient in mortality did not differ between the two cohorts (1985 vs 1992), suggesting no major increase in educational inequality during the early stage of postcommunist transition. Further research is needed to understand trends in health inequalities during socioeconomic transitions.

INTRODUCTION

The inverse association between educational attainment and health, including all cause and cause-specific mortality, has been long established.^{1–3} However, the educational gradient differs over time and between populations. For example, analysis of official data from 22 European countries found considerable differences in the magnitude of educational inequalities between countries.⁴ This suggests that sociopolitical context plays an important role in shaping social inequalities in health.^{4,5}

A relatively recent example of rapid societal change is the postcommunist transition in Central and Eastern Europe and the former Soviet Union. Macroeconomic data suggest a rapid increase in income inequalities, reappearance of unemployment and falls in living standards in the early stages of the transition.^{6,7} With democratic reforms in Central and Eastern Europe, the importance of education

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Previous studies reported an increase in social inequalities in mortality during the early stage of postcommunist transition. However, there has been less evidence from Central Europe, especially based on individual-level (linked) data. There has also been a lack of studies assessing educational gradients for men and women separately.

WHAT THIS STUDY ADDS

⇒ Although the educational gradient in mortality existed before as well as during the political transition, there was no evidence for the expected steepening of the gradient in our study. Our results suggested that the educational gradient in mortality was substantially stronger among men than in women. Further research is needed to understand trends in health inequalities during socioeconomic transitions emphasising the differential patterns between sexes.

as a route to higher income rapidly increased, contributing to increase in income inequality by education.⁸ Published studies suggest that social inequalities in mortality in the former Soviet Union and the Baltic countries increased considerably after 1990,^{9–11} but there is less evidence from Central Europe, where the transition was less dramatic than in the former Soviet Union. Geographical studies in Hungary and the Czech Republic suggest increasing socioeconomic inequality in mortality after 1990.¹² Estimates based on official unlinked data on death registration with census data in the Czech Republic and the Baltic states also show a short-term increase in all-cause mortality after 1990.^{13–15} In most reports, the educational inequalities increased very quickly, over only a few years. However, as most of available studies used unlinked routine data, the numerator-denominator bias makes them less reliable.¹⁶

In this paper, we analysed data from two Czech cohorts using identical protocol established in 1985 and 1992 to assess changes in educational inequalities using individual-level (ie, linked) data. We hypothesised that the educational gradient would become steeper in the later cohort and this would be apparent in both men and women.



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Pauls C, Dalecká A, Lu W, et al. *J Epidemiol Community Health* 2022;**76**:819–822.

METHODS

We used data from the 1985 and 1992 population studies conducted as part of the WHO MONICA Project¹⁷ in a 1% random age-sex-stratified sample of men and women aged 25–64 in of six Czech districts (more detail is available elsewhere¹⁸). In total, 2573 and 2353 participants were recruited in 1985 and 1992 (response rate 84% and 73%), respectively. Participants with missing data on education, marital status and cardiovascular risk factors were excluded, leaving 2530 and 2294, respectively, in the analytical sample.

Education was assessed by two indices. First, educational attainment was reported in categories, which were classified into primary, secondary and university education. Second, participants reported how many years they spent in education; this information was used both as continuous variable (years of schooling) and classified in tertiles by cohort. Covariates (potential confounders or mediators) included marital status (recoded into binary variable married vs single/widowed/divorced), ever smoking cigarettes (yes vs no), height (metres), body mass index (BMI, calculated as kg/m²), high-density lipoproteins cholesterol (mmol/L), total cholesterol (mmol/L), high blood pressure (yes vs no; greater than or equal to 140 mm Hg/90 mm Hg or current treatment with antihypertensive medication).

Deaths among study participants were ascertained via linkage with the national mortality register. For the present analysis, the follow-up time was restricted to 15 years for each cohort (ie, until 31 December 2000 for the 1985 cohort and until 31 December 2007 for the 1992 cohort). Cox proportional hazard regression was conducted to determine the association between education and all-cause mortality in each cohort in both sexes combined and separately for men and women. Age-adjusted models were estimated first, followed by multivariable models adjusted for covariates (age, BMI, height, cholesterol, high blood pressure,

smoking status and marital status). The HRs along with their corresponding 95% CIs were estimated using Cox regression. Terms of interaction between educational variables (one per model) and cohort were fitted to assess whether the educational gradient differed between cohorts. All analyses were performed in Stata software (V.16.1).

RESULTS

The descriptive statistics of the characteristics of the combined analytical sample and stratified by sex are reported in table 1. Over 15-year follow-up, there were 291 deaths in the 1985 cohort and 281 deaths in the 1992 cohorts. There were only minor differences in covariates between the two cohorts. However, the numbers and proportions of deaths during follow-up were substantially lower in the 1992 vs the 1985 cohort.

Table 2 presents the age-adjusted and fully adjusted HRs of all-cause mortality for both sexes combined and separately for men and women. For both sexes combined, the gradient in all-cause mortality by all educational indices was significant in age-adjusted models in both cohorts. Each additional year of schooling was associated with reduction in mortality risk by 6% (HR 0.94, 95% CI 0.90 to 0.98) and 7% (HR 0.93, 95% CI 0.90 to 0.97), respectively. After adjustment for covariates, the HRs were attenuated and only the association of years of schooling remained significant in both cohorts. When data were analysed separately by sex, a clear educational and statistically significant gradient was only seen in men. Crucially, the gradients, in both sexes, were similar between the cohorts, and there was no suggestion that the gradient became steeper in the 1992 cohort. None of the interactions between cohort and any educational indicators were statistically significant (all $p > 0.5$, not shown).

Table 1 Descriptive characteristics of the analytical sample, stratified by sex and cohort

	Men		Women	
	1985	1992	1985	1992
No of individuals	1237	1111	1293	1183
Age group				
25–34	24.4% (302)	21.8% (242)	24.7% (319)	21.6% (256)
35–44	23.4% (289)	31.3% (348)	26.1% (337)	29.5% (349)
45–54	27.2% (336)	27.1% (301)	25.7% (332)	25.9% (306)
55–64	25.1% (310)	19.8% (220)	23.6% (305)	23.0% (272)
Marital status				
Married	86.3% (1,068)	83.7% (930)	81.9% (1,059)	80.5% (952)
Single/widowed/divorced	13.7% (169)	16.3% (181)	18.1% (234)	19.5% (231)
Education				
Primary	63.3% (783)	61.8% (686)	68.1% (880)	58.7% (694)
Secondary	28.1% (347)	26.8% (298)	27.2% (351)	35.5% (420)
University	8.7% (107)	11.4% (127)	4.8% (62)	5.8% (69)
Years of schooling				
(Median)	12	11	11	11
High blood pressure	25.9% (320)	25.7% (286)	33.3% (430)	26.6% (315)
Ever smoking	44.8% (554)	39.5% (439)	23.8% (308)	23.3% (275)
BMI mean (SD)	27.04 (3.92)	27.09 (3.83)	27.31 (5.37)	26.87 (5.36)
Cholesterol mean (SD)	6.21 (1.28)	5.98 (1.28)	6.17 (1.24)	5.95 (1.28)
Height mean (SD)	1.74 (0.07)	1.75 (0.07)	1.61 (0.06)	1.62 (0.06)
HDL mean (SD)	1.35 (0.36)	1.32 (0.36)	1.57 (0.36)	1.52 (0.38)
Deaths during follow-up	15.0% (185)	16.5% (183)	8.2% (106)	8.3% (98)

BMI, body mass index; HDL, high-density lipoproteins.

Table 2 Age-adjusted and fully adjusted HRs for mortality by educational variables in 1985 and 1992

	Age adjusted		Fully adjusted*	
	1985	1992	1985	1992
Men and women combined (also adjusted for sex)				
Education				
Primary	1.0	1.0	1.0	1.0
Secondary	0.78 (0.59–1.04)	0.81 (0.62–1.07)	0.81 (0.60–1.08)	0.88 (0.66–1.17)
University	0.61 (0.35–1.05)	0.70 (0.44–1.10)	0.67 (0.39–1.17)	0.80 (0.51–1.27)
P value for trend	0.021	0.051	0.058	0.237
Years of schooling				
Lowest tertile	1.0	1.0	1.0	1.0
Medium	0.92 (0.70–1.19)	0.88 (0.65–1.20)	0.94 (0.72–1.23)	0.91 (0.67–1.23)
Highest tertile	0.65 (0.47–0.89)	0.67 (0.48–0.93)	0.71 (0.51–0.99)	0.76 (0.55–1.07)
P value for trend	0.011	0.018	0.055	0.115
Years of schooling (per year)				
P value	0.94 (0.90–0.98)	0.93 (0.90–0.97)	0.95 (0.91–0.99)	0.95 (0.91–0.99)
	0.007	0.001	0.028	0.019
Men				
Education				
Primary	1.0	1.0	1.0	1.0
Secondary	0.78 (0.55–1.10)	0.84 (0.60–1.18)	0.82 (0.58–1.16)	0.86 (0.61–1.22)
University	0.57 (0.31–1.06)	0.67 (0.40–1.12)	0.66 (0.35–1.22)	0.72 (0.43–1.20)
P value for trend	0.033	0.088	0.101	0.159
Years of schooling				
Lowest tertile	1.0	1.0	1.0	1.0
Medium	0.94 (0.67–1.30)	0.83 (0.56–1.23)	0.98 (0.70–1.37)	0.81 (0.55–1.20)
Highest tertile	0.61 (0.42–0.90)	0.66 (0.45–0.95)	0.70 (0.47–1.03)	0.72 (0.49–1.05)
P value for trend	0.016	0.025	0.090	0.068
Years of schooling (per year)				
P value	0.94 (0.89–0.99)	0.92 (0.88–0.97)	0.95 (0.91–1.00)	0.94 (0.89–0.98)
	0.012	0.001	0.058	0.008
Women				
Education				
Primary	1.0	1.0	1.0	1.0
Secondary	0.83 (0.49–1.39)	0.76 (0.47–1.23)	0.85 (0.50–1.45)	0.95 (0.58–1.55)
University	0.80 (0.25–2.54)	0.86 (0.31–2.36)	0.87 (0.27–2.80)	1.25 (0.45–3.49)
P value for trend	0.453	0.331	0.572	0.915
Years of schooling				
Lowest tertile	1.0	1.0	1.0	1.0
Medium	0.85 (0.55–1.32)	0.96 (0.60–1.55)	0.86 (0.55–1.34)	1.16 (0.71–1.90)
Highest tertile	0.82 (0.46–1.48)	0.69 (0.33–1.44)	0.88 (0.48–1.62)	0.93 (0.44–1.98)
P value for trend	0.408	0.376	0.552	0.890
Years of schooling (per year)				
P value	0.96 (0.89–1.05)	0.97 (0.89–1.05)	0.97 (0.89–1.06)	1.01 (0.93–1.10)
	0.367	0.460	0.514	0.799

*Adjusted for age, BMI, height, cholesterol, high blood pressure, smoking and marital status. BMI, body mass index.

DISCUSSION

The main objective of these analyses was to compare educational gradients in all-cause mortality in two cohorts established before and during the early stage of the postcommunist transformation in the Czech Republic. The results suggested that (1) educational gradients in mortality existed in both cohorts, (2) were pronounced only among men and (3) the gradients were of similar magnitude in both cohorts. These data, therefore, do not suggest a major divergence of educational inequalities of mortality.

Some of findings are unexpected. First, the lack of clear gradient in women was surprising, since official statistics have shown educational inequality in mortality in both sexes.¹³ The stronger association between educational level and mortality in men might be explained by differences in health behaviours among women and men (men

are more likely to follow unhealthy and risky behaviours) that might consequently contribute to higher specific mortality to, for example, accidents, lung cancer.¹⁹

Second, since the majority of published reports suggest an increase in social inequalities in mortality during the early stage of postcommunist transition,^{9 12 15} we expected a similar pattern in these cohorts. However, there was no evidence for the expected steeping of the gradient.

Previous studies in the Czech Republic generally found increase in educational differences in all-cause mortality, despite some exceptions between particular categories for both men and women.^{12 13 20} Increasing inequalities were also reported in Poland, Hungary, Estonia and Lithuania from 1990 to 2000,¹⁵ Russia from 1989 to 2001⁹ and Estonia from 1989 to 2000,¹⁴ indicating that the

postcommunist transition exacerbated educational discrepancies in mortality.

One potential explanation for the unexpected findings may be the numerator–denominator bias. A previous study conducted in Lithuania between 2001 and 2004 revealed an overestimation of the educational gradient in mortality in studies that used unlinked data rather than linked data, due to overreported deaths in lower educational groups and underreporting for those with higher education,¹⁰ although the extent of the bias seems dependent on the context and country.¹⁶ In addition, at least one prospective cohort²¹ and one retrospective study,⁹ both using individually linked observations, reported diverging mortality inequalities in Russia, suggesting that numerator–denominator bias is not a major alternative explanation.

A second potential reason for our findings may be the lower response rate in the 1992 vs 1985 survey (73% vs 84%). Given this large difference and the literature consistently suggesting that non-respondents tend to be less healthy and more likely socially disadvantaged than respondents, the potential changes in educational gradients between cohorts may be obscured by differential response rate.²² Lower response rate may partly explain the lower mortality risk in the second study.

Third, the observation period between the two compared cohorts (1985 vs 1992) might be too short (7 years) to distinguish the potential effect of the educational gradient in all-cause mortality. In order to obtain a sufficient number of deaths for the analysis, a 15-year follow-up was used and the calendar years during follow-up, therefore, partially overlap between the cohorts. Restricting analyses to non-overlapping first 5 years of follow-up reduced the number of deaths in the analyses and did not provide any additional clues.

Finally, the sample sizes of the cohorts may have been too small to detect modest changes in associations between education and mortality between cohorts; this is particularly pertinent to women, among whom the lower deaths rates further reduce the statistical power.

CONCLUSION

Notwithstanding these limitations, this first individual-based prospective investigation outside of the former Soviet Union found evidence for an educational gradient in mortality in both cohorts, which was largely driven by gradients among men. However, we found no evidence for the gradient becoming steeper in the later cohort. These findings warrant re-examination of this important question in larger individual-based datasets.

Correction notice This article has been corrected since it first published. In the last row of table 1, 'Deaths during follow-up', data has been corrected. This data has also been corrected in the main text.

Acknowledgements We are most grateful for all the participants involved in the MONICA project and the study members who participated in data collection. Authors thank the RECETOX Research Infrastructure (No LM2018121) financed by the Ministry of Education, Youth and Sports, and the Operational Programme Research, Development and Education (the CETOCOEN EXCELLENCE project No. CZ.02.1.01/0.0/0.0/17_043/0009632) for supportive background.

Contributors CP and AD contributed equally to this paper. CP, MB and HP conceived and designed the study. CP and WL analysed the data, and CP wrote the first draft of the manuscript. AD and MB contributed to writing of drafts. JH contributed on designing of the overall research study. All authors provided critical revisions. All authors read and approved the submitted manuscript.

Funding This work has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 857487, No 857560 and No 857340 and funding from the Czech NPO Systemic Risk Institute, No LX22NPO5101. The Czech MONICA study was funded by the Czech Ministry of Health through institutional support for the Institute of Clinical and Experimental

Medicine. This publication reflects only the author's view and the European Commission is not responsible for any use that may be made of the information it contains.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval The study was approved by the ethics committee at University College London, UK (99/0081) and the ethics committee at Institute of Clinical and Experimental Medicine, Prague, Czech Republic.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Andrea Dalecká <http://orcid.org/0000-0002-9808-7444>

Jaroslav Hubáček <http://orcid.org/0000-0001-6537-1353>

REFERENCES

- Lleras-Muney A. The relationship between education and adult mortality in the United States. *Rev Econ Stud* 2005;72:189–221.
- Son M, Armstrong B, Choi J-M, et al. Relation of occupational class and education with mortality in Korea. *J Epidemiol Community Health* 2002;56:798–9.
- Kohler IV, Martikainen P, Smith KP, et al. Educational differences in all-cause mortality by marital status - Evidence from Bulgaria, Finland and the United States. *Demogr Res* 2008;19:2011–42 <https://doi.org/10.4054/demres.2008.19.60>
- Mackenbach JP, Stirbu I, Roskam A-JR, et al. Socioeconomic inequalities in health in 22 European countries. *N Engl J Med* 2008;358:2468–81.
- Montez JK, Berkman LF. Trends in the educational gradient of mortality among US adults aged 45 to 84 years: bringing regional context into the explanation. *Am J Public Health* 2014;104:e82–90.
- UNICEF. A decade of transition. In: *Reg Monit Rep*. 8, 2001.
- Cornia GA. Labor market shocks, psychosocial stress and the transition's mortality crisis. *Research in Progress* 1996;4:Helsinki: UNU/WIDER.
- Simonová N, Antonowicz D. Czech and polish higher education - from bureaucracy to market competition. *Czech Sociol Rev* 2006;42:517–36.
- Murphy M, Bobak M, Nicholson A, et al. The widening gap in mortality by educational level in the Russian federation, 1980-2001. *Am J Public Health* 2006;96:1293–9.
- Shkolnikov VM, Jasilionis D, Andreev EM, et al. Linked versus unlinked estimates of mortality and length of life by education and marital status: evidence from the first record linkage study in Lithuania. *Soc Sci Med* 2007;64:1392–406.
- Shkolnikov VM, Andreev EM, Jasilionis D, et al. The changing relation between education and life expectancy in central and eastern Europe in the 1990s. *J Epidemiol Community Health* 2006;60:875–81.
- Dzúrová D. Mortality differentials in the Czech Republic during the post-1989 socio-political transformation. *Health Place* 2000;6:351–62.
- Rychtaříková J. The case of the Czech Republic. determinants of the recent favourable turnover in mortality. *Demogr Res* 2004;105:38.
- Leinsalu M, Vägerö D, Estonia KAE. Enormous increase in mortality differences by education. *Int J Epidemiol* 1989;2003:1081–7.
- Leinsalu M, Stirbu I, Vägerö D, et al. Educational inequalities in mortality in four eastern European countries: divergence in trends during the post-communist transition from 1990 to 2000. *Int J Epidemiol* 2009;38:512–25.
- Jasilionis D, Leinsalu M. Changing effect of the numerator-denominator bias in unlinked data on mortality differentials by education: evidence from Estonia 2000-2015. *J Epidemiol Community Health* 2021;75:88–91.
- World Health Organization. The world Health organization MONICA Project (monitoring trends and determinants in cardiovascular disease): a major international collaboration. *J Clin Epidemiol* 1988;41:105–14.
- Bobak M, Skodova Z, Pisa Z, et al. Political changes and trends in cardiovascular risk factors in the Czech Republic, 1985-92. *J Epidemiol Community Health* 1997;51:272–7.
- Luy M, Di Giulio P. The impact of health behaviors and life quality on gender differences in mortality. *Germany Max Planck Inst Demogr Res*; 2006: 0–36.
- Dragano N, Wahrendorf M. Consistent health inequalities in Europe: the importance of labour market disadvantage. *J Epidemiol Community Health* 2014;68:293–4.
- Plavinski SL, Plavinskaya SI, Klimov AN. Social factors and increase in mortality in Russia in the 1990s: prospective cohort study. *BMJ* 2003;326:1240–2.
- Cheung KL, ten Klooster PM, Smit C, et al. The impact of non-response bias due to sampling in public health studies: a comparison of voluntary versus mandatory recruitment in a Dutch national survey on adolescent health. *BMC Public Health* 2017;17:276.