



REVIEW

Open Access

Evolving Diagnostic and Treatment Strategies for Pancreatic Neuroendocrine Tumors

Matthew H Kulke^{1*}, Johanna Bendell², Larry Kvols³, Joel Picus⁴, Rodney Pommier⁵ and James Yao⁶

Abstract

Pancreatic neuroendocrine tumors (NET) have diverse clinical presentations. Patients with symptoms of hormone secretion may require specific medical interventions to control those symptoms prior to antitumor intervention. In some patients, tumors in the pancreas may be occult and specialized diagnostic imaging or surgery may be required for diagnosis. Other patients may present with more advanced disease, presenting with symptoms of tumor bulk rather than hormone secretion. Treatment options for patients with advanced pancreatic neuroendocrine tumors include surgical resection and hepatic directed therapies, including partial hepatectomy, hepatic artery embolization, or other ablative techniques. Streptozocin or temozolomide-based chemotherapy regimens are active against pancreatic NET, and can also play an important role in the palliation of patients with advanced disease. A number of biologically targeted agents targeting the VEGF and mTOR signaling pathways have recently shown promise, with recent trials showing treatment with the VEGFR tyrosine kinase inhibitor sunitinib or the mTOR inhibitor everolimus improves progression-free survival in patients with advanced NET.

Introduction

Pancreatic neuroendocrine tumors (NET) have been considered rare, with an estimated incidence of less than 1 per 100,000 individuals [1]. In recent years, however, the diagnosed incidence of pancreatic NET has increased, an observation that is likely due, at least in part, to improved detection and classification [2]. The diverse and sometimes non-specific clinical syndromes associated with pancreatic NET can make these malignancies difficult to diagnose at an early stage. Awareness of the clinical presentation and treatment options for patients with pancreatic NET has become increasingly relevant for both medical oncologists and other health care providers, as new treatment options emerge for patients with this disease.

Histologic Classification and Staging

Pancreatic NET have also been referred to as pancreatic islet cell tumors or pancreatic endocrine tumors. Carcinoid tumors have a similar histologic appearance to pancreatic NET, but generally arise in the bronchi, small intestine, appendix, or rectum. While the term "pancreatic carcinoid" has also sometimes been used to describe

pancreatic NET, this term is considered confusing as the clinical presentation and treatment options for pancreatic NET differ in many respects from those for carcinoid tumors.

The majority of pancreatic NET occur sporadically. However, pancreatic NET can be associated with inherited genetic syndromes; in particular, approximately 10% may be associated with multiple endocrine neoplasia type 1 (MEN1). MEN1 is an autosomal dominant syndrome associated with mutations in the tumor suppressor gene *menin*, and is characterized by the development of multiple NET involving not only the pancreas but also the parathyroid and pituitary glands [3]. Pancreatic NET have also been associated with MEN2, Von Hippel-Lindau disease, tuberous sclerosis, and neurofibromatosis.

The histologic features of pancreatic NET can vary, affecting both prognosis and treatment recommendations. An important first step following the diagnosis of a pancreatic malignancy is the differentiation of neuroendocrine cancers from the far more common pancreatic adenocarcinoma. Though the pathologic criteria for differentiating these two entities are clear, limited tissue from fine needle aspirations or endoscopic brushings may preclude accurate diagnosis. In questionable cases, repeat tissue sampling should be performed, particularly if systemic treatments are being considered.

* Correspondence: matthew_kulke@dfci.harvard.edu

¹Dana-Farber Cancer Institute, Boston MA, USA

Full list of author information is available at the end of the article

Adequate tissue sampling is also critical in differentiating the various subtypes of pancreatic NET. These tumors may fall within a broad spectrum of well-differentiated, low grade tumors to more poorly differentiated, high grade tumors. While a number of histologic classification systems have been proposed for pancreatic NET, tumors with a mitotic count $>20/10$ high powered fields or a Ki-67 proliferation index of $>20\%$ generally represent highly aggressive malignancies where treatment with platinum based regimens is generally indicated, according to small cell carcinoma guidelines [4,5].

The American Joint Committee on Cancer (AJCC) staging system for pancreatic NET is increasingly accepted as the standard staging system in North America, and is similar to the system used for pancreatic adenocarcinomas. Several other organizations, including both North-American based groups and the European Neuroendocrine Tumor Society (ENETS) have proposed similar, though not identical, staging systems for NET using the commonly accepted Tumor Node Metastasis (TNM) notation [6-10].

Clinical Presentation and Initial Management

Most pancreatic NET are considered “non-functional” in that they are not associated with symptoms of hormone hypersecretion. Such tumors are usually identified incidentally during imaging for other indications, or at an advanced stage, when patients become symptomatic from tumor bulk. Patients with hormonal hypersecretion, on the other hand, can present with diverse and sometimes puzzling clinical symptoms (Table 1). Specific recommendations for some of the more common tumors, based on the clinical presentation and hormones secreted, are described below.

Insulinoma

Insulinomas classically present with “Whipple’s Triad:” a combination of symptoms of hypoglycemia, inappropriately

high insulin levels with associated documented blood glucose levels of <50 mg/dL, and symptom relief with administration of glucose [11]. Initially, the hypoglycemia may be managed with dietary modifications or with diazoxide [12]. For these patients, octreotide or other somatostatin analogs should be used with caution, as they have the potential to worsen hypoglycemia by suppressing glucagon secretion. Treatment with the mTOR inhibitor everolimus has also been reported to be beneficial in insulinoma patients with refractory hypoglycemia [13].

Glucagonoma

Over two-thirds of patients with glucagonomas present with necrolytic migratory erythema, a rash characterized by raised erythematous patches beginning in the perineum and progressing to the trunk and extremities [14,15]. Somatostatin analogs are generally successful in the initial management of patients with the glucagonoma syndrome [16,17]. Glucagonomas may be associated with diabetes mellitus, though only half of patients experience clinically significant hyperglycemia.

Gastrinoma and Zollinger-Ellison syndrome

The gastrinoma syndrome is characterized by gastric hypersecretion [18]. In patients with non-healing peptic ulcers and a fasting gastrin level >100 pg/mL, a diagnosis of gastrinoma should be considered [19]. Moderate elevations of serum gastrin may also, however, be seen in patients receiving concomitant therapy with proton pump inhibitors, sometimes complicating efforts to confirm a diagnosis. Proton pump inhibitors are a highly effective initial treatment in controlling symptoms associated with gastric hypersecretion [20,21]. Treatment with somatostatin analogs has also been associated with improved control of serum gastrin levels and, in some cases, with tumor stabilization or regression [22].

Table 1 Clinical presentation of pancreatic neuroendocrine tumors (NET)

Tumor	Symptoms or signs	Incidence of metastases	Extrapancreatic location
Insulinoma	Hypoglycemia resulting in intermittent confusion, sweating, weakness, nausea; loss of consciousness may occur in severe cases	<15%	Rare
Glucagonoma	Rash (necrotizing migratory erythema), cachexia, diabetes, deep venous thrombosis	Majority	Rare
VIPoma, Verner-Morrison Syndrome, WDHA Syndrome	Profound secretory diarrhea, electrolyte disturbances	Majority	10%
Gastrinoma, Zollinger-Ellison Syndrome	Acid hypersecretion resulting in refractory peptic ulcer disease, abdominal pain, and diarrhea	<50%	Frequently in duodenum
Somatostatinoma	Diabetes, diarrhea, cholelithiasis	Majority	Rare
Non-functioning	May be first diagnosed due to mass effect	Majority	Rare

WDHA: Watery Diarrhea, Hypokalemia and Achlorhydria.

VIPoma

Pancreatic endocrine tumors associated with profound diarrhea, hypokalemia, and achlorhydria were first described by Verner and Morrison in 1958 [23]. This syndrome was subsequently found to be due to ectopic vasoactive intestinal peptide (VIP) secretion. Treatment with somatostatin analogs is effective in treatment of diarrhea in these patients [24].

Imaging

Patients with functioning tumors, particularly insulinomas and gastrinomas, may develop hormonal symptoms from small primary tumors, and localization of the primary lesions may be challenging. Traditional cross-sectional imaging with triple phase CT or MRI is generally the first step in attempting to localize these tumors. Endoscopic ultrasound may be more sensitive than CT or MRI for the detection of small lesions, and may also provide useful information regarding potential vessel involvement prior to planned resection. Pancreatic NET, like carcinoid tumors, frequently over express somatostatin receptors. ¹¹¹Indium-DTPA-octreotide (Octreoscan™) has been commonly used, often in combination with cross-sectional imaging, to localize and stage pancreatic NET.

Biochemical Assessment and Monitoring

In patients with symptoms of hormone hypersecretion, serial measurements of the specific hormone may be helpful in assessing treatment response or in monitoring for recurrence. The majority of patients with pancreatic NET, however, do not have clear evidence of hormone hypersecretion. Serum chromogranin A (CGA) is a neuroendocrine secretory protein that serves as a marker of disease activity in both functional and non-functional pancreatic NET [1,25-27]. CGA may decrease in patients responding to somatostatin analogs or other therapies [26,28]. In patients on stable SSA doses, consistent increases in plasma CGA levels over time may reflect loss of secretory control and/or tumor growth [25-27,29,30]. Use of CGA as a diagnostic or screening test for pancreatic NET is discouraged, as CGA may be elevated in a number of non-malignant conditions, including renal insufficiency and liver disease, and in patients taking proton pump inhibitors.

Surgical Management

In general, in the absence of distant metastases or significant comorbidities, complete surgical resection of the primary tumor should be attempted [31,32]. The primary tumors in patients who are diagnosed due to symptoms of hormone hypersecretion may be occult (Figure 1A). In contrast, patients with non-functioning pancreatic NET are commonly diagnosed at a later stage




Figure 1 Variations in size of primary pancreatic NET.

A: Insulinoma presenting as an occult nodule on the pancreas. Functional pancreatic NET may present at an early stage, and identification of the primary tumor may be challenging. B: Non-functioning pancreatic NET in the tail of the pancreas with associated hepatic metastases. Non-functioning pancreatic NET more typically present at a later stage, often as bulky lesions in the tail of the pancreas with associated liver metastases.

(Figure 1B). The prognosis following surgical resection of localized NET is often excellent. Isolated insulinomas, for example, are generally treated with enucleation; long-term survival following surgery in this patient population exceeds 90% [33]. The role of surgical resection in patients with MEN1 syndrome remains more controversial because of the risk of additional tumors within the remaining pancreas and elsewhere [34,35].

In contrast to patients with pancreatic adenocarcinoma, hepatic resection may be beneficial in patients with metastatic pancreatic NET. Resection may be

performed to render the patient free of macroscopic disease and to diminish systemic symptoms. Hepatic resection is generally favored in patients with limited hepatic disease. In a study of 170 patients, hepatic resection improved symptoms in over 90% of cases [36]. Debulking surgery in patients with more advanced disease may be recommended in selected patients if the majority (i.e., >90%) of the tumor burden can be resected [37]. The median survival for patients treated with this approach has been reported to be 7 years [38]. The reported survival rates for this surgical approach have been in excess of 60% at 5 years, which is twice that of patients with untreated liver metastases [38,39]. An important observation is that the survival of patients who have palliative hepatic debulking by 90% is indistinguishable from those who have complete resection (i.e., resection of all visible hepatic tumors) [39]. Also, the survival with this approach is the same among patients with functional and non-functional tumors [39]. Thus, surgical resection should be considered to potentially improve outcomes even though surgery may not be curative.

Liver-directed Therapy

In patients with hepatic metastases who are not candidates for surgical resection, hepatic arterial embolization may be an appropriate palliative technique, provided that their disease is primarily confined to the liver, the portal vein is patent, they have not undergone a Whipple procedure (or pancreaticoduodenectomy), and that patients have an otherwise preserved performance status [40-42]. Response rates are generally >50%, whether measured by reduced hormone secretion or radiographic regression [40-43]. A variety of techniques have been employed, including bland embolization, chemo-embolization, embolization with chemotherapy eluting beads, or embolization using radioisotopes. There are currently no data confirming superiority of any one of these techniques over the others.

Hepatic metastases can also be treated with percutaneous or laparoscopic radiofrequency ablation (RFA) and cryoablation, either alone or in conjunction with surgical debulking [40,42]. While these approaches appear to cause less morbidity than either hepatic resection or hepatic artery embolization, the clinical benefit of this approach in patients with asymptomatic, small-volume disease has not been clearly established. Ablative techniques should be considered only in carefully selected patients [40,42,44,45].

Cytotoxic Chemotherapy

Although carcinoid and pancreatic NET appear histologically similar, there is increasing evidence that pancreatic NET are more responsive to chemotherapy than are carcinoid tumors (Table 2). In an initial randomized

trial, the combination of streptozocin and doxorubicin was associated with a combined biochemical and radiologic regression of 69%, as well as a survival benefit when compared to streptozocin and fluorouracil [46]. The median overall survival duration for patients treated with streptozocin and doxorubicin was 2.2 years. Streptozocin was subsequently approved by the FDA as a treatment for patients with pancreatic NET. The very high response rates reported in this study were derived in part from the historical use of non-standard response criteria. A large retrospective analysis of 84 patients with either locally advanced or metastatic pancreatic endocrine tumors showed that a three-drug regimen of streptozocin, 5-FU, and doxorubicin was associated with an overall response rate of 39% and a median survival duration of 37 months [47].

Despite the demonstrated efficacy of streptozocin-based regimens, their potential toxicity, together with a cumbersome 5 consecutive day infusion schedule, has precluded their more widespread use in patients with advanced pancreatic NET. Recent prospective and retrospective studies have suggested that oral temozolomide-based regimens may be at least comparable in efficacy to streptozocin-based regimens, and may also be more tolerable. In retrospective series, temozolomide-based therapy has been associated with overall response rates of 8-70% [48-50] (Table 2). Temozolomide has been evaluated prospectively in combination with thalidomide, bevacizumab, or everolimus, with overall response rates of 24-45% [51-53]. Neither the optimal dosing regimen for temozolomide, nor the relative activity of temozolomide as a single agent or in combination with other therapeutic agents has been clearly established.

Somatostatin Analogs and Peptide Receptor Radiation Therapy (PRRT)

More than 90% of NET express somatostatin receptors, and somatostatin analogs (SSAs) have been shown to be effective in reducing symptoms of hormone hypersecretion associated with both carcinoid and pancreatic NET. In patients with midgut carcinoid tumors, treatment with the somatostatin analog octreotide has been shown to improve time to tumor progression over placebo. Ongoing studies are currently exploring whether somatostatin analogs have a similar antiproliferative effect in patients with pancreatic NET.

The high rate of somatostatin receptor expression in pancreatic NET also provides a rationale for peptide receptor radionuclide therapy in patients with inoperable or metastatic disease. The most frequently used radionuclides for targeted radiotherapy in NET are yttrium (⁹⁰Y), and lutetium (¹⁷⁷Lu), which differ from one another in terms of emitted particles, particle energy, and tissue penetration [54,55]. Both the yttrium

Table 2 Selected Trials of Cytotoxic Chemotherapy in Advanced Pancreatic NET

Regimen	Patients	Tumor Response Rate (%)	Median Progression-Free Survival	Median Overall Survival (Months)	Reference
Prospective Studies					
Chlorozotocin	33	30	17*	18.0	Moertel et al. 1992 [46]
Fluorouracil + Streptozocin	33	45	14*	16.8	Moertel et al. 1992
Doxorubicin + Streptozocin	36	69	18*	26.4	Moertel et al. 1992
DTIC	50	34	NR	19.3	Ramanathan et al. 2001 [67]
Temozolomide+ Thalidomide	11	45	NR	NR	Kulke et al. 2006 [52]
Temozolomide+ Bevacizumab	17	24	8.6	NR	Kulke et al. 2006 [51]
Temozolomide+ Everolimus	24	35	NR	NR	Kulke et al. 2010 [53]
Retrospective Studies					
Streptozocin+ Doxorubicin+ Fluorouracil	84	39	18	37	Kouvaraki et al. 2004 [47]
Temozolomide (diverse regimens)	53	34	13.6	35.3	Kulke et al. 2009 [49]
Temozolomide (single agent)	12	8	NR	NR	Ekeblad et al. 2007 [48]
Temozolomide+ Capecitabine	30	70	18	NR	Strosberg et al. 2010 [50]

NR: Not reported.

and the lutetium labeled compounds have demonstrated promising activity in NET patients. The radiolabeled somatostatin analog [¹⁷⁷Lu-DOTA, Tyr³] octreotate, for example, has been utilized in the treatment of 504 patients with NET, and efficacy results, reported for 310 patients, suggest single agent activity [56]. Treatment with ⁹⁰Y-DOTA tyr3-octreotide (⁹⁰Y-edotreotide) was recently reported to be associated with high rates of symptom control, though only modest tumor response rates, in a prospective, phase II study [57]. Randomized studies comparing PRRT to treatment with “cold” octreotide are anticipated to better define the relative efficacy and toxicities associated with these regimens.

Biologically Targeted Therapies for Pancreatic NET

Studies of biologically targeted therapies in pancreatic NET have, to date, focused primarily on inhibitors of the VEGF or mTOR signaling pathways. While objective RECIST-defined tumor response rates have been relatively low, recent studies have suggested that treatment with these agents is associated with improvements in progression-free survival.

VEGF pathway inhibitors

Three tyrosine kinase inhibitors—pazopanib, sorafenib, and sunitinib—all with activity against VEGFR, have been evaluated in prospective trials of patients with advanced pancreatic NET. Pazopanib was evaluated in a prospective study enrolling 51 NET patients (29 with pancreatic NET and 22 with carcinoid) on stable doses of octreotide-LAR. Patients received pazopanib at a dose of 800 mg daily. The response rate among patients with pancreatic NET was 17%; no patients with carcinoid experienced a radiographic response (by RECIST) [58]. Sorafenib is another small molecule tyrosine kinase inhibitor with activity against VEGFR. In a study of 50 patients with carcinoid and 43 patients with pancreatic NET, preliminary analysis showed responses in 7% of the carcinoid patients and 11% of the pancreatic NET patients [59].

Sunitinib malate was evaluated in a multi-institutional phase II study enrolling 109 patients with advanced NET. Patients received repeated 6-week treatment cycles of sunitinib, administered orally at 50 mg once daily for 4 weeks, followed by 2 weeks off treatment [60]. Partial responses were observed in 2% of the carcinoid cohort

and 16% of the pancreatic NET cohort. Based on evidence of activity in this study, an international randomized phase III study to confirm the activity of sunitinib in pancreatic NET was undertaken. The study was discontinued prior to a planned interim analysis after enrollment of 171 patients, 86 of whom received sunitinib and 85 of whom received placebo. The early discontinuation of the study precluded definitive hypothesis testing for differences in progression-free survival durations between the treatment and placebo groups. Nevertheless, analysis of the available data demonstrated that treatment with sunitinib was associated with a median progression-free survival (PFS) of 11.4 months, as compared with 5.5 months for placebo ($P=.0001$, Table 3) [61].

mTOR Inhibitors

Tumor cell growth, proliferation, and apoptosis are regulated in part by a serine-threonine kinase called the mammalian target of rapamycin (mTOR). This enzyme also mediates downstream signaling from a number of pathways, including the VEGF and insulin-like growth factor (IGF) signaling implicated in NET growth. Temsirolimus and everolimus are rapamycin derivatives that have been evaluated recently in NET. Weekly intravenous administration of temsirolimus was associated with a response rate of 5.6% in one study of 37 patients with advanced progressive NET. Outcomes were similar between patients with carcinoid and pancreatic NET [62].

Everolimus was initially evaluated in a single-institution study, in which 30 patients with carcinoid tumors and 30 with pancreatic NET received doses of 5 or 10 mg daily plus depot octreotide (30 mg every 4 weeks). The overall tumor response rate in evaluable patients was 17% in carcinoid and 27% in pancreatic NET [63]. A follow-up multinational phase II study (RADIANT-1) enrolled 160 patients with advanced pancreatic NET and evidence of RECIST-defined progression following chemotherapy. In this non-randomized study, treatment with everolimus was associated with an overall response rate of 4.4% and progression-free survival duration of 16.7 months in those patients receiving octreotide. Among patients not receiving octreotide, the response

rate was 9.6% and the progression-free survival duration was 9.7 months [64]. A subsequent phase III study randomized 410 patients with progressive advanced pancreatic NET (RADIANT-3) to receive treatment with everolimus or placebo; octreotide was given at the discretion of the investigator. This study demonstrated significant improvements in the primary endpoint of investigator-assess PFS associated with everolimus as compared to placebo (11 months versus 4.6 months, [$P<.0001$, Table 3]) [65]. The overall tumor response rate associated with everolimus in this study was 5%.

Ongoing studies are currently evaluating combinations of targeted agents in patients with advanced pancreatic NET. A combination of the mTOR inhibitor everolimus and the VEGF inhibitor bevacizumab was recently shown to be well tolerated and associated with antitumor activity (overall response rate 26%) in an initial phase II study enrolling patients with low or intermediate grade NET [66]. CALGB 80701 is currently randomizing patients with advanced pancreatic NET to receive either treatment with everolimus or everolimus + bevacizumab to assess the relative efficacy and toxicity of these regimens (Table 3).

Conclusions

Patients with pancreatic NET present with diverse symptoms related to hormonal hypersecretion, tumor bulk, or both. Accurate diagnosis of this condition and differentiation of pancreatic NET from the more common pancreatic adenocarcinomas is a critical first step in developing an appropriate treatment plan. Similarly, pancreatic NET should be considered separately from carcinoid tumors, which arise in other sites. Surgical resection remains the mainstay of treatment for patients with localized disease. A number of treatment options are available for patients with advanced pancreatic NET. These include hepatic-directed therapies, including surgical resection and hepatic artery embolization. Systemic treatment options include the use of SSAs for control of hormonal hypersecretion, as well as alkylating chemotherapy. Recent studies have also reported that the tyrosine kinase inhibitor sunitinib and the mTOR inhibitor everolimus improved progression-free survival in patients with pancreatic NET, further

Table 3 Randomized Trials of Biologically Targeted Therapies in Pancreatic NET

Regimen	N (total)	Overall Response Rate	Median Progression-Free Survival/TPP	P value	Reference
Sunitinib (37.5 mg po qd) Placebo (+ best supportive care)	171	9% 0%	11.4 months 5.5 months	.0001	Raymond et al, 2011 [61]
Everolimus (10 mg po qd) Placebo (+ best supportive care)	410	5% 2%	11 months 4.6 months	<.0001	Yao et al, 2011 [65]
Everolimus (10 mg po qd) Everolimus (10 mg po qd) + Bevacizumab (10 mg/kg every other week)	CALGB 80701	(Ongoing)			

expanding the therapeutic arsenal available to patients with this disease. Future studies will likely build on these results, further improving therapeutic options for patients with this disease.

Acknowledgements

The authors thank Lindy Morde (Dana-Farber Cancer Institute) for providing editorial assistance. Medical writing support was also provided by Susanne Gilbert and Keith Lantz (ACUMED®, New York, USA) and was funded by Pfizer Inc.

Author details

¹Dana-Farber Cancer Institute, Boston MA, USA. ²Sarah Cannon Research Institute, Nashville, TN, USA. ³H Lee Moffitt Cancer Center, Tampa FL, USA.

⁴Siteman Cancer Center, St Louis MO, USA. ⁵Oregon Health and Science University, Portland OR, USA. ⁶MD Anderson Cancer Center, Houston TX, USA.

Authors' contributions

All authors were involved in drafting the manuscript and revising it critically for important intellectual content. All authors have also read and approved the final version of the manuscript to be published.

Competing interests

MK has served as a consultant for Pfizer, Novartis, Ipsen, Lexicon Pharmaceuticals, and Molecular Insight Pharmaceuticals. JB has no competing interests. LK has served as a consultant and/or received honorarium from Novartis, Pfizer, and Delcath. JP has received research funding and honorarium, and served as a consultant and as a speaker for Novartis and Pfizer. JY has served as a consultant for Ipsen, Novartis, and Pfizer.

Received: 31 March 2011 Accepted: 14 June 2011

Published: 14 June 2011

References

- Metz DC, Jensen RT: Gastrointestinal neuroendocrine tumors: pancreatic endocrine tumors. *Gastroenterology* 2008, **135**:1469-1492.
- Yao JC, Hassan M, Phan A, Dagohoy C, Leary C, Mares JE, Abdalla EK, Fleming JB, Vauthhey JN, Rashid A, Evans DB: One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol* 2008, **26**:3063-3072.
- Agarwal SK, Lee BA, Sukhodolets KE, Kennedy PA, Obungu VH, Hickman AB, Mullendore ME, Whitten I, Skarulis MC, Simonds WF, et al: Molecular pathology of the MEN1 gene. *Ann N Y Acad Sci* 2004, **1014**:189-198.
- Kloppel G, Perren A, Heitz PU: The gastroenteropancreatic neuroendocrine cell system and its tumors: the WHO classification. *Ann N Y Acad Sci* 2004, **1014**:13-27.
- Rindi G, Kloppel G: Endocrine tumors of the gut and pancreas tumor biology and classification. *Neuroendocrinology* 2004, **80**(Suppl 1):12-15.
- Klimstra DS, Modlin IR, Adsay NV, Chetty R, Deshpande V, Gonem M, Jensen RT, Kidd M, Kulke MH, Lloyd RV, Moran C, Moss SF, Oberg K, O'Toole D, Rindi G, Robert ME, Suster S, Tang LH, Tzen CY, Washington MK, Wiedenmann B, Yao J: Pathology reporting of neuroendocrine tumors: application of the Delphic consensus process to the development of a minimum pathology data set. *Am J Surg Pathol* 2010, **34**:300-313.
- Klimstra DS, Modlin IR, Coppola D, Lloyd RV, Suster S: The pathologic classification of neuroendocrine tumors: a review of nomenclature, grading, and staging systems. *Pancreas* 2010, **39**:707-712.
- Rindi G, Kloppel G, Alhman H, Caplin M, Couvelard A, de Herder WW, Eriksson B, Falchetti A, Falconi M, Komminoth P, Korner M, Lopes JM, McNicol AM, Nilsson O, Perren A, Scarpa A, Scoazec JY, Wiedenmann B, All of the Frascati concensus conference participants European Neuroendocrine Tumour Society (ENTs): TNM staging of foregut (neuro) endocrine tumors: a consensus proposal including a grading system. *Virchows Arch* 2006, **449**:395-401.
- Rindi G, Kloppel G, Couvelard A, Komminoth P, Korner M, Lopes JM, McNicol AM, Nilsson O, Perren A, Scarpa A, Scoazec JY, Wiedenmann B: TNM staging of midgut and hindgut (neuro) endocrine tumors: a consensus proposal including a grading system. *Virchows Arch* 2007, **451**:757-762.
- Edge SB, Compton CC: The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM. *Ann Surg Oncol* 2010, **17**:1471-1474.
- Whipple AO, Frantz VK: Adenoma of islet cells with hyperinsulinism: a review. *Ann Surg* 1935, **101**:1299-1335.
- Goode PN, Farndon JR, Anderson J, Johnston ID, Morte JA: Diazoxide in the management of patients with insulinoma. *World J Surg* 1986, **10**:586-592.
- Kulke MH, Bergstrand EK, Yao JC: Glycemic control in patients with insulinoma treated with everolimus. *N Engl J Med* 2009, **360**:195-197.
- Horrobin DF, Cunnane SC: Interactions between zinc, essential fatty acids and prostaglandins: relevance to acrodermatitis enteropathica, total parenteral nutrition, the glucagonoma syndrome, diabetes, anorexia nervosa and sickle cell anaemia. *Med Hypotheses* 1980, **6**:277-296.
- Roth E, Muhlbacher F, Karner J, Hamilton G, Funovics J: Free amino acid levels in muscle and liver of a patient with glucagonoma syndrome. *Metabolism* 1987, **36**:7-13.
- El RZ, Partensky C, Valette PJ, Berger F, Chayvialle JA: Necrotolytic migratory erythema, first symptom of a malignant glucagonoma: treatment by long-acting somatostatin and surgical resection. Report of three cases. *Eur J Surg Oncol* 1998, **24**:562-567.
- Jockenhövel F, Lederbogen S, Olbricht T, Schmidt-Gayk H, Krenning EP, Lamberts SW, Reinwein D: The long-acting somatostatin analogue octreotide alleviates symptoms by reducing posttranslational conversion of prepro-glucagon to glucagon in a patient with malignant glucagonoma, but does not prevent tumor growth. *Clin Investig* 1994, **72**:127-133.
- Zollinger RM, Ellison EH: Primary peptic ulcerations of the jejunum associated with islet cell tumors of the pancreas. *Ann Surg* 1955, **142**:709-723.
- Jensen RT: Gastrointestinal endocrine tumours. Gastrinoma. *Baillieres Clin Gastroenterol* 1996, **10**:603-643.
- Lambers CB, Lind T, Moberg S, Jansen JB, Olbe L: Omeprazole in Zollinger-Ellison syndrome. Effects of a single dose and of long-term treatment in patients resistant to histamine H₂-receptor antagonists. *N Engl J Med* 1984, **310**:758-761.
- Frucht H, Maton PN, Jensen RT: Use of omeprazole in patients with Zollinger-Ellison syndrome. *Dig Dis Sci* 1991, **36**:394-404.
- Shojaamanesh H, Gibril F, Louie A, Ojeaburu JV, Bashir S, bou-Saif A, Jensen RT: Prospective study of the antitumor efficacy of long-term octreotide treatment in patients with progressive metastatic gastrinoma. *Cancer* 2002, **94**:331-343.
- Verner JV, Morrison AB: Islet cell tumor and a syndrome of refractory watery diarrhea and hypokalemia. *Am J Med* 1958, **25**:374-380.
- Kraenzlin ME, Ch'ng JL, Wood SM, Carr DH, Bloom SR: Long-term treatment of a VIPoma with somatostatin analogue resulting in remission of symptoms and possible shrinkage of metastases. *Gastroenterology* 1985, **88**:185-187.
- de Herder WW: Biochemistry of neuroendocrine tumours. *Best Pract Res Clin Endocrinol Metab* 2007, **21**:33-41.
- Oberg K, Eriksson B: Endocrine tumours of the pancreas. *Best Pract Res Clin Gastroenterol* 2005, **19**:753-781.
- Ardilli JE: Circulating markers for endocrine tumours of the gastroenteropancreatic tract. *Ann Clin Biochem* 2008, **45**:539-559.
- Oberg K, Kvols L, Caplin M, Delle FG, de HW, Rindi G, Ruszniewski P, Woltering EA, Wiedenmann B: Consensus report on the use of somatostatin analogs for the management of neuroendocrine tumors of the gastroenteropancreatic system. *Ann Oncol* 2004, **15**:966-973.
- Kann PH, Balakina E, Ivan D, Bartsch DK, Meyer S, Klose KJ, Behr T, Langer P: Natural course of small, asymptomatic neuroendocrine pancreatic tumours in multiple endocrine neoplasia type 1: an endoscopic ultrasound imaging study. *Endocr Relat Cancer* 2006, **13**:1195-1202.
- Janson ET, Holmberg L, Stridsberg M, Eriksson B, Theodorsson E, Wilander E, Oberg K: Carcinoid tumors: analysis of prognostic factors and survival in 301 patients from a referral center. *Ann Oncol* 1997, **8**:685-690.
- Norton JA, Fraker DL, Alexander HR, Gibril F, Liebeherr DJ, Venzon DJ, Jensen RT: Surgery increases survival in patients with gastrinoma. *Ann Surg* 2006, **244**:410-419.

32. Fraker DL, Norton JA, Alexander HR, Venzon DJ, Jensen RT: **Surgery in Zollinger-Ellison syndrome alters the natural history of gastrinoma.** *Ann Surg* 1994, **220**:320-328.
33. Service FJ, McMahon MM, O'Brien PC, Ballard DJ: **Functioning insulinoma—incidence, recurrence, and long-term survival of patients: a 60-year study.** *Mayo Clin Proc* 1991, **66**:711-719.
34. O'Riordain DS, O'Brien T, van Heerden JA, Service FJ, Grant CS: **Surgical management of insulinoma associated with multiple endocrine neoplasia type I.** *World J Surg* 1994, **18**:488-493.
35. Demeure MJ, Klonoff DC, Karam JH, Duh QY, Clark OH: **Insulinomas associated with multiple endocrine neoplasia type I: the need for a different surgical approach.** *Surgery* 1991, **110**:998-1004.
36. Sarmiento JM, Que FG: **Hepatic surgery for metastases from neuroendocrine tumors.** *Surg Oncol Clin N Am* 2003, **12**:231-242.
37. Que FG, Nagorney DM, Batts KP, Linz LJ, Kvols LK: **Hepatic resection for metastatic neuroendocrine carcinomas.** *Am J Surg* 1995, **169**:36-42.
38. Sarmiento JM, Heywood G, Rubin J, Ilstrup DM, Nagorney DM, Que FG: **Surgical treatment of neuroendocrine metastases to the liver: a plea for resection to increase survival.** *J Am Coll Surg* 2003, **197**:29-37.
39. Dousset B, Saint-Marc O, Pitre J, Soubrane O, Houssin D, Chapuis Y: **Metastatic endocrine tumors: medical treatment, surgical resection, or liver transplantation.** *World J Surg* 1996, **20**:908-914.
40. Steinmuller T, Kianmanesh R, Falconi M, Scarpa A, Taal B, Kwekkeboom DJ, Lopes JM, Perren A, Nikou G, Delle Fave GF, O'Toole D, Frascati Consensus Conference Participants: **Consensus guidelines for the management of patients with liver metastases from digestive (neuro)endocrine tumors: foregut, midgut, hindgut, and unknown primary.** *Neuroendocrinology* 2008, **87**:47-62.
41. Toumpanakis C, Meyer T, Caplin ME: **Cytotoxic treatment including embolization/chemoembolization for neuroendocrine tumours.** *Best Pract Res Clin Endocrinol Metab* 2007, **21**:131-144.
42. O'Toole D, Ruszniewski P: **Chemoembolization and other ablative therapies for liver metastases of gastrointestinal endocrine tumours.** *Best Pract Res Clin Gastroenterol* 2005, **19**:585-594.
43. Gupta S, Yao JC, Ahrra K, Wallace MJ, Morello FA, Madoff DC, Murthy R, Hicks ME, Ajani JA: **Hepatic artery embolization and chemoembolization for treatment of patients with metastatic carcinoid tumors: the M.D. Anderson experience.** *Cancer J* 2003, **9**:261-267.
44. Mazzaglia PJ, Berber E, Milas M, Siperstein AE: **Laparoscopic radiofrequency ablation of neuroendocrine liver metastases: a 10-year experience evaluating predictors of survival.** *Surgery* 2007, **142**:10-19.
45. Elvin A, Skogseid B, Hellman P: **Radiofrequency ablation of neuroendocrine liver metastases.** *Abdom Imaging* 2005, **30**:427-434.
46. Moertel CG, Lefkopoulou M, Lipsitz S, Hahn RG, Klaassen D: **Streptozocin-doxorubicin, streptozocin-fluorouracil or chlorozotocin in the treatment of advanced islet-cell carcinoma.** *N Engl J Med* 1992, **326**:519-523.
47. Kouvaraki MA, Ajani JA, Hoff P, Wolff R, Evans DB, Lozano R, Yao JC: **Fluorouracil, doxorubicin, and streptozocin in the treatment of patients with locally advanced and metastatic pancreatic endocrine carcinomas.** *J Clin Oncol* 2004, **22**:4762-4771.
48. Ekeblad S, Sundin A, Janson ET, Welin S, Granberg D, Kindmark H, Dunder K, Kozlovacki G, Orlefors H, Sigurd M, Oberg K, Eriksson B, Skogseid B: **Temozolomide as monotherapy is effective in treatment of advanced malignant neuroendocrine tumors.** *Clin Cancer Res* 2007, **13**:2986-2991.
49. Kulke MH, Hornick JL, Frauenhoffer C, Hooshmand S, Ryan DP, Enzinger PC, Meyerhardt JA, Clark JW, Stuart K, Fuchs CS, Redston MS: **O6-methylguanine DNA methyltransferase deficiency and response to temozolomide-based therapy in patients with neuroendocrine tumors.** *Clin Cancer Res* 2009, **15**:338-345.
50. Strosberg JR, Fine RL, Choi J, Nasir A, Coppola D, Chen DT, Helm J, Kvols L: **First-line chemotherapy with capecitabine and temozolomide in patients with metastatic pancreatic endocrine carcinomas.** *Cancer* 2011, **117**:268-275.
51. Kulke MH, Stuart K, Earle C: **A phase II study of temozolomide and bevacizumab in patients with advance neuroendocrine tumors [abstract].** *ASCO Annual Meeting Proc* 2006, **24**:4044a.
52. Kulke MH, Stuart K, Enzinger PC, Ryan DP, Clark JW, Muzikansky A, Vincitore M, Michelini A, Fuchs CS: **Phase II study of temozolomide and thalidomide in patients with metastatic neuroendocrine tumors.** *J Clin Oncol* 2006, **24**:401-406.
53. Kulke MH, Blaszkowsky L, Zhu A: **Phase I/II study of everolimus (RAD001) in combination with temozolomide in patients with advanced pancreatic neuroendocrine tumors [abstract].** *Gastrointestinal Cancers Symposium* 2010, 223a.
54. Breeman WA, de JM, Kwekkeboom DJ, Valkema R, Bakker WH, Kooij PP, Visser TJ, Krenning EP: **Somatostatin receptor-mediated imaging and therapy: basic science, current knowledge, limitations and future perspectives.** *Eur J Nucl Med* 2001, **28**:1421-1429.
55. Teunissen JJ, Kwekkeboom DJ, de JM, Esser JP, Valkema R, Krenning EP, **Endocrine tumours of the gastrointestinal tract: Peptide receptor radionuclide therapy.** *Best Pract Res Clin Gastroenterol* 2005, **19**:595-616.
56. Kwekkeboom DJ, de Herder WW, Kam BL, van Eijk CH, van EM, Kooij PP, Feelders RA, van Aken MO, Krenning EP: **Treatment with the radiolabeled somatostatin analog [177 Lu-DOTA 0Tyr3]octreotide: toxicity, efficacy, and survival.** *J Clin Oncol* 2008, **26**:2124-2130.
57. Bushnell DL, O'Dorisio TM, O'Dorisio MS, Menda Y, Hicks RJ, Van CE, Baulieu JL, Borson-Chazot F, Anthony L, Benson AB, Oberg K, Grossman AB, Connolly M, Bouterfa H, Li Y, Kacena KA, LaFrance N, Pauwels SA: **90Y-edotreotide for metastatic carcinoid refractory to octreotide.** *J Clin Oncol* 2010, **28**:1652-1659.
58. Phan A, Yao J, Fogelman D: **A prospective, multi-institutional phase II study of GW786034 (pazopanib) and depot octreotide in advanced low-grade neuroendocrine carcinoma [abstract].** *J Clin Oncol* 2010, **28**.
59. Hobday TJ, Rubin J, Holen K, Picus J, Donehower R, Marschke R, Maples W, Lloyd R, Mahoney M, Erlichman C: **MC044h, a phase II trial of sorafenib in patients with metastatic neuroendocrine tumors: a phase II consortium study [abstract].** *ASCO Annual Meeting Proc* 2007, **25**:4505a.
60. Kulke MH, Lenz HJ, Meropol NJ, Posey J, Ryan DP, Picus J, Bergstrand E, Stuart K, Tye L, Huang X, Li ZJ, Baum CM, Fuchs CS: **Activity of sunitinib in patients with advanced neuroendocrine tumors.** *J Clin Oncol* 2008, **26**:3403-3410.
61. Raymond E, Dahan L, Raoul JL, Bang YJ, Borbath I, Lombard-Bohas C, Valle J, Metrakos P, Smith D, Vinik A, Chen JS, Horsch D, Hammel P, Wiedenmann B, Van Cutsem E, Patyna S, Lu DR, Blanckmeister C, Chao R, Ruszniewski P: **Sunitinib malate for the treatment of pancreatic neuroendocrine tumors.** *N Engl J Med* 2011, **364**:501-513.
62. Duran I, Kortmansky J, Singh D, Hirte H, Kocha W, Goss G, Le L, Oza A, Nicklee T, Ho J, Birle D, Pond GR, Arboine D, Dancey J, Aviel-Ronan S, Tsao MS, Hedley D, Siu LL: **A phase II clinical and pharmacodynamic study of temsirolimus in advanced neuroendocrine carcinomas.** *Br J Cancer* 2006, **95**:1148-1154.
63. Yao JC, Phan AT, Chang DZ, Wolff RA, Hess K, Gupta S, Jacobs C, Mares JE, Landgraf AN, Merci-Bernstam F: **Efficacy of RAD001 (everolimus) and octreotide LAR in advanced low- to intermediate-grade neuroendocrine tumors: results of a phase II study.** *J Clin Oncol* 2008, **26**:4311-4318.
64. Yao JC, Lombard-Bohas C, Baudin E, Kvols LK, Rougier P, Ruszniewski P, Hoosen S, St PJ, Haas T, Lebwohl D, Van Cutsem E, Kulke MH, Hobday TJ, O'Dorisio TM, Shah MH, Cadiot G, Luppi G, Posey JA, Wiedenmann B: **Daily oral everolimus activity in patients with metastatic pancreatic neuroendocrine tumors after failure of cytotoxic chemotherapy: a phase II trial.** *J Clin Oncol* 2010, **28**:69-76.
65. Yao JC, Shah MH, Ito T, Bohas CL, Wolin EM, Van CE, Hobday TJ, Okusaka T, Capdevila J, de Vries EG, Tomassetti P, Pavel ME, Hoosen S, Haas T, Lincy J, Lebwohl D, Oberg K, RAD001 in Advanced Neuroendocrine Tumors Third Trial (RADIANT-3) Study Group: **Everolimus for advanced pancreatic neuroendocrine tumors.** *N Engl J Med* 2011, **364**:514-523.
66. Yao J, Phan A, Fogelman D: **Randomized run-in study of bevacizumab and everolimus in low to intermediate grade neuroendocrine tumors using perfusion CT as a functional biomarker [abstract].** *J Clin Oncol* 2011, **28**.
67. Ramanathan RK, Cnaan A, Hahn RG, Carbone PP, Haller DG: **Phase II trial of dacarbazine (DTIC) in advanced pancreatic islet cell carcinoma. Study of the Eastern Cooperative Oncology Group-E6282.** *Ann Oncol* 2001, **12**:1139-1143.

doi:10.1186/1756-8722-4-29

Cite this article as: Kulke et al.: Evolving Diagnostic and Treatment Strategies for Pancreatic Neuroendocrine Tumors. *Journal of Hematology & Oncology* 2011 4:29.