


# Antibodies against nephronectin ameliorate anti-type II collagen-induced arthritis in mice

Shigeyuki Kon<sup>1</sup> , Machiko Honda<sup>1</sup>, Kiyoshi Ishikawa<sup>2</sup>, Masahiro Maeda<sup>2</sup> and Tatsuya Segawa<sup>2</sup>

<sup>1</sup> Department of Molecular Immunology, Faculty of Pharmaceutical Sciences, Fukuyama University, Japan

<sup>2</sup> Immuno-Biological Laboratories, Fujioka, Japan

## Keywords

antibody; anti-type II collagen-induced arthritis; autoimmune diseases; ELISA; nephronectin

## Correspondence

S. Kon, Department of Molecular Immunology, Faculty of Pharmaceutical Sciences, Fukuyama University, 985, Sanzo, Higashimura-cho, Fukuyama 729-0292, Japan

Fax: +81 84 736 2024

Tel: +81 84 736 2112

E-mail: kon@fukuyama-u.ac.jp

(Received 27 August 2019, revised 25 October 2019, accepted 7 November 2019)

doi:10.1002/2211-5463.12758

The extracellular matrix protein nephronectin (Npnt) is known to be critical for kidney development, but its function in inflammatory diseases is unknown. Here, we developed a new enzyme-linked immunosorbent assay system to detect Npnt in various autoimmune diseases, which revealed that plasma Npnt levels are increased in various mouse autoimmune models. We also report that antibodies against the  $\alpha 8\beta 1$  integrin-binding region of Npnt protect mice from anti-type II collagen-induced arthritis, suggesting that Npnt may be a potential therapeutic target molecule for the prevention of autoimmune arthritis.

The extracellular matrix (ECM) primarily functions to support the intercellular space; however, recent studies suggest that the ECM regulates various cellular functions including cell differentiation, cell growth, survival, adhesion, and migration by mediating integrin receptors [1–3]. A wide variety of ECM proteins have the same primary sequence motif, a tripeptide, Arg-Gly-Asp (RGD) motif for integrin binding [4]. The expression of some ECM components is increased during autoimmune diseases [5,6]. For example, we previously reported that osteopontin (OPN), an ECM protein containing an RGD motif, is highly expressed in inflammatory diseases and critically involved in autoimmune diseases, including collagen antibody-induced arthritis (CAIA) [7,8] and concanavalin A

(ConA)-induced hepatitis [9]. Therefore, aberrant ECM expression facilitates disease development.

An ECM protein, nephronectin (Npnt), is expressed in kidneys and hair follicles [10,11]. Npnt consists of five epidermal growth factor (EGF)-like repeats in N-terminal region, a linker segment containing an RGD integrin-binding site, and a meprin/A5-protein/PTPmu (MAM) domain in C-terminal region [10]. The RGD sequence in Npnt is critical for binding to its receptor,  $\alpha 8\beta 1$  integrin, and the interaction is critically involved in kidney development [10,12]. EGF-like repeats and MAM domains are responsible for binding to chondroitin sulfate and heparan sulfate, respectively [13]. The involvement of Npnt in diseases has been reported, particularly kidney diseases [14–17], acute

## Abbreviations

CAIA, collagen antibody-induced arthritis; ConA, concanavalin A; EAE, experimental autoimmune encephalomyelitis; ECM, extracellular matrix; EGF, epidermal growth factor; ELISA, enzyme-linked immunosorbent assay; HE, hematoxylin and eosin; LPS, lipopolysaccharide; MAM, meprin/A5-protein/PTPmu; MOG, myelin oligodendrocyte glycoprotein; Npnt, nephronectin; Npnt-FD, nephronectin-functional domain; OPN, osteopontin; PLP, proteolipid protein; RA, rheumatoid arthritis; SUMO, small ubiquitin-related modifier; TLR, Toll-like receptor.

and chronic hepatitis [18], and tumor progression [19–21].

We performed real-time PCR to evaluate Npnt expression in tissues and found that Npnt expression was high in the spleen. Therefore, we hypothesize that Npnt may play an important role in autoimmune diseases. To test this hypothesis, we developed an enzyme-linked immunosorbent assay (ELISA) system for quantification of Npnt protein expression levels and generated an antibody against the  $\alpha 8\beta 1$  integrin-binding site. In the present study, we show the expression and functional importance of Npnt in autoimmune diseases.

## Materials and methods

### Reagents and cell lines

Mouse nephronectin (Npnt) protein [called Npnt (R&D) protein in this study] was from R&D systems (Minneapolis, MN, USA). RD cells (derived from a human rhabdomyosarcoma), A549 cells (derived from human lung adenocarcinoma), HepG2 (derived from human hepatocellular carcinoma), LN-229 cells (derived from human glioblastoma), B16 cells (derived from mouse skin melanoma), NIH3T3 cells, MEF cells, L929 cells (derived from mouse fibroblast) were cultured in DMEM containing 10% fetal bovine serum (FBS). Raji cells (derived from human Burkitt's lymphoma), Jurkat cells (derived from human T-cell leukemia), U937 cells (derived from human histiocytic lymphoma), HL60 cells (derived from human acute promyelocytic leukemia), Ehrlich cells (derived from mouse Ehrlich-Lette ascites carcinoma), X63 cells (derived from mouse myeloma) were cultured in RPMI containing 10% FBS.

### Animals

Mice and rats were kept under specific pathogen-free conditions and provided food and water *ad libitum*. Every effort was made to minimize suffering during injections, and all surgery was performed on humanely sacrificed animals. All animal experiments were performed in accordance with the guidelines of the Bioscience Committee of and were approved by the Animal Care and Use Committee of Immuno-biological Laboratories.

Specific pathogen-free BALB/c mice, C57BL/6, and SD rats were purchased from Japan SLC (Hamamatsu, Japan).

### Real-time PCR

Total RNA from healthy mouse tissues, arthritic joints, and synovial cells was extracted with TRIzol (Thermo Fisher, Hanover Park, IL, USA). First-strand cDNA was generated with a first-strand cDNA synthesis kit

(TOYOBO, Osaka, Japan). For human Npnt expression in human tissues, a human multiple tissue cDNA panel was used (Takara, Kusatsu, Japan). The specific primers used are shown in Table 1. The expression level of mRNA was calculated using the calibration curve method using LIGHTCYCLER software version 3 (Roche Diagnostics, Indianapolis, IN, USA). Data were standardized to G3PDH.

### Npnt preparation

The human Npnt complementary DNAs (cDNAs) were PCR-amplified from A549 cells. Primer sets for cloning of Npnt are shown in Table 2. The amplified product was cloned into a pM-secSUMOstar vector (LifeSensors, Malvern, PA, USA), which contains a His-tag and SUMO tag at the 5' end of the cloning sites for purification and enhancing expression and solubility. The cDNAs encoding domain of human Npnt was subcloned into pM-secSUMOstar in-frame with a His-tag and SUMO tag. EGF-like repeats, Linker segment, and MAM domain were amplified by PCR using primer sets described in Table 2. The Npnt expression plasmids after nucleotide sequence confirmation were transfected into 293 cells with Lipofectamine 2000 (Thermo Fisher). His-SUMO-tagged human Npnt was purified using TALON Resin (Takara) following the manufacturer's protocol.

### Production and purification of polyclonal and monoclonal antibodies against Npnt

Synthetic peptides used for immunization were as follows: Npnt3 peptide FKGEKRRGHTGEIGLDDVSL and NPNT-FD peptide POKPRGDVFPQRPTNDLFEIFEIER, which correspond to common amino acid sequences in human and mouse Npnt, and the functional domain of mouse Npnt for  $\alpha 8\beta 1$  integrin, respectively. These peptides were used to immunize rabbits after coupling with thyroglobulin. The immunoglobulin (Ig) G fraction was obtained from a sepharose column coupled with the synthetic peptide used as the immunogen.

Generation of monoclonal antibodies against Npnt was as follows. SD rats were subcutaneously immunized with mouse Npnt protein mixed with Freund's complete

**Table 1.** Real-time PCR primer sequences.

Gene	Sequence of primers
Human Npnt	5'-AAATGAGTGTGGCCTGAAGC-3' 5'-TGCCATAAATCTAGGGCAGGA-3'
Mouse Npnt	5'-GCGGATGAGGAAGTAAAGGAC-3' 5'-CCTTTGAAGATGACGCTTTTGTG-3'
Human $\alpha 8$ integrin	5'-TCTGCTGCACCCAATGATTA-3' 5'-CTCCACAGTCCACCAGAATGT-3'
Mouse $\alpha 8$ integrin	5'-AGATTTGCTTGTAGGGGCATT-3' 5'-GACAGCTTCAAGTCAGGAACG-3'

**Table 2.** Primer sequences for cloning of Npnt.

Gene	Sequence of primers
Human Npnt	5'-TATTGAGGCTCATCGCGAACAGATTGGAGGTGAGTTCGACGGGAGGTGGCCC-3' 5'-ATGCCTGCAGGTCGACTTAGCGTTCTTCAGAGCAGTG-3'
EGF-like repeats	5'-TATTGAGGCTCATCGCGAACAGATTGGAGGTGAGTTCGACGGGAGGTGGCCC-3' 5'-ATGCCTGCAGGTCGACTTACACACAAGTCAGTCCATCACC-3'
Linker segment	5'-TATTGAGGCTCATCGCGAACAGATTGGAGGTTATATCCCAAAGTTATGATT-3' 5'-ATGCCTGCAGGTCGACTTATACCAGAACACCTGGATCATC-3'
MAM domain	5'-TATTGAGGCTCATCGCGAACAGATTGGAGGTCACAGTTGTAATTTTGACCAT-3' 5'-ATGCCTGCAGGTCGACTTAGCGTTCTTCAGAGCAGTG-3'

adjuvant (Becton Dickinson, Sparks, MD, USA) (day 1) and immunized with mouse Npnt protein mixed with Freund's incomplete adjuvant (Becton Dickinson) (days 7, 14, and 21) Spleen cells were harvested and fused with X63-Ag8-653 cells in the presence of polyethylene glycol (Roche Diagnostics, Indianapolis, IN, USA). Hybridomas were screened for production of antibody specifically reactive to immobilized mouse Npnt used for the immunization, and one IgG1 clone (20D1a) from the mouse Npnt protein was obtained.

### Establishing the Npnt sandwich ELISA system

About 5  $\mu\text{g}\cdot\text{mL}^{-1}$  of rabbit anti-Npnt3 antibody was used as a capture antibody, and 2  $\text{ng}\cdot\text{mL}^{-1}$  of 20D1 conjugated with HRP by using Peroxidase labeling kit according to the manufacturer's instructions (DOJINDO, Kumamoto, Japan) was used as the detection antibody. Purified mouse Npnt (R&D) protein was used as the standard. 100  $\mu\text{L}$  of the diluted sample was applied to each well of the antibody-coated plate and incubated for 1 h at 37 °C. The wells were washed three times with PBS-T [PBS containing 0.05% (v/v) Tween-20], and then, 100  $\mu\text{L}$  of detection antibody was added and incubated for 30 min at 4 °C. Each well was washed with PBS-T and incubated with 100  $\mu\text{L}$  of 3,3',5,5'-tetramethyl benzidine (TMB) (Sigma-Aldrich, St Louis, MO, USA) for 30 min, and then, 100  $\mu\text{L}$  of 1 M sulfuric acid was added. Plates were read at a wavelength of 450 nm by an immunoreader (Thermo Fisher).

### Direct ELISA

Mouse Npnt (R&D), fibronectin (Sigma-Aldrich, St Louis, MO, USA), GRGDS peptide-coupled with BSA were coated onto 96-well plate at a concentration of 1  $\mu\text{g}\cdot\text{mL}^{-1}$  at 37 °C for 1 h, then blocked with 0.1% BSA in PBS containing 0.05%  $\text{NaN}_3$  at 37 °C for at least 1 h. The plates were washed two times with PBS-T and incubated with anti-Npnt-FD mAb at various concentrations at 37 °C for 1 h. After a further three washes, 1 : 5000 dilution of HRP-conjugated anti-mouse IgG antibody (Jackson Immuno Research, West Grove, PA, USA) was added to each well at room temperature for 30 min. Bound protein

was quantified by a colorimetric assay using TMB as a substrate for 15 min at room temperature. Plates were read at a wavelength of 450 nm by an immunoreader.

### Cell adhesion assay

The 96-well plates were precoated with hNpnt, mNpnt (R&D), NPNT-FD synthetic peptides, fibronectin, or vitronectin (2  $\mu\text{g}\cdot\text{mL}^{-1}$ ) at 37 °C for 1 h, followed by treatment with 0.5% BSA in PBS for 1 h at room temperature. Cells in the presence or absence of various concentrations of antibody were suspended in DMEM containing 0.25% BSA, and 200  $\mu\text{L}$  of cell suspension (at a cell density of  $5 \times 10^4$  cells/well) was applied to 96-well plates and incubated for 1 h at 37 °C. The medium was removed from the plates, and all wells were washed twice. The adherent cells were fixed and stained with 0.5% crystal violet in 20% methanol for 30 min. All wells were rinsed three times with water, and adherent cells were then lysed with 20% acetic acid. The resulting supernatants from each well were analyzed by an immunoreader, and the absorbance at 595 nm was measured to determine the relative number of cells adhered to wells.

### Fractionation of synovial tissues

Hind limbs were surgically removed from the ankle joint, followed by surgical separation of the dermal, subcutaneous, tendinous, and muscle tissues from joints. The remaining soft tissues were removed from the bones and homogenized roughly with dissecting scissors. Homogenates were washed once with culture medium and treated with 3  $\text{mg}\cdot\text{mL}^{-1}$  type II collagenase (Worthington Biochemical, Freehold, NJ, USA) by vigorous stirring. After washing, synovial cells were fractionated into TLR-4<sup>+</sup> or TLR-4<sup>-</sup> fractions using MACS (Miltenyi Biotec, Auburn, CA, USA). The TLR-4<sup>+</sup> fraction was considered macrophages and the TLR-4<sup>-</sup> fraction was cultured overnight, and adherent cells were used as fibroblasts.

### Arthritis model

Arthritis was induced using an arthritogenic 5-clone monoclonal antibody cocktail kit (Chondrex Inc., Redmond,

WA, USA) following the manufacturer's protocol. Briefly, 7-week-old female BALB/c mice were injected intravenously with a mixture of five anti-type II collagen monoclonal antibodies (2 mg each) on day  $-3$ , followed by an intraperitoneal injection of 50  $\mu\text{g}$  of lipopolysaccharide (LPS) (0111:B4) on day 0. About 400  $\mu\text{g}$  anti-Npnt-FD antibody or control rabbit IgG was administered intraperitoneally at doses of 400  $\mu\text{g}$  per mouse on days  $-4$  and 0. The clinical severity of arthritis was graded up to 9 days after LPS administration in each of the four paws on a 0–4 scale. The disease severity was recorded for each limb as follows: 0, normal; 1, focal slight swelling and/or redness in one digit; 2, moderate swelling and erythema of two digits; 3, marked swelling and erythema of the limb; and 4, maximal swelling, erythema, deformity, and/or ankylosis. Mice were scored in a double-blind manner. Mice were sacrificed, and then, joint sections were stained with hematoxylin and eosin (HE) and fast green/safranin-O for immunohistochemical evaluation.

### Induction of concanavalin A-induced hepatitis and experimental autoimmune encephalomyelitis

For induction of ConA-hepatitis, BALB/c mice were injected intravenously with 15 mg ConA (Sigma-Aldrich) per kilogram of body weight, dissolved in pyrogen-free PBS.

For induction of experimental autoimmune encephalomyelitis (EAE), C57BL/6 or SJL/J mice were injected subcutaneously with 100  $\mu\text{g}$  of myelin oligodendrocyte glycoprotein (MOG) 35–55 peptide (MEVG-WYRSPFSRVVHLYRNGK) or Myelin proteolipid protein (PLP) 139–151 (HSLGKWLGHDPKF) emulsified with complete Freund's adjuvant on day 0 and intravenous

injection of 400 ng of pertussis toxin (List Biological Laboratories, Campbell, CA, USA) on days 0 and 2.

### Statistical analysis

Data are presented as means  $\pm$  SEM and are representative of at least three independent experiments. The statistical significance of differences between groups was calculated with a two-tailed Student's *t*-test. Differences were considered significant for  $P < 0.05$  (\*) or 0.01 (\*\*).


## Results

### Expression of Npnt and $\alpha 8$ integrin in normal tissues


We first examined the expression level of human and mouse Npnt (Fig. 1A) and  $\alpha 8$  integrin (Fig. 1B) in healthy human and mouse tissues using real-time PCR. Subsequently, we found both Npnt and  $\alpha 8$  integrin were highly expressed in the spleen; thus, we hypothesized that the interaction between Npnt and  $\alpha 8$  integrin may be involved in immunological diseases.

### Establishing a sandwich ELISA system for Npnt

To analyze the expression level of Npnt in an autoimmune mouse model, we sought to establish an Npnt sandwich ELISA system. For ELISA construction, two antibodies against Npnt were generated, Npnt3 rabbit polyclonal antibody and 20D1 monoclonal antibody.



**Fig. 1.** Expression of Npnt and  $\alpha 8$  integrin in human and mouse tissues. Npnt (A) and  $\alpha 8$  integrin (B) expression in healthy human and mouse tissues were evaluated by real-time PCR. Data are representative of three or more independent experiments with similar results. Data are presented as means  $\pm$  SEM.



**Fig. 2.** Establishing the sandwich ELISA system for Npnt. (A) Schematic representation of Npnt and its truncated variants. Cell binding sequences (RGD sequences) in the linker segment and the epitope in the MAM domain of anti-Npnt3 antibody are indicated. (B) Specificity of Npnt3 and 20D1a antibodies by western blotting. (C) Diagram (top panel) and standard curve (lower panel) of the ELISA. Recombinant mNpnt (R&D) protein was used as the standard. (D) Expression of Npnt in cell lines by RT-PCR. Human or mouse cell lines were left or right panel, respectively. (E) Detection of Npnt in supernatants from cell lines used in (C). Data are representative of three or more independent experiments with similar results. Data are presented as means  $\pm$  SEM.


For characterization of these antibodies, we purified full length of Npnt, EGF-like repeats, linker segment, MAM domain (Fig. 2A). Both antibodies were capable of detecting mouse and human Npnt (Fig. 2B). Notably, the molecular weight of human Npnt is higher than mouse because human Npnt is fused with a SUMO tag (approximate molecular weight of 20 kDa). A link truncated variant showed higher and broader bands than the expected size (approximately 28 kDa) as reported previously [22]. Anti-Npnt3 antibody recognized the MAM domain as predicted, whereas the 20D1 Ab did not react with a panel of Npnt truncated variants, suggesting that 20D1a Ab recognizes the Npnt conformation (Fig. 2B).

We developed an ELISA combination using Npnt3 as a coating antibody and 20D1 as a detecting antibody (Fig. 2C). The standard curve of the ELISA system exhibited a linear shape ranging from 0.078 to

5.0 ng·mL<sup>-1</sup> calculated with mouse Npnt (R&D) protein (Fig. 2C). To analyze the specificity of the ELISA, Npnt mRNA expression in various human and mouse cell lines was explored by RT-PCR, and Npnt expression was detected in RD cells, A549 cells and HepG2 cells in human cell lines, and B16 cells, Ehrlich cells, and MEF cells in mouse cell lines (Fig. 2D). The ELISA we developed successfully detected secreted Npnt protein in the supernatant of the cell lines expressing Npnt by RT-PCR, but not from non-Npnt mRNA expressing cell lines (Fig. 2E).

### Elevated expression of Npnt in autoimmune diseases

The antigenic peptide of anti-Npnt3 antibody is a common sequence shared by humans, mice, and rats, and the 20D1a antibody recognizes both human and




**Fig. 3.** Npnt in plasma and urine samples. (A) Npnt in plasma and urine samples from healthy mice and humans. (B) Npnt in plasma samples from various murine autoimmune disease models. Plasma samples were tested on the day when inflammation was exacerbated as follows, for ConA-hepatitis, 24 h after ConA injection; for CAIA, 7 days after LPS injection; for EAE, 14 days after MOG or PLP peptide immunization. Data are representative of three or more independent experiments with similar results. \* $P < 0.05$  or \*\* $P < 0.01$  versus plasma samples from wild-type mice (Student's *t*-test).

mouse Npnt. Therefore, we assayed Npnt values in healthy mouse, human, and rat plasma samples. We found that the Npnt system can be used for quantification of mouse and human Npnt values with Npnt (R&D) protein as the standard. We next evaluated Npnt expression in urine since Npnt is highly expressed in kidneys. Significant expression of Npnt in human urine, but not mouse urine, was detected (Fig. 3A). Next, we tested Npnt levels in plasma samples from autoimmune mouse models, specifically ConA-hepatitis, CAIA, and EAE models. We found that Npnt levels in all of the inflammatory models we tested were significantly elevated (Fig. 3B).

### Involvement of $\alpha 8\beta 1$ integrin-binding region of Npnt in inflammatory arthritis

The RGD motif and LFEIFEIER sequence in Npnt are required for binding to  $\alpha 8\beta 1$  integrin, which is the


physiological receptor for Npnt. To assess the importance of the  $\alpha 8\beta 1$  integrin-binding region within murine Npnt, we prepared the anti-Npnt-FD antibody that recognizes the PQKPRGDVFIPRQPTNDLFEIFEIER motif used as the antigen. We first examined the specificity of the anti-Npnt-FD antibody by direct ELISA, western blotting, or cell adhesion test. We also asked whether Npnt-FD antibody cross-reacts with fibronectin or GRGDS peptide since RGD motif found in many ECM molecules is exist in antigen peptide. The anti-Npnt-FD antibody recognized mouse Npnt protein, but not RGD motif-containing ECM, fibronectin or GRGDS peptide by direct ELISA (Fig. 4A), and detected mouse Npnt, human Npnt, and the Linker segment, which contains the antigen region of the anti-Npnt-FD antibody, by western blotting (Fig. 4B). To examine the functional specificity, cell adhesion assays were performed. Anti-Npnt-FD antibody partially inhibited cellular adhesion to mouse



**Fig. 4.** Generation of neutralizing antibodies for Npnt. (A) Cross-reactivity of anti-Npnt-FD antibody to plates coated with Npnt, fibronectin, or GRGDS peptide-coupled with BSA ( $1 \mu\text{g}\cdot\text{mL}^{-1}$ ) by direct ELISA. (B) Specificity of anti-Npnt-FD antibody by western blotting. (C, D) NIH3T3 cells were allowed to adhere to 96-well plates precoated with full-length of Npnt or Npnt-Link (C), fibronectin, or vitronectin (D) in the presence of indicated concentrations of anti-Npnt-FD antibody. Normal rabbit IgG was used as a control antibody. Data are presented as mean  $\pm$  SEM. Data are representative of three or more independent experiments with similar results. \* $P < 0.05$ , anti-Npnt-FD antibody versus control antibody (Student's  $t$ -test).

and human Npnt, whereas it completely inhibited the adhesion of the  $\alpha 8\beta 1$  integrin-binding region, suggesting that an alternative Npnt exists (Fig. 4C). Anti-Npnt-FD antibody did not show any inhibitory activity toward fibronectin or vitronectin (Fig. 4D). These results indicate that anti-Npnt-FD antibody specifically recognizes and functionally inhibits the interaction of the  $\alpha 8\beta 1$  integrin-binding region of Npnt with its receptor.

An antibody against OPN, which is one of the  $\alpha 8\beta 1$  integrin ligands, inhibits an exacerbation of CAIA; thus, we assessed the function of the anti-Npnt-FD antibody in the CAIA model [23]. In the CAIA model, on day 6 the Npnt mRNA in arthritic joints was elevated. Furthermore, synovial fibroblasts and synovial macrophages, which are the main cellular components in arthritic synovia and important players in the joint pathology of rheumatoid arthritis



**Fig. 5.** Npnt neutralization by anti-Npnt-FD antibody ameliorates CAIA. (A) Protocol for CAIA with anti-Npnt-FD antibody treatment. The antibody was administered to BALB/c mice on days  $-4$  and  $0$  relative to LPS injection during CAIA induction. (B) Npnt expression in arthritic joints on day 6, synovial fibroblasts, or synovial macrophages were evaluated by real-time PCR.  $*P < 0.05$ ,  $**P < 0.01$ , wild-type versus CAIA (Student's  $t$ -test). (C) Arthritis scores of arthritic mice treated with control antibody (rabbit IgG) or an anti-Npnt-FD antibody at the indicated time points ( $n = 5$  per group).  $*P < 0.05$ ,  $**P < 0.01$ , anti-Npnt-FD antibody versus control Ig (Student's  $t$ -test). Data are presented as means  $\pm$  SEM. (D) Representative images of the forepaw and hindpaw on day 7 are shown. (E–G) Representative histology of normal joints and arthritic joints on day 14 from mice treated with control antibody or anti-Npnt-FD antibody. Sections were stained with HE (E, G) or Safranin-O (F). Magnified views of the boxed areas a and b are shown in F and G, respectively. Scale bar = 2mm (E), 200μm (F and G). Data are representative of three or more independent experiments with similar results.

(RA) [24,25], were elevated (Fig. 5B). Treatment with the anti-Npnt-FD antibody on days  $-3$  and  $0$  led to the delay of clinical onset of arthritis, and the severity of arthritis was significantly attenuated (Figure 5C). The gross appearance of arthritic joint treated with anti-Npnt-FD antibody on day 7 is shown in Fig. 5D. Finally, we evaluated the effect of anti-Npnt-FD antibody treatment on joint histology on day 14 after the LPS injection. In concert with the clinical score, histological analysis demonstrated marked differences. Cartilage destruction evaluated by Safranin-O staining was faint in the arthritic joints of

control mice. In contrast, the staining was partially rescued in anti-Npnt-FD antibody-treated mice (Fig. 5F), indicating that the destruction of proteoglycans was partially prevented by the anti-Npnt-FD antibody. The proliferation of synovial cells, evaluated by HE staining, was detected in control antibody-treated mice; however, this feature of CAIA was clearly diminished following the administration of the anti-Npnt-FD antibody (Fig. 5G). These results strongly suggested that the  $\alpha 8 \beta 1$  integrin-binding region of Npnt is involved in inflammatory arthritis.



## Discussion

In the present study, we investigated the function of Npnt in autoimmune diseases as high levels of Npnt mRNA were found in the spleens of humans and mice. We developed a sandwich ELISA system for quantifying the protein level of Npnt in autoimmune diseases and found that Npnt in blood is increased in various autoimmune diseases. The results from ConA-hepatitis are consistent with previous reports that observed elevated Npnt mRNA [18].

For clarification of Npnt function in autoimmune diseases, we generated a neutralizing antibody against Npnt (anti-Npnt-FD antibody) by immunizing rabbits with the PQKPRGDVFIPRQPTNDLFEIFEIER peptide (Npnt-FD peptide), because the RGD motif and LFEIFEIER sequence in Npnt plays a critical role in the binding to its physiological receptor,  $\alpha 8\beta 1$  integrin [22]. Anti-Npnt-FD antibody partially inhibits cell adhesion of full-length mouse and human Npnt to NIT3T3 cells, which express  $\alpha 8\beta 1$  integrin [26]. This suggests that Npnt has other receptors on NIH3T3 cells. In fact, it has been reported that the MAM domain in Npnt binds to heparan sulfate proteoglycan and EGF-like repeats domain and activates the EGF receptor [13,27]. Therefore, anti-Npnt-FD antibody specifically inhibits the interaction between Npnt and receptors bound to Npnt-FD peptide such as  $\alpha 8\beta 1$  integrin.

Previously, we reported that neutralizing antibodies for osteopontin, one of the ligands of  $\alpha 8\beta 1$  integrin, inhibits the development of inflammatory arthritis in the CAIA model. Therefore, we analyzed the function of Npnt in CAIA and determined that anti-Npnt-FD antibody attenuates the severity of arthritis. We also found higher expression of Npnt in arthritic joints than in control mice. In RA joints, proliferating fibroblasts and macrophages are clearly seen. These cells are thought to cause an exacerbation of RA by secretion of various pro-inflammatory cytokines such as tumor necrosis factor (TNF)- $\alpha$  or interleukin (IL)-1 $\beta$ . In this study, Npnt expression in synovial fibroblasts and macrophages from CAIA was also increased, suggesting a critical involvement of the PQKPRGDVFIPRQPTNDLFEIFEIER sequence of Npnt and its receptor(s) in the development of inflammatory arthritis and, perhaps, the combination of Npnt and these cytokines have a synergistic effect on the severity of inflammation.

The  $\alpha 8\beta 1$  integrin, which is part of the RGD-recognizing integrin family, is prominently expressed in smooth muscle and has a defined in vivo role in kidney development and liver and lung fibrosis

[10,11,28,29]. In this study,  $\alpha 8\beta 1$  integrin is highly expressed in human and mouse spleen, suggesting that  $\alpha 8\beta 1$  integrin has any important role in immune responses. It is conceivable that the binding of Npnt to  $\alpha 8\beta 1$  integrin is involved in  $\alpha 8\beta 1$  integrin, but we cannot exclude the possibility that other RGD-recognizing integrins, such as  $\alpha v\beta 3$  integrin, are the Npnt receptor during inflammatory arthritis. Further research is required to understand how  $\alpha 8\beta 1$  integrin is involved in inflammatory arthritis.

In conclusion, we found that Npnt is highly expressed in the spleen and upregulated in blood from various autoimmune mouse models. Furthermore, we provide evidence for a potential role of the  $\alpha 8\beta 1$  integrin-binding region of Npnt in the pathogenesis of inflammatory arthritis. Further studies will be directed at understanding the mechanism by which Npnt and its interactions with integrins affect autoimmune diseases.

## Acknowledgements

This study was supported by research grants from JSPS KAKENHI Grant Numbers 16K08221 and 19K07492, Japan Rheumatism Foundation, the SUHARA Memorial Foundation, the FUGAKU TRUST for Medical Research, academic support from Pfizer, Chugai, Astellas, and Eli Lilly to SK. We thank S. J. Win, PhD, from Edanz Group for editing a draft of this manuscript.

## Conflict of interest

SK and MH declare no conflicts of interest associated with this manuscript. KI, MM, and TS are employees of the Immuno-Biological Laboratories Co., Ltd.

## Author contributions

SK conceived and carried out experiments and wrote the manuscript; MH performed experiments; KI, MH, MM, and TS generated anti-Npnt antibody and established the Npnt ELISA system; SK and TS supervised the study. All authors read and approved the final manuscript.

## References

- 1 Hynes RO (2009) The extracellular matrix: not just pretty fibrils. *Science* **326**, 1216–1219.
- 2 Giancotti FG and Ruoslahti E (1999) Integrin signaling. *Science* **285**, 1028–1032.

- 3 Hynes RO (2014) Stretching the boundaries of extracellular matrix research. *Nat Rev Mol Cell Biol* **15**, 761–763.
- 4 Hynes RO (1987) Integrins: a family of cell surface receptors. *Cell* **48**, 549–554.
- 5 Hasegawa M, Yoshida T and Sudo A (2018) Role of tenascin-C in articular cartilage. *Mod Rheumatol* **28**, 215–220.
- 6 Schultz C (2019) Targeting the extracellular matrix for delivery of bioactive molecules to sites of arthritis. *Br J Pharmacol* **176**, 26–37.
- 7 Yamamoto N, Sakai F, Kon S, Morimoto J, Kimura C, Yamazaki H, Okazaki I, Seki N, Fujii T and Uede T (2003) Essential role of the cryptic epitope SLAYGLR within osteopontin in a murine model of rheumatoid arthritis. *J Clin Invest* **112**, 181–188.
- 8 Kon S, Nakayama Y, Matsumoto N, Ito K, Kanayama M, Kimura C, Kouro H, Ashitomi D, Matsuda T and Uede T (2014) A novel cryptic binding motif, LRSKRSRFQVSDEQY, in the C-terminal fragment of MMP-3/7-cleaved osteopontin as a novel ligand for alpha9beta1 integrin is involved in the anti-type II collagen antibody-induced arthritis. *PLoS ONE* **9**, e116210.
- 9 Diao H, Kon S, Iwabuchi K, Kimura C, Morimoto J, Ito D, Segawa T, Maeda M, Hamuro J, Nakayama T *et al.* (2004) Osteopontin as a mediator of NKT cell function in T cell-mediated liver diseases. *Immunity* **21**, 539–550.
- 10 Brandenberger R, Schmidt A, Linton J, Wang D, Backus C, Denda S, Muller U and Reichardt LF (2001) Identification and characterization of a novel extracellular matrix protein nephronectin that is associated with integrin alpha8beta1 in the embryonic kidney. *J Cell Biol* **154**, 447–458.
- 11 Fujiwara H, Ferreira M, Donati G, Marciano DK, Linton JM, Sato Y, Hartner A, Sekiguchi K, Reichardt LF and Watt FM (2011) The basement membrane of hair follicle stem cells is a muscle cell niche. *Cell* **144**, 577–589.
- 12 Miner JH (2001) Mystery solved: discovery of a novel integrin ligand in the developing kidney. *J Cell Biol* **154**, 257–259.
- 13 Sato Y, Shimono C, Li S, Nakano I, Norioka N, Sugiura N, Kimata K, Yamada M and Sekiguchi K (2013) Nephronectin binds to heparan sulfate proteoglycans via its MAM domain. *Matrix Biol* **32**, 188–195.
- 14 Cheng CW, Ka SM, Yang SM, Shui HA, Hung YW, Ho PC, Su YC and Chen A (2008) Nephronectin expression in nephrotoxic acute tubular necrosis. *Nephrol Dial Transplant* **23**, 101–109.
- 15 Muller-Deile J, Dannenberg J, Schroder P, Lin MH, Miner JH, Chen R, Brasen JH, Thum T, Nystrom J, Staggs LB *et al.* (2017) Podocytes regulate the glomerular basement membrane protein nephronectin by means of miR-378a-3p in glomerular diseases. *Kidney Int* **92**, 836–849.
- 16 Nakatani S, Ishimura E, Mori K, Fukumoto S, Yamano S, Wei M, Emoto M, Wanibuchi H and Inaba M (2012) Nephronectin expression in glomeruli of renal biopsy specimens from various kidney diseases: nephronectin is expressed in the mesangial matrix expansion of diabetic nephropathy. *Nephron Clin Pract* **122**, 114–121.
- 17 Kashani K and Kellum JA (2015) Novel biomarkers indicating repair or progression after acute kidney injury. *Curr Opin Nephrol Hypertens* **24**, 21–27.
- 18 Inagaki FF, Tanaka M, Inagaki NF, Yagai T, Sato Y, Sekiguchi K, Oyaizu N, Kokudo N and Miyajima A (2013) Nephronectin is upregulated in acute and chronic hepatitis and aggravates liver injury by recruiting CD4 positive cells. *Biochem Biophys Res Commun* **430**, 751–756.
- 19 Dilmac S, Erin N, Demir N and Tanriover G (2018) Nephronectin is decreased in metastatic breast carcinoma and related to metastatic organs. *Pathol Oncol Res* **24**, 679–688.
- 20 Steigedal TS, Toraskar J, Redvers RP, Valla M, Magnussen SN, Bofin AM, Opdahl S, Lundgren S, Eckhardt BL, Lamar JM *et al.* (2018) Nephronectin is correlated with poor prognosis in breast cancer and promotes metastasis via its integrin-binding motifs. *Neoplasia* **20**, 387–400.
- 21 Wang D, Zhao C, Gao L, Wang Y, Gao X, Tang L, Zhang K, Li Z, Han J and Xiao J (2018) NPNT promotes early-stage bone metastases in breast cancer by regulation of the osteogenic niche. *J Bone Oncol* **13**, 91–96.
- 22 Sato Y, Uemura T, Morimitsu K, Sato-Nishiuchi R, Manabe R, Takagi J, Yamada M and Sekiguchi K (2009) Molecular basis of the recognition of nephronectin by integrin alpha8beta1. *J Biol Chem* **284**, 14524–14536.
- 23 Kanayama M, Kurotaki D, Morimoto J, Asano T, Matsui Y, Nakayama Y, Saito Y, Ito K, Kimura C, Iwasaki N *et al.* (2009) Alpha9 integrin and its ligands constitute critical joint microenvironments for development of autoimmune arthritis. *J Immunol* **182**, 8015–8025.
- 24 Muller-Ladner U, Ospelt C, Gay S, Distler O and Pap T (2007) Cells of the synovium in rheumatoid arthritis. *Synovial fibroblasts, Arthritis Res Ther* **9**, 223.
- 25 Falconer J, Murphy AN, Young SP, Clark AR, Tiziani S, Guma M and Buckley CD (2018) Review: synovial cell metabolism and chronic inflammation in rheumatoid arthritis. *Arthritis Rheumatol* **70**, 984–999.
- 26 Denda S, Reichardt LF and Muller U (1998) Identification of osteopontin as a novel ligand for the

- integrin alpha8 beta1 and potential roles for this integrin-ligand interaction in kidney morphogenesis. *Mol Biol Cell* **9**, 1425–1435.
- 27 Kahai S, Lee SC, Seth A and Yang BB (2010) Nephronectin promotes osteoblast differentiation via the epidermal growth factor-like repeats. *FEBS Lett* **584**, 233–238.
- 28 Hung CF, Wilson CL, Chow YH and Schnapp LM (2018) Role of integrin alpha8 in murine model of lung fibrosis. *PLoS ONE* **13**, e0197937.
- 29 Levine D, Rockey DC, Milner TA, Breuss JM, Fallon JT and Schnapp LM (2000) Expression of the integrin alpha8beta1 during pulmonary and hepatic fibrosis. *Am J Pathol* **156**, 1927–1935.