



Can ivermectin mass drug administrations to control scabies also reduce skin and soft tissue infections? Hospitalizations and primary care presentations lower after a large-scale trial in Fiji

Jo Middleton^{a,b*}

^aNIHR Global Health Research Unit on Neglected Tropical Diseases and Department of Primary Care and Public Health, Brighton and Sussex Medical School, Falmer, United Kingdom

^bEvolution, Behavior and Environment, School of Life Sciences, University of Sussex, Falmer, East Sussex, United Kingdom

Evaluations of Mass Drug Administrations (MDAs) targeting scabies have generally not assessed their secondary impacts on skin and soft tissue infections. These are major health burdens in the global south, with sequelae including glomerulonephritis, rheumatic fever and chronic kidney disease.^{1–3} In a major step forward Li Jun Thean and colleagues⁴ report in *The Lancet Regional Health Western Pacific* on the impact on skin infections of an oral ivermectin MDA targeting scabies in Fiji. Importantly, they show in the largest before-after trial of its kind that not only did scabies and impetigo prevalence reduce, but so too did hospitalisations with severe skin and soft tissue infections and attendance at primary care with scabies and skin infections.

Scabies is a highly irritant skin infestation with the mite *Sarcoptes scabiei* and is primarily transmitted by touch. It is classed by WHO as a Neglected Tropical Disease (NTD) in recognition of its high burden and the health risks associated with secondary infections of impetigo (i.e., cellulitis, necrotising fasciitis, skin abscesses, rheumatic fever, chronic kidney disease).^{1–3} Evidence indicates prevalence is not influenced by personal washing.⁵ Neglected tropical skin diseases are significant health problems across Oceania.^{6,7} Some of the world's highest prevalence rates of scabies have been measured in the region (e.g., Papua New Guinea, 71%; Fiji, 32%).² The situation is similar for impetigo, particularly in children (e.g., Samoa, 57%; Solomon Islands, 52%; Fiji, 36%; Vanuatu, 16%).^{2,8} The first trial in Oceania of an ivermectin MDA for scabies (outside of institutional outbreaks) was carried out in a Papua New

Guinea village in 1996,⁹ followed on a greater scale in the Solomon Islands 1997–2000,¹⁰ Fiji in 2004,¹¹ and in aboriginal communities in Australia.¹² A 15-year follow-up of the first Solomon Islands MDA found very low scabies prevalence, in fact just one case. Marks et al.¹⁰ attributed this to MDA and an associated period of active case finding. Subsequently, in the Solomon Islands¹³ and Fiji¹⁴ MDAs targeting scabies have been associated with decreased impetigo prevalence (respectively: from 24.7% to 9.6%; 24.6% to 8.0%). Influenced by these developments in Oceania, similar interventions are increasingly being conducted elsewhere.¹ The largest adoption of this approach has been in Ethiopia where a 2018 scabies MDA involved over nine million people,¹⁵ and the secondary impacts on scabies of an ivermectin MDA targeting onchocerciasis is also being assessed.¹⁶

Against this regional and global background Thean and colleagues⁴ report a 2018–2020 before-after MDA trial covering Fiji's Northern Division: first dose, 135,744 people (97% coverage); second dose, 121,760 people (87%). The prior MDA in Fiji had 716 participants enrolled in the ivermectin arm,¹⁴ in the Solomon Islands 26,188.¹³ This then is by far the largest study to date to report the impact of a scabies MDA on impetigo. Scabies prevalence decreased from 14.2% to 7.7%, and impetigo from 15.3% to 6.1%. Whilst this valuably supports the conclusions of the earlier smaller trials, what is most important in this study is the observed reductions in primary and secondary care burdens. Incidence of childhood invasive infections and post-streptococcal sequelae were unchanged. However, hospitalisations with severe skin and soft tissue infections reduced 17%, attendance at primary care with scabies and skin infections 21%. Thean and colleagues note the reduction in scabies prevalence was lower than expected compared to other in-region studies. They are likely correct in linking this in part to the intervention not being distributed and overseen directly by the study team, as many smaller scabies MDA trials have been, but instead by the national health system. A similar reduction in

The Lancet Regional Health - Western Pacific
2022;22: 100454
Published online xxx
<https://doi.org/10.1016/j.lanwpc.2022.100454>

DOI of original article: <http://dx.doi.org/10.1016/j.lanwpc.2022.100443>

*Correspondence to: Department of Primary Care and Public Health, Brighton and Sussex Medical School, 318a Watson Building, Falmer BN1 9PH, United Kingdom.

E-mail address: j.middleton@bsms.ac.uk

© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

expected effectiveness has been observed in a non-researcher run MDA in Ethiopia.¹⁶ Rather than a weakness of the study, this underlines its value as the important effects shown may more realistically indicate benefits to be expected when operationalising MDAs at large-scales beyond smaller researcher-run trials. These results strongly support (1) expansion of ivermectin MDAs targeting scabies and impetigo, (2) collection of similar hospitalisation and primary care data as MDAs expand, and (3) development of further research and interventions that capitalise on how NTD control can secondarily reduce wider burdens to health systems. Given resource scarcity such synergistic integrations are important clinically not only in Oceania⁶, but worldwide.¹

Declaration of interests

The author declares that they have no competing interests.

Funding

The author is a co-investigator of SURFACES Papua New Guinea, which is funded by Sussex Sustainability Research Project (University of Sussex, UK), and the Darwin Initiative (Department for Environment, Food & Rural Affairs, UK government). Their work on scabies in Ethiopia is funded by the National Institute for Health Research (NIHR) Global Health Research Unit on Neglected Tropical Diseases. The views expressed are those of the author and not necessarily those of the NHS, the NIHR or the UK government Department of Health and Social Care.

Role of the funding source

This piece is not a report of study data, but a commentary on a paper of which JM is not an author. JM's funders had no role in writing this commentary.

References

- 1 Engelman D, Marks M, Steer AC, et al. A framework for scabies control. *PLoS Negl Trop Dis*. 2021;15: e0009661. <https://doi.org/10.1371/journal.pntd.0009661>.

- 2 Romani L, Steer AC, Whitfield MJ, Kaldor JM. Prevalence of scabies and impetigo worldwide: a systematic review. *Lancet Infect Dis*. 2015;15:960–967. [https://doi.org/10.1016/S1473-3099\(15\)00132-2](https://doi.org/10.1016/S1473-3099(15)00132-2).
- 3 Chung SD, Wang KH, Huang CC, Lin HC. Scabies increased the risk of chronic kidney disease: a 5-year follow-up study. *JEADV*. 2014;28:286–292. <https://doi.org/10.1111/jdv.12099>.
- 4 Thean LJ, Romani L, Engelman D, et al. Prevention of bacterial complications of scabies using mass drug administration, a population-based, before-after trial in Fiji, 2018–2020. *Lancet Reg Health West Pac*. 2022. <https://doi.org/10.1016/j.lanwpc.2022.100433>.
- 5 Middleton J, Cassell JA, Jones CI, Lanza S, Head MG, Walker SL. Scabies control the forgotten role of personal hygiene – authors reply. *Lancet Infect Dis*. 2018;18:1068–1069. [https://doi.org/10.1016/S1473-3099\(18\)30558-9](https://doi.org/10.1016/S1473-3099(18)30558-9).
- 6 Kline K, McCarthy JS, Pearson M, Loukas A, Hotez PJ. Neglected tropical diseases of Oceania: review of their prevalence, distribution, and opportunities for control. *PLoS Negl Trop Dis*. 2013;7: e1755. <https://doi.org/10.1371/journal.pntd.0001755>.
- 7 Middleton J, Cassell JA, Colthart G, et al. Rationale, experience and ethical considerations underpinning integrated actions to further global goals for health and land biodiversity in Papua New Guinea. *Sustain Sci*. 2020;15:1653–1664. <https://doi.org/10.1007/s11625-020-00805-x>.
- 8 Taiaroa G, Matalavea B, Tafuna'i M, et al. Scabies and impetigo in Samoa: a school-based clinical and molecular epidemiological study. *Lancet Reg Health West Pac*. 2021;6:100081. <https://doi.org/10.1016/j.lanwpc.2020.100081>.
- 9 Bockarie MJ, Alexander NDE, Kazura JW, Bockarie F, Griffin L, Alpers MP. Treatment with ivermectin reduces the high prevalence of scabies in a village in Papua New Guinea. *Acta Trop*. 2000;75:127–130. [https://doi.org/10.1016/S0001-706X\(99\)00087-X](https://doi.org/10.1016/S0001-706X(99)00087-X).
- 10 Marks M, Taotao-Wini B, Satorara L, et al. Long term control of scabies fifteen years after an intensive treatment programme. *PLoS Negl Trop Dis*. 2015;9: e0004246. <https://doi.org/10.1371/journal.pntd.0004246>.
- 11 Haar K, Romani L, Filimone R. Scabies community prevalence and mass drug administration in two Fijian villages. *Int J Dermatol*. 2014;53:739–745. <https://doi.org/10.1111/ijd.12353>.
- 12 Kearns TM, Speare R, Cheng AC, et al. Impact of an Ivermectin mass drug administration on scabies prevalence in a remote Australian aboriginal community. *PLoS Negl Trop Dis*. 2015;9: e0004151. <https://doi.org/10.1371/journal.pntd.0004151>.
- 13 Marks M, Lucia Romani L, Oliver Sokana O, et al. Prevalence of scabies and impetigo 3 years after mass drug administration with ivermectin and azithromycin. *Clin Infect Dis*. 2020;70:1591–1595. <https://doi.org/10.1093/cid/ciz444>.
- 14 Romani L, Whitfield MJ, Koroiveuta J, et al. Mass drug administration for scabies control in a population with endemic disease. *N Engl J Med*. 2015;373:2305–2313. <https://doi.org/10.1056/NEJMoar1500987>.
- 15 Enbiale W, Bayniem TB, Ayalew A, et al. Stopping the itch": mass drug administration for scabies outbreak control covered for over nine million people in Ethiopia. *J Infect Dev Ctries*. 2020;14:28S–35S. <https://doi.org/10.3855/jidc.11701>.
- 16 Yirgu R, Middleton J, Fekadu A, et al. The secondary impact of ivermectin MDA for the control of onchocerciasis on the prevalence of scabies in the Amhara region of northern Ethiopia. In: *International Alliance for the Control of Scabies Annual Scientific Meeting*. 2021. November 2021, online.