



Distribution of *PLD* and *FagA*, *B*, *C* and *D* genes in *Corynebacterium pseudotuberculosis* isolates from sheep and goats with caseous lymphadenitis

Maria da Conceição Aquino de Sá¹, Gisele Veneroni Gouveia², Carina da Costa Krewer³,
Josir Laine Aparecida Veschi⁴, Ana Luiza de Mattos-Guaraldi⁵ and Mateus Matiuuzzi da Costa¹

¹Programa de Pós-Graduação em Ciência Animal, Universidade Federal do Vale do São Francisco, Petrolina, PE, Brazil.

²Laboratório de Microbiologia e Imunologia Animal, Universidade Federal do Vale do São Francisco, Petrolina, PE, Brazil.

³Programa de Pós-Graduação em Biociência Animal, Universidade Federal Rural de Pernambuco, Recife, PE, Brazil.

⁴Embrapa Semiárido, Petrolina, PE, Brazil.

⁵Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brazil.

Abstract

Caseous lymphadenitis (CL) is a chronic and subclinical disease that affects goats and sheep and, consequently, causes economic losses, especially to small producers. The purpose of this study, through use of Polymerase Chain Reaction (PCR), was to verify the presence of virulence genes of phospholipase D (*PLD*), integral membrane protein (*FagA*), iron enterobactin transporter (*FagB*), ATP binding cytoplasmic membrane protein (*FagC*) and iron siderophore binding protein (*FagD*) in 168 isolates of *C. pseudotuberculosis* obtained from cases of caseous lymphadenitis in goats and sheep. *FagA*, *FagB* and *PLD* genes were detected in all 145 strains isolated from abscesses in superficial lymph nodes and in 23 strains isolated from viscera. The *FagC* gene was positive in 167 (99.40%) isolates. The *FagD* gene was detected in 160 (95.23%) isolates. All virulence factors analyzed were found more frequently among isolates collected in the viscera of animals with CL, indicating a multifactorial nature, as well as variations, in the invasive potential of *C. pseudotuberculosis* strains.

Keywords: virulence genes, *C. pseudotuberculosis*, caseous lymphadenitis.

Received: October 5, 2012; Accepted: February 26, 2013.

Corynebacterium pseudotuberculosis is a facultative intracellular pathogen that mainly infects sheep and goats, causing the disease called caseous lymphadenitis (CL). In disease manifestations, the main characteristic is abscessing of the lymph nodes (Fontaine and Baird, 2008). Considering the current status of CL prevalence worldwide, including Brazil (Connor *et al.*, 2000), there is a pressing need for more efficient alternatives for disease control that do not only cure sick animals, but also minimize or even prevent the onset of the disease in herds (Çetinkaya *et al.*, 2002; D'Afonseca *et al.*, 2008). The high prevalence of CL in sheep and goats has made studies on ways to detect *C. pseudotuberculosis* virulence factors increasingly important. Although the mechanisms by which the disease is caused remain poorly

understood, advances in genomics have allowed the discovery of new genes, especially ones related to *C. pseudotuberculosis* pathogenicity and lifestyle (Quinn *et al.*, 1994; Dorella *et al.*, 2006; Ruiz *et al.*, 2011).

Hemolysis is an important characteristic of the virulence of *C. pseudotuberculosis* because it has been associated with the uptake of iron, which is a micronutrient important for energy gain and bacterial growth (Jost and Billington, 2004). The gene product for phospholipase D (*PLD*) is regarded as the main virulence factor in *C. pseudotuberculosis* (Hodgson *et al.*, 1999). The *PLD* gene encodes the phospholipase D - *PLD* exotoxin, an enzyme that catalyzes the dissociation of sphingomyelin and increases vascular permeability. This leads to the spread (Hodgson *et al.*, 1994) and survival of *C. pseudotuberculosis* in the cells, and consequently the invasion of the body and transport by phagocytes to regional lymph nodes (Baird and Fontaine, 2007). Although it is not considered to be directly hemolytic, it has been reported as capable of producing synergistic hemolysis (Jost and Billington, 2004).

Send correspondence to Gisele Veneroni Gouveia. Laboratório de Microbiologia e Imunologia Animal, Campus de Ciências Agrárias, Universidade Federal do Vale do São Francisco, Rodovia BR-407 km 12, Lote 543 - Projeto de Irrigação Nilo Coelho, C1, s/n°, Zona Rural, 56300-000 Petrolina, PE, Brazil. E-mail: giseleveneroni@yahoo.com.br.

Other important virulence factors include integral membrane protein (*FagA*), iron enterobactin transporter (*FagB*), ATP binding cytoplasmic membrane protein (*FagC*) and iron siderophore binding protein (*FagD*). These are organized in an operon involved in iron uptake, which provides persistence to *C. pseudotuberculosis* in infections in goats. This operon is located downstream from the *PLD* gene, contributing to the virulence of *C. pseudotuberculosis* (Billington *et al.*, 2002). The finding of seven putative pathogenicity islands containing the classical virulence elements, including genes for iron uptake, fimbrial subunits, insertional elements and secreted toxins, probably mostly acquired through horizontal transfer, contributes to the understanding of how this species causes disease (Ruiz *et al.*, 2011).

The aim of this study was to verify the presence of *PLD*, *FagA*, *FagB*, *FagC* and *FagD* virulence genes in strains of *C. pseudotuberculosis* obtained from caseous lymphadenitis in goats and sheep.

A total of 168 isolates of *C. pseudotuberculosis* were used in this study. The strains were obtained from caseous lymphadenitis samples of goats and sheep originating from farms located in different municipalities of the state of Pernambuco, Brazil: Afranio (n = 2), Floresta (n = 4), Jatobá (n = 2), Petrolândia (n = 4) and Petrolina (n = 38), as well as from animals slaughtered in slaughterhouses located in Petrolina, PE (n = 116) and Juazeiro, BA (n = 2). Of these isolates, 151 were obtained from abscesses in superficial lymph nodes and 17 from abscesses in viscera. Isolates of *C. pseudotuberculosis* were previously identified by means of their morphology and biochemical characteristics (Quinn *et al.*, 1994). The cultures were subcultured in a Tryptone Soy Agar (TSA) medium, and then used for molecular characterization.

The DNA from *C. pseudotuberculosis* isolates was heat extracted in a final volume of 500 µL. Initially the strains were grown in BHI (Brain Heart Infusion) medium for 48 h at 37 °C. Subsequently, 10 colony forming units (CFU) were placed in 500 µL of sterile ultrapure water. Thermal lysis of DNA was done at a temperature of 100 °C for 11 min and cooling at 4 °C for 4 min. After centrifugation at 14,000 g for 2 min the supernatant was removed.

PCR was used to check for the presence of *FagA*, *FagB*, *FagC*, *FagD* and *PLD* genes with the use of the specific primers shown in Table 1. The primers for amplification of *FagA*, *FagB*, *FagC*, *FagD* and *PLD* genes were designed with the Primer3 Plus software. The quality of the primers was checked online with the IDT Oligo Analyzer. The primers used for amplification of the *PLD* gene have been described by Hodgson *et al.*, 1990.

The PCR consisted of a final volume of 25 µL containing 4 µL of genomic DNA, 0.4 µM of each primer, 50 mM KCl, 10 mM Tris-HCl pH 8.4, 0.1 mM dNTPs, 2.5 U *Taq* polymerase (Centibiot) and 1 mM of MgCl₂ for *PLD* primer, 1.4 mM for *FagA* and 1.3 mM for *FagB*, *FagC*

Table 1 - Primers used for the detection of virulence genes.

Primer	Sequence	Product size in base pairs
<i>PLD_F</i>	5'ATGAGGGAGAAAGTTGTTTAA3'	924
<i>PLD_R</i>	5'TCACCACGGGTTATCCGC3'	
<i>Fag_A_F</i>	5'AGCAAGACCAAGAGACATGC3'	245
<i>Fag_A_R</i>	5'AGTCTCAGCCCAACGTACAG3'	
<i>Fag_B_F</i>	5'GTGAGAAGAACCCCGGTATAAG3'	291
<i>Fag_B_R</i>	5'TACCGCACTTATTCTGACACTG3'	
<i>Fag_C_F</i>	5'GTTTGGCTATCTCCTTGGTATG3'	173
<i>Fag_C_R</i>	5'CGACCTTAGTGTGACATACCC3'	
<i>Fag_D_F</i>	5'GAGACTATCGACCAGGCAGA3'	226
<i>Fag_D_R</i>	5'ACTTCTTGGGGAGCAGTTCT3'	

and *FagD* primers. The amplification reaction for the *PLD* gene consisted of the following steps: initial denaturation at 94 °C for 30 s, followed by 40 cycles of denaturation at 94 °C for 40 s, annealing at 61 °C for 40 s and extension at 72 °C for 40 s, terminating with a final extension step at 72 °C for 10 min. For the other genes, the difference was in relation to the annealing temperature, which was 58 °C, 55 °C, 60 °C and 61 °C for the *FagA*, *FagB*, *FagC* and *FagD* genes, respectively. The amplification products were then separated in 1.5% agarose gels, stained with ethidium bromide and viewed under ultraviolet light. The specificity of the primers used was confirmed in 1.5% agarose gels, revealing a specific band of the estimated size.

PCR is an accurate, efficient and highly sensitive method that can rapidly lead to the identification of a pathogen in material collected directly from lesions, as well as isolates in culture (Pacheco *et al.*, 2007). The use of PCR also permits to detect the presence of virulence genes, thus demonstrating the potential of a disease causing microorganism.

The *FagA*, *FagB* and *PLD* genes were found in all isolates. The *FagC* gene was present in 99.40% of the isolates tested, and the *FagD* gene had a frequency of 95.23%. Frequencies of virulence genes in *C. pseudotuberculosis* isolates according to their origin (viscera or superficial lymph nodes) are shown in Table 2. The combinations of virulence factors observed in 168 isolates of *C.*

Table 2 - Frequencies of virulence genes in *C. pseudotuberculosis* isolates from viscera or superficial lymph nodes.

Gene	Percentage (%) of positive isolates from viscera	Percentage (%) of positive isolates from superficial lymph nodes
<i>FagA</i>	10.11	89.89
<i>FagB</i>	10.11	89.89
<i>FagC</i>	10.18	89.82
<i>FagD</i>	10.62	89.38
<i>PLD</i>	10.11	89.89

Table 3 - Combinations of virulence genes in *C. pseudotuberculosis* isolates.

Gene combinations	Number of positive isolates from superficial lymph nodes	Number of positive isolates from viscera	Total percentage (%)
<i>FagA/FagB/PLD</i>	1	0	0.60
<i>FagA/FagB/FagC/PLD</i>	7	0	4.17
<i>FagA/FagB/FagC/FagD/PLD</i>	143	17	95.23

pseudotuberculosis are shown in Table 3. All samples collected in the viscera were positive for the five genes tested. The *FagD* gene was less frequent in samples collected from slaughtered animals.

In this study, the *PLD* gene was found in all clinical isolates, which is in agreement with other studies that show the spread of the *PLD* gene among *C. pseudotuberculosis* strains (Hodgson *et al.*, 1990; Çetinkaya *et al.*, 2002; Pacheco *et al.*, 2007). All isolates in this study showed a positive reverse CAMP test, indicating expression of *PLD* (data not shown). The mechanisms by which *PLD* is expressed are not well understood, but it can be regulated by a variety of environmental factors that may involve the binding of repressors or activators or structural changes in DNA (McKean *et al.*, 2007). Pacheco *et al.* (2011) compared the proteomes of two strains and demonstrated that *PLD* was expressed only in the virulent isolate. Attenuation of the *PLD* gene reduces the virulence of *C. pseudotuberculosis* isolates and prevents the development of caseous lymphadenitis (McNamara *et al.*, 1994). Knockout isolates for *PLD* may represent new candidates for the production of live attenuated vaccines (Simmons *et al.*, 1998; Meyer *et al.*, 2002).

The *FagA*, *B*, *C* and *D* putative genes are related to regulation and uptake of iron by *C. pseudotuberculosis* and are located downstream of the *PLD* gene. These genes are rarely expressed *in vitro*, but when expressed *in vivo*, they increase the virulence of *C. pseudotuberculosis*, which indicates that host factors may be necessary for the expression of these genes (Billington *et al.*, 2002). In the present paper, the *FagA*, *B* and *PLD* genes exhibited a rate of 100% in the isolates tested. In a study by Billington *et al.* (2002), a knockout mutant for the *Fag* gene showed reduced virulence in inoculation in goats. This information is important, as all isolates evaluated in this study were virulent. The *FagC* and *FagD* genes were detected less frequently; but still, this was more than 95%. Interestingly, all isolates negative for the *FagD* gene were obtained from superficial abscesses, suggesting differences in the virulence potential of the clinical isolates and the possibility of participation of *FagD* in the invasive mechanisms expressed by some *C. pseudotuberculosis* strains.

Since treatment of caseous lymphadenitis is difficult, vaccination would be an important alternative (Meyer *et al.*, 2002). The potential of interference of the *PLD* and *Fag* genes in bacterial fitness, and their ability to multiply and survive within the host (Jost and Billington, 2004), place

these genes as candidates for production of live vaccines (Hodgson *et al.*, 1999; Billington *et al.*, 2002). Since most studies published so far investigated only a small number of *C. pseudotuberculosis* strains (Billington *et al.*, 2002; Pacheco *et al.*, 2011), the present analysis of a large number of virulent isolates should contribute to mapping the distribution of virulence factors in *C. pseudotuberculosis*.

We could show that genes encoding *PLD* and *FagA-D* virulence factors were present in most of the *C. pseudotuberculosis* strains isolated from animals with CL, indicating a wide distribution in the field and a high pathogenic potential.

Acknowledgments

We thank FACEPE (Fundação de Amparo e Pesquisa de Pernambuco) and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for providing fellowships to MCAS and GVG. This study was financially supported by the CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico N° - 479624/2010-0). This study was authorized by the Ethics Committee in Research in Human and Animal Studies at Univasf under number 27091054 of October 6, 2010.

References

- Baird GJ and Fontaine MC (2007) *Corynebacterium pseudotuberculosis* and its role in ovine caseous lymphadenitis. J Comp Pathol 137:179-210.
- Billington JS, Esmay PA, Songer JG and Jost HB (2002) Identification and role in virulence of putative iron acquisition genes from *Corynebacterium pseudotuberculosis*. FEMS Microbiol Lett 208:41-45.
- Çetinkaya B, Karahan M, Atıl E, Kalın R, De Baere T and Vaneechoutte M (2002) Identification of *Corynebacterium pseudotuberculosis* isolates from sheep and goats by PCR. Vet Microbiol 88:75-83.
- Connor KM, Quirie MM, Baird G and Donachie W (2000) Characterization of United Kingdom isolates of *Corynebacterium pseudotuberculosis* using pulsed-field gel electrophoresis. J Clin Microbiol 38:2633-2637.
- D'Afonseca V, Moraes PM, Dorella FA, Pacheco LGC, Meyer R, Portela RW, Miyoshi A and Azevedo V (2008) A description of genes of *Corynebacterium pseudotuberculosis* useful in diagnostics and vaccine applications. Genet Mol Res 7:252-260.
- Dorella FA, Pacheco LGC, Oliveira SC, Miyoshi A and Azevedo V (2006) *Corynebacterium pseudotuberculosis*: Microbiol-

- ogy, biochemical properties, pathogenesis and molecular studies of virulence. *Vet Rec* 37:201-218.
- Fontaine MC and Baird GJ (2008) Caseous lymphadenitis. *Small Rumin Res* 76:42-48.
- Hodgson ALM, Bird P and Nisbet T (1990) Cloning, nucleotide sequence, and expression in *Escherichia coli* of the phospholipase D gene from *Corynebacterium pseudotuberculosis*. *J Bacteriol* 172:1256-1261.
- Hodgson ALM, Tachedjian M, Corner LA and Radford AJ (1994) Protection of sheep against Caseous Lymphadenitis by use of a single oral dose of live recombinant *Corynebacterium pseudotuberculosis*. *Infect Immun* 62:5275-5280.
- Hodgson ALM, Carter K, Tachedjian M, Krywult J, Corner LA, McColl M and Cameron A (1999) Efficacy of an ovine caseous lymphadenitis vaccine formulated using a genetically inactive form of the *Corynebacterium pseudotuberculosis* Phospholipase D. *Vaccine* 17:802-808.
- Jost BH and Billington SJ (2004) *Corynebacterium* and *Arcanobacterium* in: Gyles CL, Prescott JF, Songer G and Thoen CO (eds) *Pathogenesis of Bacterial Infections in Animals*. 3rd edition. John Wiley & Sons, Yowa, pp 77-86.
- McKean SC, Davies JK and Moore RJ (2007) Expression of phospholipase D, the major virulence factor of *Corynebacterium pseudotuberculosis*, is regulated by multiple environmental factors and plays a role in macrophage death. *Microbiology* 153:2203-2211.
- McNamara PJ, Bradley GA and Songer JG (1994) Targeted mutagenesis of the phospholipase D gene results in decreased virulence of *Corynebacterium pseudotuberculosis*. *Mol Microbiol* 12:921-930.
- Meyer R, Carminati R, Cerqueira RB, Vale V, Viegas S, Martinez T, Nascimento I, Schaer R, Silva JAH, Ribeiro M, et al. (2002) Evaluation of the goats humoral immune response induced by the *Corynebacterium pseudotuberculosis* lyophilized live vaccine. *Cienc Méd Biol* 1:42-48.
- Pacheco LGC, Pena RR, Castro TLP, Dorella FA, Bahia RC, Carminati R, Frota MNL, Oliveira SC, Meyer R, Alves FSF, et al. (2007) Multiplex PCR assay for identification of *Corynebacterium pseudotuberculosis* from pure cultures and for rapid detection of this pathogen in clinical samples. *J Med Microbiol* 56:480-486.
- Pacheco LGC, Slade SE, Seyffert N, Santos AR, Castro TLP, Silva WM, Santos AV, Santos SG, Farias LM, Carvalho MAR, et al. (2011) A combined approach for comparative exoproteome analysis of *Corynebacterium pseudotuberculosis*. *BMC Microbiology* 11:e12.
- Quinn PJ, Carter ME, Markey B and Carter GR (1994) *Clinical Veterinary Microbiology*. Wolfe, London, 648 pp.
- Ruiz JC, D'Afonseca V, Silva A, Ali A, Pinto AC, Santos AR, Rocha AAM, Lopes DO, Dorella FA, Pacheco LGC, et al. (2011) Evidence for reductive genome evolution and lateral acquisition of virulence functions in two *Corynebacterium pseudotuberculosis* strains. *PLoS ONE* 6:e18551.
- Simmons CP, Dunstan SJ, Tachedjian M, Krywult J, Hodgson ALM and Strugnell RA (1998) Vaccine potential of attenuated mutants of *Corynebacterium pseudotuberculosis* in sheep. *Infect Immun* 66:474-479.

Internet Resources

- Primer3 Plus software, http://www.genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi (September 10, 2010).
- IDT Oligo Analyzer, <http://www.idtdna.com/analyzer/Applications/OligoAnalyzer/> (September 10, 2010).

Associate Editor: Célia Maria de Almeida Soares

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.