

Waist–Calf Circumference Ratio Is an Independent Risk Factor of HRQoL in Centenarians

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Shanshan Yang ^{1,2,*}
Miao Liu ^{1,*}
Shengshu Wang¹
Wangping Jia¹
Ke Han¹
Yao He¹

¹Institute of Geriatrics, State Key Laboratory of Kidney Disease, Beijing Key Laboratory of Aging and Geriatrics, The 2nd Clinical Center, Chinese PLA General Hospital, Beijing 100853, People's Republic of China; ²Department of Disease Control, Northern Military Area Center for Disease Control and Prevention, Jinan, People's Republic of China

*These authors contributed equally to this work

Purpose: To analyze the associations between waist circumference (WC), body mass index (BMI), waist–hip ratio (WHR), waist–height ratio (WHtR), calf circumference, waist–calf circumference ratio (WCR), and quality of life in Hainan centenarians.

Patients and Methods: A total of 1002 centenarians in Hainan were selected by a full sample survey. The EQ-5D visual analogue scale (EQ-5D-VAS) was used to investigate the quality of life. Restricted cubic splines were used to analyze and visualize the linear relationships.

Results: After adjustment, the standard β values for BMI, WC, WHR, WHtR, calf circumference, and WCR associated with EQ-5D score were 0.101, 0.126, -0.018 , 0.100, 0.302, and -0.219 , respectively; all associations except for WHR were significant ($P < 0.01$). With increasing BMI, WC, and calf circumference, the risk of EQ-5D score <1 decreased (odds ratios [ORs] 0.91 [95% confidence interval (CI): 0.86–0.97], 0.97 [95% CI: 0.95–0.99], and 0.87 [95% CI: 0.82–0.92] after adjustment, respectively). With increasing WCR, the risk also increased (OR 2.70 [95% CI: 1.54–4.75]).

Conclusion: After excluding nutritional and muscle retention factors, fat central distribution negatively impacted the health-related quality of life of the oldest old population.

Keywords: centenarians, waist–calf circumference ratio, quality of life, obesity

Introduction

Health-related quality of life (HRQoL) is a comprehensive reflection of health-related factors such as physical health, psychological status, independence, social relations, and environmental factors in the elderly population that has received increasing attention.¹ Population aging is the inevitable result of social and economic development. Given the increasing proportions of elderly populations, successful aging (SA) is a cornerstone of healthy societal development. SA refers to the comprehensive bio-psycho-social aspects of elderly populations to not only be healthy in their daily lives and physiological functions but also their moods and social lives.^{2,3} Centenarians may be the templates for SA,⁴ and researchers in the United States, Japan, and Denmark have studied these populations.^{5–7} Studies have also been performed on centenarians in Bama and Rugao in China, but the sample sizes were relatively small.^{8,9}

Obesity is a common risk factor for cardiovascular and cerebrovascular disease, type 2 diabetes mellitus (T2DM), cancer, and other diseases.^{10,11} Obesity is also a risk factor for poor HRQoL.^{12,13} However, the results of studies on the effects of obesity on HRQoL among elderly populations are inconsistent.^{14–16} The “obesity paradox” has been especially prevalent among the elderly older than 80 years and individuals with chronic diseases in recent decades; that is, unlike the general population, obese

Correspondence: Yao He
Institute of Geriatrics State Key
Laboratory of Kidney Disease, Beijing Key
Laboratory of Aging and Geriatrics, The
2nd Clinical Center, Chinese PLA
General Hospital, 28 Fuxing Road, Beijing
100853, People's Republic of China
Tel +86-10-66876411
Fax +86-10-68219351
Email yhe301@x263.net

elderly and obese patients with chronic diseases have better HRQoL and prognosis as well as lower disability and mortality than those with normal weight/body mass index (BMI).^{17,18}

Similar to the general population, body mass index (BMI) and waist circumference (WC) are also generally used as the criteria for obesity in the elderly. However, due to the natural aging progress, it is difficult to measure the height of the elderly and the accuracy of BMI or waist–height ratio (WHtR) in elderly populations is difficult to guarantee, especially those older than 80 years. Furthermore, unlike abdominal obesity (indicated by WC and WHtR) and general obesity (indicated by BMI), peripheral adiposity and larger hip circumference may offer protection from T2DM, cerebrovascular disease, and premature death.¹⁹ Among the elderly, the mechanism of the obesity paradox is largely due to better nutritional status and higher muscle retention.^{17,18} Calf circumference is often used to represent the degree of muscle retention and nutritional status in the elderly.^{20,21} Considering the opposing effects of central obesity and peripheral adiposity, an indicator that assesses both masses simultaneously may better evaluate the risk of obesity on HRQoL than indicators that separately estimate either central obesity or peripheral adiposity; for example, waist–hip ratio (WHR). However, WHR may mask central obesity if both hip circumference and WC increase.²² Recent studies have used the waist–calf ratio (WCR) an index to assess the disproportional relationships between abdominal fat and leg muscle mass and showed it to be an independent predictor of cardiovascular disease and hepatic steatosis and fibrosis.^{23,24} Therefore, the obesity paradox in elderly populations may be related to the use of BMI and WC, which are prone to measurement accuracies and mixed nutritional factors and are, thus, not suitable for evaluating obesity in elderly populations, especially in the oldest old. Furthermore, research on the relationship between obesity and QOL in centenarians is very limited. If the obesity paradox on QOL in centenarians is also due to nutritional factors, the present centenarian study assumed that WC and calf circumference were positively correlated with HRQoL, while WCR and HRQoL were negatively correlated or unrelated.

This study used a series of common obesity evaluation indicators (BMI, WC, WHR, and WHtR) as well as calf circumference and WCR to evaluate the obesity status and analyzed the correlation between obesity status and HRQoL in a cluster sample of centenarians in China.

Materials and Methods

The China Hainan Centenarian Cohort Study (CHCCS)

The present analysis used CHCCS baseline data. The details of the methods have been reported elsewhere.²⁵ Briefly, according to the list of centenarians provided by the Hainan Provincial Civil Affairs Department, a full sample household survey was conducted among all centenarians of Hainan Province between June 2014 and June 2016, excluding those who had died, who failed to pass age verification, and centenarians or their family members who did not or could not cooperate with the examination. Among 1,811 living centenarians according to the household register provided by the civil affairs bureau in 2014, 1,473 were contacted after age verification and address survey and 1,002 centenarians were included in the present survey after excluding those who declined survey participation and those unable to take the physical exam.²⁵ The baseline data of these centenarians were collected by household survey, including questionnaire interviews, physical examinations, and laboratory blood sample testing. Questionnaire interviews including domains such as sociodemographics, functional capacity, cognitive function, behaviors, sleep quality, and quality of life²⁵ and human body indicators (height, weight, WC, hip circumference, calf circumference, blood pressure) were measured by trained local Hainan nurses who spoke the local language and could communicate with the centenarians without linguistic obstacles.

Ethics

The CHCCS was conducted in accordance with the Declaration of Helsinki and was approved by the Medical Ethics Committee of the Chinese PLA General Hospital (301hn11-206-01). All participants provided written informed consent before joining the study.

Exposures

The obesity indicators included BMI, WC, WHR, waist–height ratio (WHtR), calf circumference, and WCR. The body measurements, including height (measured by a unified scale), weight (measured by a unified scale with the participants wearing light clothing), WC, hip, and calf circumference (measured by a tape measure with the participants standing and wearing light clothing), were performed by trained nurses. For the measurements, the centenarians were required to remove their shoes, hats,

and coats and to remove personal belongings such as keys, mobile phones, etc. The height measurements were accurate to 0.5 cm, while the weight measurement required two consecutive results with an error of less than 0.5 kg. The waist and calf circumference measurements were accurate to 0.1 cm.²⁶ BMI = height/weight², WHR= WC/hip circumference, WHtR= WC/height, WCR = WC/calf circumference. According to the Guidelines for Prevention and Control of Overweight and Obesity in Chinese Adults,²⁷ obesity was defined as (1) BMI ≥ 28 kg/m²; (2) WC ≥ 90 cm in men or ≥ 85 cm in women; (3) WHR ≥ 0.9 in men or ≥ 0.8 in women; and (4) WHtR ≥ 0.6 .

Study Outcomes

The EuroQol five dimensions questionnaire-visual analogue scale (EQ-5D-VAS)²⁸ was used to measure HRQoL. The EQ-5D is a validated and extensively used general health questionnaire that covers five health domains (mobility, self-care, usual activities, pain/discomfort, and anxiety/depression) that was also validated in the Chinese elderly population before data collection. The EQ-5D index was calculated using the Japanese population-based time trade-off (TTO) model which was the most suitable tool for Chinese individuals at the time of the present study.²⁹ The EQ-5D scores after TTO conversion ranged from -0.11 to 1. Interviewees without any problems in the five domains of EQ-5D scored 1, and were defined as having a high QOL, while interviewees with EQ-5D scores < 1 were defined as having a low QOL. The VAS is a self-rating tool used to assess health status on a 20-cm vertical scale in which values of 100 (at the top) 0 (at the bottom) indicate the best and worst health statuses, respectively. Information from the EQ-5D and VAS were collected by trained nurses.

Measurements

Information on demographic characteristics and lifestyle including age, sex, ethnicity (Han, Li, or others), education, marital status (married, widowed, divorced, or never married), residential type (living with family or living alone), smoking (never, former, or current), alcohol use (never, former, or current) and physical activity (low, medium, or high)²⁵ was collected via questionnaires administered by trained nurses. Education was assessed and classified into three groups: illiterate (0 years), primary school (1–6 years), or middle school or higher (> 6 years).

Statistical Analysis

IBM SPSS Statistics for Windows, version 24.0 and R 3.5.2 were used for the data analyses. The significance level for all tests was set at a two-tailed α value of 0.05. The differences in the means and proportions were evaluated using Student's t- and chi-square tests, respectively. Linear and logistic regression models were used to identify the associations between obesity indicators and EQ-5D/VAS scores. Restricted cubic splines were used to display and test the relationships between the risk of low QOL (EQ-5D score < 1) and obesity-related indicators (WC, BMI, WHR, WHtR, calf circumference, and WCR). The linear and logistic regression models were repeated in men and women to determine if the associations differed by sex. The graphs were created in R 3.5.2 and were arranged into the final format in Adobe Illustrator.

Patient Involvement

No patients were involved in setting the research question or the outcome measures, nor were they involved in the study design and implementation. There are no plans to involve patients in the dissemination of these findings.

Results

Socio-Demographic Characteristics

Of 1,002 centenarians of CHCCS recruited from June 2014 to June 2016, 822 (82.2%) were females. The mean age was 102.77 (standard deviation [SD] 2.75) years. Compared to that in men, more women more commonly had central obesity as defined by WC (12.8 vs 6.8%, $P=0.021$) and the group with central obesity had similar demographic and lifestyle characteristics including age, ethnicity, education, marital status, residential type, smoking, alcohol use, and physical activity ($P>0.05$, Table 1).

The average EQ-5D index was 0.62 ± 0.25 (range: -0.11–1.00) and the average VAS score was 61.60 ± 15.56 (range: 0–100). The average EQ-5D indices for centenarians with and without central obesity were 0.61 ± 0.25 and 0.65 ± 0.22 , respectively. The overall coefficient of EQ-5D and VAS was 0.414 ($P<0.01$). Scores > 1 were observed in 873 centenarians (143 men and 730 women).

The average WC, BMI, WHR, WHtR, calf circumference, and WCR were 75.27 ± 8.79 cm, 18.11 ± 3.22 kg/m², 0.90 ± 0.14 , 0.52 ± 0.07 , 24.73 ± 3.67 cm, and 3.08 ± 0.41 , respectively (Table 1).

Table 1 Sociodemographic Variables of the CHCCS Participants

	All (n=1002)	Normal (n=885)	Central Obesity (n=117)	P
Mean±SD				
Age (years)	102.77±2.75	102.82±2.79	102.38±2.41	0.110
Height (cm)	144.52±8.95	144.40±9.05	145.45±8.12	0.230
Weight (kg)	37.85±7.69	37.09±7.16	43.61±8.99	<0.001
Waist circumference (cm)	75.27±8.79	73.23±6.84	90.68±6.2	<0.001
BMI	18.11±3.22	17.78±2.92	20.66±4.13	<0.001
WHR	0.90±0.14	0.89±0.14	0.98±0.09	<0.001
WHtR	0.52±0.07	0.51±0.05	0.63±0.05	<0.001
Calf circumference	24.73±3.67	24.39±3.21	27.09±3.15	<0.001
WCR	3.08±0.41	3.04±0.39	3.39±0.43	<0.001
EQ5D	0.62±0.25	0.61±0.25	0.65±0.22	0.113
VAS	61.60±15.56	61.41±15.76	63.05±13.93	0.283
N (%)				
Gender				0.021
Male	180 (18.0)	168 (93.3)	12 (6.7)	
Female	822 (82.0)	717 (87.2)	105 (12.8)	
Ethnic				0.175
Han	883 (88.1)	777 (88.0)	106 (12.0)	
Li	106 (10.6)	98 (92.5)	8 (7.5)	
Others	13 (1.3)	10 (76.9)	3 (23.1)	
Marital Status				0.269
Married	100 (10.0)	90 (90.0)	10 (10.0)	
Widowed	836 (83.4)	733 (87.7)	103 (12.3)	
Divorced or never married	66 (6.6)	62 (93.9)	4 (6.1)	
Education Level				0.603
Illiterate	915 (91.3)	806 (88.1)	109 (11.9)	
Primary school	67 (6.7)	60 (89.6)	7 (10.4)	
Middle school or higher	20 (2.0)	19 (95.0)	1 (5.0)	
Residential Type				0.358
Living together with families	863 (86.1)	759 (87.9)	104 (12.1)	
Living alone at home	139 (13.9)	126 (90.6)	13 (9.4)	
Body Mass Index (kg/m ²)				<0.001
Underweight (<18.5)	575 (57.4)	538 (93.6)	37 (6.4)	
Normal (18.5–24.0)	391 (39.0)	327 (83.6)	64 (16.4)	
Overweight (24.0–27.9)	28 (2.8)	20 (71.4)	8 (28.6)	
Obese (≥28.0)	8 (0.8)	0 (0.0)	8 (100.0)	
WHR				<0.001
Normal	177 (17.7)	176 (99.4)	1 (0.6)	
Obese	825 (82.3)	709 (85.9)	116 (14.1)	
WHtR				<0.001
Normal	128 (12.8)	841 (96.2)	33 (3.8)	
Obese	874 (87.2)	44 (34.4)	84 (65.6)	
Smoking Status				0.574
Non-smoker	919 (91.7)	809 (88.0)	110 (12.0)	
Former	52 (5.2)	47 (90.4)	5 (9.6)	

(Continued)

Table 1 (Continued).

	All (n=1002)	Normal (n=885)	Central Obesity (n=117)	P
Current	31 (3.1)	29 (93.5)	2 (6.5)	
Drinking Status				0.129
Non-drinker	872 (87.0)	764 (87.6)	108 (12.4)	
Former	81 (8.1)	77 (95.1)	4 (4.9)	
Current	49 (4.9)	44 (89.8)	5 (10.2)	
Physical Activity				0.449
Low	874 (87.2)	768 (87.9)	106 (12.1)	
Medium	37 (3.7)	33 (89.2)	4 (10.8)	
High	91 (9.1)	84 (92.3)	7 (7.7)	

Results of Linear Regression Models

Obesity-related indicators including WC, BMI, WHR, WHtR, calf circumference, and WCR were included in the linear regression models as continuous variables. Demographic and lifestyle characteristics including age, sex, ethnicity, education, marital status, residential type, smoking, alcohol use, and physical activity were included as adjust variables by the stepwise method.

As shown in Table 2, with the EQ-5D index as the dependent variable, WC, BMI, WHtR, and calf circumference had positive impacts on the EQ-5D index. After adjustment, the standard β values were 0.101, 0.126, 0.100, and 0.302, respectively (P <0.001), while WCR negatively impacted the EQ-5D index (standard β=-0.219 after adjustment, P<0.001). Similar results were observed in women. In men, only calf circumference and WCR significantly impacted EQ-5D after adjustment.

A shown in Table 3, with VAS as the dependent variable, WC, BMI, WHtR, and calf circumference positively impacted the VAS score. The standard β values were 0.082, 0.076, and 0.156 after adjustment, respectively (P =0.009, 0.014, and <0.001, respectively), while WCR negatively impacted the VAS score (standard β=-0.117 after adjustment, P<0.001). Similar results were found in women. The associations were not significant in men.

Results of Logistic Regression Models

Table 4 model C shows that, after adjustment, compared to lower BMI, WC, and calf circumference, the odds ratios (AORs) of EQ-5D score <1 for centenarians with 1 kg/m² or 1 cm increases were 0.91 (95% CI: 0.86–0.97), 0.97 (95% CI: 0.95–0.99) and 0.87 (0.82–0.92), respectively (Table 4). WCR was a risk factor for low QOL ((EQ-5D score <1) (OR=2.70,

Table 2 Association Between BMI, WC, WHR, WHtR, Calf Circumference, WCR, and EQ-5D

	Model A				Model B				Model C				P		
	β	95% CI		Standard β	P	β	95% CI		Standard β	P	β	95% CI		Standard β	
		Lower	Upper				Lower	Upper				Lower			Upper
Male															
BMI	-0.003	-0.016	0.010	-0.034	0.652	-0.003	-0.016	0.010	-0.036	0.629	-0.002	-0.014	0.010	-0.028	0.696
WC	-0.002	-0.006	0.002	-0.064	0.392	-0.003	-0.007	0.002	-0.097	0.203	-0.003	-0.007	0.002	-0.087	0.230
WHR	-0.333	-0.806	0.139	-0.104	0.166	-0.313	-0.782	0.155	-0.098	0.188	-0.230	-0.680	0.219	-0.072	0.313
WHtR	-0.468	-1.079	0.143	-0.113	0.133	-0.527	-1.135	0.082	-0.127	0.089	-0.417	-1.005	0.172	-0.100	0.164
Calf circumference	0.015	0.005	0.025	0.212	0.004	0.013	0.023	0.190	0.011	0.011	0.015	0.005	0.024	0.213	0.003
WCR	-0.166	-0.256	-0.076	-0.264	<0.001	-0.168	-0.257	-0.078	-0.266	<0.001	-0.168	-0.252	-0.083	-0.266	<0.001
Female															
BMI	0.010	0.005	0.015	0.132	0.000	0.010	0.005	0.015	0.132	0.000	0.009	0.004	0.014	0.123	0.000
WC	0.004	0.003	0.006	0.159	0.000	0.004	0.003	0.006	0.158	0.000	0.005	0.003	0.007	0.168	0.000
WHR	-0.030	-0.142	0.082	-0.018	0.597	-0.031	-0.144	0.081	-0.019	0.583	-0.006	-0.114	0.102	-0.004	0.916
WHtR	0.483	0.229	0.737	0.129	0.000	0.483	0.229	0.738	0.129	0.000	0.507	0.264	0.750	0.136	0.000
Calf circumference	0.024	0.019	0.029	0.342	0.000	0.024	0.020	0.029	0.343	0.000	0.022	0.017	0.026	0.310	0.000
WCR	-0.151	-0.191	-0.110	-0.246	<0.001	-0.151	-0.192	-0.110	-0.246	<0.001	-0.123	-0.163	-0.083	-0.200	<0.001
Total															
BMI	0.009	0.005	0.014	0.122	<0.001	0.008	0.003	0.013	0.107	0.001	0.008	0.003	0.012	0.101	0.001
WC	0.004	0.002	0.005	0.130	<0.001	0.003	0.002	0.005	0.117	0.000	0.004	0.002	0.005	0.126	<0.001
WHR	-0.054	-0.163	0.055	-0.031	0.330	-0.046	-0.154	0.063	-0.026	0.411	-0.032	-0.136	0.072	-0.018	0.551
WHtR	0.267	0.033	0.501	0.071	0.025	0.342	0.107	0.576	0.090	0.004	0.377	0.152	0.602	0.100	0.001
Calf circumference	0.023	0.019	0.027	0.338	<0.001	0.022	0.018	0.026	0.327	<0.001	0.020	0.017	0.024	0.302	<0.001
WCR	-0.161	-0.198	-0.125	-0.268	<0.001	-0.154	-0.191	-0.117	-0.255	<0.001	-0.132	-0.168	-0.096	-0.219	<0.001

Notes: Model A: Crude model; Model B: Adjusted for gender, age, ethnic, education level, residential type; Model C: Adjusted for gender, age, ethnic, education level, residential type, smoking, drinking, physical activity.

Table 3 Association Between BMI, WC, WHR, WHtR, Calf Circumference, WCR, and VAS

	Model A				Model B				Model C				P		
	β	95% CI		Standard β	P	β	95% CI		Standard β	P	β	95% CI		Standard β	
		Lower	Upper				Lower	Upper				Lower			Upper
Male															
BMI	0.182	-0.573	0.937	0.036	0.634	0.219	-0.532	0.971	0.043	0.565	0.230	-0.511	0.971	0.045	0.541
WC	0.044	-0.211	0.299	0.025	0.735	0.021	-0.237	0.279	0.012	0.874	0.028	-0.226	0.282	0.016	0.828
WHR	-1.1311	-39.527	16.906	-0.059	0.430	-9.447	-37.436	18.541	-0.049	0.506	-6.248	-33.936	21.440	-0.033	0.657
WHtR	-2.051	-38.636	34.535	-0.008	0.912	-2.717	-39.227	33.793	-0.011	0.883	0.113	-36.260	36.486	0.000	0.995
Calf circumference	0.569	-0.035	1.172	0.138	0.065	0.490	-0.118	1.099	0.119	0.113	0.542	-0.056	1.139	0.131	0.075
WCR	-4.381	-9.888	1.125	-0.117	0.118	-4.100	-9.592	1.391	-0.109	0.142	-4.177	-9.561	1.208	-0.111	0.128
Female															
BMI	0.450	0.121	0.779	0.093	0.007	0.454	0.126	0.782	0.094	0.007	0.434	0.110	0.758	0.090	0.009
WC	0.155	0.034	0.276	0.087	0.012	0.150	0.029	0.271	0.084	0.015	0.161	0.041	0.280	0.091	0.008
WHR	-1.302	-8.391	5.787	-0.013	0.719	-1.426	-8.504	5.652	-0.014	0.693	-1.118	-8.147	5.911	-0.011	0.755
WHtR	15.457	-0.720	31.635	0.065	0.061	15.391	-0.746	31.528	0.065	0.062	16.433	0.499	32.367	0.069	0.043
Calf circumference	0.761	0.461	1.061	0.171	<0.001	0.772	0.472	1.071	0.174	<0.001	0.700	0.401	0.998	0.158	<0.001
WCR	-5.312	-7.946	-2.678	-0.137	<0.001	-5.396	-8.023	-2.768	-0.139	<0.001	-4.406	-7.052	-1.760	-0.114	0.001
Total															
BMI	0.485	0.187	0.783	0.100	0.001	0.409	0.109	0.708	0.085	0.008	0.396	0.100	0.692	0.082	0.009
WC	0.152	0.043	0.262	0.086	0.006	0.124	0.015	0.234	0.070	0.026	0.135	0.027	0.243	0.076	0.014
WHR	-2.321	-9.156	4.514	-0.021	0.505	-1.845	-8.632	4.942	-0.017	0.594	-1.525	-8.220	5.170	-0.014	0.655
WHtR	8.443	-6.223	23.110	0.036	0.259	12.319	-2.404	27.042	0.052	0.101	13.720	-0.817	28.256	0.058	0.064
Calf circumference	0.802	0.543	1.060	0.189	<0.001	0.713	0.444	0.981	0.168	<0.001	0.661	0.395	0.927	0.156	<0.001
WCR	-5.855	-8.169	-3.541	-0.155	<0.001	-5.231	-7.599	-2.862	-0.139	<0.001	-4.401	-6.763	-2.039	-0.117	<0.001

Notes: Model A: Crude model; Model B: Adjusted for gender, age, ethnic, education level, residential type; Model C: Adjusted for gender, age, ethnic, education level, residential type, smoking, drinking, physical activity.

Table 4 Odds Ratios (ORs) for Having a Low QOL

	Model A	Model B	Model C
Male			
BMI	0.95 (0.83–1.08)	0.95 (0.83–1.09)	0.93 (0.81–1.08)
WC	0.99 (0.95–1.03)	1.00 (0.95–1.04)	0.99 (0.95–1.04)
WHR	0.24 (0.00–23.95)	0.23 (0.00–24.81)	0.08 (0.00–11.62)
WHtR	0.60 (0.00–304.59)	1.43 (0.00–942.95)	0.38 (0.00–392.32)
Calf circumference	0.87 (0.78–0.98)	0.88 (0.79–0.99)	0.87 (0.77–0.98)
WCR	2.76 (0.93–8.20)	3.16 (1.03–9.69)	3.39 (1.06–10.89)
BMI obese			
WC obese	3.00 (0.37–24.01)	3.31 (0.41–27.05)	2.20 (0.26–18.49)
WHR obese	0.80 (0.38–1.66)	0.83 (0.39–1.76)	0.73 (0.33–1.61)
WHtR obese	1.58 (0.18–13.52)	1.41 (0.16–12.52)	1.34 (0.15–12.34)
Female			
BMI	0.91 (0.86–0.97)	0.91 (0.86–0.97)	0.90 (0.84–0.97)
WC	0.97 (0.95–1.00)	0.97 (0.95–1.00)	0.97 (0.94–0.99)
WHR	3.59 (0.30–43.03)	3.93 (0.31–49.41)	2.94 (0.20–42.26)
WHtR	0.13 (0.01–3.22)	0.13 (0.01–3.31)	0.06 (0.00–2.05)
Calf circumference	0.85 (0.80–0.92)	0.86 (0.80–0.92)	0.87 (0.81–0.94)
WCR	3.43 (1.84–6.39)	3.46 (1.85–6.46)	2.45 (1.28–4.70)
BMI obese	0.88 (0.11–7.24)	0.92 (0.11–7.61)	0.67 (0.08–5.83)
WC obese	0.56 (0.32–0.98)	0.56 (0.32–0.98)	0.48 (0.26–0.87)
WHR obese	1.44 (0.71–2.93)	1.46 (0.71–2.97)	1.49 (0.70–3.15)
WHtR obese	0.80 (0.45–1.42)	0.82 (0.46–1.47)	0.76 (0.41–1.40)
Total			
BMI	0.91 (0.86–0.96)	0.92 (0.87–0.97)	0.91 (0.86–0.97)
WC	0.97 (0.95–0.99)	0.98 (0.96–0.99)	0.97 (0.95–0.99)
WHR	2.41 (0.33–17.48)	2.26 (0.31–16.69)	1.82 (0.24–13.88)
WHtR	0.42 (0.03–6.90)	0.20 (0.01–3.60)	0.09 (0.00–1.88)
Calf circumference	0.85 (0.80–0.90)	0.86 (0.81–0.91)	0.87 (0.82–0.92)
WCR	3.74 (2.21–6.33)	3.39 (1.96–5.85)	2.70 (1.54–4.75)
BMI obese	1.04 (0.13–8.48)	0.90 (0.11–7.45)	0.77 (0.09–6.49)
WC obese	0.73 (0.43–1.24)	0.68 (0.40–1.17)	0.59 (0.34–1.03)
WHR obese	1.58 (1.01–2.45)	1.12 (0.65–1.92)	1.08 (0.61–1.92)
WHtR obese	0.96 (0.56–1.66)	0.86 (0.49–1.50)	0.81 (0.45–1.45)

Notes: Model A: Crude model; Model B: Adjusted for gender, age, ethnic, education level, residential type; Model C: Adjusted for gender, age, ethnic, education level, residential type, smoking, drinking, physical activity. Bold Values: P<0.05.

95% CI: 1.54–4.75) Table 4). When BMI, WC, WHR, and WHtR included involved as binary variables (obesity or not), the association was no longer significant (Table 4). Similar results were observed in women. In men, only calf circumference and WCR significantly impacted low QOL.

Further division of BMI, WC, WHR, WHtR, calf circumference, and WCR by quintile and inclusion as categorical variables in the logistic regression models showed that, compared to Q1, an increase in calf circumference decreased the risk of low QOL, while an increase in WCR increased the risk (P for trend <0.01, OR of Q5 calf circumference: 0.11 [95% CI: 0.05–0.27], OR of Q5 WCR: 3.73 [95% CI: 1.76–7.93] after adjustment) (Table 5).

Restricted Cubic Splines

Restricted cubic splines were used to display and test the relationships between the risk of QOL and obesity-related indicators (WC, BMI, WHR, WHtR, calf circumference, and WCR) (Figure 1) (non-linear P >0.05).

Discussion

The results of this study showed that calf circumference and WCR is a risk factor for HRQoL and that calf circumference was a protective factor among centenarians in China. To our knowledge, this study was the first to show the association between obesity-related indicators

Table 5 Odds Ratios (ORs) for Having a Low QOL (Quintile)

	Model A	Model B	Model C
BMI			
Q1	1	1	1
Q2	0.45(0.23–0.90)	0.45(0.22–0.91)	0.41 (0.20–0.86)
Q3	0.52(0.26–1.06)	0.53(0.26–1.08)	0.52 (0.25–1.09)
Q4	0.34(0.17–0.67)	0.35(0.18–0.70)	0.34 (0.17–0.70)
Q5	0.36(0.19–0.72)	0.36(0.18–0.71)	0.33 (0.16–0.67)
WC			
Q1	1	1	1
Q2	1.21(0.63–2.32)	1.28(0.66–2.49)	0.99 (0.50–1.98)
Q3	0.87(0.48–1.59)	0.85(0.47–1.57)	0.83 (0.44–1.57)
Q4	0.61(0.36–1.05)	0.63(0.36–1.10)	0.54 (0.30–0.97)
Q5	0.56(0.32–0.99)	0.60(0.34–1.06)	0.48 (0.26–0.88)
WHR			
Q1	1	1	1
Q2	1.33(0.76–2.32)	1.39(0.79–2.44)	1.50 (0.83–2.71)
Q3	1.32(0.73–2.40)	1.31(0.72–2.40)	1.46 (0.78–2.76)
Q4	1.78(0.97–3.23)	1.80(0.98–3.31)	1.70 (0.90–3.21)
Q5	1.09(0.62–1.92)	1.15(0.65–2.03)	1.16 (0.64–2.12)
WHtR			
Q1	1	1	1
Q2	0.77(0.41–1.44)	0.82(0.44–1.56)	0.82 (0.42–1.61)
Q3	0.90(0.48–1.71)	0.97(0.51–1.86)	0.92 (0.46–1.81)
Q4	0.57(0.31–1.06)	0.60(0.32–1.11)	0.55 (0.29–1.05)
Q5	0.68(0.37–1.26)	0.71(0.38–1.33)	0.60 (0.31–1.16)
Calf Circumference			
Q1	1	1	1
Q2	0.25 (0.11–0.60)	0.25 (0.11–0.60)	0.28 (0.12–0.68)
Q3	0.18 (0.07–0.46)	0.19 (0.08–0.48)	0.23 (0.09–0.59)
Q4	0.14 (0.06–0.31)	0.14 (0.06–0.33)	0.19 (0.08–0.44)
Q5	0.09 (0.04–0.21)	0.10 (0.04–0.24)	0.11 (0.05–0.27)
WCR			
Q1	1	1	1
Q2	1.56 (0.94–2.60)	1.45 (0.86–2.43)	1.26 (0.73–2.17)
Q3	1.91 (1.11–3.26)	1.75 (1.01–3.04)	1.58 (0.88–2.82)
Q4	2.67 (1.50–4.77)	2.43 (1.34–4.38)	2.03 (1.09–3.76)
Q5	5.53 (2.7–11.34)	5.02 (2.4–10.46)	3.73 (1.76–7.93)

Notes: Model A: Crude model; Model B: Adjusted for gender, age, ethnic, education level, residential type; Model C: Adjusted for gender, age, ethnic, education level, residential type, smoking, drinking, physical activity. Bold Values: P<0.05.

(WC, BMI, WHR, WHtR, calf circumference, and WCR) and HRQoL in centenarians. The major strength of our study was its comprehensive control and adjustment for a wide range of potential confounders using different statistical models. The similarity in results demonstrated their robustness. Moreover, as the accuracy of height measurement in the oldest old population is difficult to guarantee, in addition to commonly used indicators such as BMI, WC, WHR, and WHtR, this study also included calf circumference and WCR as indicators of nutrition/muscle

retention and central obesity. Calf circumference and WCR were more powerful protective and risk factors, respectively, for HRQoL.

The correlation coefficient of EQ-5D and VAS in this centenarian sample was 0.414, slightly lower than that reported in the fifth National Health Service Survey of China in 2013³⁰ and higher than that in the British population.³¹ This discordance may be due to differences in social structures and culture as well as different ages of the study samples. A study of elderly people over 72 years of age in German communities reported an EQ-5D index of 78.3+15.8; they used the original value from 0 to 100 and did not report the EQ-5D index calculated by TTO.³² The average VAS in our population was 61.60 +15.56, which was higher than that in a previous survey based on cognitive impairment in the elderly. In this study, 12.9% of participants had no problems based on EQ-5D score (score=1), a rate higher than that among elderly with cognitive impairment (6.1%). This observation indicates that the centenarians in the present study were relatively healthy.³³

Studies on the impact of obesity on HRQoL in centenarians are limited. A study on this impact in Spanish adults (aged above 18 years) showed that HRQoL decreased along with BMI.³⁴ Worldwide, most studies in adults have confirmed the significant negative correlation between BMI and HRQoL using the SF-36 or EQ-5D to evaluate HRQoL.^{35–37} The obesity paradox in elderly populations shows that, compared to normal-weight elderly and patients with chronic disease, obese elderly people and obese patients had higher HRQoL, better prognosis, lower disability, and lower mortality.^{17,18} However, the criteria for evaluating obesity in these studies were usually BMI or WC; due to the natural aging progress, it is difficult to accurately measure the height of the elderly. Secondly, it is difficult to exclude the impact of nutrition and muscle retention in WC measurements; thus, the conclusion that obesity was beneficial to maintain a better HRQoL was biased. Furthermore, WHR is generally used to balance the relationship between fat distribution and nutrition or muscle retention; however, this measurement may mask central obesity if both hip circumference and WC increase.²² In this study, increased BMI and WC resulted in increased HRQoL (reflected by EQ-5D-VAS) and decreased risk of low QOL in centenarians. The WHR was not significantly correlated with HRQoL in this population. However, the analysis of calf circumference and WCR showed that the protective effect of calf




Figure 1 Restricted cubic splines of the relationship between (A) BMI, (B) WC, (C) WHR, (D) WHtR, (E) calf circumference, (F) WCR and the risk of EQ-5D scored less than 1 after adjusting for gender, age, ethnic, education level, residential type, smoking, drinking, physical activity.

circumference (the indicator of nutrition and muscle retention^{20,21}) on HRQoL compared to those for BMI and WC. We also observed that WCR (the indicator simultaneously assessing both central obesity and nutrition and muscle retention and excluding the effects of nutrition and muscle retention to reflect the central distribution of fat),

was a risk factor for HRQoL in centenarians. The results of the restrictive cubic spline analysis also showed the linear correlation between WCR and the risk of low QoL.

The major strength of our study was the comprehensive control and adjustment for a wide range of potential confounders using different statistical models. Moreover, to our

knowledge, this study is the first to analyze the correlations between obesity-related indicators (especially calf circumference and WCR) and HRQoL in a large sample of Chinese centenarians.

This study has several limitations. Firstly, this study used baseline data from the CHCCS; thus, these cross-sectional data do not permit causal deduction. Secondly, the large sample of centenarians of this study had lived in the island environment for nearly their entire lives; therefore, caution is required in extrapolating the conclusions of the present study. Further, as the results and implications were based on centenarians, caution is required regarding the external validity. Thirdly, due to the natural aging of the elderly population, there may be errors in height measurement; thus, BMI and WHtR may have had corresponding errors that may have consequently affected the correlation analysis of BMI/WHtR and EQ-5D-VAS.

Conclusions

Despite the limitations, this study analyzed the correlations between obesity-related indicators (especially calf circumference and WCR) and the HRQoL in a large sample of Chinese centenarians. In this study, calf circumference (the indicator of nutrition and muscle retention) was a protective factor for the HRQoL of centenarians, while WCR was a risk factor. These results showed the negative impact of central fat distribution on HRQoL in the oldest old population after excluding nutritional and muscle retention factors. As the height of elderly individuals is difficult to measure, calf circumference and WCR can be used as measurement indicators of nutrition and obesity, respectively, in these populations.

Abbreviations

HRQoL, Health-related quality of life; WC, Waist circumference; BMI, Body mass index; WCR, Waist-calf circumference ratio; WHR, Waist-hip ratio; WHtR, Waist-height ratio; CHCCS, The China Hainan Centenarian Cohort Study; EQ-5D-VAS, EuroQol five dimensions questionnaire-visual analogue scale; TTO, time trade-off.

Data Sharing Statement

All data relevant to the study are included in the article.

Acknowledgments

The abstract for this paper was presented at the Eighth Chinese Congress on Gerontology and Health Industry and was published in the Journal of the American Geriatrics

Society, Vol 67, Issue S4. Shanshan Yang and Miao Liu are co-first authors for this study.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

This study was supported by the National Natural Science Foundation of China (81773502, 81703308, and 81703285).

Disclosure

The authors report no conflicts of interest in this work.

References

- Rahman AR. Development of the World Health Organization WHOQOL-BREF quality of life assessment. The WHOQOL Group. *Psychol Med.* 1998;28(3):551–558. doi:10.1017/S0033291798006667
- Depp CA, Jeste DV. Definitions and predictors of successful aging: a comprehensive review of larger quantitative studies. *Am J Geriatric Psychiatry Off J Am Assoc Geriatric Psychiatry.* 2006;14(1):6–20. doi:10.1097/01.JGP.0000192501.03069.bc
- Schulz R, Heckhausen J. A life span model of successful aging. *Am Psychol.* 1996;51(7):702–714. doi:10.1037/0003-066X.51.7.702
- Motta M, Bennati E, Ferlito L, Malaguamera M, Motta L. Successful aging in centenarians: myths and reality. *Arch Gerontol Geriatrics.* 2005;40(3):241–251.
- Arnold J, Dai J, Nahapetyan L, et al. Predicting successful aging in a population-based sample of Georgia centenarians. *Current Gerontol Geriatrics Res.* 2014;2010(3):9.
- Willcox BJ, Willcox DC, Suzuki M. Demographic, phenotypic, and genetic characteristics of centenarians in Okinawa and Japan: part 1? Centenarians in Okinawa. *Mech Ageing Devel.* 2017;165:75–79.
- Andersen-Ranberg K, Schroll M, Jeune B. Healthy centenarians do not exist, but autonomous centenarians do: a population-based study of morbidity among Danish centenarians. *J Am Geriatr Soc.* 2001;49(7):900–908. doi:10.1046/j.1532-5415.2001.49180.x
- Liu Z, Wang Y, Zhang Y, et al. Cohort profile: the Rugao Longevity and Ageing Study (RuLAS). *Int J Epidemiol.* 2015;45(4):1064–1073.
- Yang X, Wang X, Yao H, et al. Mitochondrial DNA polymorphisms are associated with the longevity in the Guangxi Bama population of China. *Mol Biol Rep.* 2012;39(9):9123–9131. doi:10.1007/s11033-012-1784-8.
- Skinner AC, Perrin EM, Moss LA, Skelton JA. Cardiometabolic risks and severity of obesity in children and young adults. *N Engl J Med.* 2015;373(14):1307–1317. doi:10.1056/NEJMoa1502821
- Calle EE, Kaaks R. Overweight, obesity and cancer: epidemiological evidence and proposed mechanisms. *Nat Rev Cancer.* 2004;4(8):579–591. doi:10.1038/nrc1408
- Hassan MK, Joshi AV, Madhavan SS, Amonkar MM. Obesity and health-related quality of life: a cross-sectional analysis of the US population. *Int J Obesity Related Metab Disord.* 2003;27(10):1227–1232. doi:10.1038/sj.ijo.0802396

13. McDonough C, Dunkley AJ, Aujla N, Morris D, Davies MJ, Khunti K. The association between body mass index and health-related quality of life: influence of ethnicity on this relationship. *Diabetes Obes Metab.* 2013;15(4):342–348. doi:10.1111/dom.2013.15.issue-4
14. Wang L, Crawford JD, Reppermund S, et al. Body mass index and waist circumference predict health-related quality of life, but not satisfaction with life, in the elderly. *Qual Life Res.* 2018;27(10):2653–2665. doi:10.1007/s11136-018-1904-6
15. Xu KY, Wisnivesky JP, Martynenko M, et al. Assessing the association of obesity and asthma morbidity in older adults. *Ann Allergy Asthma Immunol.* 2016;117(1):33–37. doi:10.1016/j.anai.2016.04.027
16. Yin Z, Shi X, Kraus VB, et al. Gender-dependent association of body mass index and waist circumference with disability in the Chinese oldest old. *Obesity (Silver Spring, Md).* 2014;22(8):1918–1925. doi:10.1002/oby.v22.8
17. Lavie CJ, De Schutter A, Parto P, et al. Obesity and prevalence of cardiovascular diseases and prognosis—the obesity paradox updated. *Prog Cardiovasc Dis.* 2016;58(5):537–547. doi:10.1016/j.pcad.2016.01.008
18. Neeland IJ, Das SR, Simon DN, et al. The obesity paradox, extreme obesity, and long-term outcomes in older adults with ST-segment elevation myocardial infarction: results from the NCDR. *Eur Heart J Qual Care Clin Outcomes.* 2017;3(3):183–191. doi:10.1093/ehjqcco/qcx010
19. Heitmann BL. Thigh circumference and risk of heart disease and premature death: prospective cohort study. *BMJ.* 2009;339(7723):704–705. doi:10.1136/bmj.b3292
20. Chumlea WC, Guo SS, Vellas B, Guigoz Y. Techniques of assessing muscle mass and function (sarcopenia) for epidemiological studies of the elderly. *J Gerontol Ser A Biol Sci Med Sci.* 1995;50 Spec No:45–51. doi:10.1093/gerona/50a.special_issue.45
21. Bonnefoy M, Jauffret M, Kostka T, Jusot JF. Usefulness of calf circumference measurement in assessing the nutritional state of hospitalized elderly people. *Gerontology.* 2002;48(3):162–169. doi:10.1159/000052836
22. Despres JP, Lemieux I, Prudhomme D. Treatment of obesity: need to focus on high risk abdominally obese patients. *BMJ Chin Edn.* 2001;322(7288):716–720. doi:10.1136/bmj.322.7288.716
23. Kim SK, Choi YJ, Huh BW, et al. Ratio of waist-to-calf circumference and carotid atherosclerosis in Korean patients with Type 2 diabetes. *Diabetes Care.* 2011;34(9):2067–2071. doi:10.2337/dc11-0743
24. Choe EY, Lee YH, Choi YJ, et al. Waist-to-calf ratio is an independent predictor of hepatic steatosis and fibrosis patients with type 2 diabetes. *J Gastroenterol Hepatol.* 2017;33:5.
25. He Y, Zhao Y, Yao Y, et al. Cohort profile: the China Hainan Centenarian Cohort Study (CHCCS). *Int J Epidemiol.* 2018. doi:10.1093/ije/dyy017
26. Batsis JA, Singh S, Lopezjimenez F. Anthropometric measurements and survival in older Americans: results from the third National Health and Nutrition Examination Survey. *J Nutri Health Aging.* 2014;18(2):123–130. doi:10.1007/s12603-013-0366-3
27. WGO. Guidelines for prevention and control of overweight and obesity in Chinese adults. *Acta Nutrimenta Sinica.* 2004;26(1):1–4.
28. Krabbe P, Weijnen T. Guidelines for analysing and reporting EQ-5D outcomes. In: Brooks R, Rabin R, de Charro F, editors. *The Measurement and Valuation of Health Status Using EQ-5D: A European Perspective: Evidence from the EuroQol BIOMED Research Programme.* Dordrecht: Springer Netherlands; 2003:7–19.
29. Wu YQ, Liu K, Tang X, et al. Empirical research of measuring elderly health utility in the outskirts of Beijing by using European quality of life 5-dimensions. *J Peking Univ.* 2012;44(3):397.
30. Zhang Yaoguang YQ, Lin X. Study on the relationship between health-related quality of life and its relevance from the individual and group perspective. *Chin Health Econ.* 2018;37(1):77–80.
31. Whyne DK. Correspondence between EQ-5D health state classifications and EQ VAS scores. *Health Qual Life Outcomes.* 2008;6(1):94. doi:10.1186/1477-7525-6-94
32. Thiem U, Klaaßen-Mielke R, Trampisch U, Moschny A, Pientka L, Hinrichs T. Falls and EQ-5D rated quality of life in community-dwelling seniors with concurrent chronic diseases: a cross-sectional study. *Health Qual Life Outcomes.* 2014;12(1):2. doi:10.1186/1477-7525-12-2
33. Wolfs CA, Dirksen CD, Kessels A, Willems DC, Verhey FR, Severens JL. Performance of the EQ-5D and the EQ-5D+C in elderly patients with cognitive impairments. *Health Qual Life Outcomes.* 2007;5(1):33. doi:10.1186/1477-7525-5-33
34. Busutil R, Espallardo O, Torres A, Martínez-Galdeano L, Zozaya N, Hidalgo-Vega Á. The impact of obesity on health-related quality of life in Spain. *Health Qual Life Outcomes.* 2017;15(1):197. doi:10.1186/s12955-017-0773-y
35. Kearns B, Ara R, Young T, Relton C. Association between body mass index and health-related quality of life, and the impact of self-reported long-term conditions - Cross-sectional study from the south Yorkshire cohort dataset. *BMC Public Health.* 2013;13(1):1009. doi:10.1186/1471-2458-13-1009
36. McLaughlin L, Hinyard LJ. The relationship between health-related quality of life and body mass index. *West J Nurs Res.* 2014;36(8):989–1001. doi:10.1177/0193945913520415
37. Ul-Haq Z, Mackay DF, Fenwick E, Pell JP. Meta-analysis of the association between body mass index and health-related quality of life among adults, assessed by the SF-36. *Obesity.* 2013;21(3):E322–E327. doi:10.1002/oby.20107.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>