



**Cite this article:** Peng S, Gerasimenko JV, Tsugorka T, Gryshchenko O, Samarasinghe S, Petersen OH, Gerasimenko OV. 2016 Calcium and adenosine triphosphate control of cellular pathology: asparaginase-induced pancreatitis elicited via protease-activated receptor 2. *Phil. Trans. R. Soc. B* **371**: 20150423. <http://dx.doi.org/10.1098/rstb.2015.0423>

Accepted: 20 April 2016

One contribution of 15 to a Theo Murphy meeting issue 'Evolution brings Ca<sup>2+</sup> and ATP together to control life and death'.

**Subject Areas:**

physiology

**Keywords:**

calcium, adenosine triphosphate, asparaginase, pancreatitis, leukaemia, calcium channels

**Authors for correspondence:**

Ole H. Petersen

e-mail: [petersenoh@cardiff.ac.uk](mailto:petersenoh@cardiff.ac.uk)

Oleg V. Gerasimenko

e-mail: [gerasimenkoov@cardiff.ac.uk](mailto:gerasimenkoov@cardiff.ac.uk)

# Calcium and adenosine triphosphate control of cellular pathology: asparaginase-induced pancreatitis elicited via protease-activated receptor 2

Shuang Peng<sup>1,2</sup>, Julia V. Gerasimenko<sup>1</sup>, Tatiana Tsugorka<sup>1</sup>, Olesiy Gryshchenko<sup>1,3</sup>, Sujith Samarasinghe<sup>4</sup>, Ole H. Petersen<sup>1,5</sup> and Oleg V. Gerasimenko<sup>1</sup>

<sup>1</sup>Cardiff School of Biosciences, Cardiff University, Cardiff CF10 3AX, Wales, UK

<sup>2</sup>Department of Pathophysiology, Medical College, Jinan University, Guangzhou 510632, People's Republic of China

<sup>3</sup>Bogomoletz Institute of Physiology, Kiev 01024, Ukraine

<sup>4</sup>Great Ormond Street Hospital for Children NHS Foundation Trust, Great Ormond Street, London WC1N 3JH, UK

<sup>5</sup>Systems Immunity Research Institute, Cardiff University, Cardiff CF14 4XN, Wales, UK

OHP, 0000-0002-6998-0380

Exocytotic secretion of digestive enzymes from pancreatic acinar cells is elicited by physiological cytosolic Ca<sup>2+</sup> signals, occurring as repetitive short-lasting spikes largely confined to the secretory granule region, that stimulate mitochondrial adenosine triphosphate (ATP) production. By contrast, sustained global cytosolic Ca<sup>2+</sup> elevations decrease ATP levels and cause necrosis, leading to the disease acute pancreatitis (AP). Toxic Ca<sup>2+</sup> signals can be evoked by products of alcohol and fatty acids as well as bile acids. Here, we have investigated the mechanism by which L-asparaginase evokes AP. Asparaginase is an essential element in the successful treatment of acute lymphoblastic leukaemia, the most common type of cancer affecting children, but AP is a side-effect occurring in about 5–10% of cases. Like other pancreatitis-inducing agents, asparaginase evoked intracellular Ca<sup>2+</sup> release followed by Ca<sup>2+</sup> entry and also substantially reduced Ca<sup>2+</sup> extrusion because of decreased intracellular ATP levels. The toxic Ca<sup>2+</sup> signals caused extensive necrosis. The asparaginase-induced pathology depended on protease-activated receptor 2 and its inhibition prevented the toxic Ca<sup>2+</sup> signals and necrosis. We tested the effects of inhibiting the Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> entry by the Ca<sup>2+</sup> channel inhibitor GSK-7975A. This markedly reduced asparaginase-induced Ca<sup>2+</sup> entry and also protected effectively against the development of necrosis.

This article is part of the themed issue 'Evolution brings Ca<sup>2+</sup> and ATP together to control life and death'.

## 1. Background

The importance of Ca<sup>2+</sup> for the control of secretion has been known for a long time [1]. However, it was only through a detailed analysis—using permeabilized adrenal chromaffin cells—that the intracellular requirements for exocytotic secretion were clarified [2]. This work established that secretion occurs when the intracellular free Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>i</sub>) increases from the basic level (around 100 nM) to the low μM range if, and only if, Mg-adenosine triphosphate (ATP; 0.5–5 mM) is present [2]. Under physiological conditions, the rise in [Ca<sup>2+</sup>]<sub>i</sub> that initiates secretion from intact chromaffin cells, as well as nerve endings, is owing to Ca<sup>2+</sup> entering the cells through voltage-gated Ca<sup>2+</sup> channels [2–4]. This is, however, fundamentally different from what happens in the electrically

non-excitabile exocrine gland cells [5,6], where neurotransmitter- or hormone-evoked enzyme and fluid secretion are initiated by release of  $\text{Ca}^{2+}$  from the endoplasmic reticulum (ER) [7], mediated by the intracellular messenger inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) [8]. The primary intracellular  $\text{Ca}^{2+}$  release is followed by secondary  $\text{Ca}^{2+}$  entry from the extracellular solution, and this  $\text{Ca}^{2+}$  influx does not occur through voltage-gated  $\text{Ca}^{2+}$  channels, but via  $\text{Ca}^{2+}$  release-activated  $\text{Ca}^{2+}$  (CRAC) channels [9].

The physiologically relevant  $\text{Ca}^{2+}$  signals that control secretion in the pancreatic acinar cells occur as repetitive local  $[\text{Ca}^{2+}]_i$  spikes in the apical secretory granule region [10,11], owing to primary  $\text{Ca}^{2+}$  release from thin strands of ER penetrating into this region [12,13], which is dominated by zymogen granules (ZG). These ER strands are fully functionally connected (ER tunnels) to the principal ER store dominating the basal region [12,13]. The  $\text{IP}_3$  receptors ( $\text{IP}_3\text{Rs}$ ) are concentrated in the apical area [11,14,15] and the functional ER tunnels effectively allow  $\text{Ca}^{2+}$  to be mobilized from the main basal store into the apical secretory granule area [12]. Plasma membrane  $\text{Ca}^{2+}$  pumps, concentrated in the apical (secretory) membrane [16], extrude  $\text{Ca}^{2+}$  from the cell during each  $\text{Ca}^{2+}$  spike, and compensatory  $\text{Ca}^{2+}$  entry and uptake into the ER are therefore necessary. This occurs through conventional [17] CRAC channels [9] localized in the baso-lateral membrane [12,18], and  $\text{Ca}^{2+}$  entering through these channels is immediately taken up into the ER by powerful  $\text{Ca}^{2+}$  pumps in the ER membrane [12]. The physiological apical  $\text{Ca}^{2+}$  release via  $\text{IP}_3\text{Rs}$  in that region results in apically confined  $[\text{Ca}^{2+}]_i$  spikes owing to the mitochondrial firewall separating the ZG-containing part of the cell from the ER-dominated baso-lateral part [19]. During each apical  $\text{Ca}^{2+}$  spike,  $\text{Ca}^{2+}$  is taken up into the peri-granular mitochondria and then slowly released [13,20]. This mitochondrial  $\text{Ca}^{2+}$  uptake, activating several  $\text{Ca}^{2+}$ -sensitive dehydrogenases in the Krebs cycle [21], results in an increased cytosolic ATP level, despite the increased ATP utilization [22]. This is helpful for powering the exocytotic secretory process.

Whereas the  $\text{Ca}^{2+}$  release evoked by the neurotransmitter acetylcholine (ACh) is primarily mediated by  $\text{IP}_3$  acting on  $\text{IP}_3\text{Rs}$  [23,24], the  $\text{Ca}^{2+}$  release elicited by physiological concentrations of the hormone cholecystokinin (CCK) is primarily mediated by the  $\text{Ca}^{2+}$ -releasing messenger nicotinic acid adenine dinucleotide phosphate (NAADP) [25–27]. The action of NAADP depends on functional ryanodine receptors (RyRs) and two-pore channels (TPCs) and also involves acid  $\text{Ca}^{2+}$  stores [27]. Physiological  $\text{Ca}^{2+}$  spiking, irrespective of whether it is evoked by ACh or CCK, depends on both functional  $\text{IP}_3\text{Rs}$  and RyRs [13,25].

Although the acinar cells in the pancreas are quantitatively dominant and perform the most crucial physiological role in the exocrine pancreas, by synthesizing and secreting the digestive (pro-) enzymes that are essential for the digestion of food in the intestine, there are other cells that need to be considered. The acinar cells secrete fluid together with the enzymes [5], but most of the fluid carrying the enzymes into the gut is produced by the duct cells, principally regulated by secretin-evoked cyclic AMP formation rather than by  $\text{Ca}^{2+}$  signals [28,29]. The acinar–ductal system is functionally integrated and regulation of one cell type has influence on the other [30]. The physiological role of the more recently discovered peri-acinar stellate cells has not yet been fully clarified, but they

generate substantial  $\text{Ca}^{2+}$  signals in response to concentrations of bradykinin that have been measured in plasma during exercise and pancreatitis [31,32]. They do not respond to the principal controllers of acinar cell function, namely ACh or CCK [31,32].

In the normal digestion process, the inactive acinar pancreatic pro-enzymes, which are secreted by exocytosis into the acinar lumen, are carried into the gut by acinar and ductal fluid secretion and then activated in the gut [33]. In acute pancreatitis (AP), a potentially fatal human disease, the inactive pancreatic pro-enzymes become active enzymes inside the acinar cells, digesting the pancreas and its surroundings. The main causes of AP are biliary disease (gallstone complications) and alcohol abuse [33]. More than 20 years ago, it was proposed that AP is essentially a disease brought about by excessive cytosolic  $\text{Ca}^{2+}$  signals [34] and since then a substantial amount of evidence in favour of this hypothesis has accumulated [35–38]. Pathological stimuli—for example, combinations of alcohol and fatty acids or bile acids—can evoke excessive release of  $\text{Ca}^{2+}$  from internal stores followed by excessive  $\text{Ca}^{2+}$  entry through store-operated CRAC channels, resulting in sustained global elevations of  $[\text{Ca}^{2+}]_i$  [38]. This causes intracellular protease activation [38] and excessive mitochondrial  $\text{Ca}^{2+}$  uptake, leading to opening of the mitochondrial permeability transition pore (MPTP) [20,39]. The MPTP opening causes depolarization of the inner mitochondrial membrane, resulting in failure of the normal mitochondrial ATP production [20,39]. The lack of ATP prevents the acinar cells from disposing of the excess  $\text{Ca}^{2+}$  in the cytosol and, in combination with the abnormal intracellular protease activity, this leads to necrosis. It is the acinar necrosis that generates the damaging inflammatory response [39–42]. Since the primary pathological event in AP is the excessive and sustained  $[\text{Ca}^{2+}]_i$  elevation and as this depends on excessive  $\text{Ca}^{2+}$  entry through CRAC channels, it would be logical, as a therapy, to target these  $\text{Ca}^{2+}$  entry channels. Gerasimenko *et al.* [9,38] demonstrated, in experiments on isolated mouse acinar cell clusters, that the intracellular protease activation and necrosis evoked by fatty acid ethyl esters—non-oxidative combinations of ethanol and fatty acids, which are important mediators of alcohol-related pancreatitis [33]—could be effectively inhibited by specific pharmacological blockade of CRAC channels. These results have recently been confirmed *in vivo*, in three different mouse models of AP [43], giving hope that CRAC channel blockade may become the first rational and effective therapy against AP [44].

As both alcohol-related and bile-induced AP are owing to toxic  $\text{Ca}^{2+}$  signal generation, it seems possible that all types of AP are  $\text{Ca}^{2+}$  toxicity diseases. We were therefore interested in exploring the mechanism underlying pancreatitis caused by a side-effect of L-asparaginase, which is used for the treatment of acute lymphoblastic leukaemia (ALL). L-asparaginase is a well-known anticancer agent effective against lymphoid malignancies. Since 1971, it has been an essential element in the successful treatment of ALL, the most common type of cancer affecting children [45,46]. However, asparaginase treatment can result in AP (asparaginase-associated pancreatitis (AAP)). This occurs in about 5–10% of cases and AAP is the most frequent cause of discontinuing the asparaginase treatment [47–50]. The mechanism underlying the development of AAP has so far not been explored [48].

The aim of the present study was to define the mechanism underlying AAP and then to identify potential steps for

therapeutic intervention. As outlined above, the history of our path to the current understanding of AP shows that studies on isolated cells or cell clusters have been enormously helpful and our approach has therefore been to study, for the first time, the effects of asparaginase on isolated mouse acinar cells or cell clusters. As *in vitro* results concerning the effects of, for example, fatty acid ethyl esters or bile acids [9,38,43] have turned out to be excellent predictors of the outcome of studies of real AP *in vivo* [39,43,51], acute studies on isolated cells are a natural starting point for investigations of the mechanism of AAP.

Our results show that asparaginase acts on protease-activated receptor 2 (PAR2) to evoke sustained elevations of  $[Ca^{2+}]_i$  owing to release of  $Ca^{2+}$  from internal stores, followed by  $Ca^{2+}$  entry from the extracellular solution. The sustained  $[Ca^{2+}]_i$  elevation reduces ATP formation. These effects can be markedly reduced by specific pharmacological blockade of CRAC channels, which also markedly reduces the extent of necrosis.

## 2. Results

### (a) Asparaginase increases $[Ca^{2+}]_i$ in pancreatic acinar cells

Pancreatitis-inducing agents, combinations of ethanol and fatty acids or bile acids, are able to elevate  $[Ca^{2+}]_i$  in isolated acinar cell clusters causing intracellular  $Ca^{2+}$  overload. This is owing to  $Ca^{2+}$  release from internal stores triggering excessive store-operated  $Ca^{2+}$  entry [9,38]. To identify the mechanism of action of asparaginase, we therefore started out by testing the effects of asparaginase on  $[Ca^{2+}]_i$  over a wide concentration range. One of the challenges inherent in this approach is that the time course needed for a study on normal freshly isolated cells is quite different from that of the development of AAP in the clinical situation. AAP typically develops several days after several injections (over many weeks) of asparaginase [49], whereas studies on isolated cells in the laboratory require observations of the effects of asparaginase within hours. In the present study, we have worked with the lowest concentration of asparaginase that reliably evoked cellular changes that are similar to those previously found to be associated with AP initiated by alcohol metabolites or bile acids.

We found that only in some cells did a low concentration of asparaginase (20 IU ml<sup>-1</sup>) induce  $[Ca^{2+}]_i$  oscillations (9 out of 42 cells). Figure 1a shows a representative positive trace with repetitive  $Ca^{2+}$  spikes induced by 20 IU ml<sup>-1</sup>, whereas figure 1b represents the more typical negative response (33 out of 42 cells), in which the same concentration of asparaginase failed to cause any change in  $[Ca^{2+}]_i$ . A higher concentration of asparaginase (200 IU ml<sup>-1</sup>) elicited repetitive  $[Ca^{2+}]_i$  oscillations, mostly on top of a sustained elevated  $[Ca^{2+}]_i$  (43 out of 55 cells; figure 1c). The development of a sustained  $[Ca^{2+}]_i$  elevation has previously been shown to be a distinctive characteristic of  $[Ca^{2+}]_i$  changes induced by pathological concentrations of alcohol metabolites or bile acids [9,38], and we found that an elevated  $[Ca^{2+}]_i$  plateau, although often small, was seen in the vast majority of cells (52 out of 55) stimulated by asparaginase (200 IU ml<sup>-1</sup>). In some cases (12 out of 55), there were no, or very few, spikes superimposed on the elevated  $[Ca^{2+}]_i$  plateau (figure 1d).

### (b) The asparaginase-elicited sustained increase in $[Ca^{2+}]_i$ depends on the presence of external $Ca^{2+}$

Removal of external  $Ca^{2+}$  always terminated the elevated  $[Ca^{2+}]_i$  plateau (figure 1c,d), but did not significantly ( $p > 0.27$ ) change the amplitudes of the asparaginase-induced  $Ca^{2+}$  spikes ( $n = 28$ ) within the time frame of our experiments (figure 1e).  $Ca^{2+}$  entry clearly plays an important role in the formation of the asparaginase-induced elevated  $[Ca^{2+}]_i$  plateau and the role of  $Ca^{2+}$  entry was further demonstrated by increasing the external  $Ca^{2+}$  concentration to 2 mM during asparaginase stimulation, which caused a marked and sustained increase in  $[Ca^{2+}]_i$  ( $n = 18$ ; figure 1f). Subsequent removal of external  $Ca^{2+}$  and addition of the  $Ca^{2+}$  chelator EGTA (200  $\mu$ M) abolished the elevated  $[Ca^{2+}]_i$  plateau, whereas the  $Ca^{2+}$  oscillations continued for some time (figure 1f).

It has previously been shown that the excessive  $Ca^{2+}$  entry into pancreatic acinar cells induced by alcohol metabolites or bile acids, as well as their pathological effects, can be markedly inhibited by the CRAC channel blockers GSK-7975A and CM-128 [9,43,52]. We have therefore tested the effect of CRAC blockade (GSK-7975A, 10  $\mu$ M) on the asparaginase-induced sustained  $[Ca^{2+}]_i$  elevation and found that it was abolished in the presence of the inhibitor, although repetitive  $Ca^{2+}$  spiking was still observed within the time frame of our experiments (figure 1g,  $n = 32$ ). Figure 1h summarizes the degree of inhibition, caused by removal of external  $Ca^{2+}$  or by GSK-7975A, of the integrated  $Ca^{2+}$  signal evoked by asparaginase.


### (c) Asparaginase-elicited $Ca^{2+}$ release involves inositol 1,4,5-trisphosphate receptors and ryanodine receptors as well as nicotinic acid adenine dinucleotide phosphate signalling

To investigate the involvement of intracellular  $Ca^{2+}$  release channels in asparaginase-induced  $[Ca^{2+}]_i$  elevations, we have used caffeine, a substance known to reliably inhibit IP<sub>3</sub>-mediated intracellular  $Ca^{2+}$  release in pancreatic acinar cells [23]. Caffeine (20 mM) substantially reduced the asparaginase-induced  $[Ca^{2+}]_i$  elevations in a  $Ca^{2+}$ -free solution ( $n = 8$ ; figure 2a; compare with figure 1e as the appropriate control). The phospholipase C (PLC) inhibitor U73122 (10  $\mu$ M) also significantly blocked the asparaginase-induced  $Ca^{2+}$  release as well as the response to 1  $\mu$ M ACh ( $n = 11$ ; figure 2b). Ryanodine (100  $\mu$ M), inhibiting RyRs, also substantially reduced the asparaginase-induced  $[Ca^{2+}]_i$  elevations ( $n = 13$ ; figure 2c).

It has previously been shown that NAADP signalling in pancreatic acinar cells can be inactivated by the cell-permeable NAADP analogue, and selective antagonist, Ned-19 [27]. After pre-treatment of cells with 100  $\mu$ M Ned-19, the asparaginase (200 IU ml<sup>-1</sup>)-induced  $[Ca^{2+}]_i$  elevations were profoundly inhibited (virtually abolished; figure 2d,  $n = 8$ ), whereas ACh (1  $\mu$ M), eliciting  $Ca^{2+}$  release via IP<sub>3</sub>Rs independently of NAADP [25], was still able to evoke a typical  $[Ca^{2+}]_i$  rise ( $n = 8$ ; figure 2d).

### (d) Protease-activated receptor 2 is involved

PAR2 is widely expressed in human and animal tissues, including the pancreas, and has previously been implicated




**Figure 1.** Asparaginase (ASNase) induces cytosolic  $\text{Ca}^{2+}$  signals in pancreatic acinar cells. A low concentration of asparaginase ( $20 \text{ IU ml}^{-1}$ ) induces repetitive  $\text{Ca}^{2+}$  spikes in a minority (9 out of 42) of experiments (a), but elicits no response in the majority (33 out of 42) of cases (b). A higher concentration of asparaginase ( $200 \text{ IU ml}^{-1}$ ) elicits an elevated  $[\text{Ca}^{2+}]_i$  plateau in practically all cases (52 out of 55), often with repetitive  $\text{Ca}^{2+}$  transients on top of the plateau (43 out of 55) as shown in the representative trace (c). An elevated  $[\text{Ca}^{2+}]_i$  plateau is the main type of response to asparaginase ( $200 \text{ IU ml}^{-1}$ ; 52 out of 55), but in some cases (12 out of 55) with no or very few spikes (d). The sustained  $[\text{Ca}^{2+}]_i$  elevation depends on the continued presence of  $\text{Ca}^{2+}$  in the external solution (c,d). In the absence of external  $\text{Ca}^{2+}$ , there is no elevated  $[\text{Ca}^{2+}]_i$  plateau (e,  $n = 28$ ). When the external  $\text{Ca}^{2+}$  concentration is increased from 1 to 2 mM, there is a marked transient  $[\text{Ca}^{2+}]_i$  rise and a continuing elevated  $[\text{Ca}^{2+}]_i$  plateau (f). When external  $\text{Ca}^{2+}$  is subsequently removed and a  $\text{Ca}^{2+}$  chelator (EGTA) added, the plateau gradually disappears (f;  $n = 18$ ). In the presence of the CRAC channel blocker GSK-7975A ( $10 \mu\text{M}$ ), asparaginase ( $200 \text{ IU ml}^{-1}$ ) evokes repetitive  $\text{Ca}^{2+}$  spikes without an elevated  $[\text{Ca}^{2+}]_i$  plateau (g,  $n = 32$ ). Comparison of the integrated  $\text{Ca}^{2+}$  signals ('area under the curve' from start of the  $\text{Ca}^{2+}$  signal until 1800 s later), in the presence of 1 mM external  $\text{Ca}^{2+}$ , in the absence of external  $\text{Ca}^{2+}$  and in the presence of 1 mM external  $\text{Ca}^{2+}$  with the addition of  $10 \mu\text{M}$  GSK-7975A, shows that the absence of external  $\text{Ca}^{2+}$  ( $n = 28$ ) or the addition of the CRAC blocker GSK-7975A ( $n = 32$ ) significantly ( $p < 0.0001$  in both cases) reduced the responses to asparaginase ( $200 \text{ IU ml}^{-1}$ ; h). Bars represent mean  $\pm$  s.e.m. (Online version in colour.)

in the pathology of AP [53,54]. The activation of PAR family members is coupled to multiple heteromeric G proteins that lead to PLC activation and production of  $\text{IP}_3$  and diacylglycerol [55]. Therefore, we tested the possibility that asparaginase could activate PAR2 by pre-treating the cells

with the PAR2 blocker FSLRY-NH<sub>2</sub> ( $10 \mu\text{M}$ ) before the addition of asparaginase ( $200 \text{ IU ml}^{-1}$ ). The PAR2 blocker reduced significantly (virtually abolished) the asparaginase-induced  $[\text{Ca}^{2+}]_i$  oscillations as well as the sustained  $[\text{Ca}^{2+}]_i$  elevation (figure 2e;  $n = 32$ ).





**Figure 2.** The primary intracellular  $\text{Ca}^{2+}$  release induced by asparaginase depends on  $\text{IP}_3$  and NAADP signalling mechanisms. The  $\text{IP}_3$ R blocker caffeine (20 mM) inhibited very markedly the asparaginase-induced  $[\text{Ca}^{2+}]_i$  elevations in the absence of external  $\text{Ca}^{2+}$  (*a*, averaged trace,  $n = 8$ ). The PLC inhibitor U73122 (10  $\mu\text{M}$ ) blocked the asparaginase-induced  $[\text{Ca}^{2+}]_i$  elevation in the absence of external  $\text{Ca}^{2+}$  (*b*, averaged trace,  $n = 11$ ). ACh (1  $\mu\text{M}$ ) was applied at the end of each experiment and, as expected, did not elicit any change in  $[\text{Ca}^{2+}]_i$  (*b*) as its effect depends on  $\text{IP}_3$  formation and  $\text{IP}_3$  receptors [23,24]. Ryanodine (100  $\mu\text{M}$ ) markedly inhibited the asparaginase-induced  $\text{Ca}^{2+}$  signals in the absence of external  $\text{Ca}^{2+}$  (*c*, averaged trace,  $n = 13$ ). Ned-19 (100  $\mu\text{M}$ ) prevented the asparaginase-induced  $[\text{Ca}^{2+}]_i$  elevation in the absence of external  $\text{Ca}^{2+}$  (*d*, averaged trace,  $n = 8$ ). ACh (1  $\mu\text{M}$ ) was applied at the end of each experiment and, as expected, caused a major  $[\text{Ca}^{2+}]_i$  rise as its effect does not depend on NAADP receptors [25,27]. The asparaginase-induced  $\text{Ca}^{2+}$  signals were virtually eliminated by the PAR2 inhibitor FSLRY-NH<sub>2</sub> (10  $\mu\text{M}$ ; *e*, averaged trace,  $n = 32$ ).


### (e) $\text{Ca}^{2+}$ extrusion mechanisms are affected by asparaginase

To study  $\text{Ca}^{2+}$  movements in more detail, we have applied a specific protocol routinely used to assess  $\text{Ca}^{2+}$  entry and extrusion [9,38]. In these experiments,  $\text{Ca}^{2+}$  stores were emptied using the ER  $\text{Ca}^{2+}$  pump inhibitor cyclopiazonic acid (CPA) in a nominally  $\text{Ca}^{2+}$ -free solution. Thereafter, 2 mM of  $\text{Ca}^{2+}$  was added to the external solution for a short period and then removed (figure 3*a*). In the presence of asparaginase, the  $\text{Ca}^{2+}$  entry following external  $\text{Ca}^{2+}$  admission was significantly increased (figure 3*a*), assessed by both amplitude of the  $[\text{Ca}^{2+}]_i$  change (figure 3*b*) and the rate of  $[\text{Ca}^{2+}]_i$  increase (half-time of the increase, figure 3*c*). However, the quantitatively most important effect of asparaginase was to slow down the rate of  $\text{Ca}^{2+}$  extrusion after removal of external  $\text{Ca}^{2+}$  (figure 3*a*). The half-time of  $[\text{Ca}^{2+}]_i$  recovery was more than three times longer than in the control cells (figure 3*d*).

### (f) Asparaginase depletes intracellular adenosine triphosphate

$\text{Ca}^{2+}$  extrusion is an energy-demanding process and has previously been found to be abnormal in pancreatic acinar cell pathologies owing to disruption of mitochondrial metabolism and, therefore, reduction of ATP levels [42]. We have conducted indirect assessments of intracellular changes in ATP concentration, using Magnesium Green (MgGreen) fluorescence measurements. As most of the intracellular ATP will be in the form of Mg-ATP, a reduction of the ATP concentration will increase the fluorescence intensity of MgGreen owing to the inevitable increase of  $[\text{Mg}^{2+}]_i$  [56–58]. By this measure, asparaginase induced a substantial reduction of intracellular ATP levels (figure 4*a*) superseded only by the full ATP depletion caused by a mixture of the protonophore CCCP, oligomycin and iodoacetate [42].

Because the presence of external  $\text{Ca}^{2+}$  was important for the cytoplasmic and mitochondrial effects induced by



**Figure 3.** Asparaginase accelerates  $\text{Ca}^{2+}$  entry and substantially slows down  $\text{Ca}^{2+}$  extrusion. (a) In the absence of external  $\text{Ca}^{2+}$ , CPA, a specific inhibitor of the  $\text{Ca}^{2+}$  pump in the ER membrane, causes a modest and largely transient  $[\text{Ca}^{2+}]_i$  rise. When subsequently 2 mM  $\text{Ca}^{2+}$  is added to the external solution, there is a marked rise in  $[\text{Ca}^{2+}]_i$ , which then declines after removal of external  $\text{Ca}^{2+}$ . In the presence of asparaginase (red averaged trace,  $n = 32$ ), the amplitude (b) and the rate of rise of  $[\text{Ca}^{2+}]_i$  (c) are somewhat increased ( $p < 0.048$  and  $p < 0.0001$ , respectively) when compared with control (a, blue averaged trace,  $n = 34$ ). The extrusion of  $\text{Ca}^{2+}$  by the plasma membrane  $\text{Ca}^{2+}$  pumps, observed as the decline in  $[\text{Ca}^{2+}]_i$  following removal of external  $\text{Ca}^{2+}$ , is very markedly and significantly ( $p < 0.0001$ ) reduced in the presence of asparaginase (a,d). (Online version in colour.)

asparaginase, we decided to check if inhibition of  $\text{Ca}^{2+}$  entry channels [9,43] could affect the ATP loss evoked by asparaginase. Blocking  $\text{Ca}^{2+}$  entry by GSK-7975A (10  $\mu\text{M}$ ) substantially reduced the ATP loss evoked by asparaginase (figure 4b).

### (g) The effect of pyruvate


It has been shown previously that ethyl pyruvate (aliphatic ester derived from pyruvic acid [59]) attenuates the severe AP induced by sodium taurocholate in rats [59]. When we included 1 mM pyruvate in the external solution, we found that the asparaginase-induced ATP loss was substantially reduced compared with control experiments (figure 4c,d). The protective effect was significant and similar to what was achieved by GSK-7975A (figure 4b,d), although it did not give complete protection.

### (h) Necrosis

We have previously shown that the cytosolic  $\text{Ca}^{2+}$  overload and ATP deprivation induced by fatty acid ethyl esters and bile acids lead to necrosis [38], and we have therefore now tested whether asparaginase can also induce necrosis, the hallmark of AP [33,38]. The extent of necrosis induced by asparaginase (200 IU ml<sup>-1</sup>) treatment ( $17.4 \pm 0.4\%$  of the cells; figure 5a) was comparable to, but somewhat smaller than, the level of necrosis induced by pamitoic acid ethyl ester (POAEE;  $29 \pm 3.1\%$ ) or the bile acid taurocholic acid sulphate (TLC-S;  $27.6 \pm 1.9\%$ ; figure 5a), whereas a lower concentration of asparaginase (20 IU ml<sup>-1</sup>) did not increase the level of necrosis above that seen in control experiments (figure 5b). The CRAC channel

inhibitor GSK-7975A (10  $\mu\text{M}$ ) [9,43] reduced the asparaginase-induced necrosis to the control level ( $4.5 \pm 0.7\%$ ; figure 5b). Pyrazole compounds have generally been thought to inhibit other types of cation channels, namely the relatively non-selective TRP (transient receptor potential) channels, which have significant  $\text{Ca}^{2+}$  permeability [60,61], but pyrazole6 (Pyr6) has been shown to have more of an effect on the very  $\text{Ca}^{2+}$ -selective CRAC channels [61], which are the ones specifically inhibited by GSK-7975A and CM-128 [9,43]. In our experiments, Pyr6 partially inhibited asparaginase-induced necrosis to  $8.4 \pm 0.9\%$ . Both caffeine and Ned-19 inhibited asparaginase-induced necrosis to control levels ( $4.2 \pm 0.5$  and  $6.6 \pm 1.2\%$ , respectively). A PAR2 inhibitor (FSLLRY-NH<sub>2</sub>) significantly blocked asparaginase-induced necrosis (figure 5b). We also tested the effect of pyruvate on asparaginase-induced necrosis. As seen in figure 5b, this gave significant protection against necrosis. Figure 5c shows representative images of some of the cells under the treatment protocols, together with the results of staining for propidium iodide (PI). It is seen that asparaginase (200 IU ml<sup>-1</sup>) elicited strong intracellular PI staining and that GSK-7975A provided protection against this.

Asparaginase kills lymphoblastic cells by depriving them of asparagine, which they—unlike normal cells—cannot produce themselves [62]. The effects of asparaginase on normal pancreatic acinar cells described in this study are therefore unlikely to be owing to asparagine deprivation. We tested whether there was any difference between the ability of asparaginase to induce necrosis in the absence or presence of asparagine. As seen in figure 5d, there was no difference in the necrosis levels evoked by asparaginase in the presence or absence of asparagine.




**Figure 4.** Asparaginase reduces intracellular ATP levels as assessed by increases in  $[Mg^{2+}]_i$ . In these experiments, changes in  $[Mg^{2+}]_i$  were assessed by monitoring changes in MgGreen fluorescence. Most of the  $Mg^{2+}$  inside cells is bound to ATP, so when ATP declines—for example owing to interference with mitochondrial function— $[Mg^{2+}]_i$  will increase. A rise in  $[Mg^{2+}]_i$  can therefore be taken to represent a reduction in the intracellular ATP level [58]. Asparaginase (200 IU ml<sup>-1</sup>) markedly increased  $[Mg^{2+}]_i$  and a further rise occurred after poisoning mitochondrial function with a cocktail of CCCP (5 nM), oligomycin (10  $\mu$ M) and sodium iodoacetate (2 mM; *a*, averaged trace,  $n = 39$ ). The asparaginase effect was markedly reduced by GSK-7975A (*b*: orange averaged trace,  $n = 21$ ; blue averaged trace,  $n = 39$ ) and was also markedly reduced by supplementing the external medium with pyruvate (1 mM; *c*,  $n = 8$ ). Comparisons of the integrated responses ('areas under the curve' from the start of the responses until 1800 s later) show that GSK-7975A and pyruvate significantly reduced the asparaginase-induced ATP depletion ( $p < 0.0001$  for both treatments).

### 3. Discussion

The results presented here, on the asparaginase-elicited injury to pancreatic acinar cells, provide fresh evidence for the hypothesis that all types of AP are owing to toxic  $Ca^{2+}$  signal generation and explain how asparaginase could cause AP (figure 6). Asparaginase, like fatty acid ethyl esters and bile acids, can evoke sustained  $[Ca^{2+}]_i$  elevation owing to release of  $Ca^{2+}$  from intracellular stores followed by store-operated  $Ca^{2+}$  entry through CRAC channels. Qualitatively, the effects

of asparaginase fit well with those induced by fatty acid ethyl esters and bile acids, which we have described previously [33,38]. However, the sustained  $[Ca^{2+}]_i$  elevations evoked by asparaginase are somewhat smaller than those evoked by bile acids or fatty acid ethyl esters. Nevertheless asparaginase evokes significant reductions in the intracellular ATP levels and extensive necrosis. Further studies on mitochondrial  $Ca^{2+}$  handling during the action of asparaginase are warranted because the regulation of mitochondrial  $Ca^{2+}$  uptake under different conditions may be a critical issue [63,64].




**Figure 5.** Asparaginase induces necrosis in pancreatic acinar cells. (a) The level of necrosis induced by 200 IU ml<sup>-1</sup> of asparaginase ( $17.4 \pm 0.4\%$ , eight series with  $n > 250$ ) is comparable with, but somewhat lower than, that caused by the alcohol metabolite POAEE (100  $\mu$ M;  $29 \pm 3.1\%$ , three series of experiments with  $n > 100$ ) or the bile acid TLC-S (200  $\mu$ M;  $27.6 \pm 1.9\%$ , three series of experiments with  $n > 120$ ), whereas a lower concentration of asparaginase (20 IU ml<sup>-1</sup>) did not induce any significant change in the level of necrosis (b) when compared with control ( $4.5 \pm 0.4$  and  $4.5 \pm 0.7\%$ , respectively,  $p > 0.8$ , each test had eight series of experiments with  $n > 130$  of tested cells in each group). The CRAC channel inhibitor GSK-7975A (10  $\mu$ M) essentially abolished asparaginase-induced necrosis to a level not significantly different from control ( $p > 0.24$ ), while significantly ( $p < 0.0001$ , eight series,  $n > 70$ ) lower than that caused by 200 IU ml<sup>-1</sup> of asparaginase alone (b). The PAR2 inhibitor FSLLRN-NH<sub>2</sub> (10  $\mu$ M) significantly blocked the asparaginase-induced necrosis ( $p < 0.0001$ ; four series of experiments with  $n > 150$  cells in each sample). Pyruvate (1 mM) reduced the asparaginase-induced necrosis to near the control level (b;  $p < 0.0001$ , three series of experiments with the number of tested cells in each group more than 130). (c) Representative images of cells from some of the experiments shown in (b). Transmitted light images (upper row) and PI-stained fluorescence images (lower row). (d) The presence or absence of asparagine (50  $\mu$ M) made no difference to the level of asparaginase-induced necrosis ( $p > 0.75$ , three series of experiments with more than 70 cells in each sample).

The findings presented in this study provide the first mechanistic insights into the process by which asparaginase treatment of ALL may cause AAP (figure 6). These insights also provide the first pointers to rational therapies (figure 6) that may prevent the currently necessary cessation of

asparaginase treatment of ALL in cases of severe pancreatitis. The most accessible therapeutic target is the Ca<sup>2+</sup> entry route, namely the previously characterized CRAC channels [9,43,44,65]. We have now shown that the asparaginase-induced Ca<sup>2+</sup> elevations depend on CRAC-mediated Ca<sup>2+</sup>





**Figure 6.** Schematic diagram illustrating the likely effects of asparaginase on pancreatic acinar cells, based partly on the new data shown in this study and partly on analogy with processes identified in previous studies on the mechanisms by which fatty acid ethyl esters and bile acids cause AP [38]. Potential sites for therapeutic intervention are also indicated.

entry and, therefore, are strongly inhibited by the CRAC channel inhibitor GSK-7975A and also that, consequently, asparaginase-induced necrosis is dramatically reduced to near control levels by GSK-7975A (figure 5*b*). On the basis of the previously documented protective effects of Ca<sup>2+</sup> entry channel inhibition against alcohol-related pancreatic pathology in isolated cell clusters [9,38], the recent confirmation of its effectiveness *in vivo* in three different mouse pancreatitis models [43] and the very recently demonstrated inhibition of prolonged Ca<sup>2+</sup> signal generation in pancreatic stellate cells [31,32], our new data indicate that this therapeutic approach is also likely to be successful against asparaginase-induced pancreatitis. Clearly, the next—but challenging—step would be to test the effectiveness of CRAC channel blockade against AAP in an *in vivo* mouse model.

The mechanism by which asparaginase induces pancreatitis is fundamentally different from its therapeutic action on the lymphoblastic cells in ALL. The asparaginase effect on cancer cells relies on depletion of asparagine, which the malignant cells cannot produce themselves, in contrast to normal cells [62], whereas the side-effect of asparaginase, inducing pancreatitis, is owing to activation of a signal transduction mechanism involving PAR2, the intracellular messengers IP<sub>3</sub> and NAADP, and the intracellular receptors IP<sub>3</sub>Rs, RYRs and possibly TPCs (figure 6). The asparaginase effect on the pancreas is therefore independent of the presence or absence of asparagine (figure 5*d*). This means that there are several potential intervention points available for treating the side-effect of asparaginase (figure 6), without interfering with its primary effect on the cancer cells. The primary site of action of

asparaginase on pancreatic acinar cells seems to be PAR2. This receptor has previously been implicated in AP, although its exact role is still debated [53,54]. Blocking PAR2 in our experiments inhibited both the pathological [Ca<sup>2+</sup>]<sub>i</sub> elevations (figure 2*e*) and the asparaginase-induced necrosis (figure 5*b*), suggesting that PAR2 inhibitors in addition to, or in combination with, CRAC channel inhibitors could be a useful tool to supplement asparaginase ALL treatment in AAP cases.

Both Ca<sup>2+</sup> entry and extrusion are significantly affected by asparaginase, leading to formation of the pathological elevated [Ca<sup>2+</sup>]<sub>i</sub> plateau, and this sustained elevation of [Ca<sup>2+</sup>]<sub>i</sub> would appear to be responsible for the necrosis. The demonstrated reduction of the intensity of Ca<sup>2+</sup> extrusion (figure 3*a*) is clearly a key element, and the simplest explanation for this is the reduction in the intracellular ATP level (figure 4*a*) limiting the energy supply to the Ca<sup>2+</sup> ATPase in the plasma membrane. When energy supply is partially restored by the addition of pyruvate (figure 4*d*), it provides a similar degree of protection against pancreatic necrosis to PAR2 inhibition or GSK-7975A (figure 5*b*). Clearly, both Ca<sup>2+</sup> and ATP play key roles in pancreatic pathology, as indeed they do in physiological regulation of secretion, and therapeutic strategies must take both into account.

## 4. Material and methods

### (a) Materials

All fluorescent dyes were purchased from ThermoFisher Scientific (Invitrogen, UK), and CPA was from Merck Millipore (Calbiochem,

UK). Collagenase was obtained from Worthington (USA), asparaginase was from Abcam (UK), the PAR2 inhibitor FSLRY-NH2 from TOCRIS (UK) and GSK-7975A from GlaxoSmithKline (UK). All other chemicals were purchased from Sigma. C57BL/6J mice were from Charles River UK Ltd.

### (b) Isolation of pancreatic acinar cells

Pancreatic acinar cells were isolated as previously described [9]. Briefly, animals were sacrificed according to the Animal Scientific Procedures Act, 1986 and approved by the Ethical Review Committee of Cardiff University. After dissection, the pancreas was digested using collagenase-containing solution (200 U ml<sup>-1</sup>, Worthington, UK) and incubated in a 37°C water bath for 14–15 min. The extracellular solution contained: 140 mM NaCl, 4.7 mM KCl, 10 mM Hepes, 1 mM MgCl<sub>2</sub>, 10 mM glucose, pH 7.2, and CaCl<sub>2</sub> (0–2 mM as described in the text).

### (c) Fluorescence measurements

For measurements of [Ca<sup>2+</sup>]<sub>i</sub>, isolated pancreatic acinar cells were loaded with Fluo-4-AM (5 μM; excitation 488 nm) or Fura-2-AM (2.5 μM; excitation 365 and 385 nm) following the manufacturer's instruction. The cells were then washed and re-suspended in extracellular solution containing 1 mM CaCl<sub>2</sub>.

Measurement of intracellular ATP was performed with MgGreen, which senses changes in [Mg<sup>2+</sup>]<sub>i</sub> at concentrations around the resting [Mg<sup>2+</sup>]<sub>i</sub> (approx. 0.9 mM). This was used as an indirect approach to detect cytosolic ATP depletion [58]. Pancreatic acinar cells were incubated with 4 μM MgGreen for 30 min at room temperature (excitation 488 nm). ATP depletion mixture (4 μM CCCP, 10 μM oligomycin and 2 mM iodoacetate) was applied for a final 10 min to induce maximum ATP depletion [42].

Necrotic cell death was assessed with PI uptake as previously described [9].

All experiments were performed at room temperature using freshly isolated cells attached to coverslips of the perfusion chamber. Fluorescence was imaged over time using an Leica TCS SPE confocal microscope.

### (d) Statistical analysis

Data are presented as mean ± SEM. Statistical significance and *p*-values were calculated using *t*-test or ANOVA, with *p* < 0.05 considered significant.

**Ethics.** All animal-related research was run according to the Animal Scientific Procedures Act, 1986 and approved by the Ethical Review Committee of Cardiff University.

**Data accessibility.** Data sets for this article have been obtained and processed using Leica LAS AF software (<http://www.leica-microsystems.com/products/microscope-software/>) needed to open the original files. Supporting files will be made available in the Research section via: <https://www.cardiff.ac.uk/people/view/108900-gerasimenko-oleg>.

**Authors' contributions.** S.P., J.V.G., T.T., O.G., S.S., O.H.P. and O.V.G. designed the research; S.P., J.V.G., T.T. and O.G. performed the research; S.P., J.V.G., T.T., O.G. and O.V.G. analysed data; J.V.G., O.H.P. and O.V.G. wrote the study.

**Competing interests.** We have no competing interests.

**Funding.** This work was funded by a Medical Research Council Programme grant no. (MR/J002771/1) and a Children with Cancer UK grant no. (2014/167). Work was also supported by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and University College London. S.P. is supported by China Scholarship Council (201406780021). O.H.P. is a Medical Research Council Professor (G19/22/2).

## References

- Harvey AM, MacIntosh FC. 1940 Calcium and synaptic transmission in a sympathetic ganglion. *J. Physiol.* **97**, 408–416. (doi:10.1113/jphysiol.1940.sp003818)
- Baker PF, Knight DE. 1981 Calcium control of exocytosis and endocytosis in bovine adrenal medullary cells. *Phil. Trans. R. Soc. Lond. B* **296**, 83–103. (doi:10.1098/rstb.1981.0174)
- Katz B, Miledi R. 1967 The timing of calcium action during neuromuscular transmission. *J. Physiol.* **189**, 535–544. (doi:10.1113/jphysiol.1967.sp008183)
- Neher E, Augustine GJ. 1992 Calcium gradients and buffers in bovine chromaffin cells. *J. Physiol.* **450**, 273–301. (doi:10.1113/jphysiol.1992.sp019127)
- Petersen OH, Maruyama Y, Graf J, Laugier R, Nishiyama A, Pearson GT. 1981 Ionic currents across pancreatic acinar cell membranes and their role in fluid secretion. *Phil. Trans. R. Soc. Lond. B* **296**, 151–166. (doi:10.1098/rstb.1981.0179)
- Schulz I, Kimura T, Wakasugi H, Haase W, Kribben A. 1981 Analysis of Ca<sup>2+</sup> fluxes and pools in pancreatic acini. *Phil. Trans. R. Soc. Lond. B* **296**, 105–113. (doi:10.1098/rstb.1981.0175)
- Nielsen SP, Petersen OH. 1972 Transport of calcium in the perfused submandibular gland of the cat. *J. Physiol.* **223**, 685–697. (doi:10.1113/jphysiol.1972.sp009869)
- Streb H, Irvine RF, Berridge MJ, Schulz I. 1983 Release of Ca<sup>2+</sup> from a nonmitochondrial intracellular store in pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* **306**, 67–69. (doi:10.1038/306067a0)
- Gerasimenko JV *et al.* 2013 Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> channel blockade as a potential tool in antipancreatitis therapy. *Proc. Natl. Acad. Sci. USA* **110**, 13 186–13 191. (doi:10.1073/pnas.1300910110)
- Wakui M, Potter BVL, Petersen OH. 1989 Pulsatile intracellular calcium release does not depend on fluctuations in inositol trisphosphate concentration. *Nature* **339**, 317–320. (doi:10.1038/339317a0)
- Thorn P, Lawrie AM, Smith PM, Gallacher DV, Petersen OH. 1993 Local and global cytosolic Ca<sup>2+</sup> oscillations in exocrine cells evoked by agonists and inositol trisphosphate. *Cell* **74**, 661–668. (doi:10.1016/0092-8674(93)90513-P)
- Mogami H, Nakano K, Tepikin AV, Petersen OH. 1997 Ca<sup>2+</sup> flow via tunnels in polarized cells: recharging of apical Ca<sup>2+</sup> stores by focal Ca<sup>2+</sup> entry through basal membrane patch. *Cell* **88**, 49–55. (doi:10.1016/S0092-8674(00)81857-7)
- Petersen OH, Tepikin AV. 2008 Polarized calcium signaling in exocrine gland cells. *Annu. Rev. Physiol.* **70**, 273–299. (doi:10.1146/annurev.physiol.70.113006.100618)
- Kasai H, Li YX, Miyashita Y. 1993 Subcellular distribution of Ca<sup>2+</sup> release channels underlying Ca<sup>2+</sup> waves and oscillations in exocrine pancreas. *Cell* **74**, 669–677. (doi:10.1016/0092-8674(93)90514-Q)
- Nathanson MH, Fallon MB, Padfield PJ, Maranto AR. 1994 Localization of the type 3 inositol 1,4,5-trisphosphate receptor in the Ca<sup>2+</sup> wave trigger zone of pancreatic acinar cells. *J. Biol. Chem.* **269**, 4693–4696.
- Belan PV, Gerasimenko OV, Tepikin AV, Petersen OH. 1996 Localization of Ca<sup>2+</sup> extrusion sites in pancreatic acinar cells. *J. Biol. Chem.* **271**, 7615–7619. (doi:10.1074/jbc.271.13.7615)
- Parekh AB. 2010 Store-operated CRAC channels: function in health and disease. *Nat. Rev. Drug Discov.* **9**, 399–410. (doi:10.1038/nrd3136)
- Lur G, Haynes LP, Prior IA, Gerasimenko OV, Feske S, Petersen OH, Burgoyne RD, Tepikin AV. 2009 Ribosome-free terminals of rough ER allow formation of STIM1 puncta and segregation of STIM1 from IP<sub>3</sub> receptors. *Curr. Biol.* **19**, 1648–1653. (doi:10.1016/j.cub.2009.07.072)
- Tinel H, Cancela JM, Mogami H, Gerasimenko JV, Gerasimenko OV, Tepikin AV, Petersen OH. 1999

- Active mitochondria surrounding the pancreatic acinar granule region prevent spreading of inositol trisphosphate-evoked local cytosolic  $\text{Ca}^{2+}$  signals. *EMBO J.* **18**, 4999–5008. (doi:10.1093/emboj/18.18.4999)
20. Gerasimenko OV, Gerasimenko JV. 2012 Mitochondrial function and malfunction in the pathophysiology of pancreatitis. *Pflugers Arch.* **464**, 89–99. (doi:10.1007/s00424-012-1117-8)
  21. Denton RM, McCormack JG. 1986 The calcium-sensitive dehydrogenases of vertebrate mitochondria. *Cell Calcium* **7**, 377–386. (doi:10.1016/0143-4160(86)90040-0)
  22. Voronina SG, Barrow SL, Simpson AWM, Gerasimenko OV, Da Silva Xavier G, Rutter GA, Petersen OH, Tepikin AV. 2010 Dynamic changes in cytosolic and mitochondrial ATP levels in pancreatic acinar cells. *Gastroenterology* **138**, 1976–1987. (doi:10.1053/j.gastro.2010.01.037)
  23. Wakui M, Osipchuk YV, Petersen OH. 1990 Receptor-activated cytoplasmic  $\text{Ca}^{2+}$  spiking mediated by inositol trisphosphate is due to  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release. *Cell* **63**, 1025–1032. (doi:10.1016/0092-8674(90)90505-9)
  24. Futatsugi A *et al.* 2005  $\text{IP}_3$  receptor types 2 and 3 mediate exocrine secretion underlying energy metabolism. *Science* **309**, 2232–2234. (doi:10.1126/science.1114110)
  25. Cancela JM, Gerasimenko OV, Gerasimenko JV, Tepikin AV, Petersen OH. 2000 Two different but converging messenger pathways to intracellular  $\text{Ca}^{2+}$  release: the roles of NAADP, cADPR and  $\text{IP}_3$ . *EMBO J.* **19**, 2549–2557. (doi:10.1093/emboj/19.11.2549)
  26. Yamasaki M, Thomas JM, Churchill GC, Garnham C, Lewis AM, Cancela JM, Patel S, Galione A. 2005 Role of NAADP and cADPR in the induction and maintenance of agonist-evoked  $\text{Ca}^{2+}$  spiking in mouse pancreatic acinar cells. *Curr. Biol.* **9**, 874–878. (doi:10.1016/j.cub.2005.04.033)
  27. Gerasimenko JV, Charlesworth RM, Sherwood MW, Ferdek PE, Mikoshiba K, Parrington J, Petersen OH, Gerasimenko OV. 2015 Both RyRs and TPCs are required for NAADP-induced intracellular  $\text{Ca}^{2+}$  release. *Cell Calcium* **58**, 237–245. (doi:10.1016/j.ceca.2015.05.005)
  28. Scratcherd T, Hutson D, Case RM. 1981 Ionic transport mechanisms underlying fluid secretion by the pancreas. *Phil. Trans. R. Soc. Lond. B* **296**, 167–178. (doi:10.1098/rstb.1981.0180)
  29. Maléth J, Hegyi P. 2016  $\text{Ca}^{2+}$  toxicity and mitochondrial damage in acute pancreatitis: translational overview. *Phil. Trans. R. Soc. B* **371**, 20150425. (doi:10.1098/rstb.2015.0425)
  30. Hegyi P, Petersen OH. 2013 The exocrine pancreas: the acinar-ductal tango in physiology and pathophysiology. *Rev. Physiol. Biochem. Pharmacol.* **165**, 1–30. (doi:10.1007/112\_2013\_14)
  31. Gryshchenko O, Gerasimenko JV, Gerasimenko OV, Petersen OH. 2016  $\text{Ca}^{2+}$  signals mediated by bradykinin type 2 receptors in normal pancreatic stellate cells can be inhibited by specific  $\text{Ca}^{2+}$  channel blockade. *J. Physiol.* **594**, 281–293. (doi:10.1113/JP271468)
  32. Gryshchenko O, Gerasimenko JV, Gerasimenko OV, Petersen OH. 2016 Calcium signalling in pancreatic stellate cells: mechanisms and potential roles. *Cell Calcium* **59**, 140–144. (doi:10.1016/j.ceca.2016.02.003)
  33. Petersen OH, Sutton R. 2006  $\text{Ca}^{2+}$  signalling and pancreatitis: effects of alcohol, bile and coffee. *Trends Pharmacol. Sci.* **27**, 113–120. (doi:10.1016/j.tips.2005.12.006)
  34. Ward JB, Petersen OH, Jenkins SA, Sutton R. 1995 Is an elevated concentration of acinar cytosolic free ionised calcium the trigger for acute pancreatitis? *Lancet* **346**, 1016–1019. (doi:10.1016/S0140-6736(95)91695-4)
  35. Raraty M, Ward J, Erdemli G, Vaillant C, Neoptolemos JP, Sutton R, Petersen OH. 2000 Calcium-dependent enzyme activation and vacuole formation in the apical granular region of pancreatic acinar cells. *Proc. Natl. Acad. Sci. USA* **97**, 13 126–13 131. (doi:10.1073/pnas.97.24.13126)
  36. Kruger B, Albrecht E, Lerch MM. 2000 The role of intracellular calcium signaling in premature protease activation and the onset of pancreatitis. *Am. J. Pathol.* **157**, 43–50. (doi:10.1016/S0002-9440(10)64515-4)
  37. Voronina S, Longbottom R, Sutton R, Petersen OH, Tepikin AV. 2002 Bile acids induce calcium signals in mouse pancreatic acinar cells: implications for bile-induced pancreatic pathology. *J. Physiol.* **540**, 49–55. (doi:10.1113/jphysiol.2002.017525)
  38. Gerasimenko JV, Gerasimenko OV, Petersen OH. 2014 The role of  $\text{Ca}^{2+}$  in the pathophysiology of pancreatitis. *J. Physiol.* **592**, 269–280. (doi:10.1113/jphysiol.2013.261784)
  39. Mukherjee R *et al.* 2016 Mechanism of mitochondrial permeability transition pore induction and damage in the pancreas: inhibition prevents acute pancreatitis by protecting production of ATP. *Gut*. In press. (doi:10.1136/gutjnl-2014-308553)
  40. Maléth J, Hegyi P, Rakonczay Jr, Z, Venglovecz V. 2015 Breakdown of bioenergetics evoked by mitochondrial damage in acute pancreatitis: mechanisms and consequences. *Pancreatology* **15**, S18–S22. (doi:10.1016/j.pan.2015.06.002)
  41. Criddle DN. 2015 The role of fat and alcohol in acute pancreatitis: a dangerous liaison. *Pancreatology* **15**, S6–S12. (doi:10.1016/j.pan.2015.02.009)
  42. Baggaley EM, Elliott AC, Bruce JL. 2008 Oxidant-induced inhibition of the plasma membrane  $\text{Ca}^{2+}$ -ATPase in pancreatic acinar cells: role of the mitochondria. *Am. J. Physiol. Cell Physiol.* **295**, C1247–C1260. (doi:10.1152/ajpcell.00083.2008)
  43. Wen L *et al.* 2015 Inhibitors of ORAI1 prevent cytosolic calcium-associated injury of human pancreatic acinar cells and acute pancreatitis in 3 mouse models. *Gastroenterology* **149**, 481–492. (doi:10.1053/j.gastro.2015.04.015)
  44. Hegyi P. 2016 Blockade of calcium entry provides a therapeutic window in acute pancreatitis. *J. Physiol.* **594**, 257. (doi:10.1113/JP271710)
  45. Jaffe N, Traggis D, Das L, Moloney WC, Hann HW, Kim BS, Nair R. 1971 L-asparaginase in the treatment of neoplastic diseases in children. *Cancer Res.* **31**, 942–949.
  46. Silverman LB *et al.* 2001 Improved outcome for children with acute lymphoblastic leukemia: results of Dana-Farber Consortium Protocol 91-01. *Blood* **97**, 1211–1218. (doi:10.1182/blood.V97.5.1211)
  47. Kearney SL, Dahlberg SE, Levy DE, Voss SD, Sallan SE, Silverman LB. 2009 Clinical course and outcome in children with acute lymphoblastic leukemia and asparaginase-associated pancreatitis. *Pediatr. Blood Cancer* **53**, 162–167. (doi:10.1002/pbc.22076)
  48. Raja RA, Schmiegelow K, Frandsen TL. 2012 Asparaginase-associated pancreatitis in children. *Br. J. Haematol.* **159**, 18–27. (doi:10.1111/bjh.12016)
  49. Raja RA *et al.* 2014 Asparaginase-associated pancreatitis in children with acute lymphoblastic leukaemia in the NOPHO ALL2008 protocol. *Br. J. Haematol.* **165**, 126–133. (doi:10.1111/bjh.12733)
  50. Samarasinghe S, Dhir S, Slack J, Iyer P, Wade R, Clack R, Vora A, Goulden N. 2013 Incidence and outcome of pancreatitis in children and young adults with acute lymphoblastic leukaemia treated on a contemporary protocol, UKALL 2003. *Br. J. Haematol.* **162**, 710–713. (doi:10.1111/bjh.12407)
  51. Huang W *et al.* 2014 Fatty acid ethyl ester synthase inhibition ameliorates ethanol-induced  $\text{Ca}^{2+}$ -dependent mitochondrial dysfunction and acute pancreatitis. *Gut* **63**, 1313–1324. (doi:10.1136/gutjnl-2012-304058)
  52. Voronina S *et al.* 2015 The role of  $\text{Ca}^{2+}$  influx in endocytic vacuole formation in pancreatic acinar cells. *Biochem. J.* **465**, 405–412. (doi:10.1042/BJ20140398)
  53. Namkund W, Han W, Luo X, Muallem S, Cho KH, Kim KH, Lee MG. 2004 Protease-activated receptor 2 exerts local protection and mediates some systemic complications in acute pancreatitis. *Gastroenterology* **126**, 1844–1859. (doi:10.1053/j.gastro.2004.03.019)
  54. Laukkarinen JM, Weiss ER, van Acker GJD, Steer ML, Perides G. 2008 Protease-activated receptor-2 exerts contrasting model-specific effects on acute experimental pancreatitis. *J. Biol. Chem.* **283**, 20 703–20 712. (doi:10.1074/jbc.M801779200)
  55. Soh UJ, Dores MR, Chen B, Trejo J. 2010 Signal transduction by protease-activated receptors. *Br. J. Pharmacol.* **160**, 191–203. (doi:10.1111/j.1476-5381.2010.00705.x)
  56. Leyssens A, Nowicky AV, Patterson L, Crompton M, Duchon MR. 1996 The relationship between mitochondrial state, ATP hydrolysis,  $[\text{Mg}^{2+}]_i$  and  $[\text{Ca}^{2+}]_i$  studied in isolated rat cardiomyocytes. *J. Physiol.* **496**, 111–128. (doi:10.1113/jphysiol.1996.sp021669)
  57. Inoue M, Fujishiro N, Imanaga I, Sakamoto Y. 2002 Role of ATP decrease in secretion induced by mitochondrial dysfunction in guinea-pig adrenal chromaffin cells. *J. Physiol.* **539**, 145–155. (doi:10.1113/jphysiol.2001.012936)
  58. Criddle D, Murphy J, Fistetto G, Barrow S, Tepikin AV, Neoptolemos JP, Sutton R, Petersen OH. 2006

- Fatty acid ethyl esters cause pancreatic calcium toxicity via inositol trisphosphate receptors and loss of ATP synthesis. *Gastroenterology* **130**, 781–793. (doi:10.1053/j.gastro.2005.12.031)
59. Luan ZG, Ma XC, Zhang H, Zhang C, Guo RX. 2013 Protective effect of ethyl pyruvate on pancreas injury in rats with severe acute pancreatitis. *J. Surg. Res.* **181**, 76–84. (doi:10.1016/j.jss.2012.05.066)
60. Kiyonaka S *et al.* 2009 Selective and direct inhibition of TRPC3 channels underlies biological activities of a pyrazole compound. *Proc. Natl. Acad. Sci. USA* **106**, 5400–5405. (doi:10.1073/pnas.0808793106)
61. Schleifer H *et al.* 2012 Novel pyrazole compounds for pharmacological discrimination between receptor-operated and store-operated  $\text{Ca}^{2+}$  entry pathways. *Br. J. Pharmacol.* **167**, 1712–1722. (doi:10.1111/j.1476-5381.2012.02126.x)
62. Broome JD. 1968 Studies on the mechanism of tumor inhibition by L-asparaginase. Effects of the enzyme on asparagine levels in the blood, normal tissues, and 6C3HED lymphomas of mice: differences in asparagine formation and utilization in asparaginase-sensitive and -resistant lymphoma cells. *J. Exp. Med.* **127**, 1055–1072. (doi:10.1084/jem.127.6.1055)
63. Spät A, Szanda G. 2012 Special features of mitochondrial  $\text{Ca}^{2+}$  signalling in adrenal glomerulosa cells. *Pflügers Arch.* **464**, 43–50. (doi:10.1007/s00424-012-1086-y)
64. Vais H, Mallilankaraman K, Mak DO, Hoff H, Payne R, Tanis JE, Foskett JK. 2016 EMRE is a matrix  $\text{Ca}^{2+}$  sensor that governs gatekeeping of the mitochondrial  $\text{Ca}^{2+}$  uniporter. *Cell Rep.* **14**, 403–410. (doi:10.1016/j.celrep.2015.12.054)
65. Kar P, Samanta K, Kramer H, Morris O, Bakowski D, Parekh AB. 2014 Dynamic assembly of a membrane signaling complex enables selective activation of NFAT by Orai1. *Curr. Biol.* **24**, 1361–1368. (doi:10.1016/j.cub.2014.04.046)