

The ability to enter into an avirulent viable but non-culturable (VBNC) form is widespread among *Listeria monocytogenes* isolates from salmon, patients and environment

Toril LINDBÄCK^{1*}, Martin E. ROTTENBERG², Sylvie M. ROCHE³, Liv Marit RØRVIK¹

¹ Department of Food Safety and Infection Biology, Norwegian School of Veterinary Science, P.O. Box 8146 Dep., N-0033 Oslo, Norway

² Microbiology & Tumorbiology Center, Karolinska Institute, Stockholm, Sweden

³ INRA, UR 1282, Infectiologie Animale et Santé Publique, IFR136, Agents transmissibles et Infectiologie, F-37380 Nouzilly, France

(Received 12 May 2009; accepted 30 September 2009)

Abstract – Media-based bacteriological testing will fail to detect non-culturable organisms and the risk of consuming viable but non-culturable (VBNC) *Listeria monocytogenes* is unknown. We have here studied whether *L. monocytogenes* obtained from seafoods, processing environment and clinical cases enter the VBNC state and assessed the virulence of the non-culturable forms of the bacteria. A number of 16 *L. monocytogenes* strains were starved in microcosm water at 4 °C until loss of culturability. Metabolic activity in the VBNC form was measured as ATP generation using a luciferase assay and membrane integrity was examined using the LIVE/DEAD BacLight assay. All tested *L. monocytogenes* strains entered the VBNC state after starvation in microcosm water. Ongoing mRNA synthesis of *hly* in VBNC *L. monocytogenes* cells re-incubated in culture medium indicated a potential virulence of these forms. Sodium pyruvate and replenishment of nutrient were used in attempts to resuscitate VBNC cells. However, VBNC *L. monocytogenes* were not resuscitated under these conditions. VBNC *L. monocytogenes* were tested for virulence in a cell plaque assay and by intraperitoneally inoculation in immunodeficient RAG1^{-/-} mice. Inoculation of VBNC *L. monocytogenes* in immunodeficient mice did not cause morbidity, and plaque assay on HT-29 cells in culture indicated that the VBNC cells were avirulent. The results indicate that the risk of non-culturable *L. monocytogenes* in foods, when the VBNC state is induced by starvation, is negligible.

Listeria monocytogenes / VBNC / ATP / mRNA

1. INTRODUCTION

The occurrence of *Listeria monocytogenes* in foods has been a growing concern over the last decades. The bacteria cause listeriosis, a rare but serious illness, especially for the risk groups; the elderly, unborn children and

immunocompromised persons. In 2007, 0.3 cases per 100 000 were reported in the EU states, with a mortality of 20% [2]. Several small outbreaks of listeriosis have been associated with seafood [11, 22, 23, 30], and some countries have zero tolerance for *L. monocytogenes* in imported fish. Producers of seafood products demand *L. monocytogenes* free raw material for further processing, and the surveillance of

* Corresponding author: toril.lindback@nvh.no

L. monocytogenes during seafood-processing is becoming increasingly important.

Viable but non-culturable (VBNC) bacteria have been described as organisms that fail to grow and develop colonies on routine bacteriological media, but which are to be considered alive because of their capability of metabolic activity [34]. A number of pathogens including *Vibrio vulnificus* [33], *Legionella pneumophila* [45], *Salmonella typhi* [16] and *Campylobacter jejuni* [46] have been described to enter the state of VBNC after exposure to different stress factors such as starvation, shifts in temperature, high NaCl concentration and visible light [36]. *L. monocytogenes* has been shown to enter the VBNC state in response to starvation and exposure to high NaCl concentrations [6, 8]. However, the studies included only four culture collection strains, of which only two were able to enter the VBNC state after starvation in microcosm water.

ATP levels, declining rapidly in dead cells, have been reported to remain high in VBNC cells, thus presence of ATP is a good viability marker of cells in the VBNC state [9, 24, 27]. The LIVE/DEAD BacLight assay, utilizing a mixture of the stains SYTO 9 and propidium iodide to evaluate cell membrane integrity, has been used to confirm viability in non-cultural bacteria [1, 27, 37]. Detection of mRNA, due to its central role in cell metabolism and its very short half-life, has previously been used as a viability marker for several pathogens including *Vibrio cholerae* [4], *Escherichia coli* [43], *L. monocytogenes* [26] and *Vibrio parahaemolyticus* [18]. Since Listeriolysin O (*hly*) is an important virulence factor in *L. monocytogenes*, expression of *hly* in the VBNC state will indicate a virulence potential present in the bacteria in the VBNC state.

The ability of *L. monocytogenes* to enter the VBNC state is of concern since media-based bacteriological tests used in routine will fail to detect non-culturable organisms from the processing environments and the products, and the hazard risks of consuming VBNC *L. monocytogenes* is still unknown. VBNC *L. monocytogenes* in foods might theoretically be able to resuscitate and account for virulence. Most studies on resuscitation have been performed

on *Vibrio* sp., where a simple temperature up-shift or supplementing media with H₂O₂-degrading compounds such as Na-pyruvate results in resuscitation [31, 33, 35]. VBNC *L. monocytogenes* cannot be resuscitated directly in culture medium [8, 20], nevertheless they have been shown to resuscitate under appropriate conditions such as in embryonated eggs [14].

To our knowledge, little is known about how VBNC *L. monocytogenes* may resuscitate in the human organism. In the present study, we sought to investigate whether *L. monocytogenes* isolated from salmon and salmon-processing environment and clinical isolates were able to enter the VBNC state during starvation, and if the VBNC forms were virulent towards immunodeficient mice and human colon adenocarcinoma cells in culture.

2. MATERIALS AND METHODS

2.1. Bacterial strains and culture conditions

Fourteen strains of *L. monocytogenes* comprising six clinical isolates, five strains isolated from salmon and three strains isolated from processing plants [42] were studied. In addition, the Scott A (Pasteur Institute Culture Collection, France) and the EGDe strain [32] were included. Unless otherwise stated, bacteria were cultured in Brain Heart Infusion broth (BHI) (Oxoid, Basingstoke, UK) at 32 °C. Colony forming units (CFU) were determined by plating appropriate dilutions on 5% bovine blood in blood agar base (Oxoid), followed by incubation at 37 °C for 24 or 48 h.

2.2. Starvation in microcosm water

Overnight BHI cultures of *L. monocytogenes* were washed three times in microcosm water (filtered, autoclaved MQ water adjusted to pH 6.0) [41], and diluted to 5×10^7 organisms/mL in 100 mL microcosm water in 250 mL flasks. The flasks were incubated at 4 °C with gentle shaking at 100 rpm. At inoculation and each week samples from the flasks were plated on blood agar for determination of CFU. The plates were incubated at 37 °C for 48 h. When no CFU were detected in 1 mL microcosm water, the organisms were examined for metabolic activity (ATP generation), damage to membranes (LIVE/DEAD BacLight™ viability

staining) and accumulation of *hly* mRNA. The flasks were kept at 4 °C for 12 months for further examinations.

2.3. ATP assay

In order to test for ATP generation, 3 mL of the *L. monocytogenes* in microcosm water was pelleted for 5 min at $10\,000 \times g$, and re-suspended in 220 μ L BHI. The samples were divided in two, and ATP was quantified in half of the sample immediately while the other half was incubated at 32 °C for 1 h before ATP quantification. ATP was quantified using the BacTiter-Glo Microbial Cell Viability assay kit (Promega, Madison, WI, USA) where 100 μ L sample was mixed with 100 μ L luciferase reagent and incubated for 5 min at room temperature. Luminescence (relative light units (RLU)) was recorded by a GloMax Integrated Luminescence System (Promega). The remaining 10 μ L of each sample was plated on blood agar to exclude that quantified ATP was generated by a limited number of cells that have not lost culturability. Previous experiments had shown a detection limit of 1×10^3 culturable *L. monocytogenes* cells by the BacTiter-Glo Microbial Cell Viability assay kit¹, thus plating of 10 μ L would reveal culturable cells above the detection limit. ATP was quantified in BHI without VBNC cells as negative control. Each strain was tested in at least three experiments. An ATP standard curve was used for conversion of RLU values into ATP concentration (Roche Diagnostics, Mannheim, Germany).

Four randomly chosen strains (L839, L1069, L1372 and L2842) were grown to stationary phase, and diluted samples from the cultures were tested for ATP content, and plated on blood agar for determination of CFU. A curve was created by plotting ATP concentration against CFU, to estimate the number of *L. monocytogenes* cells corresponding to a known ATP concentration.

2.4. LIVE/DEAD BacLight™ bacterial viability fluorescence microscopy

Viability of non-culturable *L. monocytogenes* was examined using the LIVE/DEAD BacLight™ bacteria viability kit L-7007 (Molecular Probes, Eugene, OR, USA) according to the manufacturer's directions, and visualized with an Olympus IX81 inverted fluorescence microscope. The LIVE/DEAD BacLight™ assay is based on cell membrane integrity. The fluorescent SYTO 9 stain viable bacteria cells with intact

membranes green, while propidium iodide stains dead organisms with compromised membranes red. For each of the 16 tested strains, viable bacteria were observed in five random fields in each of two independent experiments from the same VBNC culture.

2.5. Transcriptional analysis by reverse transcriptase PCR (RT-PCR)

To demonstrate de novo mRNA synthesis of *hly* in VBNC *L. monocytogenes*, VBNC cells were incubated in BHI with and without rifampicin (Sigma-Aldrich), which inhibits mRNA synthesis. Five mL of VBNC solution was spun down at $10\,000 \times g$ for 5 min, re-suspended in BHI with 10 μ g/mL rifampicin (Sigma-Aldrich) and incubated at 32 °C for 2–10 h. The RNeasy Mini Kit (Qiagen, Hilden, Germany) was used for isolation of total RNA. The cDNA synthesis was performed using 0.5 μ g of DNase treated RNA; 0.2 μ M reverse *hly* primer, 1 mM dNTP mix; 10 mM DTT, 1 \times First Strand Synthesis Buffer (Invitrogen, Carlsbad, CA, USA) and 200 U SuperScript III Reverse Transcriptase (Invitrogen), and incubated at 55 °C for 60 min. RNA was incubated in cDNA synthesis reaction buffer without reverse transcriptase to serve as a control for DNA contamination. Eight μ L of the RT reaction was used in the following PCR amplification using 2 U DyNAzyme (Finnzymes Oy, Espoo, Finland) under following conditions; 94 °C for 4 min followed by a temperature cycling routine of denaturation at 94 °C for 1 min, annealing at 52 °C for 1 min and extension at 72 °C for 1 min. After 36 cycles a final extension at 72 °C for 5 min was performed and the tubes were cooled to 4 °C. The primers used in the cDNA synthesis and the PCR reactions were *hly* forward (5'-ATTTCCCTCACTGATTGC-3') and *hly* reverse (5'-CACTCAGCATTGATTGCCA-3') [17].

2.6. Resuscitation by replenishment of nutrition

Attempts to resuscitate the VBNC *L. monocytogenes* cells by replenishment of nutrients were done at 12 °C and 32 °C on 4 strains; Scott A (reference strain), L839 (salmon), L2842 (smoked salmon) and L3365 (clinical isolate). Erlenmeyer flasks containing three different concentrations of BHI medium (0.1%, 1% and 10% of the concentration recommended by manufacturer) were incubated at the two different temperatures for each strain. Five mL diluted BHI medium was inoculated with 100 μ L each from the flasks with non-culturable *L. monocytogenes* strains, and incubated for 3 days at 12 °C and

¹ Lindbäck T., unpublished results.

32 °C. After 3 days, samples of 100 µL were plated on blood agar. The concentration of BHI in the flasks was then increased to 1%, 10% and 50%, respectively. The flasks were further incubated for 3 days and samples of 100 µL were plated on blood agar. A third sample was plated on blood agar after 9 days of incubation. The blood agar plates were incubated overnight at 32 °C. The experiment was performed twice.

2.7. Resuscitation with Na-pyruvate

H₂O₂-degrading agents such as sodium pyruvate are previously shown to facilitate resuscitation [47] of VBNC organisms. Samples of 0.1 mL microcosms with VNBC *L. monocytogenes* cells were plated on BHI agar with 0.1% sodium pyruvate. Plates were checked for colonies after 24 and 48 h incubation.

2.8. Plaque-forming cell assay

The virulence of *L. monocytogenes* strains and their VBNC forms was estimated by a plaque-forming assay [40]. The virulent strain EGDe BUG1600 (Pasteur Institute culture Collection, France) was used as control. Human colon adenocarcinoma cell line HT-29 (ECACC N° 85061109, Salisbury, UK) was used between passages 86 and 95. Cells were grown in 75-cm² plastic tissue culture flasks (Falcon, VWR) in Dulbecco's Modified Eagle Medium with 4.5 g/L glucose (DMEM; GIBCO, Invitrogen) supplemented with 10% (v/v) foetal calf serum (PAA) and 2 mM L-glutamine (PAA). Antibiotics (100 IU/mL penicillin, 100 µg/mL streptomycin (PAA)) were routinely added to the culture medium during maintenance of the cells, but not for the virulence assays. Cells were maintained in culture in a humidified incubator (NuAire DH Autoflow, MN, USA) at 37 °C under 5% (v/v) CO₂. Confluent HT-29 cell monolayers were cultured in DMEM supplemented with antibiotics in 96-well tissue culture plates for 4 or 5 days (Falcon, VWR) after inoculum of 3.2×10^4 cells or 2.4×10^4 cells in 100 µL culture medium per well, respectively. One day before infection, cells were incubated in culture medium without antibiotics. Numbers of 10^2 to 10^7 *L. monocytogenes* cells from an overnight BHI culture at 37 °C or 10^7 VBNC *L. monocytogenes* cells were added to each well of the monolayers. After 2 h at 37 °C, the extracellular bacteria were killed by incubation with 100 µg/mL gentamicin for 1 h 30 min. The HT-29 cell monolayers were then overlaid with medium containing 0.8% SeaPlaque[®] GTG[®] Agarose (Lonza, Rockland, ME,

USA) in the culture medium, supplemented with 10 µg/mL gentamicin, and further on with the latter culture medium without agarose. After overnight incubation, the wells were examined for formation of plaques using an inverted microscope. Duplicate wells were used and the experiment was repeated for each strain. Results were expressed as the number of plaques obtained for 7 log *L. monocytogenes* or VBNC *L. monocytogenes* per well. The mean of the results from all the experiments was used for determination the virulence of the EGDe control strain. Strains with values higher than the lower CI limit of the EGDe value were considered to be virulent.

2.9. Injection of immunodeficient mice with VBNC *L. monocytogenes* Scott A

L. monocytogenes Scott A starved to non-culturability for 12 weeks in microcosm water were concentrated by centrifugation to approximately 1×10^7 VBNC cells per 100 µL (according to the LIVE/DEAD BacLight[™] assay) and injected intraperitoneally in PBS into four RAG1^{-/-} mice. Mice were routinely weighted, sacrificed 2 weeks after injection and liver and spleen lysates prepared in 5 and 2 mL 0.1% triton containing PBS. One hundred µL of the lysates were then plated on BHI plates. The experiments in mice comply with the Karolinska Institute's regulations for care and use of laboratory animals.

3. RESULTS

3.1. Starvation in microcosm water

All tested strains of *L. monocytogenes* ($n = 16$) entered a non-culturable state after 5–12 weeks of incubation in microcosm water (mean 7.9 ± 2.6 SD). Evolution of the culturable count of *L. monocytogenes* Scott A starved in microcosm water is shown in Figure 1. The decrease in CFU varied substantially among the strains, and no correlation was observed between the source of isolation and the rate of decrease in CFU/mL.

3.2. Viability of non-culturable *L. monocytogenes* cells

Non-culturable cells of all the 16 tested strains generated ATP, although they did not




Figure 1. Evolution of the culturable count (●) and the viable count (■) of *L. monocytogenes* Scott A in microcosm water. The culturable cells were enumerated by spread plate counts, while viable counts were carried out using the LIVE/DEAD BacLight™ bacteria viability kit. The standard deviation is indicated.

grow on blood agar. Figure 2 shows the average ATP detected in non-culturable cells of the 16 tested strains after incubation in BHI for 1 h (3.4 ± 1.5 nM SD). The ATP concentration remained at the approximately same level (2.3 ± 1.5 nM SD) when measured after 3 days of incubation in BHI. The production of ATP indicated that the non-culturable cells were viable, and confirmed that they had entered the VBNC state. ATP generation measured in the strains maintained in microcosm water for 12 weeks, 6 and 12 months were similar, indicating that the VBNC *L. monocytogenes* were stable for at least one year in the microcosm water.

An ATP concentration of 3.4 nM (the mean ATP concentration in the VBNC cells measured in 1.5 mL microcosm water) correspond to the ATP concentration in approximately 8×10^5 culturable *L. monocytogenes* cells. The total numbers of *L. monocytogenes* cells originally inoculated in the microcosm water were 5×10^7 cells/mL, thus 7.5×10^7 cells were tested for ATP content in each sample. This indicates that approximately 1% of the starved cells entered the VBNC state.




Figure 2. Average generation of ATP by 16 *L. monocytogenes* strains in the VBNC state were monitored by measuring ATP concentration in the cells directly from microcosm water (T0) and after 1 h of incubation in BHI (T1). BHI as negative control showed no variation in ATP concentration. The standard deviation is indicated.

Despite the loss of culturability, LIVE/DEAD BacLight™ assay of *L. monocytogenes* strains starved to non-culturability showed that $68 \pm 16\%$ SD of the cells remained viable (average of the 16 tested strains). Figure 3 shows microscopic pictures of LIVE/DEAD BacLight™ assay of VBNC *L. monocytogenes* ScottA.

3.3. Transcriptional analysis of *hly*

Transcripts of *hly* were detected in VBNC cells of all the 16 strains by RT-PCR. To confirm that detected *hly* mRNA was a result of ongoing transcription, VBNC *L. monocytogenes* were incubated in BHI containing rifampicin which inhibits de novo mRNA synthesis, thus degradation of present mRNA can be studied. The variation in degradation of *hly* transcript with time for four of the strains is shown in Figure 4. In two of the tested strains, Scott A and L839, no *hly* transcripts could be detected 8 h after incubation with rifampicin. For the EGDe strain and strain L1069 a weak RT-PCR product indicated small amounts of *hly* mRNA 10 h after incubation with rifampicin.




Figure 3. Microscopic pictures of LIVE/DEAD BacLight™ assay of VBNC *L. monocytogenes* Scott A. Left: directly from BHI culture, day 0. Right: after 12 weeks in microcosm water. (A color version of this figure is available on line at www.vetres.org.)




Figure 4. Expression of *hly* in five *L. monocytogenes* strains in the VBNC state after 2, 4, 6, 8 and 10 h incubation at 32 °C in BHI containing rifampicin detected by RT-PCR. The positive control (lane pos) was incubated in BHI without rifampicin for 10 h. Experiments were performed at least twice.

3.4. Resuscitation

Attempts to resuscitate the VBNC *L. monocytogenes* cells by replenishment of nutrients were done at 12 °C and 32 °C on 4 of the 16 strains (L1069, L2842, L3365 and Scott A). None of the tested strains did resume growing in any concentration of BHI at any tested temperature. Neither plating VBNC cells directly on BHI plates containing 0.1% Na-pyruvate

nor incubating VBNC cells in BHI cultures containing 0.1% Na-pyruvate, prior to plating, did result in resuscitation.

3.5. Virulence in plaque-forming assay and immunodeficient mice

All the 16 tested *L. monocytogenes* strains expressed virulence in the plaque-forming assay in their culturable form, except for L1338 (4.28 ± 0.24 log). This strain was therefore tested in a mouse assay, and was shown to be virulent (data not shown). In contrast, none of the 16 tested VBNC *L. monocytogenes* expressed virulence (Tab. I).

VBNC *L. monocytogenes* Scott A were injected i.p. in RAG1^{-/-} mice. Despite the high sensitivity of these B and T cell deficient animals to infection with *L. monocytogenes*, bacteria were not detected in spleen or livers from mice inoculated with 10⁷ VBNC *L. monocytogenes*.

4. DISCUSSION

In this study we show that all examined strains (16) of *L. monocytogenes* isolated from salmon, salmon-processing environment and patients, as well as two reference strains, entered the state of VBNC after starvation in microcosm water. We examined numerous

Table I. Expression of virulence of 16 *L. monocytogenes* strains and their VBNC forms, by formation of plaques in the human adenocarcinoma cell line HT-29. The results are expressed as the number of plaques obtained for 7 log CFU deposited per well.

Strain	Source	Virulence (mean log + SD) culturable strains	Virulence (mean log + SD) VBNC
EGDe ^a (control)	Reference strain	5.23 ± 0.40	ND
Scott A	Reference strain	6.38 ± 0.03	0
EGDe ^b	Reference strain	5.18 ± 0.11	0
L1069	Patient	5.75 ± 0.34	0
L1371	Patient	5.18 ± 0.27	0
L1372	Patient	6.70 ± 0.13	0
L3365	Patient	6.18 ± 0.47	0
L3368	Patient	5.35 ± 0.23	0
L3371	Patient	5.66 ± 0.14	0
L839	Salmon	5.44 ± 0.37	0
L1272	Salmon	5.81 ± 0.26	0
L1338 ^c	Salmon	4.28 ± 0.24	0
L2663	Salmon	4.98 ± 0.16	0
L2735	Salmon	5.56 ± 0.21	0
L1317	Drainpipe	4.88 ± 0.27	0
L2842	Salmon waste	5.61 ± 0.31	0
L1232	Seawater	5.88 ± 0.13	0

ND: Not done.

^a Control used in the plaque cell test.

^b Used in the VBNC experiments.

^c Examined in a immunocompetent mouse assay and shown to be virulent.

strains from a wide selection of sources, and strains from different sources had the same ability to enter the VBNC state. All the tested strains entered the VBNC state, in contrast to results obtained by Besnard et al. [7, 8] where only 2 out of 4 culture collection *L. monocytogenes* strains entered the VBNC state after starvation. The difference in results may be due the use of environmental and patient isolates instead of laboratory reference strains, and the fact that the VBNC state was induced by starvation at different temperatures.

The time taken to reach the VBNC state broadly is in agreement with other studies, spanning between 5 and 12 weeks before no CFU could be detected in 1 mL microcosm. It lapsed 2, 4 and 6 weeks, respectively for *Enterococcus faecalis*, *Edwardsiella tarda* and *Aeromonas hydrophila* to become non-culturable in microcosm water at 4 °C [21, 29, 47], however, only a few strains were examined in each of those studies.

ATP quantification, LIVE/DEAD *BacLight* viable counts and mRNA transcription are included in this study to demonstrate viability in the non-culturable *L. monocytogenes* strains. ATP measurements has previously been used as a viability marker [9, 24, 27], and RT-PCR have been used for investigating viability and potential virulence in VBNC *V. parahaemolyticus*, VBNC *Enterococcus faecalis* and VBNC *V. vulnificus* [18, 25, 28]. The so far most common used technique to confirm the VBNC state is the microscopic method using intracellular hydrolysis of CTC (5-cyano-2,3-ditoyl tetrazolium chloride) into an insoluble fluorescent salt by enzymes in the respiratory chain, and LIVE/DEAD *BacLight* viable counts are reported to be comparable to results obtained with the CTC staining assay [10].

When ATP in the VBNC cells was measured after incubation in BHI for 3 days the concentration remained at the same level as for 1 h (Fig. 2), although the VBNC did not regain

the ability to divide and grow on blood agar. This is consistent with results reported previously; *L. monocytogenes* cannot be resuscitated in rich medium such as BHI [8, 20], although the metabolic activity seems to be maintained. When *L. monocytogenes* had entered the state of VBNC and the cells were maintained in microcosm water at 4 °C, we observed that the VBNC cells were stable for at least 12 months, corresponding with the observation presented by Bunker et al. [12] that *Pseudomonas fluorescens* can remain in the VBNC state for over a year. According to the LIVE/DEAD BacLight assay, the number of viable cells after 12 weeks in microcosm water varied between the strains, from 45 to 90% (mean $68 \pm 16\%$ SD).

Our estimation of the proportion of viable cells differed markedly with the two methods employed. If one assumes that the ATP concentration is equivalent in culturable cells and VBNC cells, then the amount of ATP generated by VBNC *L. monocytogenes* suggests that 1% of the starved cells had entered the VBNC state. If, however, the ATP concentration is higher in growing cells than in VBNC cells, the percentage of cells entering the state of VBNC will be correspondingly greater. Even so, this appears to be substantially lower than that predicted from the LIVE/DEAD BacLight assay (68%).

RT-PCR is not a quantitative method and may not be ideal for detection of mRNA half-life, but the results comparing accumulation of transcripts in presence or absence of rifampicin indicated a very long half-life (6–10 h) in VBNC *L. monocytogenes* compared to the mRNA half-life in culturable cells (2–4 min) [5, 38, 39]. Using RT-PCR, Smith and Oliver concluded that detected mRNA in VBNC *V. vulnificus* was a result of de novo synthesis, with a half-life of 60 min [44]. The results shown in Figure 4 clearly indicate an effect of rifampicin in three of the strains, Scott A, L839 and L1069, indicating de novo mRNA synthesis in VBNC *L. monocytogenes*.

Ongoing transcription of *hly* described in the present study together with the fact that VBNC *L. monocytogenes* could be resuscitated in embryonated eggs in a previous study [14], suggest maintenance of potential virulence. However, in this study they appeared avirulent

to immunodeficient RAG1^{-/-} mice and HT-29 cells in culture. This is supported by Cappelier et al. [13] finding that VBNC *L. monocytogenes* LO 28, ATCC 19115, CNL 895807 and Scott A, were negative for virulence in the HT-29 cell assay and in immunocompetent mice.

In this study we have tried to resuscitate the VBNC *L. monocytogenes* by gradually adaptation to nutrients at different temperatures attempting to emulate real life conditions. However, none of the tested strains regained the ability to grow and form colonies by gradual replenishment of nutrients. Neither did addition of the H₂O₂-degrading compound Na-pyruvate facilitate resuscitation of the VBNC *L. monocytogenes* cells. This is in contrast to *Vibrio* spp. [44, 48], *Enterococci* [29] and *Citrobacter* [19] that have been successfully resuscitated in culture medium. *Campylobacter* required animal models such as mice and embryonated eggs to regain culturability [3, 15]. However, *L. monocytogenes* did neither resuscitate after injection in immunocompetent mice in the study by Cappelier et al. [13], nor in immunodeficient mice in the present study.

In conclusion, the entrance of *L. monocytogenes* into the VBNC state is of interest for the food industry since routine bacteriological testing will fail to detect non-culturable organisms in the processing environment and in the products. We found that *L. monocytogenes* isolated from salmon, salmon-processing environment and patients entered the VBNC state after starvation. Despite viability according to ATP generation, membrane integrity and *hly* de novo transcription, the VBNC cells were neither resuscitated by addition of the H₂O₂-degrading compound Na-pyruvate nor by a gradual adaptation to nutrients. In view of the fact that VBNC *L. monocytogenes* did not show virulence in immunodeficient mice and HT-29 cells in culture, we propose that VBNC *L. monocytogenes* induced by starvation are avirulent.

Acknowledgements. The study was supported by grant 169123-NKJ123 from the Norwegian Research Council. We thank Annette Fagerlund, Simon Hardy and Hannah Jørgensen for fruitful discussions and for critically reading the manuscript.

REFERENCES

- [1] Adams B.L., Bates T.C., Oliver J.D., Survival of *Helicobacter pylori* in a natural freshwater environment, Appl. Environ. Microbiol. (2003) 69:7462–7466.
- [2] Anonymous, The Community Summary Report on Trends and sources of zoonoses and zoonotic agents in the European Union in 2007, The EFSA Journal (2009) 223:136–158.
- [3] Baffone W., Casaroli A., Citterio B., Pierfelici L., Campana R., Vittoria E., et al., *Campylobacter jejuni* loss of culturability in aqueous microcosms and ability to resuscitate in a mouse model, Int. J. Food Microbiol. (2006) 107:83–91.
- [4] Bej A.K., Ng W.-Y., Morgan S., Jones D.D., Mahbubani M.H., Detection of viable *Vibrio cholerae* by reverse-transcriptase polymerase chain reaction (RT-PCR), Mol. Biotechnol. (1996) 5:1–10.
- [5] Bernstein J.A., Khodursky A.B., Lin P.H., Lin-Chao S., Cohen S.N., Global analysis of mRNA decay and abundance in *Escherichia coli* at single-gene resolution using two-color fluorescent DNA microarrays, Proc. Natl. Acad. Sci. USA (2002) 99:9697–9702.
- [6] Besnard V., Federighi M., Cappelier J.M., Development of a direct viable count procedure for the investigation of VBNC state in *Listeria monocytogenes*, Lett. Appl. Microbiol. (2000) 31:77–81.
- [7] Besnard V., Federighi M., Cappelier J.-M., Evidence of viable but non-culturable state in *Listeria monocytogenes* by direct viable count and CTC-DAPI double staining, Food Microbiol. (2000) 17:697–704.
- [8] Besnard V., Federighi M., Declercq E., Jugiau F., Cappelier J.-M., Environmental and physico-chemical factors induce VBNC state in *Listeria monocytogenes*, Vet. Res. (2002) 33:359–370.
- [9] Beumer R.R., deVries J., Rombouts F.M., *Campylobacter jejuni* non-culturable coccoid cells, Int. J. Food Microbiol. (1992) 15:153–163.
- [10] Boulos L., Prevost M., Barbeau B., Coallier J., Desjardins R., LIVE/DEAD BacLight : application of a new rapid staining method for direct enumeration of viable and total bacteria in drinking water, J. Microbiol. Methods (1999) 37:77–86.
- [11] Brett M.S.Y., Short P., McLauchlin J., A small outbreak of listeriosis associated with smoked mussels, Int. J. Food Microbiol. (1998) 43:223–229.
- [12] Bunker S.T., Bates T.C., Oliver J.D., Effects of temperature on detection of plasmid or chromosomally encoded *gfp*- and *lux*-labeled *Pseudomonas fluorescens* in soil, Environ. Biosafety Res. (2004) 3:83–90.
- [13] Cappelier J.M., Besnard V., Roche S., Garrec N., Zundel E., Velge P., Federighi M., Avirulence of viable but non-culturable *Listeria monocytogenes* cells demonstrated by in vitro and in vivo models, Vet. Res. (2005) 36:589–599.
- [14] Cappelier J.-M., Besnard V., Roche S.M., Velge P., Federighi M., Avirulent viable but non culturable cells of *Listeria monocytogenes* need the presence of an embryo to be recovered in egg yolk and regain virulence after recovery, Vet. Res. (2007) 38:573–583.
- [15] Chaveerach P., ter Huurne A.A., Lipman L.J., van Knapen F., Survival and resuscitation of ten strains of *Campylobacter jejuni* and *Campylobacter coli* under acid conditions, Appl. Environ. Microbiol. (2003) 69:711–714.
- [16] Cho J.C., Kim S.J., Viable, but non-culturable, state of a green fluorescence protein-tagged environmental isolate of *Salmonella typhi* in groundwater and pond water, FEMS Microbiol. Lett. (1999) 170:257–264.
- [17] Cooray K.J., Nishibori T., Xiong H., Matsuyama T., Fujita M., Mitsuyama M., Detection of multiple virulence-associated genes of *Listeria monocytogenes* by PCR in artificially contaminated milk samples, Appl. Environ. Microbiol. (1994) 60:3023–3026.
- [18] Coutard F., Pommepuy M., Loaec S., Hervio-Heath D., mRNA detection by reverse transcription-PCR for monitoring viability and potential virulence in pathogenic strain of *Vibrio parahaemolyticus* in viable but nonculturable state, J. Appl. Microbiol. (2005) 98:951–961.
- [19] Dhiaf A., Bakhrouf A., Witzel K.P., Resuscitation of eleven-year VBNC *Citrobacter*, J. Water Health (2008) 6:565–568.
- [20] Dreux N., Albagnac C., Federighi M., Carlin F., Morris C.E., Nguyen-The C., Viable but non-culturable *Listeria monocytogenes* on parsley leaves and absence of recovery to a culturable state, J. Appl. Microbiol. (2007) 103:1272–1281.
- [21] Du M., Chen J., Zhang X., Li A., Li Y., Wang Y., Retention of virulence in a viable but nonculturable *Edwardsiella tarda* isolate, Appl. Environ. Microbiol. (2007) 73:1349–1354.
- [22] Ericsson H., Eklöv A., Danielsson-Tham M.-L., Loncarevic S., Mentzing L.-O., Persson I., et al., An outbreak of listeriosis suspected to have been caused by rainbow trout, J. Clin. Microbiol. (1997) 35:2904–2907.
- [23] Farber J.M., Daley E.M., MacKie M.T., Limerick B., A small outbreak of listeriosis potentially linked to the consumption of imitation crab meat, Lett. Appl. Microbiol. (2000) 31:100–104.
- [24] Federighi M., Tholozan J.L., Cappelier J.M., Tissier J.P., Jouve J.L., Evidence of non-coccoid viable but non-culturable *Campylobacter jejuni* cells in microcosm water by direct viable count, CTC-DAPI double staining, and scanning electron microscopy, Food Microbiol. (1998) 15:539–550.
- [25] Fischer-Le Saux M., Hervio-Heath D., Loaec S., Colwell R.R., Pommepuy M., Detection of

cytotoxin-hemolysin mRNA in nonculturable populations of environmental and clinical *Vibrio vulnificus* strains in artificial seawater, Appl. Environ. Microbiol. (2002) 68:5641–5646.

[26] Klein P.G., Juneja V.K., Sensitive detection of viable *Listeria monocytogenes* by reverse-transcription-PCR, Appl. Environ. Microbiol. (1997) 63: 4441–4448.

[27] Li J., Kolling G.L., Matthews K.R., Chikindas M.L., Cold and carbon dioxide used as multi-hurdle preservation do not induce appearance of viable but non-culturable *Listeria monocytogenes*, J. Appl. Microbiol. (2003) 94:48–53.

[28] Lleo M.M., Pierobon S., Tafi M.C., Signoretto C., Canepari P., mRNA detection by reverse transcription-PCR for monitoring viability over time in an *Enterococcus faecalis* viable but nonculturable population maintained in a laboratory microcosm, Appl. Environ. Microbiol. (2000) 66:4564–4567.

[29] Lleo M.M., Bonato B., Tafi M.C., Signoretto C., Boaretti M., Canepari P., Resuscitation rate in different enterococcal species in the viable but non-culturable state, J. Appl. Microbiol. (2001) 91:1095–1102.

[30] Miettinen M.K., Siitonen A., Heiskanen P., Haajanen H., Björkroth K.J., Korkeala H.J., Molecular epidemiology of an outbreak of febrile gastroenteritis caused by *Listeria monocytogenes* in cold-smoked rainbow trout, J. Clin. Microbiol. (1999) 37: 2358–2360.

[31] Mizunoe Y., Wai S.N., Ishikawa T., Takade A., Yoshida S., Resuscitation of viable but nonculturable cells of *Vibrio parahaemolyticus* induced at low temperature under starvation, FEMS Microbiol. Lett. (2000) 186:115–120.

[32] Murray E.G.D., Webb R.A., Swann M.B.R., A disease of rabbit characterised by a large mononuclear leucocytosis, caused by a hitherto undescribed bacillus; *Bacterium monocytogenes* (n.sp.), J. Pathol. Bacteriol. (1926) 29:407–439.

[33] Oliver J.D., Hite D., McDougald D., Andon N.L., Simpson L.M., Entry into, and resuscitation from, the viable but nonculturable state by *Vibrio vulnificus* in an estuarine environment, Appl. Environ. Microbiol. (1995) 61:2624–2630.

[34] Oliver J.D., The public health significance of viable but nonculturable bacteria, in: Colwell R.R., Grimes D.J. (Eds.), Nonculturable Microorganisms in the Environment, American Society for Microbiology Press, Washington, DC, 2000, pp. 277–299.

[35] Oliver J.D., Bockian R., In vivo resuscitation, and virulence towards mice, of viable but nonculturable cells of *Vibrio vulnificus*, Appl. Environ. Microbiol. (2005) 61:2620–2623.

[36] Oliver J.D., The viable but nonculturable state in bacteria, J. Microbiol. (2006) 43:93–100.

[37] Ordax M., Marco-Noales E., Lopez M.M., Biosca E.G., Survival strategy of *Erwinia amylovora*

against copper: induction of the viable-but-nonculturable state, Appl. Environ. Microbiol. (2006) 72:3482–3488.

[38] Pease A.J., Roa B.R., Luo W., Winkler M.E., Positive growth rate-dependent regulation of the *pdxA*, *ksgA*, and *pdxB* genes of *Escherichia coli* K-12, J. Bacteriol. (2002) 184:1359–1369.

[39] Rieu A., Weidmann S., Garmyn D., Piveteau P., Guzzo J., agr System of *Listeria monocytogenes* EGD-e: Role in adherence and differential expression pattern, Appl. Environ. Microbiol. (2007) 73:6125–6133.

[40] Roche S.M., Velge P., Bottreau E., Durier C., Marquet-van der Mee N., Pardon P., Assessment of the virulence of *Listeria monocytogenes*: agreement between a plaque-forming assay with HT-29 cells and infection of immunocompetent mice, Int. J. Food Microbiol. (2001) 68:33–44.

[41] Rollins D.M., Colwell R.R., Viable but nonculturable stage of *Campylobacter jejuni* and its role in survival in the natural aquatic environment, Appl. Environ. Microbiol. (1986) 52:531–538.

[42] Rørvik L.M., Aase B., Alvestad T., Caugant D.A., Molecular epidemiological survey of *Listeria monocytogenes* in seafoods and seafood-processing plants, Appl. Environ. Microbiol. (2000) 66: 4779–4784.

[43] Sheridan G.E.C., Masters C.I., Shallcross J.A., Mackey B.M., Detection of mRNA by reverse transcription-PCR as an indicator of viability in *Escherichia coli* cells, Appl. Environ. Microbiol. (1998) 64:1313–1318.

[44] Smith B., Oliver J.D., In situ and in vitro gene expression by *Vibrio vulnificus* during entry into, persistence within, and resuscitation from the viable but nonculturable state, Appl. Environ. Microbiol. (2006) 72:1445–1451.

[45] Steinert M., Emody L., Amann R., Hacker J., Resuscitation of viable but nonculturable *Legionella pneumophila* Philadelphia JR32 by *Acanthamoeba castellanii*, Appl. Environ. Microbiol. (1997) 63:2047–2053.

[46] Tholozan J.L., Cappelletti J.-M., Tissier J.P., Delattre G., Federighi M., Physiological characterization of viable-but-nonculturable *Campylobacter jejuni* cells, Appl. Environ. Microbiol. (1999) 65: 1110–1116.

[47] Wai S., Mizunoe Y., Takade A., Yoshida S., A comparison of solid and liquid media for resuscitation of starvation- and low-temperature-induced nonculturable cells of *Aeromonas hydrophila*, Arch. Microbiol. (2000) 173:307–310.

[48] Wong H.C., Wang P., Chen S.Y., Chiu S.W., Resuscitation of viable but non-culturable *Vibrio parahaemolyticus* in a minimum salt medium, FEMS Microbiol. Lett. (2004) 233:269–275.