

Review



OPEN ACCESS

Received: Aug 13, 2019

Revised: Oct 28, 2019

Accepted: Nov 11, 2019

Correspondence to

Hideaki Morita, MD, PhD

Department of Allergy and Clinical Immunology, National Research Institute for Child Health and Development, 2-10-1 Okura, Setagaya-ku, Tokyo 157-8535, Japan.

Tel: +81-3-3416-0181

Fax: +81-3-5494-7173

E-mail: morita-hi@ncchd.go.jp

Copyright © 2020 The Korean Academy of Asthma, Allergy and Clinical Immunology · The Korean Academy of Pediatric Allergy and Respiratory Disease

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Keisuke Orimo

<https://orcid.org/0000-0002-6496-0508>

Hirohisa Saito

<https://orcid.org/0000-0002-6630-8337>

Hideaki Morita

<https://orcid.org/0000-0003-0928-8322>

Disclosure

There are no financial or other issues that might lead to conflict of interest.

Innate Lymphoid Cells in the Airways: Their Functions and Regulators

Keisuke Orimo , Hirohisa Saito , Kenji Matsumoto, Hideaki Morita

Department of Allergy and Clinical Immunology, National Research Institute for Child Health and Development, Tokyo, Japan

ABSTRACT

Since the airways are constantly exposed to various pathogens and foreign antigens, various kinds of cells in the airways—including structural cells and immune cells—interact to form a precise defense system against pathogens and antigens that involve both innate immunity and acquired immunity. Accumulating evidence suggests that innate lymphoid cells (ILCs) play critical roles in the maintenance of tissue homeostasis, defense against pathogens and the pathogenesis of inflammatory diseases, especially at body surface mucosal sites such as the airways. ILCs are activated mainly by cytokines, lipid mediators and neuropeptides that are produced by surrounding cells, and they produce large amounts of cytokines that result in inflammation. In addition, ILCs can change their phenotype in response to stimuli from surrounding cells, which enables them to respond promptly to microenvironmental changes. ILCs exhibit substantial heterogeneity, with different phenotypes and functions depending on the organ and type of inflammation, presumably because of differences in microenvironments. Thus, ILCs may be a sensitive detector of microenvironmental changes, and analysis of their phenotype and function at local sites may enable us to better understand the microenvironment in airway diseases. In this review, we aimed to identify molecules that either positively or negatively influence the function and/or plasticity of ILCs and the sources of the molecules in the airways in order to examine the pathophysiology of airway inflammatory diseases and facilitate the issues to be solved.

Keywords: Respiratory tract diseases; innate immune response; cellular microenvironment; lymphocytes; phenotype; cytokines

INTRODUCTION

The airways are organs that are constantly exposed to airborne pathogens and foreign antigens throughout life. Therefore, various kinds of cells in the airways, including structural cells and immune cells, interact to form a precise defense system against pathogens and foreign antigens that involves both innate immunity and acquired immunity. In recent years, innate lymphoid cells (ILCs), which are widely conserved in humans and non-human vertebrates, such as the mouse and zebra fish, have attracted attention as cells involved in inflammatory diseases as well as defense against pathogens and foreign antigens.

Although ILCs have a lymphoid morphology, unlike T cells and B cells they do not have antigen receptors that recognize the microstructures of antigens; thus, they are not directly activated by specific antigens. They are activated mainly by antigen non-specific mechanisms, suggesting that they are involved in innate immunity. Some ILCs express toll-like receptors (TLRs) that recognize microbes, and the cells may be directly activated by the pathogen-associated molecular patterns of microbes. However, some recent articles have shown that ILCs express various kinds of receptors for cytokines, danger signals, neuropeptides and lipid mediators that are more dominant than TLRs. These findings suggest that ILCs may be activated by signals from other cells around it upon exposure to foreign antigens (including microbes), rather than by being directly activated by foreign antigens.

ILCs exhibit substantial heterogeneity, with different phenotypes and functions depending on the organ and type of inflammation, presumably because of differences in microenvironments. Thus, ILCs may be a sensitive detector of microenvironmental changes at our body surfaces, such as mucosal sites and the skin. Conversely, by analyzing the phenotype and function of ILCs, we can better understand the details of the microenvironment at local sites in airway diseases. In this review, we aimed to identify molecules that either positively or negatively influence the function and/or plasticity of ILCs and the sources of molecules in the airways in order to examine the pathophysiology of airway inflammatory diseases and facilitate the issues to be solved.

ILC SUBSETS AND THEIR PLASTICITY

ILCs have a lymphoid morphology, but they have no cell-surface lineage markers such as those that define other immune cell subsets. Instead, all ILCs express a cytokine receptor subunit called interleukin (IL)-7 receptor α (CD127). ILCs are currently divided into 3 different subtypes, according to their expression of cytokines and transcription factors: group 1 ILCs (ILC1s), group 2 ILCs (ILC2s) and group 3 ILCs (ILC3s). ILC1s are defined as ILCs that express T-box-expressed-in-T cells (T-bet) and produce interferon (IFN)- γ ; they include conventional natural killer cells (cNK) and are considered to be involved in anti-viral immunity, like T helper (Th)1 cells. In addition to IFN- γ , cNK cells release perforin and granzymes to induce apoptosis of target cells. ILC2s are defined as ILCs that express GATA-binding protein 3 and produce such cytokines as IL-4, IL-5, IL-9 and IL-13 as well as the epidermal growth factor, amphiregulin; like Th2 cells, they are considered to be involved in anti-helminth immunity. ILC3s are defined as ILCs that express retinoic acid receptor-related orphan receptor- γ t and produce cytokines such as IL-17A, IL-22 and granulocyte macrophage colony-stimulating factor (GM-CSF); they include both natural cytotoxicity receptor (NCR)⁻ ILC3s and NCR⁺ ILC3s, and are considered to be involved in anti-bacterial immunity, like Th17 cells.

Recently, another ILC subset called regulatory ILCs (ILCregs), which resemble regulatory T cells (Tregs) and have regulatory functions, has been reported.^{1,3} ILCregs produce regulatory cytokines such as IL-10 and/or transforming growth factor- β (TGF- β), but they do not express Forkhead box protein P3 which is the canonical transcription factor of Tregs. Therefore, it remains controversial whether ILCregs represent an independent effector subset, like ILC1s, ILC2s and ILC3s, or just a temporary state of ILCs.

ILCs are generally thought to be tissue-resident cells that differentiate into mature effector cells in tissues and show minimal movement between organs.⁴ Instead, they have functional




Fig. 1. Plasticity of ILCs in the airways. IL-4 induces conversion of ILC1s and ILC3s to ILC2s. IL-12, in conjunction with IL-1β or IL-18, induces conversion of ILC2s to ILC1s. IL-1β and IL-23, in conjunction with TGF-β, induce conversion of ILC2s to ILC3s. IL-1β and IL-23 also induce conversion of ILC1s to ILC3s in the intestines, but this has not been demonstrated in the airways. Retinoic acid, in conjunction with IL-33, induces conversion of ILC2s to ILCregs. ILC, innate lymphoid cell; ILC1, group 1 innate lymphoid cell; ILC2, group 2 innate lymphoid cell; ILC3, group 3 innate lymphoid cell; IL, interleukin; TGF-β, transforming growth factor-β; ILCreg, regulatory innate lymphoid cell; IFN, interferon; T-bet, T-box-expressed-in-T cells; RORγt, express retinoic acid receptor-related orphan receptor-γt; GATA3, GATA binding protein 3.

plasticity that enables them to respond promptly to microenvironmental changes, thereby precluding any need for differentiation and/or migration of new ILC subsets adapted to a new environment. For instance, transdifferentiation has already been shown between ILC1s and ILC3s,^{5,6} between ILC1s and ILC2s,^{7,9} and between ILC2s and ILC3s (**Fig. 1**).^{10,11}

DISTRIBUTION, HETEROGENEITY AND TISSUE-SPECIFIC SIGNATURES OF ILCs

ILCs are present in various organs throughout the body, and the proportions of the subsets differ depending on the organ. In humans, ILC3s are the predominant population in mucosal tissues, including the lung and gut,⁷ whereas the proportion of ILC2s is a little higher in the skin compared to mucosal tissues.⁷ The proportions of the ILC subsets are also influenced by age. For instance, although ILC3s (including NCR⁻ ILC3s and NCR⁺ ILC3s) are the predominant population in the fetal human lung, their proportion decreases while the proportions of ILC1s and ILC2s increase with age in the adult human lung.⁷ Another important point is that there is substantial heterogeneity in each subset of ILCs. Moreover, ILCs show different phenotypes depending on the organ.¹² For instance, although ILC2s from different organs share canonical markers such as GATA3 and IL-7R, their expressions of IL-33R, IL-25R and IL-18R1 differ depending on the organ.¹² Such heterogeneity is probably due to the high plasticity of ILCs, which enables them to adjust to their microenvironment.




Fig. 2. Regulators of ILCs in the airways and their sources. In the airways, ILCs are regulated by various stimuli that are produced by other cells. The red arrows indicate stimuli that activate ILCs, and the blue ones indicate stimuli that suppress ILCs.

ILC, innate lymphoid cell; ILC1, group 1 innate lymphoid cell; ILC2, group 2 innate lymphoid cell; ILC3, group 3 innate lymphoid cell; IL, interleukin; ILCreg, regulatory innate lymphoid cell; IFN, interferon; DC, dendritic cell; Th, T helper; Treg, regulatory T cell; SCC, solitary chemosensory cell; TSLP, thymic stromal lymphopoietin; CysLT, cysteinyl leukotriene; PGD₂, prostaglandin D₂; NMU, neuromedin U; VIP, vasoactive intestinal peptide; CGRP, calcitonin gene-related peptide; PNEC, pulmonary neuroendocrine cell.

CYTOKINES AND OTHER FACTORS ASSOCIATED WITH EACH ILC SUBSET IN THE AIRWAYS ILC1S

IL-12

IL-12 is known as a major activator of ILC1s and promotes their secretion of IFN- γ .⁶ The major physiological producers of IL-12 are antigen-presenting cells (APCs) such as dendritic cells (DCs) and macrophages. In the mouse lung, IFN- γ produced by ILC1s in response to DC-derived IL-12 during viral infection suppresses early viral growth,¹³ suggesting that the IL-12–ILC1 axis may be involved in anti-viral immunity (**Fig. 2**). Furthermore, IL-12 mediates the transdifferentiation of ILC2s⁷⁻⁹ and ILC3s⁵ into IFN- γ -producing ILC1s (**Fig. 1**), a mechanism that may be involved in immune responses to viral infections and in the pathophysiology of chronic obstructive pulmonary disease (COPD).

IL-15

Like IL-12, IL-15 activates ILC1s to produce IFN- γ . IL-15 is known to be produced by APCs, a subset of thymic epithelial cells, and by stromal cells. In the airways, human bronchial epithelial cells produce IL-15 in response to respiratory syncytial virus infection (**Fig. 2**).¹⁴ In human airway diseases, IL-15–positive cells have been reported to be increased in patients with sarcoidosis, tuberculosis or chronic bronchitis compared to asthmatic patients and healthy subjects,¹⁵ suggesting the involvement of IL-15 in the pathophysiology of these diseases.

IL-18

IL-18 was once thought to be just an activator of ILC1s that induces IFN- γ production. However, recent studies have shown that IL-18 also activates ILC2s and ILC3s to produce their signature cytokines,^{12,16} suggesting that IL-18 may be a pan-activator of ILCs (**Fig. 2**). Furthermore, IL-18 and IL-12 together promote conversion of ILC2s to ILC1s (**Fig. 1**).⁸ IL-18 is produced by APCs such as macrophages and DCs. In regard to the airways, IL-18 was shown to be released from human bronchial epithelial cells upon human rhinovirus infection¹⁷ and *Alternaria* extract stimulation¹⁸ *in vitro*. In addition, cigarette smoke exposure induced IL-18 production by alveolar macrophages in the mouse lungs (**Fig. 2**).¹⁹ In humans, the levels of IL-18 in bronchoalveolar lavage fluids (BALFs) were significantly higher in patients with COPD than in healthy subjects, and even higher in patients with acute exacerbation of COPD.²⁰ In addition, the expression of IL-18 in lung epithelial cells was significantly increased in patients with severe COPD compared to healthy individuals who never smoked.¹⁷ These findings suggest that IL-18 may be involved in the pathophysiology of COPD.

ILC2s

IL-25

IL-25 activates ILC2s and promotes type 2 cytokine production.²¹ Various kinds of immune cells, such as macrophages, eosinophils and T cells, have been shown to produce IL-25. Recently, bottle-shaped epithelial-lineage cells expressing taste receptors, named tuft cells—including intestinal tuft cells, brush cells in the lower airways and solitary chemosensory cells (SCCs) in nasopharyngeal tissue—have attracted broad attention as major sources of IL-25.²¹ In mice, intestinal tuft cells produce IL-25 after sensing microbial metabolites through succinate receptors or taste receptors during protozoan and helminth infections, which results in activation of ILC2s and promotion of an anti-helminth response.²¹ Similarly, recent findings have suggested that SCCs in the human upper respiratory tract²² and brush cells in the murine lower respiratory tract²³ are major producers of IL-25 in the airways (**Fig. 2**). In humans, IL-25 messenger RNA (mRNA) expression in bronchial epithelial cells was significantly increased in a subgroup of asthmatic patients with a Th2-high phenotype.²⁴ In addition, the levels of IL-25 were significantly increased in nasal mucosal fluid from asthmatic patients during RV infection compared to healthy individuals.²⁵ In mice, IL-25-producing brush cells were increased in the lungs of mice challenged with leukotriene (LT) E4, *Alternaria* or house dust mites, resulting in asthma-like airway inflammation.²³ Moreover, epithelial cell-derived IL-25 has been shown to be crucial in an ovalbumin (OVA)-induced asthma model,²⁶ suggesting that IL-25 may play a critical role in the development of asthma. IL-25-producing SCCs²² and ILC2s²⁷⁻²⁹ were increased in nasal polyps from chronic rhinosinusitis with nasal polyps (CRSwNP) compared to nasal tissue from healthy individuals, and the level of IL-25 correlated with infiltration of inflammatory cells and expression of inflammatory markers.³⁰ These findings suggest that the IL-25–ILC2 axis may be involved in the pathophysiology of CRSwNP. Besides allergic disorders, the concentration of IL-25 and the number of ILC2s were increased in BALF from patients with idiopathic pulmonary fibrosis and in the lungs of mice with helminth-induced lung fibrosis compared to controls,³¹ suggesting possible involvement of the IL-25–ILC2 axis in lung fibrosis as well.

IL-33

Unlike other cytokines that are newly synthesized upon stimulation and secreted via the endoplasmic reticulum/Golgi pathway, IL-33 is constitutively expressed in cells at the mucosal barrier and released from the nucleus in active form in response to tissue damage.³² Therefore, it is believed to be one of the “alarmins” that gather components of

the repair response to the sites of injury. However, several studies suggest that IL-33 may be actively secreted from live cells, including bronchial epithelial cells³³ and fibroblasts, even in the absence of necrosis (**Fig. 2**). Although the mechanisms of IL-33 secretion are not fully understood, adenosine triphosphate-induced purinoceptor-dependent activation of epithelial nicotinamide adenine dinucleotide phosphate oxidase, *i.e.*, dual oxidase 1, may be involved.³³ IL-33 is recognized as one of the major activators of ILC2s that induce production of type 2 cytokines. In mice, IL-33 is released from alveolar epithelial cells in response to tissue damage³² caused by fungi such as *Alternaria* and *Aspergillus* and viruses such as RSV and RV.³² Meanwhile, in humans, IL-33 is released from bronchial epithelial cells located more centrally,³² similar to IL-25 and thymic stromal lymphopoietin (TSLP) (**Fig. 2**).

The expression of IL-33 in the lungs peaks during infancy, and declines with age. In line with that, the number of ILC2s in the lungs also peaks in infancy.³⁴ These findings suggest that IL-33 may play a major role in the developing phase of acquired immunity and that epithelial damage may induce more severe allergic airway inflammation during infancy than during adulthood through the IL-33–ILC2s axis. Furthermore, the expression of IL-33 in the nucleus of airway epithelial cells is enhanced by exposure to atmospheric substances such as cigarette smoke³² and diesel exhaust particles.³⁵ This suggests that damage to epithelial cells after exposure to cigarette smoke and diesel exhaust particles evokes release of larger amounts of IL-33, which results in aggravation of allergic inflammation. In addition to epithelial cells, stromal cells,³⁶ endothelial cells, fibroblasts³² and platelets³⁷ may produce IL-33 (**Fig. 2**). In agreement with the finding that the IL-33 gene was strongly associated with asthma susceptibility in several genome-wide association studies,³⁸ the level of IL-33 was higher in BALF from asthmatic patients than in controls and showed an inverse correlation with forced expiratory volume in one second (FEV1), regardless of the history of steroid treatment.³⁹ The levels of IL-33 in serum⁴⁰ as well as in exhaled breath condensate were higher in COPD patients than in healthy individuals and correlated with the eosinophil counts in the peripheral blood.⁴¹ Although it remains unclear whether IL-33 is involved in the pathogenesis of COPD, these findings suggest such involvement, especially in patients with eosinophilic COPD.

TSLP

Like other epithelial-derived cytokines such as IL-33 and IL-25, TSLP is recognized as a major activator of ILC2s that induces production of type 2 cytokines. However, unlike other epithelial-derived cytokines, TSLP was shown to induce corticosteroid resistance in murine ILC2s through activation of an intracellular signaling molecule, signal transducer and activator of transcription 5.⁴² TSLP is produced by various kinds of cells including DCs, vascular endothelial cells, macrophages and mast cells. In the airways, similar to IL-25 and IL-33, TSLP is produced mainly by airway epithelial cells in response to exposure to bacteria, fungi and viruses.⁴³ Adventitial stromal cells localize with ILC2s in adventitial niche around the lung bronchi and large vessels, and support ILC2s through constitutive expression of TSLP and IL-33.³⁶ In humans, the number of type 2 cytokine-producing ILC2s was significantly increased in sputum from patients with corticosteroid-resistant severe asthma compared those with mild asthma, whereas the number of type 2 cytokine-producing Th2 cells was comparable in both groups. The levels of TSLP in BALF from patients with asthma correlate inversely with steroid-mediated inhibition of IL-5-producing ILC2s in the BALF. Furthermore, a clinical trial of an anti-TSLP antibody, tezepelumab, showed that, in moderate to severe steroid-resistant asthma, anti-TSLP treatment significantly reduced asthma exacerbation as well as important biologic markers such as the blood eosinophil count and fractional exhaled nitric oxide (FeNO).⁴⁴ These findings suggest that the TSLP–ILC2 axis

may play critical roles in steroid resistance in asthmatic patients and that TSLP is likely to be continuously produced and released in the airways of moderate to severe asthmatic patients.

IL-27

IL-27 is generally produced by DCs and macrophages (**Fig. 2**). In mice, IL-27 suppresses the proliferation and cytokine production of ILC2 cells *in vitro*,^{4,45} and it also suppresses *Alternaria*-induced eosinophilic airway inflammation by regulating ILC2 activation *in vivo*.⁴ However, in humans, IL-27 mRNA levels and the percentage of IL-27-expressing cells in BALF were increased in patients with asthma, especially those with a severe phenotype.⁴⁶ Therefore, the involvement of IL-27-ILC2s in the pathophysiology of asthma in humans remains to be elucidated.

IFNs

IFNs are divided into types 1 (α/β), 2 (γ) and 3 (λ). Type 1 and 2 IFNs have been shown to suppress type 2 cytokine production by ILC2s, both *in vitro* and *in vivo*.^{4,45} The major producers of IFN- α and - β are macrophages and DCs. IFN- γ is produced by activated Th1 cells and ILC1s, including NK cells, which are activated mainly through TLRs (**Fig. 2**). In mice, the deficiency of type 1 IFN during influenza virus and helminth infections results in severe or prolonged eosinophilic airway inflammation mediated by activated ILC2s. In humans, dozens of reports have shown impaired production of type 1 and 3 IFNs by cultured primary bronchial epithelial cells, BAL cells, peripheral blood mononuclear cells (PBMCs) and plasmacytoid DCs in response to infection with viruses such as RSV and RV in patients with asthma compared to healthy individuals.⁴⁷ Therefore, dysregulation of ILC2 activity by type 1 and 3 IFNs during viral infection in asthmatic patients may result in the development and exacerbation of allergic airway inflammation.

Lipid inflammatory mediators

Lipids are primarily involved in the formation of cell membranes of organs. However, some reports have shown that certain lipids also play crucial roles in immune responses and the maintenance of homeostasis. These lipids are called bioactive lipids or lipid mediators. Cysteinyl leukotrienes (CysLTs) as well as prostaglandin (PG) D₂ are products of arachidonic acid and were known to be major pro-inflammatory lipid mediators of allergic disorders from early days. Mast cells activated by immunoglobulin (Ig) E-crosslinking are the major source of PGD₂ in terms of quantity, but other leukocytes, including eosinophils, Th2 cells, DCs and cytokine-activated ILC2s,⁴⁸ also produce PGD₂ (**Fig. 2**). Since human ILC2s are identified as lineage-negative cells expressing chemoattractant receptor-homologous molecules on Th2 cells (CRTH2),²⁹ which is the PGD₂ receptor, PGD₂ influences ILC2s in a variety of ways, including their migration⁴⁹ and production of IL-13.⁵⁰ CysLTs are generally produced by leukocytes such as eosinophils, mast cells, macrophages and basophils. CysLTs act directly on ILC2s to enhance their ability to produce type 2 cytokines, both *in vivo* and *in vitro*.⁵⁰ On the other hand, there are some lipid molecules that inhibit ILC2 activation. PGI₂, PGE₂ and lipoxin A₄—also products of arachidonic acid—suppress ILC2s' cytokine production and proliferation, *in vitro* and *in vivo*.⁵⁰ Interestingly, however, LTE₄ and PGD₂ reportedly induce Th2 cytokines, including IL-4, synergistically in purified human peripheral blood ILC2s.⁵¹

Neuropeptides

Neuropeptides are peptides that are expressed in the nervous system and exhibit physiological activity. They are present not only in the central nervous system, but also in the nervous system of peripheral tissues such as the lungs, and they function as signal transmitters between cells. Among several neuropeptides known to act on ILC2s, vasoactive intestinal peptide (VIP) was

the first one shown to modulate ILC2 activation. VIP belongs to the glucagon/secretin family and is highly expressed in intestinal neurons, where it coordinates pancreatic secretion with smooth muscle relaxation in response to feeding. Both lung and intestinal ILC2s express VIP receptors, including VIP receptor type 1 and type 2, and VIP stimulation induces IL-5 production by the cells. The IL-5 produced in turn activates sensory neurons to produce VIP,⁵² which may exacerbate allergic airway inflammation (**Fig. 2**). Lung ILC2s also express receptors for another neuropeptide, called neuromedin U (NMU), whereas ILC1s and ILC3s do not. NMU is thought to directly activate lung ILC2s to proliferate and produce type 2 cytokines.⁵³ Calcitonin gene-related peptide (CGRP) is a calcitonin gene product, like the thyroid hormone calcitonin and it is involved in the regulation of blood calcium levels. CGRP is widely distributed in the central and peripheral nervous systems; It was also produced by non-neuronal cells in the airways—called pulmonary neuroendocrine cells (PNECs)—after OVA challenge in an OVA-sensitized mouse model (**Fig. 2**).⁵⁴ Interestingly, a recent study has shown that ILC2s are localized in close proximity to PNECs and that CGRP enhances type 2 cytokine production by lung ILC2s in the presence of IL-33 or IL-25,⁵⁴ suggesting that interaction between PNECs and ILC2s may be involved in allergic airway inflammation. Besides the neuropeptides that induce activation of ILC2s, there is also a neuropeptide that regulates activation of ILC2s. Both lung and intestinal ILC2s express the β 2-adrenergic receptor (β 2-AR), which is a receptor for epinephrine released by sympathetic nerve stimulation. Treatment with a β 2-AR agonist, salmeterol, suppressed proliferation and type 2 cytokine production by ILC2s in an IL-33-induced airway inflammation model (**Fig. 2**).⁵⁵ These findings suggest that β 2-AR agonists used as therapeutic agents for asthma may work not only as a bronchodilator, but also as a suppressor of type 2 inflammation induced by ILC2s.

Sex steroids

Sex steroids, such as estrogen and androgen, are steroid hormones that are produced mainly by the reproductive organs and modulate reproductive functions. In addition to their effects on the reproductive organs, sex steroids have recently been shown to have effects on immune cells, including ILC2s in peripheral tissues. Androgen receptors are expressed on lung ILC2s⁵⁶ as well as ILC2 progenitors (ILC2Ps) in bone marrow (BM),⁵⁷ whereas estrogen receptors are expressed on lung ILC2s and uterine ILC2s,⁵⁸ but not on ILC2Ps in BM.⁵⁷ These findings indicate that androgens may influence both the development of ILC2s in BM and the activation of ILC2s in peripheral tissues, whereas estrogens may influence mainly ILC2s in peripheral tissues. Androgens and estrogens are thought to exert opposite effects on ILC2s. Androgen signaling inhibits differentiation of ILC2Ps into ILC2s⁵⁷ and also activation of ILC2s.^{56,57} In contrast, estrogen has been suggested to have supportive effects on ILC2s (**Fig. 2**).⁵⁸ Indeed, the numbers of lung ILC2s⁵⁷ and BM ILC2Ps are significantly lower in adult male mice than in adult female mice in the steady state.⁵⁷ In addition, the number of ILC2s in the peripheral blood of adult male patients with severe asthma,⁵⁶ as well as their number in the lungs of *Alternaria*-exposed adult male mice,⁵⁶ was significantly lower than in females. In humans, although males are more susceptible to asthma than females before puberty, the asthma incidence reverses after puberty.⁵⁹ These findings suggest that ILC2s are involved in the sex difference in asthma prevalence in humans.

ILC3s

IL-23

IL-23 is a major activator of ILC3s that induces production of inflammatory cytokines such as IL-17 and IL-22. IL-23 also induces conversion of ILC1s to ILC3s in conjunction with IL-1 β and retinoic acid,⁵ and ILC2s to ILC3s in conjunction with IL-1 β and TGF- β (**Fig. 1**).^{10,11} IL-

23 is generally produced by DCs and macrophages, but it was also produced by bronchial epithelial cells in a house dust mite-induced asthma model (**Fig. 2**).⁶⁰ The serum level of IL-23 was higher in asthmatic children than in healthy children, which shows a strong inverse correlation with FEV1.⁶¹ In addition, IL-23-positive cells were increased in the airway mucosa of stable COPD patients.⁶² However, the roles of IL-23 in the pathogenesis of asthma and COPD remain unclear.

IL-1 β

IL-1 β is a major activator of ILC3s that induces IL-17A production.⁶³ While IL-1 β is a potent activator of ILC2s that induce type 2 cytokine production,⁷ it also induces conversion of ILC2s to ILC1s together with IL-12,^{7,9} and to ILC3s together with IL-23 and TGF- β (**Fig. 1**).¹¹ In the airways, IL-1 β is produced by DCs in response to exposure to chitin and IL-33,⁶⁴ and by nasal epithelial cells exposed to *Staphylococcus aureus* or *Pseudomonas aeruginosa* (**Fig. 2**).¹¹ The levels of IL-1 β in BALF and expression of mRNA for IL-1 β and NLRP3 in sputum cells⁶⁵ were significantly higher in steroid-resistant severe asthma patients with neutrophilic inflammation than in healthy subjects and mild asthmatic patients. In mice, steroid-resistant severe asthma-like airway inflammation was attenuated by inhibiting IL-1 β or inflammasome activation.⁶⁵ These findings suggest that IL-1 β is involved in the pathogenesis of steroid-resistant severe asthma. In contrast, the levels of IL-1 β in BALF and IL-1 β mRNA expression in the bronchial mucosa were comparable between severe COPD patients and control subjects.⁶⁶ In addition, a clinical trial of anti-IL-1R1 monoclonal antibody (mAb) showed no improvement in the QOL score or acute exacerbation rate in COPD patients.⁶⁷ Therefore, the question of involvement of IL-1 β in the pathogenesis of COPD should be further studied.

Vitamins

Retinoic acid (RA)—which is a metabolite of vitamin A (Vit A)—and vitamin D (Vit D) are known to be molecules that regulate ILCs.^{1,3,5,11,68} RA is synthesized from a Vit A metabolite, retinal, by cells having enzymes such as retinaldehyde dehydrogenase (ALDH)1A1, ALDH1A2 and ALDH1A3. RA is generally synthesized by CD103⁺ DCs, intestinal epithelial cells and lamina propria stromal cells in the gut that express ALDHs. In the airways, bronchial epithelial cells express ALDHs in response to IL-13 stimulation,¹ suggesting that these cells could be the source of RA during allergic airway inflammation. Vit D can be absorbed by oral intake, but it is synthesized mainly in the skin upon exposure to ultraviolet light from the sun. RA enhances activation of ILC3s by IL-1 β and IL-23 to increase production of IL-22 (**Fig. 2**), and it also induces conversion of ILC1s to ILC3s in conjunction with IL-1 β and IL-23 (**Fig. 1**).⁵ In addition, RA inhibits development of ILC2s from ILC2Ps in mouse BM⁶⁹ and induces conversion of ILC2s to IL-10–producing ILCregs in both humans and mice (**Fig. 1**).^{1,3} In contrast to the positive effects of RA on ILC3s, Vit D suppresses production of cytokines such as IL-22, IL-17F and GM-CSF by ILC3s by down-regulating the IL-23/IL-23R pathway,⁶⁸ and it also prevents IL-1 β -, IL-23- and TGF- β -induced conversion of ILC2s to ILC3s (**Fig. 1**).¹¹

Involvement of ILCs in airway inflammatory diseases

Asthma

Asthma is a chronic airway inflammatory disease characterized by wheezing, prolonged exhalation and dyspnea due to reversible airflow obstruction. It is considered to be a heterogeneous disease comprising a number of subtypes.⁷⁰ ILC2s might be involved in the pathogenesis of asthma in patients who have severe eosinophilic inflammation induced by type 2 cytokines. Indeed, the frequencies of ILC2s in BALFs, sputum and PBMCs were significantly increased in asthmatic patients, especially in uncontrolled patients, compared

to healthy individuals, and correlated negatively with airway function.⁷¹ ILC2s may be involved in the pathogenesis of asthma via 2 different mechanisms: antigen non-specific and antigen specific. The former involves direct activation of ILC2s by mediators such as IL-33, TSLP, IL-25 and neuropeptides that are produced by tissue structural cells in response to microenvironmental changes such as viral infection, allergen exposure and stress. The latter involves migration and activation of ILC2s by lipid mediators such as PGD₂ and CysLTs, which are synthesized by mast cells activated upon antigen exposure in an IgE-dependent manner. The number of ILC2s in BALFs of asthmatic patients is increased after segmental allergen challenge, whereas the number of ILC2s in the blood is decreased.⁴⁹ Moreover, the levels of PGD₂ in BALFs correlated with decreased ILC2s in the blood.⁴⁹ Although ILCs are recognized as tissue-resident cells that show minimal migration between organs in mice, these findings suggest that, at least in humans, some ILC2s dynamically migrate into the airways from the peripheral blood in response to PGD₂ that is synthesized by mast cells upon allergen exposure.

As the importance of ILC2s and its activators in the pathogenesis of asthma becomes apparent, new therapeutic strategies—including the use of biologics to suppress their activity—are attracting attention. Treatment with an anti-TSLP mAb, tezepelumab, was shown to reduce bronchoconstriction and airway inflammation induced by allergen challenge in mild asthmatic patients.⁷² That mAb also significantly reduced the annualized rate of asthma exacerbations in steroid-resistant severe asthmatic patients treated with long-acting β -agonists and medium-to-high doses of inhaled glucocorticoids in a phase 2 clinical trial.⁴⁴ Of note, biomarkers such as the blood eosinophil count and FeNO were significantly decreased as early as 4 weeks after treatment with tezepelumab,⁴⁴ suggesting that TSLP is continuously synthesized and released in the airways of asthmatic patients. Since TSLP is a key molecule inducing steroid resistance of ILC2s⁴² as described in the previous section, treatment with tezepelumab may be the most logical strategy to treat steroid-resistant severe asthmatic patients. Another biologic directed at an epithelial cell-derived cytokine, anti-IL-33 antibody, is currently undergoing a phase 2a clinical trial in asthmatic patients being treated with high-dose inhaled corticosteroid and long-acting β 2-agonist. Fevipiprant, a CRTH2 antagonist that targets CRTH2 on ILC2s as well as on eosinophils, basophils and Th2 cells, significantly reduced eosinophilic inflammation in the sputum and bronchial mucosa and improved the lung function and clinical outcome in week 12 in moderate to severe eosinophilic asthmatic patients.⁷³ However, 2 studies using different CRTH2 antagonists, AZD 1981⁷⁴ and OC000459,⁷⁵ showed no significant improvement in lung function in moderate, persistent asthma patients in week 4. Therefore, further studies are needed to determine the patient phenotypes likely to benefit from use of a CRTH2 agonist, and the duration of treatment. In contrast to the involvement of ILC2s in eosinophilic asthma, ILC3s may be involved in non-allergic asthma, including obesity-related asthma. In mice, airway hyperresponsiveness (AHR) was induced in high-fat-diet-fed obese mice even in the absence of antigen challenge compared to normal-diet-fed mice.⁶³ AHR was induced by IL-17-producing ILC3s activated by IL-1 β produced upon obesity-related inflammasome activation.⁶³ However, in humans, the details of any involvement of ILC3s in non-eosinophilic asthma remain unclear.

Allergic rhinitis (AR)

AR is an allergic disorder of the nasal mucosa that is characterized by paroxysmal recurrent sneezing, serous rhinorrhea and nasal congestion. AR is thought to be induced mainly by an immediate hypersensitivity reaction that involves IgE-mediated release of histamine

and other mediators from mast cells in the nose. However, a delayed allergic reaction accompanied by production of type 2 cytokines that is triggered by inhalation of a sensitizing antigen may also be involved in the development of the symptoms and pathology of AR. It is thus assumed that ILC2s may also be involved via type 2 cytokine production. Indeed, the proportion of ILC2s in the peripheral blood of patients with seasonal AR is significantly higher in the pollen season than out of season.⁷⁶ In addition, ILC2s sorted from the peripheral blood of AR patients produce larger amounts of type 2 cytokines than those from healthy individuals.⁷⁷ These findings suggest that ILC2s in patients with AR may be activated upon allergen exposure. Like in asthma, ILC2s may be involved in the pathogenesis of AR via 2 different mechanisms: antigen non-specific and specific. The former involves direct activation of ILC2s by mediators such as TSLP and IL-25, which are increased in the nasal lavage of patients with AR. The latter involves migration and activation of ILC2s in response to lipid mediators such as PGD₂ and CysLTs, synthesized by mast cells activated in an IgE-dependent manner upon antigen exposure. The proportion of ILC2s in the peripheral blood of patients who underwent allergen-specific immunotherapy did not increase even during the pollen season.⁷⁶ That may be due to desensitization of mast cells by the immunotherapy, resulting in decreased release of PGD₂ and CysLTs by the mast cells.

Chronic rhinosinusitis (CRS)

CRS is a heterogeneous disease characterized by chronic inflammation of the sinus and nasal mucosa, with nasal congestion and a decreased sense of smell. CRS can be roughly divided into 2 major subtypes based on the presence/absence of nasal polyps, and each subset shows different pathological conditions.⁷⁸ In particular, CRSwNP is generally considered to involve mainly eosinophilic inflammation caused by typical type 2 cytokines, with high expression of IL-5 and IL-13 in polyps. The percentage of ILC2s is significantly higher in nasal polyps from patients with CRSwNP than in nasal tissues from controls and those with chronic rhinosinusitis without nasal polyps (CRSsNP).²⁷⁻²⁹ The percentage correlates with the nasal symptom score.²⁷ Furthermore, unlike ILC2s in the peripheral blood, ILC2s sorted from nasal polyps of CRSwNP spontaneously produce type 2 cytokines such as IL-5 and IL-13, even without additional stimulation.²⁸ These findings indicate that ILC2s in CRSwNP patients are continuously exposed to activating signals in the nasal cavity.

Although factors that continuously activate ILC2s in nasal polyps remain unclear, recent findings highlighted a substantial role of IL-25 in CRSwNP, especially in polyps having high eosinophil infiltration.^{22,30} The expression of IL-25³⁰ and the number of IL-25-producing SCCs²² were significantly higher in nasal polyps from CRSwNP patients than in nasal tissue from controls and CRSsNP patients. In addition, production of IL-25 by SCCs in nasal polyps was increased in response to stimulation with IL-13,²² suggesting that a vicious loop of IL-25-ILC2s-IL-13 may be present. Recent studies have shown that the pathophysiology of CRSwNP may differ depending on the race, implying the existence of subtypes in CRSwNP.⁷⁸ Indeed, it has been shown that CRSwNP patients can be divided into 2 subtypes depending on the expression level of IL-25 in nasal tissues. The IL-25-high subtype had a type 2-prone phenotype accompanied by high eosinophil infiltration and steroid sensitivity.⁷⁹ TSLP was also constantly up-regulated in nasal tissue from CRSwNP patients compared to healthy subjects.⁸⁰ In contrast, there is still debate regarding the expression level of IL-33 in nasal polyps.^{80,81} In regard to treatment, a recent case report demonstrated that an IL-4R mAb (dupilumab) significantly improved CRSwNP disease outcomes, along with asthma control and lung function, in patients with CRSwNP and aspirin-exacerbated respiratory disease.⁸² A clinical phase 2 trial of ANB020, an anti-IL-33 antibody, is currently ongoing.

COPD

COPD is a chronic airway inflammatory disease characterized by dyspnea due to irreversible airway obstruction. Cigarette smoking is considered to be the primary cause of COPD, but continuous inhalation of air pollutants and occupational dust may also be involved. COPD is thought to be a heterogeneous disease that includes patients with chronic bronchitis, emphysema and/or asthma components. Although the pathogenesis of COPD may also be heterogeneous, type 1 and 3 immunity accompanied by neutrophil and macrophage infiltration is thought to be involved.⁸³ Indeed, the frequency of ILC1s in the peripheral blood is increased in COPD patients and correlates inversely with lung function.⁸ In addition, the frequencies of ILC1s and NCR⁺ ILC3s were increased in lung tissue of COPD patients compared to control subjects, while the frequency of ILC2s was decreased.⁷ Given that the total ILC numbers are comparable between healthy individuals and COPD patients, the increase in ILC1 frequency and decrease in ILC2 frequency have been proposed to be caused by conversion of ILC2s to ILC1s in the lung rather than influx of ILCs from other organs.⁷ With regard to ILC3s, a subset of ILC3s that express neuropilin 1 was increased in the lung tissue of COPD patients, and these cells were suggested to be involved in ectopic lymphocyte accumulation and angiogenesis.⁸⁴ In line with these findings in humans, the numbers of ILC1s and ILC3s were reportedly increased in lungs from a cigarette smoke-induced COPD mouse model.⁸⁵ These findings suggest that ILC1s and ILC3s, but not ILC2s, are involved in the pathophysiology of COPD. However, as mentioned above, COPD is thought to be a heterogeneous disease, and accumulating evidence suggests the existence of subtypes involving type 2 immunity. For instance, it has been shown that some COPD patients have eosinophilic inflammation during acute exacerbation.⁸⁶ Furthermore, some patients have features of both asthma and COPD, called asthma-COPD overlap, with increased FeNO and eosinophils in the peripheral blood.⁸⁷ Although type 2 immunity may be involved in the pathophysiology of these patients, any possible involvement of ILC2s in it remains unclear.

CONCLUSIONS

Many articles regarding the functions and regulators of ILC2s have been published in recent years, resulting in those features being relatively well characterized. On the other hand, articles regarding ILC1s and ILC3s have been scanty in number for characterizing their functions and regulators. One of the reasons why analysis of ILC2s is advanced compared to the other ILC subsets may be that ILC2 activation leads to a clear outcome: eosinophil infiltration of tissues. Indeed, many of the studies showing involvement of ILC2s consist of patients with severe eosinophil infiltration. For such patients, treatments that target such upstream regulators of ILC2s as TSLP, IL-25 and IL-33, and downstream effector molecules of ILC2s, such as IL-4, IL-5 and IL-13, can be thought to have promise. However, in clinical practice, there are many cases that involve not only pure eosinophil infiltration but also neutrophil infiltration, and such “mixed-type” patients often exhibit more severe and treatment-resistant phenotypes. Since not only ILC2s but also ILC1s and ILC3s may be involved in the pathophysiology of such cases, we need to further elucidate functions and molecules that modulate activation of ILC1s and ILC3s in the airways. Recently, IL-17-producing ILC2s expressing the canonical ILC2 surface marker ST2 were reportedly induced in the lungs of papain- or IL-33-treated mice,⁸⁸ suggesting the existence of cells intermediate between 2 different ILC subsets *in vivo*. Conventional analysis using canonical surface markers and cytokines would not be enough to clarify the characteristics and functions of

ILCs involved in the pathophysiology of “mixed-type” airway inflammatory disease. Analysis of airway ILCs at the single-cell levels using a recently developed technology, single cell RNA-seq may help determine the ILC phenotype and local microenvironment in individual cases.

ACKNOWLEDGMENTS

We thank Mr. Lawrence W. Stiver (Tokyo, Japan; Quality Translation Co., Ltd.; qualityt@gol.com) for editing our manuscript. This work was supported in part by a Grants-in-Aid for Young Scientists (B) (to H.M.), a Grant of National Center for Child Health and Development (#29-2; to H.M.) and a Grant from Nipponham Foundation for the Future of Food (to H.M.).

REFERENCES

1. Morita H, Kubo T, Rückert B, Ravindran A, Soyka MB, Rinaldi AO, et al. Induction of human regulatory innate lymphoid cells from group 2 innate lymphoid cells by retinoic acid. *J Allergy Clin Immunol* 2019;143:2190-2201.e9.
[PUBMED](#) | [CROSSREF](#)
2. Wang S, Xia P, Chen Y, Qu Y, Xiong Z, Ye B, et al. Regulatory innate lymphoid cells control innate intestinal inflammation. *Cell* 2017;171:201-216.e18.
[PUBMED](#) | [CROSSREF](#)
3. Seehus CR, Kadavallore A, Torre B, Yeckes AR, Wang Y, Tang J, et al. Alternative activation generates IL-10 producing type 2 innate lymphoid cells. *Nat Commun* 2017;8:1900.
[PUBMED](#) | [CROSSREF](#)
4. Moro K, Kabata H, Tanabe M, Koga S, Takeno N, Mochizuki M, et al. Interferon and IL-27 antagonize the function of group 2 innate lymphoid cells and type 2 innate immune responses. *Nat Immunol* 2016;17:76-86.
[PUBMED](#) | [CROSSREF](#)
5. Bernink JH, Krabbendam L, Germar K, de Jong E, Gronke K, Kofoed-Nielsen M, et al. Interleukin-12 and -23 control plasticity of CD127(+) group 1 and group 3 innate lymphoid cells in the intestinal lamina propria. *Immunity* 2015;43:146-60.
[PUBMED](#) | [CROSSREF](#)
6. Bernink JH, Peters CP, Munneke M, te Velde AA, Meijer SL, Weijer K, et al. Human type 1 innate lymphoid cells accumulate in inflamed mucosal tissues. *Nat Immunol* 2013;14:221-9.
[PUBMED](#) | [CROSSREF](#)
7. Bal SM, Bernink JH, Nagasawa M, Groot J, Shikhagaie MM, Golebski K, et al. IL-1 β , IL-4 and IL-12 control the fate of group 2 innate lymphoid cells in human airway inflammation in the lungs. *Nat Immunol* 2016;17:636-45.
[PUBMED](#) | [CROSSREF](#)
8. Silver JS, Kearley J, Copenhaver AM, Sanden C, Mori M, Yu L, et al. Inflammatory triggers associated with exacerbations of COPD orchestrate plasticity of group 2 innate lymphoid cells in the lungs. *Nat Immunol* 2016;17:626-35.
[PUBMED](#) | [CROSSREF](#)
9. Ohne Y, Silver JS, Thompson-Snipes L, Collet MA, Blanck JP, Cantarel BL, et al. IL-1 is a critical regulator of group 2 innate lymphoid cell function and plasticity. *Nat Immunol* 2016;17:646-55.
[PUBMED](#) | [CROSSREF](#)
10. Bernink JH, Ohne Y, Teunissen MB, Wang J, Wu J, Krabbendam L, et al. c-Kit-positive ILC2s exhibit an ILC3-like signature that may contribute to IL-17-mediated pathologies. *Nat Immunol* 2019;20:992-1003.
[PUBMED](#) | [CROSSREF](#)
11. Golebski K, Ros XR, Nagasawa M, van Tol S, Heesters BA, Aglmous H, et al. IL-1 β , IL-23, and TGF- β drive plasticity of human ILC2s towards IL-17-producing ILCs in nasal inflammation. *Nat Commun* 2019;10:2162.
[PUBMED](#) | [CROSSREF](#)
12. Ricardo-Gonzalez RR, Van Dyken SJ, Schneider C, Lee J, Nussbaum JC, Liang HE, et al. Tissue signals imprint ILC2 identity with anticipatory function. *Nat Immunol* 2018;19:1093-9.
[PUBMED](#) | [CROSSREF](#)

13. Weizman OE, Adams NM, Schuster IS, Krishna C, Pritykin Y, Lau C, et al. ILC1 confer early host protection at initial sites of viral infection. *Cell* 2017;171:795-808.e12.
[PUBMED](#) | [CROSSREF](#)
14. Zdrengeha MT, Telcian AG, Laza-Stanca V, Bellettato CM, Edwards MR, Nikonova A, et al. RSV infection modulates IL-15 production and MICA levels in respiratory epithelial cells. *Eur Respir J* 2012;39:712-20.
[PUBMED](#) | [CROSSREF](#)
15. Muro S, Taha R, Tscopoulos A, Olivenstein R, Tonnel AB, Christodoulopoulos P, et al. Expression of IL-15 in inflammatory pulmonary diseases. *J Allergy Clin Immunol* 2001;108:970-5.
[PUBMED](#) | [CROSSREF](#)
16. Simoni Y, Fehlings M, Kløverpris HN, McGovern N, Koo SL, Loh CY, et al. Human innate lymphoid cell subsets possess tissue-type based heterogeneity in phenotype and frequency. *Immunity* 2017;46:148-61.
[PUBMED](#) | [CROSSREF](#)
17. Briend E, Ferguson GJ, Mori M, Damera G, Stephenson K, Karp NA, et al. IL-18 associated with lung lymphoid aggregates drives IFN γ production in severe COPD. *Respir Res* 2017;18:159.
[PUBMED](#) | [CROSSREF](#)
18. Murai H, Okazaki S, Hayashi H, Kawakita A, Hosoki K, Yasutomi M, et al. *Alternaria* extract activates autophagy that induces IL-18 release from airway epithelial cells. *Biochem Biophys Res Commun* 2015;464:969-74.
[PUBMED](#) | [CROSSREF](#)
19. Kang MJ, Homer RJ, Gallo A, Lee CG, Crothers KA, Cho SJ, et al. IL-18 is induced and IL-18 receptor alpha plays a critical role in the pathogenesis of cigarette smoke-induced pulmonary emphysema and inflammation. *J Immunol* 2007;178:1948-59.
[PUBMED](#) | [CROSSREF](#)
20. Wang H, Lv C, Wang S, Ying H, Weng Y, Yu W. NLRP3 inflammasome involves in the acute exacerbation of patients with chronic obstructive pulmonary disease. *Inflammation* 2018;41:1321-33.
[PUBMED](#) | [CROSSREF](#)
21. Schneider C, O'Leary CE, Locksley RM. Regulation of immune responses by tuft cells. *Nat Rev Immunol* 2019;19:584-93.
[PUBMED](#) | [CROSSREF](#)
22. Kohanski MA, Workman AD, Patel NN, Hung LY, Shtraks JP, Chen B, et al. Solitary chemosensory cells are a primary epithelial source of IL-25 in patients with chronic rhinosinusitis with nasal polyps. *J Allergy Clin Immunol* 2018;142:460-469.e7.
[PUBMED](#) | [CROSSREF](#)
23. Bankova LG, Dwyer DF, Yoshimoto E, Ualiyeva S, McGinty JW, Raff H, et al. The cysteinyl leukotriene 3 receptor regulates expansion of IL-25-producing airway brush cells leading to type 2 inflammation. *Sci Immunol* 2018;3:eaat9453.
[PUBMED](#) | [CROSSREF](#)
24. Cheng D, Xue Z, Yi L, Shi H, Zhang K, Huo X, et al. Epithelial interleukin-25 is a key mediator in Th2-high, corticosteroid-responsive asthma. *Am J Respir Crit Care Med* 2014;190:639-48.
[PUBMED](#) | [CROSSREF](#)
25. Beale J, Jayaraman A, Jackson DJ, Macintyre JD, Edwards MR, Walton RP, et al. Rhinovirus-induced IL-25 in asthma exacerbation drives type 2 immunity and allergic pulmonary inflammation. *Sci Transl Med* 2014;6:256ra134.
[PUBMED](#) | [CROSSREF](#)
26. Suzukawa M, Morita H, Nambu A, Arae K, Shimura E, Shibui A, et al. Epithelial cell-derived IL-25, but not Th17 cell-derived IL-17 or IL-17F, is crucial for murine asthma. *J Immunol* 2012;189:3641-52.
[PUBMED](#) | [CROSSREF](#)
27. Ho J, Bailey M, Zaunders J, Mrad N, Sacks R, Sewell W, et al. Group 2 innate lymphoid cells (ILC2s) are increased in chronic rhinosinusitis with nasal polyps or eosinophilia. *Clin Exp Allergy* 2015;45:394-403.
[PUBMED](#) | [CROSSREF](#)
28. Puposki JA, Klingler AI, Tan BK, Soroosh P, Banie H, Lewis G, et al. Group 2 innate lymphoid cells are elevated and activated in chronic rhinosinusitis with nasal polyps. *Immun Inflamm Dis* 2017;5:233-43.
[PUBMED](#) | [CROSSREF](#)
29. Mjösberg JM, Trifari S, Crellin NK, Peters CP, van Drunen CM, Piet B, et al. Human IL-25- and IL-33-responsive type 2 innate lymphoid cells are defined by expression of CCR2 and CD161. *Nat Immunol* 2011;12:1055-62.
[PUBMED](#) | [CROSSREF](#)
30. Shin HW, Kim DK, Park MH, Eun KM, Lee M, So D, et al. IL-25 as a novel therapeutic target in nasal polyps of patients with chronic rhinosinusitis. *J Allergy Clin Immunol* 2015;135:1476-1485.e7.
[PUBMED](#) | [CROSSREF](#)

31. Hams E, Armstrong ME, Barlow JL, Saunders SP, Schwartz C, Cooke G, et al. IL-25 and type 2 innate lymphoid cells induce pulmonary fibrosis. *Proc Natl Acad Sci U S A* 2014;111:367-72.
[PUBMED](#) | [CROSSREF](#)
32. Cayrol C, Girard JP. Interleukin-33 (IL-33): a nuclear cytokine from the IL-1 family. *Immunol Rev* 2018;281:154-68.
[PUBMED](#) | [CROSSREF](#)
33. Hristova M, Habibovic A, Veith C, Janssen-Heininger YM, Dixon AE, Geiszt M, et al. Airway epithelial dual oxidase 1 mediates allergen-induced IL-33 secretion and activation of type 2 immune responses. *J Allergy Clin Immunol* 2016;137:1545-1556.e11.
[PUBMED](#) | [CROSSREF](#)
34. de Kleer IM, Kool M, de Bruijn MJ, Willart M, van Moorlegghem J, Schuijs MJ, et al. Perinatal activation of the interleukin-33 pathway promotes type 2 immunity in the developing lung. *Immunity* 2016;45:1285-98.
[PUBMED](#) | [CROSSREF](#)
35. Weng CM, Wang CH, Lee MJ, He JR, Huang HY, Chao MW, et al. Aryl hydrocarbon receptor activation by diesel exhaust particles mediates epithelium-derived cytokines expression in severe allergic asthma. *Allergy* 2018;73:2192-204.
[PUBMED](#) | [CROSSREF](#)
36. Dahlgren MW, Jones SW, Cautivo KM, Dubinin A, Ortiz-Carpena JF, Farhat S, et al. Adventitial stromal cells define group 2 innate lymphoid cell tissue niches. *Immunity* 2019;50:707-722.e6.
[PUBMED](#) | [CROSSREF](#)
37. Takeda T, Unno H, Morita H, Futamura K, Emi-Sugie M, Arae K, et al. Platelets constitutively express IL-33 protein and modulate eosinophilic airway inflammation. *J Allergy Clin Immunol* 2016;138:1395-1403.e6.
[PUBMED](#) | [CROSSREF](#)
38. Kim KW, Ober C. Lessons learned from GWAS of asthma. *Allergy Asthma Immunol Res* 2019;11:170-87.
[PUBMED](#) | [CROSSREF](#)
39. Li Y, Wang W, Lv Z, Li Y, Chen Y, Huang K, et al. Elevated expression of IL-33 and TSLP in the airways of human asthmatics *in vivo*: a potential biomarker of severe refractory disease. *J Immunol* 2018;200:2253-62.
[PUBMED](#) | [CROSSREF](#)
40. Xia J, Zhao J, Shang J, Li M, Zeng Z, Zhao J, et al. Increased IL-33 expression in chronic obstructive pulmonary disease. *Am J Physiol Lung Cell Mol Physiol* 2015;308:L619-27.
[PUBMED](#) | [CROSSREF](#)
41. Tworek D, Majewski S, Szewczyk K, Kiszalkiewicz J, Kurmanowska Z, Górski P, et al. The association between airway eosinophilic inflammation and IL-33 in stable non-atopic COPD. *Respir Res* 2018;19:108.
[PUBMED](#) | [CROSSREF](#)
42. Kabata H, Moro K, Fukunaga K, Suzuki Y, Miyata J, Masaki K, et al. Thymic stromal lymphopoietin induces corticosteroid resistance in natural helper cells during airway inflammation. *Nat Commun* 2013;4:2675.
[PUBMED](#) | [CROSSREF](#)
43. Varricchi G, Pecoraro A, Marone G, Criscuolo G, Spadaro G, Genovese A, et al. Thymic stromal lymphopoietin isoforms, inflammatory disorders, and cancer. *Front Immunol* 2018;9:1595.
[PUBMED](#) | [CROSSREF](#)
44. Corren J, Parnes JR, Wang L, Mo M, Roseti SL, Griffiths JM, et al. Tezepelumab in adults with uncontrolled asthma. *N Engl J Med* 2017;377:936-46.
[PUBMED](#) | [CROSSREF](#)
45. Duerr CU, McCarthy CD, Mindt BC, Rubio M, Meli AP, Pothlichet J, et al. Type I interferon restricts type 2 immunopathology through the regulation of group 2 innate lymphoid cells. *Nat Immunol* 2016;17:65-75.
[PUBMED](#) | [CROSSREF](#)
46. Xie M, Mustovich AT, Jiang Y, Trudeau JB, Ray A, Ray P, et al. IL-27 and type 2 immunity in asthmatic patients: association with severity, CXCL9, and signal transducer and activator of transcription signaling. *J Allergy Clin Immunol* 2015;135:386-94.
[PUBMED](#) | [CROSSREF](#)
47. Edwards MR, Strong K, Cameron A, Walton RP, Jackson DJ, Johnston SL. Viral infections in allergy and immunology: how allergic inflammation influences viral infections and illness. *J Allergy Clin Immunol* 2017;140:909-20.
[PUBMED](#) | [CROSSREF](#)
48. Maric J, Ravindran A, Mazzurana L, Van Acker A, Rao A, Kokkinou E, et al. Cytokine-induced endogenous production of prostaglandin D₂ is essential for human group 2 innate lymphoid cell activation. *J Allergy Clin Immunol* 2019;143:2202-2214.e5.
[PUBMED](#) | [CROSSREF](#)

49. Winkler C, Hochdörfer T, Israelsson E, Hasselberg A, Cavallin A, Thörn K, et al. Activation of group 2 innate lymphoid cells after allergen challenge in asthmatic patients. *J Allergy Clin Immunol* 2019;144:61-69.e7.
[PUBMED](#) | [CROSSREF](#)
50. Doherty TA, Broide DH. Lipid regulation of group 2 innate lymphoid cell function: moving beyond epithelial cytokines. *J Allergy Clin Immunol* 2018;141:1587-9.
[PUBMED](#) | [CROSSREF](#)
51. Salimi M, Stöger L, Liu W, Go S, Pavord I, Klenerman P, et al. Cysteinyl leukotriene E₄ activates human group 2 innate lymphoid cells and enhances the effect of prostaglandin D₂ and epithelial cytokines. *J Allergy Clin Immunol* 2017;140:1090-1100.e11.
[PUBMED](#) | [CROSSREF](#)
52. Talbot S, Abdunour RE, Burkett PR, Lee S, Cronin SJ, Pascal MA, et al. Silencing nociceptor neurons reduces allergic airway inflammation. *Neuron* 2015;87:341-54.
[PUBMED](#) | [CROSSREF](#)
53. Wallrapp A, Riesenfeld SJ, Burkett PR, Abdunour RE, Nyman J, Dionne D, et al. The neuropeptide NMU amplifies ILC2-driven allergic lung inflammation. *Nature* 2017;549:351-6.
[PUBMED](#) | [CROSSREF](#)
54. Sui P, Wiesner DL, Xu J, Zhang Y, Lee J, Van Dyken S, et al. Pulmonary neuroendocrine cells amplify allergic asthma responses. *Science* 2018;360:eaan8546.
[PUBMED](#) | [CROSSREF](#)
55. Moriyama S, Brestoff JR, Flamar AL, Moeller JB, Klose CS, Rankin LC, et al. β_2 -adrenergic receptor-mediated negative regulation of group 2 innate lymphoid cell responses. *Science* 2018;359:1056-61.
[PUBMED](#) | [CROSSREF](#)
56. Cephus JY, Stier MT, Fuseini H, Yung JA, Toki S, Bloodworth MH, et al. Testosterone attenuates group 2 innate lymphoid cell-mediated airway inflammation. *Cell Reports* 2017;21:2487-99.
[PUBMED](#) | [CROSSREF](#)
57. Laffont S, Blanquart E, Savignac M, Cénac C, Laverny G, Metzger D, et al. Androgen signaling negatively controls group 2 innate lymphoid cells. *J Exp Med* 2017;214:1581-92.
[PUBMED](#) | [CROSSREF](#)
58. Bartemes K, Chen CC, Iijima K, Drake L, Kita H. IL-33-responsive group 2 innate lymphoid cells are regulated by female sex hormones in the uterus. *J Immunol* 2018;200:229-36.
[PUBMED](#) | [CROSSREF](#)
59. Almqvist C, Worm M, Leynaert B; working group of GA2LEN WP 2.5 Gender. Impact of gender on asthma in childhood and adolescence: a GA2LEN review. *Allergy* 2008;63:47-57.
[PUBMED](#)
60. Lee HS, Park DE, Lee JW, Chang Y, Kim HY, Song WJ, et al. IL-23 secreted by bronchial epithelial cells contributes to allergic sensitization in asthma model: role of IL-23 secreted by bronchial epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2017;312:L13-21.
[PUBMED](#) | [CROSSREF](#)
61. Ciprandi G, Cuppari C, Salpietro AM, Tosca MA, Rigoli L, Grasso L, et al. Serum IL-23 strongly and inversely correlates with FEV1 in asthmatic children. *Int Arch Allergy Immunol* 2012;159:183-6.
[PUBMED](#) | [CROSSREF](#)
62. Di Stefano A, Caramori G, Gnemmi I, Contoli M, Vicari C, Capelli A, et al. T helper type 17-related cytokine expression is increased in the bronchial mucosa of stable chronic obstructive pulmonary disease patients. *Clin Exp Immunol* 2009;157:316-24.
[PUBMED](#) | [CROSSREF](#)
63. Kim HY, Lee HJ, Chang YJ, Pichavant M, Shore SA, Fitzgerald KA, et al. Interleukin-17-producing innate lymphoid cells and the NLRP3 inflammasome facilitate obesity-associated airway hyperreactivity. *Nat Med* 2014;20:54-61.
[PUBMED](#) | [CROSSREF](#)
64. Arae K, Morita H, Unno H, Motomura K, Toyama S, Okada N, et al. Chitin promotes antigen-specific Th2 cell-mediated murine asthma through induction of IL-33-mediated IL-1 β production by DCs. *Sci Rep* 2018;8:11721.
[PUBMED](#) | [CROSSREF](#)
65. Rossios C, Pavlidis S, Hoda U, Kuo CH, Wiegman C, Russell K, et al. Sputum transcriptomics reveal upregulation of IL-1 receptor family members in patients with severe asthma. *J Allergy Clin Immunol* 2018;141:560-70.
[PUBMED](#) | [CROSSREF](#)
66. Di Stefano A, Caramori G, Barczyk A, Vicari C, Brun P, Zanini A, et al. Innate immunity but not NLRP3 inflammasome activation correlates with severity of stable COPD. *Thorax* 2014;69:516-24.
[PUBMED](#) | [CROSSREF](#)

67. Calverley PM, Sethi S, Dawson M, Ward CK, Finch DK, Penney M, et al. A randomised, placebo-controlled trial of anti-interleukin-1 receptor 1 monoclonal antibody MEDI8968 in chronic obstructive pulmonary disease. *Respir Res* 2017;18:153.
[PUBMED](#) | [CROSSREF](#)
68. Konya V, Czarnewski P, Forkel M, Rao A, Kokkinou E, Villablanca EJ, et al. Vitamin D downregulates the IL-23 receptor pathway in human mucosal group 3 innate lymphoid cells. *J Allergy Clin Immunol* 2018;141:279-92.
[PUBMED](#) | [CROSSREF](#)
69. Spencer SP, Wilhelm C, Yang Q, Hall JA, Bouladoux N, Boyd A, et al. Adaptation of innate lymphoid cells to a micronutrient deficiency promotes type 2 barrier immunity. *Science* 2014;343:432-7.
[PUBMED](#) | [CROSSREF](#)
70. Kim MA, Shin SW, Park JS, Uh ST, Chang HS, Bae DJ, et al. Clinical characteristics of exacerbation-prone adult asthmatics identified by cluster analysis. *Allergy Asthma Immunol Res* 2017;9:483-90.
[PUBMED](#) | [CROSSREF](#)
71. van Rijt L, von Richthofen H, van Ree R. Type 2 innate lymphoid cells: at the cross-roads in allergic asthma. *Semin Immunopathol* 2016;38:483-96.
[PUBMED](#) | [CROSSREF](#)
72. Gauvreau GM, O'Byrne PM, Boulet LP, Wang Y, Cockcroft D, Bigler J, et al. Effects of an anti-TSLP antibody on allergen-induced asthmatic responses. *N Engl J Med* 2014;370:2102-10.
[PUBMED](#) | [CROSSREF](#)
73. Gonem S, Berair R, Singapuri A, Hartley R, Laurencin MF, Bacher G, et al. Fevipiprant, a prostaglandin D2 receptor 2 antagonist, in patients with persistent eosinophilic asthma: a single-centre, randomised, double-blind, parallel-group, placebo-controlled trial. *Lancet Respir Med* 2016;4:699-707.
[PUBMED](#) | [CROSSREF](#)
74. Kuna P, Bjermer L, Tornling G. Two Phase II randomized trials on the CRTh2 antagonist AZD1981 in adults with asthma. *Drug Des Devel Ther* 2016;10:2759-70.
[PUBMED](#) | [CROSSREF](#)
75. Barnes N, Pavord I, Chuchalin A, Bell J, Hunter M, Lewis T, et al. A randomized, double-blind, placebo-controlled study of the CRTH2 antagonist OC000459 in moderate persistent asthma. *Clin Exp Allergy* 2012;42:38-48.
[PUBMED](#) | [CROSSREF](#)
76. Lao-Araya M, Steveling E, Scadding GW, Durham SR, Shamji MH. Seasonal increases in peripheral innate lymphoid type 2 cells are inhibited by subcutaneous grass pollen immunotherapy. *J Allergy Clin Immunol* 2014;134:1193-1195.e4.
[PUBMED](#) | [CROSSREF](#)
77. Zhong H, Fan XL, Yu QN, Qin ZL, Chen D, Xu R, et al. Increased innate type 2 immune response in house dust mite-allergic patients with allergic rhinitis. *Clin Immunol* 2017;183:293-9.
[PUBMED](#) | [CROSSREF](#)
78. Kim DW, Cho SH. Emerging endotypes of chronic rhinosinusitis and its application to precision medicine. *Allergy Asthma Immunol Res* 2017;9:299-306.
[PUBMED](#) | [CROSSREF](#)
79. Hong H, Chen F, Sun Y, Yang Q, Gao W, Cao Y, et al. Nasal IL-25 predicts the response to oral corticosteroids in chronic rhinosinusitis with nasal polyps. *J Allergy Clin Immunol* 2018;141:1890-2.
[PUBMED](#) | [CROSSREF](#)
80. Ogasawara N, Klingler AI, Tan BK, Poposki JA, Hulse KE, Stevens WW, et al. Epithelial activators of type 2 inflammation: elevation of thymic stromal lymphopoietin, but not IL-25 or IL-33, in chronic rhinosinusitis with nasal polyps in Chicago, Illinois. *Allergy* 2018;73:2251-4.
[PUBMED](#) | [CROSSREF](#)
81. Lam EP, Kariyawasam HH, Rana BM, Durham SR, McKenzie AN, Powell N, et al. IL-25/IL-33-responsive TH2 cells characterize nasal polyps with a default TH17 signature in nasal mucosa. *J Allergy Clin Immunol* 2016;137:1514-24.
[PUBMED](#) | [CROSSREF](#)
82. Laidlaw TM, Mullol J, Fan C, Zhang D, Amin N, Khan A, et al. Dupilumab improves nasal polyp burden and asthma control in patients with CRSwNP and AERD. *J Allergy Clin Immunol Pract* 2019;7:2462-2465.e1.
[PUBMED](#) | [CROSSREF](#)
83. Barnes PJ. Targeting cytokines to treat asthma and chronic obstructive pulmonary disease. *Nat Rev Immunol* 2018;18:454-66.
[PUBMED](#) | [CROSSREF](#)

84. Shikhagaie MM, Björklund AK, Mjösberg J, Erjefält JS, Cornelissen AS, Ros XR, et al. Neuropilin-1 is expressed on lymphoid tissue residing LT α i-like group 3 innate lymphoid cells and associated with ectopic lymphoid aggregates. *Cell Reports* 2017;18:1761-73.
[PUBMED](#) | [CROSSREF](#)
85. Donovan C, Starkey MR, Kim RY, Rana BM, Barlow JL, Jones B, et al. Roles for T/B lymphocytes and ILC2s in experimental chronic obstructive pulmonary disease. *J Leukoc Biol* 2019;105:143-50.
[PUBMED](#) | [CROSSREF](#)
86. Bafadhel M, McKenna S, Terry S, Mistry V, Reid C, Haldar P, et al. Acute exacerbations of chronic obstructive pulmonary disease: identification of biologic clusters and their biomarkers. *Am J Respir Crit Care Med* 2011;184:662-71.
[PUBMED](#) | [CROSSREF](#)
87. Yanagisawa S, Ichinose M. Definition and diagnosis of asthma-COPD overlap (ACO). *Allergol Int* 2018;67:172-8.
[PUBMED](#) | [CROSSREF](#)
88. Cai T, Qiu J, Ji Y, Li W, Ding Z, Suo C, et al. IL-17-producing ST2 $^{+}$ group 2 innate lymphoid cells play a pathogenic role in lung inflammation. *J Allergy Clin Immunol* 2019;143:229-244.e9.
[PUBMED](#) | [CROSSREF](#)