



EGb761 Provides a Protective Effect against A β ₁₋₄₂ Oligomer-Induced Cell Damage and Blood-Brain Barrier Disruption in an *In Vitro* bEnd.3 Endothelial Model

Wen-bin Wan¹, Lan Cao², Lu-mei Liu³, Bill Kalionis⁴, Chuan Chen⁵, Xian-tao Tai⁶, Ya-ming Li^{3*}, Shijin Xia^{7*}

1 Department of Neurology, Zhongshan Hospital, Fudan University, Shanghai, China, **2** State Key Laboratory of Medical Neurobiology, School of Basic Medical Sciences, Fudan University, Shanghai, China, **3** Geriatrics Department of Traditional Chinese Medicine, Huadong Hospital, Fudan University, Shanghai, China, **4** Department of Perinatal Medicine Pregnancy Research Centre and University of Melbourne Department of Obstetrics and Gynaecology, Royal Women's Hospital, Parkville, VIC, Australia, **5** Shanghai Geriatric Institute of Chinese Medicine, Shanghai, China, **6** School of Acupuncture, Massage and Rehabilitation, Yunnan University of Traditional Chinese Medicine, Kunming, China, **7** Shanghai Institute of Geriatrics, Huadong Hospital, Fudan University, Shanghai, China

Abstract

Alzheimer's disease (AD) is the most common form of senile dementia which is characterized by abnormal amyloid beta (A β) accumulation and deposition in brain parenchyma and cerebral capillaries, and leads to blood-brain barrier (BBB) disruption. Despite great progress in understanding the etiology of AD, the underlying pathogenic mechanism of BBB damage is still unclear, and no effective treatment has been devised. The standard *Ginkgo biloba* extract EGb761 has been widely used as a potential cognitive enhancer for the treatment of AD. However, the cellular mechanism underlying the effect remain to be clarified. In this study, we employed an immortalized endothelial cell line (bEnd.3) and incubation of A β ₁₋₄₂ oligomer, to mimic a monolayer BBB model under conditions found in the AD brain. We investigated the effect of EGb761 on BBB and found that A β ₁₋₄₂ oligomer-induced cell injury, apoptosis, and generation of intracellular reactive oxygen species (ROS), were attenuated by treatment with EGb761. Moreover, treatment of the cells with EGb761 decreased BBB permeability and increased tight junction scaffold protein levels including ZO-1, Claudin-5 and Occludin. We also found that the A β ₁₋₄₂ oligomer-induced upregulation of the receptor for advanced glycation end-products (RAGE), which mediates A β cytotoxicity and plays an essential role in AD progression, was significantly decreased by treatment with EGb761. To our knowledge, we provide the first direct *in vitro* evidence of an effect of EGb761 on the brain endothelium exposed to A β ₁₋₄₂ oligomer, and on the expression of tight junction (TJ) scaffold proteins and RAGE. Our results provide a new insight into a possible mechanism of action of EGb761. This study provides a rational basis for the therapeutic application of EGb761 in the treatment of AD.

Citation: Wan W-b, Cao L, Liu L-m, Kalionis B, Chen C, et al. (2014) EGb761 Provides a Protective Effect against A β ₁₋₄₂ Oligomer-Induced Cell Damage and Blood-Brain Barrier Disruption in an *In Vitro* bEnd.3 Endothelial Model. PLoS ONE 9(11): e113126. doi:10.1371/journal.pone.0113126

Editor: Wanli Smith, University of Maryland School of Pharmacy, United States of America

Received: May 25, 2014; **Accepted:** October 20, 2014; **Published:** November 26, 2014

Copyright: © 2014 Wan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

Funding: The authors gratefully acknowledge the financial support of this study by National Natural Science Foundation of China (Grant No. 81473739), and Shanghai Committee of Science and Technology, China (Grant No. 12401904500), YML received the funding, (<http://www.nsf.gov.cn/>) and (<http://www.stcsm.gov.cn/>), respectively; by National Natural Science Foundation of China (Grant No. 31171129, Grant No. 81460748), SJX received the funding, (<http://www.nsf.gov.cn/>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: doctorymli@163.com (YML); xiashijinhd@163.com (SJX)

Introduction

Ginkgo biloba leaves are a type of medicinal herb and their extract has been shown to have neuroprotective properties and enhance cognitive functions [1,2]. EGb761 is the standardized extract of *Ginkgo biloba* produced by Dr. Willar Schwabe Pharmaceuticals, which contains 22–27% flavonol glycosides, 5.4–6.6% terpene trilactones, 2.8–3.4% ginkgolides (A, B and C), 2.6–3.2% bilobalide, and less than 5 ppm ginkgolic acids [1]. Recently, EGb761 has received significant attention as a potential cognitive enhancer for the treatment of Alzheimer's disease (AD) [1–4]. Substantial clinical and preclinical evidence indicates that EGb761 limits vascular and neural damage and has many beneficial effects that support its use in treating AD individuals

[5–7]. However, the cellular and molecular mechanisms underlying these effects remain to be elucidated.

AD is the most common neurodegenerative disease that causes progressive cognitive and behavioral deterioration in the elderly [8,9]. Extracellular deposition of the amyloid beta (A β) is widely accepted as an important event in the pathogenesis of AD [10,11]. A β is considered to be one of the most acute neurotoxins in the central nervous system [10–12]. Very recently, cerebrovascular changes leading to blood-brain barrier (BBB) leakiness have been associated with A β deposition in the brains of AD individuals, and this may be involved in AD progression [13–15]. Despite great progress in understanding the etiology of AD, the process of deposition of A β aggregates in cerebral capillaries and the brain is still poorly understood and the underlying pathogenic mechanisms

of BBB leakage remain unclear. Furthermore, no effective treatment has been devised.

The receptor for advanced glycation end-products (RAGE) is an essential transmembrane cell-signaling receptor, which binds free A β and mediates pathophysiological cellular responses, including oxidative stress, neurodegeneration, transport of circulating plasma A β across the BBB into the brain, and brain endothelial cell (EC) damage [16–19]. RAGE expression is increased in cells of the neurovascular unit in the brains of AD individuals, and in disease models of AD both *in vivo* and *in vitro* [19,20]. This is particularly the case in models associated with an A β -rich environment [21]. More importantly, antagonizing RAGE expression, or RAGE-knockout studies, show that blocking the RAGE-A β interaction at the BBB suppresses the accumulation of A β in brain parenchyma [22], prevents A β -induced BBB disruption and ameliorates tight junction (TJ) scaffold protein expression [20]. These data suggest that RAGE is related to A β accumulation as well as disruption of BBB integrity, and that RAGE might be a potential therapeutic target for AD.

Recently, an *in vitro* study in a cell monolayer BBB model reported that EGb761 diminished cell injury induced by chronic hypoxia and hypoglycemia (CHH), and significantly reversed CHH-induced upregulation of RAGE expression [23]. Considering the protective properties of EGb761 and its therapeutic potential, we speculated that EGb761 treatment might have a protective effect on A β -induced BBB disruption by inhibition of RAGE. To testify our hypothesis, we employed an *in vitro* BBB model comprising an immortalized mouse brain capillary endothelial cell line (bEnd.3). Our study assessed the effects of A β _{1–42} oligomer treatment of bEnd.3 endothelial cells with respect to changes in the expression of RAGE, and TJ scaffold proteins including ZO-1, Claudin-5 and Occludin. Finally, we investigated the effect of EGb761 on A β _{1–42} oligomer treatment of bEnd.3 endothelial cells.

Materials and Methods

Reagents and antibodies

Lyophilized human A β _{1–42}, purified by HPLC, was purchased from GL Biochem (Shanghai, China). EGb761 powder, a standardized *Ginkgo biloba* extract that contains two major active constituents 24% flavonol glycosides and 6% terpene trilactones, was purchased from Dr. Willmar Schwabe (Karlsruhe, Germany). The rabbit anti-ZO-1, anti-Claudin-5 and anti-Occludin antibodies were purchased from Invitrogen (CA, USA), whilst the rabbit anti-RAGE antibody was purchased from Millipore (MA, USA). The rabbit anti-GAPDH antibody was purchased from Santa Cruz Biotechnology (CA, USA) and the IRDye 680LT goat anti-rabbit IgG was purchased from LI-COR (CA, USA). MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide] was purchased from Sigma (CA, USA). Sodium fluorescein (Na-F, MW: 376 Da) powder was purchased from Kayon Bio-tech Co.(Shanghai, China).

Reagents preparation

Lyophilized human A β _{1–42} was used to prepare A β _{1–42} oligomer as described previously [24,25]. A β _{1–42} was initially dissolved to 1 mM in hexafluoroisopropanol (HFIP, Sigma, USA) and aliquoted into sterile microcentrifuge tubes. Then, HFIP was removed under vacuum in a Speed Vac, and the peptide stored at –20°C. For oligomer preparation, 2 mM A β _{1–42} peptide that dissolved in dry dimethyl sulfoxide (DMSO, Sigma, USA) was subsequently diluted into ice-cold Opti-MEM (Gibco, USA) to bring the peptide to a final concentration of 100 μ M. The solution

was vortexed for 30 seconds, centrifuged for 1 minute, and incubated at 4°C for 24 h before use. EGb761 was dissolved in DMSO at a concentration of 200 mg/ml and stored at room temperature. The required concentrations of EGb761 were made by further dilution of the concentrated stock solution with Opti-MEM.

Cell culture and treatments

Murine brain capillary endothelial cells (bEnd.3 obtained from ATCC, USA) were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco, USA) containing 4.5 g/l glucose supplemented with 10% fetal bovine serum (Gibco, USA), 100 U/ml penicillin and 100 μ g/ml streptomycin (Gibco, USA) at 37°C in a humidified atmosphere containing 5% CO₂ and subcultured every 3 days.

Cells were grown to 70–80% confluence prior to treatment. Before the treatments were applied, cells were rinsed in PBS and then the medium was replaced with Opti-MEM (Invitrogen, USA). For treatment of the cells exposed to A β _{1–42} oligomer and EGb761, the cells were pretreated with EGb761 for 2 h and then treated with A β _{1–42} oligomer.

Measurement of cell viability

Cell viability was measured using MTT assay. bEnd.3 cells were seeded onto 96-well plates and treated with EGb761 at different concentrations. MTT (20 μ L of a 5 mg/ml stock, diluted in PBS) was added to each cell culture well containing 100 μ L of medium. After 4 h incubation at 37°C, the medium was gently aspirated. Deposited formazan crystals were lysed in 100 μ L DMSO by gently shaking the plate. Absorbance at 570 nm was measured using a micro plate reader (Bio-Rad). The cell viability (%) was expressed as a percentage relative to the untreated control cells.

Detection of cell apoptosis

Apoptosis was observed by Hoechst-33258 staining (Apoptosis Hoechst staining kit; Beyotime, China). Briefly, cells were fixed in 0.5 mL of methanol for 15 min, followed by two washes with PBS. Cells were stained with 1 μ g/mL Hoechst 33258 in a dark chamber at room temperature for 10 min and again washed twice in PBS. Cells were analyzed by fluorescence microscopy using excitation at 350 nm and emission at 460 nm. Apoptotic cells were identified on the basis of nuclear morphology changes such as chromatin condensation and fragmentation. In each group, ten fields of view were selected randomly and counted.

Detection of intracellular ROS

The level of intracellular reactive oxygen species (ROS) was quantified using the Reactive Oxygen Species Assay Kit (Beyotime, China). DCFH-DA is oxidized by reactive oxygen species in viable cells to 2',7'-dichlorofluorescein (DCF) which is highly fluorescent at 530 nm. Cells were washed three times with PBS and then DCFH-DA, diluted to a final concentration of 10 μ M, was added and the cells were incubated for 30 min at 37°C in the dark. After washing three times with PBS, the stained cells in the 6-well plate were analyzed by inverted fluorescence microscopy (CKX41, OLYMPUS, Japan). The relative levels of fluorescence in cells were quantified by a multi-detection microplate reader (Bio-Rad) with excitation at 488 nm and emission at 525 nm. The level of intracellular ROS was expressed as the percentage of the control cells.

BBB permeability assay

Transendothelial permeability was measured using Na-F as described previously [26,27] with the following modifications. bEnd.3 cells (5×10^4 cells/cm²) were cultured in the apical compartment, on a 0.4 μ m pore size, 6.5 mm diameter polycarbonate membrane Transwell permeable insert (Corning). After the cells achieved confluence, 1.5 ml HHBS assay buffer (136 mM NaCl, 0.9 mM CaCl₂, 0.5 mM MgCl₂, 2.7 mM KCl, 1.5 mM KH₂PO₄, 10 mM Na H₂PO₄, 25 mM glucose, and 10 mM HEPES, pH 7.4) was added to the basolateral compartment. Culture medium in the apical compartment was replaced by 0.5 ml HHBS assay buffer containing 10 μ g/ml Na-F (Kayon Biology). After 30 min, the medium from the basolateral compartments was removed and fluorescence in this medium was determined by a multiwell plate reader (Bio-Rad) at the wavelengths of 485 nm (excitation) and 535 nm (emission).

Western blotting

Cells extracts were prepared by washing cells twice with PBS and resuspending in RIPA buffer (150 mM NaCl, 1% NP-40, 0.5% Sodium Deoxycholate, 0.1% SDS, 50 mM Tris-HCl (pH 7.4), 20 mM NaF, 20 mM EGTA, 1 mM DTT, 1 mM Na₃VO₄) with PMSF containing protease and phosphatase inhibitors. The extracts were then subjected to ultrasonication. Western blotting was performed to measure the change in tight junction protein levels including ZO-1, Claudin-5 and Occludin, and RAGE. Protein samples (30 μ g total protein per lane) were subjected to 10% SDS-PAGE. After electrophoresis, protein was transferred onto a nitrocellulose (NC) blotting membrane (Millipore). Membranes were blocked with 5% fat-free milk for 1 h at room temperature, and then incubated overnight at 4°C with the following rabbit primary antibodies diluted to 1:1000; anti-ZO-1 (Invitrogen, USA), anti-Claudin-5 (Invitrogen, USA), anti-Occludin (Invitrogen, USA), anti-RAGE (Millipore, USA) and anti-GAPDH (Santa Cruz, USA). Secondary goat anti-rabbit antibody (LI-COR, USA) was incubated with the filters for 1 h at room temperature. The images were captured using Odyssey infrared fluorescence imaging system (LI-COR, USA).

Statistical analysis

All results are expressed as the mean \pm S.E.M. Statistical analysis was performed using GraphPad Prism 5.0 software (GraphPad Software, Inc.). All experiments were repeated three times independently. Statistical significance of differences among different groups was analyzed by one-way analysis of variance (ANOVA) or student *t* test. A *p*-value < 0.05 was considered statistically significant.

Results

EGb761 diminished A β ₁₋₄₂ oligomer-induced cell injury of bEnd.3 cells

In this study, we first investigated whether EGb761 influenced the cell viability of bEnd.3 cells by MTT analysis. The results showed that incubation with various concentrations of EGb761 (25–200 μ g/ml) in Opti-MEM did not lead to any significant changes in cell viability (**Fig. 1A**). However, at a concentration of 300 μ g/ml, EGb761-treatment resulted in a significant decrease in cell viability (*p* = 0.0008, **Fig. 1A**). Therefore, concentration of EGb761 between 25–200 μ g/ml was used in the subsequent experiments. This concentration range of EGb761 includes the 100 μ g/ml concentration, which was showed to be effective in bEnd.3 cells in a related study [23].

The viability of bEnd.3 cells, pretreated with 25–200 μ g/ml EGb761 and then incubated with A β ₁₋₄₂ oligomer was determined. The concentration of A β ₁₋₄₂ oligomer (10 μ M) was based on the optimization data as described previously [20,28] with some modifications. The results showed that cells treated with A β ₁₋₄₂ oligomer alone had significantly reduced viability compared with untreated controls. Pretreatment with EGb761 for 2 h prior to addition of A β ₁₋₄₂ oligomer resulted in a significant increase in cell viability in a dose-dependent manner from 25 μ g/ml to 100 μ g/ml EGb761. Fold changes in cell viability following EGb761 and A β ₁₋₄₂ oligomer treatment, relative to A β ₁₋₄₂ oligomer alone, were 1.07, 1.19, 1.48 and 1.41-fold at 25, 50, 100 and 200 μ g/ml EGb761 respectively (*#p* < 0.01, **Fig. 1B**).

EGb761 prevented A β ₁₋₄₂ oligomer-triggered apoptosis in bEnd.3 cells

To investigate the effect of EGb761 on bEnd.3 cell apoptosis, cells were incubated with or without EGb761 for 2 h, followed by treatment with 10 μ M A β ₁₋₄₂ oligomer for another 24 h. We used a concentration of 100 μ g/ml EGb761 since this was most effective in the MTT assay (**Fig. 1B**). In the untreated (Control) group, cell nuclei were uniformly stained with the Hoechst-33258 dye (**Fig. 2A, Control**), whilst in the group treated with A β ₁₋₄₂ oligomer alone, bright chromatin condensation and nuclear fragmentation were observed, which is typical of apoptotic nuclei (**Fig. 2A, A β**). In the EGb761 and A β ₁₋₄₂ treated group, the nuclei were stained uniformly and the intensity of staining matched the untreated (Control) group (**Fig. 2A, EGb761+A β**). Apoptotic nuclei were quantitated and the results showed a significant increase in the percentage of apoptotic cells following treatment with A β ₁₋₄₂ oligomer alone (*p* < 0.01, A β versus Control, **Fig. 2B**). Treatment with EGb761 prior to addition of A β ₁₋₄₂ oligomer significantly reduced the percentage of apoptotic cells (*p* < 0.01, EGb761+A β versus A β , **Fig. 2B**).

EGb761 attenuated A β ₁₋₄₂ oligomer-induced ROS generation in bEnd.3 cells

Oxidative stress plays an important role in A β -induced cytotoxicity. Therefore, we examined the effect of EGb761 on A β ₁₋₄₂ oligomer-induced ROS generation in bEnd.3 endothelial cells. A marked increase in ROS generation was detected after treatment with A β ₁₋₄₂ oligomer alone, with 4.05-fold higher levels of oxidized DCF detected compared with untreated control cells (**p* < 0.01, A β versus Control, **Fig. 3A**). Treatment with EGb761 prior to addition of A β ₁₋₄₂ oligomer significantly reduced ROS formation induced by the A β ₁₋₄₂ oligomer (*#p* < 0.01, EGb761+A β versus A β , **Fig. 3A**). These data suggest that EGb761 attenuated A β ₁₋₄₂ oligomer-induced ROS generation in bEnd.3 cells.

EGb761 reduced BBB leakage induced by the A β ₁₋₄₂ oligomer

The BBB is a specialized barrier that controls the transport of various molecules and maintains the integrity of brain by restricting permeability across the brain endothelium [17]. We found that A β ₁₋₄₂ oligomer increased permeability in cultured bEnd.3 cells (**p* < 0.01, **Fig. 4**). Pretreatment with EGb761 reversed the barrier permeability damaged induced by A β ₁₋₄₂ oligomer (*#p* < 0.01, **Fig. 4**), and the effect was detected in a dose-dependent manner from 25 μ g/ml to 100 μ g/ml.




Figure 1. EGb761 increased cell viability against A β ₁₋₄₂ oligomer toxicity. Panel A, bEnd.3 cells were incubated with various concentrations of EGb761 for 24 h and then cell viability was analyzed according to MTT assay by detecting the absorbance at 490 nm. Panel B, cells were incubated with or without various concentrations of EGb761 for 2 h, followed by incubation with 10 μ M A β ₁₋₄₂ oligomer for 24 h. Subsequently, cell viability was determined by MTT assay. Results are shown as the Mean \pm S.E.M. (* p <0.01, A β versus Control; # p <0.01, EGb761+A β versus A β ; ** p <0.05, versus A β). doi:10.1371/journal.pone.0113126.g001

EGb761 increased protein levels of ZO-1, Claudin-5 and Occludin in A β ₁₋₄₂ oligomer-induced bEnd.3 cells

TJs are the most prominent feature of the brain endothelium and are key structures that ensure the integrity of the BBB [28,29]. On the basis of the above results, we determined the effect of EGb761-pretreatment of bEnd.3 cells on the expression of TJ scaffold proteins ZO-1, Claudin-5 and Occludin. Cells were pretreated with or without EGb761 for 2 h, at concentrations from 25 μ g/ml to 200 μ g/ml, then exposed to 10 μ M A β ₁₋₄₂ oligomer. Western blot and semi-quantitative analysis showed that the treatment with A β ₁₋₄₂ oligomer alone significantly decreased the

levels of ZO-1, Claudin-5 and Occludin in bEnd.3 cells relative to the control (Ctrl) (* p <0.01, **Fig. 5**). Pretreatment with EGb761 significantly increased the levels of those proteins (# p <0.01, **Fig. 5**). The protective effect of EGb761 on ZO-1 and Claudin-5 was in a concentration dependent manner from 25 μ g/ml to 100 μ g/ml, whereas Occludin levels increased in a concentration dependent manner from 25 μ g/ml to 200 μ g/ml.




Figure 2. EGb761 prevented A β ₁₋₄₂ oligomer-induced apoptosis. bEnd.3 cells were incubated with or without EGb761 (100 μ g/ml), followed by incubation with 10 μ M A β ₁₋₄₂ oligomer for 24 h. Cells were then subjected to Hoechst-33258 staining and viewed under a fluorescence microscope. Arrows indicate apoptotic nuclei. The apoptotic nuclei appear shrunken, irregular and fragmented (arrows). Panel B shows the percentage of apoptotic cells. In each group, ten microscopic fields were selected randomly and counted. Results are shown as the Mean \pm S.E.M. (* p <0.01, A β versus Control; # p <0.01, EGb761+A β versus A β). doi:10.1371/journal.pone.0113126.g002




Figure 3. EGb761 attenuated the A β_{1-42} oligomer-induced increase of ROS. Panel A: ROS generation in bEnd.3 cells was evaluated by the oxidation of H₂DCF-DA to DCF (Fig. 3A, Control) and assessed by inverted fluorescent microscopy (100 \times). Following treatment for 24 h with 10 μ M A β_{1-42} oligomer, an increase in fluorescence was detected (Fig. 3A, A β). Cells treated with 100 μ g/mL EGb761 for 2 h prior A β_{1-42} oligomer treatment for 24 h, showed a decrease in fluorescence (Fig. 3A, EGb761+A β). Panel B shows the relative levels of intracellular ROS quantified by a microplate reader (488 nm excitation and 525 nm emission), with the results normalized to the control (set at 100). Results are shown as the Mean \pm S.E.M. (* p < 0.01, A β versus Control; # p < 0.01, EGb761+A β versus A β). doi:10.1371/journal.pone.0113126.g003




Figure 4. EGb761 decreased BBB permeability in A β_{1-42} oligomer-induced bEnd.3 cells. BBB permeability, evaluated by the Na-F leakage test, was assessed after incubation with 10 μ M A β_{1-42} oligomer. Cells were incubated with or without various concentrations of EGb761 for 2 h, followed by incubation with A β_{1-42} oligomer for 24 h. Then, the absorbance of Na-F was determined by fluorescence spectrophotometry. Results are shown as the Mean \pm S.E.M. (* p < 0.01, A β versus Control; # p < 0.01, EGb761+A β versus A β). doi:10.1371/journal.pone.0113126.g004




Figure 5. EGb761 increased the expression ZO-1, Claudin-5 and Occludin in A β ₁₋₄₂ oligomer-induced bEnd.3 cells. Panel A, cells were incubated with or without various concentrations of EGb761 for 2 h, followed by incubation with A β ₁₋₄₂ oligomer for 24 h. Then, the expression levels of ZO-1, Claudin-5 and Occludin were determined by Western Blot. Panels B, C and D show the quantitation of the data from Panel A. The Western blots of target proteins were semi-quantitatively analyzed by Image J software and the sum optical density was obtained. Protein levels relative to GAPDH were determined and then normalized to the Control value (i.e. untreated cells), which was set to 1.0. Results are shown as the Mean \pm S.E.M. (* p <0.01, A β versus Control; # p <0.01, EGb761+A β versus A β). doi:10.1371/journal.pone.0113126.g005

EGb761 reversed A β ₁₋₄₂ oligomer-induced upregulation of RAGE expression in bEnd.3 cells

In this study, we hypothesized that EGb761 would protect against A β -induced BBB disruption through inhibition of RAGE. To test the hypothesis, we determined the effect on the expression of RAGE in A β ₁₋₄₂ oligomer-induced bEnd.3 cells. Western blot and semi-quantitative analysis revealed that after incubation with A β ₁₋₄₂ oligomer for 24 h, the expression of RAGE was

significantly increased by 1.97-fold when compared with the unexposed Control bEnd.3 cells (* p <0.01, **Fig. 6**). Whereas, treatment of A β ₁₋₄₂ oligomer-induced bEnd.3 cells with various concentrations of EGb761 led to a significant decrease in the expression of RAGE (# p <0.01, **Fig. 6**). Furthermore, the findings suggest that the protective effect of EGb761 on RAGE was in a dose-dependent manner from 25 μ g/ml to 100 μ g/ml. A further decrease in RAGE expression after pretreated with




Figure 6. EGb761 reversed A β ₁₋₄₂ oligomer-induced up-regulation of RAGE expression in bEnd.3 cells. Panel A, cells were incubated with or without various concentrations of EGb761 for 2 h, followed by incubation with A β ₁₋₄₂ oligomer for 24 h. Then, RAGE was determined by Western Blot. Panel B is the bar graph based on the result of Panel A. The blots of target proteins were semi-quantitatively analyzed by Image J software and the sum optical density was obtained. Results are shown as the Mean \pm S.E.M. (* p <0.01, A β versus Control; # p <0.01, EGb761+A β versus A β). doi:10.1371/journal.pone.0113126.g006

200 $\mu\text{g/ml}$ EGb761 was not detectable, when compared with 100 $\mu\text{g/ml}$ EGb761 (**Fig. 6**).

Discussion

According to the vascular hypothesis of AD, initial vascular damage plays a critical role in the disease development [30]. The origin of BBB dysfunction during AD is not known. However, in a number of AD transgenic animal models, accumulation of A β in blood vessels results in the disruption of the BBB [15,20,31]. The hypothesis is that BBB breakdown leads to accumulation in the brain of multiple vasculotoxic and neurotoxic macromolecules, and this can initiate functional and structural changes in neurons before A β deposition occurs [30]. More importantly, BBB damage impairs vascular clearance of brain A β and increases RAGE-mediated influx of blood A β into the brain [22,30]. In this study, we treated cultured immortalized mouse cerebral microvessel endothelial cells with A β to model the conditions of the BBB in AD, and subsequently observed the effect of EGb761 on this cell monolayer model of BBB. bEnd.3 cell viability was significantly decreased in response to incubation with A β_{1-42} oligomer (**Fig. 1**). There was also a qualitative increase in the number of apoptotic bEnd.3 cells (**Fig. 2**) and an increase in ROS generation (**Fig. 3**). Treatment of EGb761 restored cell viability and reduced both A β_{1-42} oligomer-induced cell apoptosis and ROS production *in vitro*.

Intercellular TJs are the most prominent feature of brain endothelium and are responsible for BBB integrity [32]. The physical seal of the BBB is maintained by several different interendothelial TJ complexes that are composed of connecting transmembrane proteins (Occludin and Claudins). These proteins form the primary seal and are linked to accessory cytoplasmic proteins of Zona Occludens family members (ZO-1/2/3 etc), which can also independently link other types of transmembrane proteins to the actin cytoskeleton [33,34]. Studies have shown that TJ breakdown contributes to the deficiency in BBB function, and abnormal expression of TJ scaffold proteins results in loss of TJ integrity and increased BBB permeability [35,36]. In this study, we demonstrated that treatment with A β_{1-42} oligomer caused significant BBB leakage (**Fig. 4**) and downregulations of ZO-1, Claudin-5 and Occludin (**Fig. 5**). These effects were reduced by EGb761 treatment.

RAGE is a pattern recognition receptor that binds to number of ligands including A β [37]. With the exception of the lungs, the basal expression of RAGE is low in physiological conditions but increases with the levels of its ligands [37,38]. Further, RAGE-ligand interaction and the subsequent up-regulation of RAGE through a positive feedback loop are associated with various diseases, including AD [39]. Accumulating evidence suggests that A β plays an essential role in BBB disruption, however, the exact mechanism leading to BBB alteration has not been determined. Recently, A β treatment was shown to induce RAGE expression in an *in vitro* study, and furthermore, interaction between A β and RAGE triggered an intercellular cascade that disrupted TJ leading to the breakdown of BBB integrity [20,33]. When pathogenic A β species accumulated in the AD brain, either in transgenic models of β -amyloidosis or in the human brain, RAGE expression was increased in affected cerebral vessels, neurons or microglia [40]. This mechanism provides the potential for exacerbating cellular dysfunction due to RAGE-A β interactions. The activation of RAGE expressed in neuronal cells promotes synaptic dysfunction

and as well leads to neurodegeneration by inducing inflammation in glial cells [9,41]. Moreover, RAGE-A β interaction is implicated in the development of Alzheimer's neurovascular disorder through various mechanisms. These include mediation of transcytosis of circulating A β across the BBB, induction of inflammatory responses in the endothelium, brain endothelial nuclear factor- κB (NF- κB) dependent apoptosis and suppression of cerebral blood flow (CBF), all of which culminate in BBB disruption [19,42]. In our present study we demonstrated that A β_{1-42} oligomer exposure led to a significant increase in the expression level of RAGE in bEnd.3 cells (**Fig. 6**).

Accumulating evidence suggests that RAGE is a potential target for therapies to lower brain A β burden, prevent BBB damage, and improve both CBF and behavioral performance [19,20]. These data suggest RAGE is a potential therapeutic target for AD. A recent study showed that EGb761 markedly reversed the up-regulation of RAGE induced by a CHH condition in a BBB *in vitro* model at both the RAGE mRNA and protein level [23]. These data suggest a rational basis for the therapeutic application of EGb761 in the treatment of AD [23]. Thus, we hypothesized that EGb761 would protect brain ECs against A β toxicity via inhibition of RAGE expression. The results indicated that the up-regulation of RAGE expression induced by A β_{1-42} oligomer was reversed by treatment with EGb761 (**Fig. 6**).

EGb761 has received a great many attentions because it exerts beneficial effects in conditions which are associated with impaired cognitive function [1,3,7]. In the present study, we found that 100 $\mu\text{g/ml}$ of EGb61 showed maximal protection in mainly detection indexes including cell viability, apoptosis, ROS, and the expression levels of ZO-1 and Claudin-5. However, the results also showed that 200 $\mu\text{g/ml}$ of EGb761 resulted in maximal protection with regard to the expression of Occludin. Furthermore, the data indicated that the difference was not significant between 100 $\mu\text{g/ml}$ and 200 $\mu\text{g/ml}$ of EGb761 at the BBB permeability and the expression level of RAGE after incubation with A β .

In conclusion, we have presented novel evidence to show that EGb761 effectively prevented A β_{1-42} oligomer-induced brain EC damage, which was characterized by reduced cell viability injury, increased cell apoptosis and increased intracellular ROS generation. Furthermore, we found that EGb761 reduced BBB leakage, reversed A β_{1-42} oligomer-induced down-regulation of TJ scaffold proteins and prevented the A β_{1-42} oligomer-induced up-regulation of RAGE in bEnd.3 cells. To our knowledge, this is the first direct evidence for an effect of EGb761 on brain endothelial cells, and for an effect of EGb761 on the expression of RAGE and TJ scaffold proteins exposed to A β_{1-42} oligomer. Our results provide a rational basis for the therapeutic application of EGb761 in the treatment of AD.

Acknowledgments

We highly appreciate the help from the members in State Key Laboratory of Medical Neurobiology, School of Basic Medical Sciences, Fudan University.

Author Contributions

Conceived and designed the experiments: WBW YML SJX. Performed the experiments: WBW. Analyzed the data: LC BK CC XTT. Contributed reagents/materials/analysis tools: LC LML. Contributed to the writing of the manuscript: WBW.

References

- Muller WE, Heiser J, Leuner K (2012) Effects of the standardized Ginkgo biloba extract EGb 761(R) on neuroplasticity. *Int Psychogeriatr* 24 Suppl 1: S21–24.
- Zhang C, Ren C, Chen H, Geng R, Fan H, et al. (2013) The analog of Ginkgo biloba extract 761 is a protective factor of cognitive impairment induced by chronic fluorosis. *Biol Trace Elem Res* 153: 229–236.
- Kehr J, Yoshitake S, Ijiri S, Koch E, Noldner M, et al. (2012) Ginkgo biloba leaf extract (EGb 761(R)) and its specific acylated flavonol constituents increase dopamine and acetylcholine levels in the rat medial prefrontal cortex: possible implications for the cognitive enhancing properties of EGb 761(R). *Int Psychogeriatr* 24 Suppl 1: S25–34.
- Jahanshahi M, Nikmahzar E, Yadollahi N, Ramazani K (2012) Protective effects of Ginkgo biloba extract (EGB 761) on astrocytes of rat hippocampus after exposure with scopolamine. *Anat Cell Biol* 45: 92–96.
- Mashayekh A, Pham DL, Yousem DM, Dizon M, Barker PB, et al. (2011) Effects of Ginkgo biloba on cerebral blood flow assessed by quantitative MR perfusion imaging: a pilot study. *Neuroradiology* 53: 185–191.
- Weinmann S, Roll S, Schwarzbach C, Vauth C, Willich SN (2010) Effects of Ginkgo biloba in dementia: systematic review and meta-analysis. *BMC Geriatr* 10: 14.
- Evans JR (2013) Ginkgo biloba extract for age-related macular degeneration. *Cochrane Database Syst Rev* 1: CD001775.
- Chen Y, Wang C, Hu M, Pan J, Chen J, et al. (2012) Effects of ginkgolide A on okadaic acid-induced tau hyperphosphorylation and the PI3K-Akt signaling pathway in N2a cells. *Planta Med* 78: 1337–1341.
- Perrone L, Sbai O, Nawroth PP, Bierhaus A (2012) The Complexity of Sporadic Alzheimer's Disease Pathogenesis: The Role of RAGE as Therapeutic Target to Promote Neuroprotection by Inhibiting Neurovascular Dysfunction. *Int J Alzheimers Dis* 2012: 734956.
- Meyer-Luehmann M, Spiess-Jones TL, Prada C, Garcia-Alloza M, de Calignon A, et al. (2008) Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. *Nature* 451: 720–724.
- Pomara N, Sidtis JJ (2010) Alzheimer's disease. *N Engl J Med* 362: 1844; author reply 1844–1845.
- Xiao Q, Wang C, Li J, Hou Q, Li J, et al. (2010) Ginkgolide B protects hippocampal neurons from apoptosis induced by beta-amyloid 25–35 partly via up-regulation of brain-derived neurotrophic factor. *Eur J Pharmacol* 647: 48–54.
- Love S, Miners S, Palmer J, Chalmers K, Kehoe P (2009) Insights into the pathogenesis and pathogenicity of cerebral amyloid angiopathy. *Front Biosci (Landmark Ed)* 14: 4778–4792.
- Marco S, Skaper SD (2006) Amyloid beta-peptide 1–42 alters tight junction protein distribution and expression in brain microvessel endothelial cells. *Neurosci Lett* 401: 219–224.
- Biron KE, Dickstein DL, Gopaul R, Jefferies WA (2011) Amyloid triggers extensive cerebral angiogenesis causing blood brain barrier permeability and hypervascularity in Alzheimer's disease. *PLoS One* 6: e23789.
- Wan W, Chen H, Li Y (2014) The potential mechanisms of Abeta-receptor for advanced glycation end-products interaction disrupting tight junctions of the blood-brain barrier in Alzheimer's disease. *Int J Neurosci* 124: 75–81.
- Kook SY, Seok Hong H, Moon M, Mook-Jung I (2013) Disruption of blood-brain barrier in Alzheimer disease pathogenesis. *Tissue Barriers* 1: e23993.
- Choi BR, Cho WH, Kim J, Lee HJ, Chung C, et al. (2014) Increased expression of the receptor for advanced glycation end products in neurons and astrocytes in a triple transgenic mouse model of Alzheimer's disease. *Exp Mol Med* 46: e75.
- RJ D (2012) Is RAGE still a therapeutic target for Alzheimer's disease? *Future Med Chem* 4: 915–925.
- Kook SY, Hong HS, Moon M, Ha CM, Chang S, et al. (2012) Abeta(1–42)-RAGE interaction disrupts tight junctions of the blood-brain barrier via Ca(2+)-calcineurin signaling. *J Neurosci* 32: 8845–8854.
- Silverberg GD, Miller MC, Messier AA, Majmudar S, Machan JT, et al. (2010) Amyloid deposition and influx transporter expression at the blood-brain barrier increase in normal aging. *J Neuropathol Exp Neurol* 69: 98–108.
- Deane R, Du Yan S, Subramanian RK, LaRue B, Jovanovic S, et al. (2003) RAGE mediates amyloid-beta peptide transport across the blood-brain barrier and accumulation in brain. *Nat Med* 9: 907–913.
- Yan FL, Zheng Y, Zhao FD (2008) Effects of ginkgo biloba extract EGB761 on expression of RAGE and LRP-1 in cerebral microvascular endothelial cells under chronic hypoxia and hypoglycemia. *Acta Neuropathol* 116: 529–535.
- Li B, Zhong L, Yang X, Andersson T, Huang M, et al. (2011) WNT5A signaling contributes to Abeta-induced neuroinflammation and neurotoxicity. *PLoS One* 6: e22920.
- Dahlgren KN, Manelli AM, Stine WB, Jr., Baker LK, Krafft GA, et al. (2002) Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 277: 32046–32053.
- Kis B, Deli MA, Kobayashi H, Abraham CS, Yanagita T, et al. (2001) Adrenomedullin regulates blood-brain barrier functions in vitro. *Neuroreport* 12: 4139–4142.
- Nakagawa S, Deli MA, Nakao S, Honda M, Hayashi K, et al. (2007) Pericytes from brain microvessels strengthen the barrier integrity in primary cultures of rat brain endothelial cells. *Cell Mol Neurobiol* 27: 687–694.
- Zehendner CM, Librizzi L, Hedrich J, Bauer NM, Angamo EA, et al. (2013) Moderate hypoxia followed by reoxygenation results in blood-brain barrier breakdown via oxidative stress-dependent tight-junction protein disruption. *PLoS One* 8: e82823.
- Zhou T, He Q, Tong Y, Zhan R, Xu F, et al. (2014) Phospholipid transfer protein (PLTP) deficiency impaired blood-brain barrier integrity by increasing cerebrovascular oxidative stress. *Biochemical and biophysical research communications* 445: 352–356.
- Deane R, Singh I, Sagare AP, Bell RD, Ross NT, et al. (2012) A multimodal RAGE-specific inhibitor reduces amyloid beta-mediated brain disorder in a mouse model of Alzheimer disease. *J Clin Invest* 122: 1377–1392.
- Paul J, Strickland S, Melchor JP (2007) Fibrin deposition accelerates neurovascular damage and neuroinflammation in mouse models of Alzheimer's disease. *J Exp Med* 204: 1999–2008.
- Chen X, Threlkeld SW, Cummings EE, Juan I, Makeyev O, et al. (2012) Ischemia-reperfusion impairs blood-brain barrier function and alters tight junction protein expression in the ovine fetus. *Neuroscience* 226: 89–100.
- Carrano A, Hoozemans JJ, van der Vies SM, Rozemuller AJ, van Horssen J, et al. (2011) Amyloid Beta induces oxidative stress-mediated blood-brain barrier changes in capillary amyloid angiopathy. *Antioxid Redox Signal* 15: 1167–1178.
- Gonzalez-Mariscal L, Quiros M, Diaz-Coranz M (2011) ZO proteins and redox-dependent processes. *Antioxid Redox Signal* 15: 1235–1253.
- Bednarczyk J, Lukasiuk K (2011) Tight junctions in neurological diseases. *Acta Neurobiol Exp (Wars)* 71: 393–408.
- Liu WY, Wang ZB, Zhang LC, Wei X, Li L (2012) Tight junction in blood-brain barrier: an overview of structure, regulation, and regulator substances. *CNS Neurosci Ther* 18: 609–615.
- Slowik A, Merres J, Elfgen A, Jansen S, Mohr F, et al. (2012) Involvement of formyl peptide receptors in receptor for advanced glycation end products (RAGE) - and amyloid beta1-42-induced signal transduction in glial cells. *Mol Neurodegener* 7: 55.
- Ramasamy R, Yan SF, Schmidt AM (2012) Advanced glycation endproducts: from precursors to RAGE: round and round we go. *Amino Acids* 42: 1151–1161.
- Han SH, Kim YH, Mook-Jung I (2011) RAGE: the beneficial and deleterious effects by diverse mechanisms of actions. *Mol Cells* 31: 91–97.
- Deane R, Bell RD, Sagare A, Zlokovic BV (2009) Clearance of amyloid-beta peptide across the blood-brain barrier: implication for therapies in Alzheimer's disease. *CNS Neurol Disord Drug Targets* 8: 16–30.
- Yan SD, Roher A, Chaney M, Zlokovic B, Schmidt AM, et al. (2000) Cellular cofactors potentiating induction of stress and cytotoxicity by amyloid beta-peptide. *Biochim Biophys Acta* 1502: 145–157.
- Deane R, Zlokovic BV (2007) Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res* 4: 191–197.