






Article

Population Pharmacokinetic/Pharmacodynamic Modelling of Daptomycin for Schedule Optimization in Patients with Renal Impairment

Teresa García-Martínez ^{1,2}, María Dolores Bellés-Medall ² , María García-Cremades ³ ,
Raúl Ferrando-Piqueres ² , Víctor Mangas-Sanjuán ^{1,4,*}  and Matilde Merino-Sanjuán ^{1,4} 

¹ Department of Pharmacy and Pharmaceutical Technology and Parasitology, University of Valencia, 46100 Valencia, Spain

² Department of Pharmacy, University Hospital of Castellon, 12004 Castellon, Spain

³ Department of Pharmaceutics and Food Technology, School of Pharmacy, Complutense University of Madrid, 28040 Madrid, Spain

⁴ Interuniversity Research Institute for Molecular Recognition and Technological Development, 46022 Valencia, Spain

* Correspondence: victor.mangas@uv.es; Tel.: +34-963-543351

Abstract: The aims of this study are (i) to develop a population pharmacokinetic/pharmacodynamic model of daptomycin in patients with normal and impaired renal function, and (ii) to establish the optimal dose recommendation of daptomycin in clinical practice. Several structural PK models including linear and non-linear binding kinetics were evaluated. Monte Carlo simulations were conducted with a fixed combination of creatinine clearance (30–90 mL/min/1.73 m²) and body weight (50–100 kg). The final dataset included 46 patients and 157 daptomycin observations. A two-compartment model with first-order peripheral distribution and elimination kinetics assuming non-linear protein-binding kinetics was selected. The bactericidal effect for Gram+ strains with MIC ≤ 0.5 mg/L could be achieved with 5–12 mg/kg daily daptomycin based on body weight and renal function. The administration of 10–17 mg/kg q48 h daptomycin allows to achieve bactericidal effect for Gram+ strains with MIC ≤ 1 mg/L. Four PK samples were selected as the optimal sampling strategy for an accurate AUC estimation. A quantitative framework has served to characterize the non-linear binding kinetics of daptomycin in patients with normal and impaired renal function. The impact of different dosing regimens on the efficacy and safety outcomes of daptomycin treatment based on the unbound exposure of daptomycin and individual patient characteristics has been evaluated.

Keywords: daptomycin; pharmacokinetic; AUC; optimal dose selection; creatinine clearance



Citation: García-Martínez, T.; Bellés-Medall, M.D.; García-Cremades, M.; Ferrando-Piqueres, R.; Mangas-Sanjuán, V.; Merino-Sanjuán, M. Population Pharmacokinetic/Pharmacodynamic Modelling of Daptomycin for Schedule Optimization in Patients with Renal Impairment. *Pharmaceutics* **2022**, *14*, 2226. <https://doi.org/10.3390/pharmaceutics14102226>

Academic Editor: Hye Suk Lee

Received: 26 August 2022

Accepted: 15 October 2022

Published: 18 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The clinical use of daptomycin in adult patients includes complicated skin and soft-tissue infections, right-sided infective endocarditis and bacteremia due to *Staphylococcus aureus* (*S. aureus*), with an approved posology ranging from 4 to 6 mg/kg once daily for 7–14 days or until the resolution of the infection [1–6]. The emergence of more resistant Gram-positive strains is leading to the recommendation and use of daptomycin at higher than approved dosage levels [7–15], especially in critically ill patients, which increases the uncertainty of clinicians when managing patients who also present renal-type disorders or extreme weight values.

The pharmacokinetic (PK) properties of daptomycin have been investigated in multiple publications in recent years, both in adult and pediatric patients [16–18]. So far, linear PK processes of daptomycin have been proposed, which show a rapid disposition phase followed by a slower and sustained disposition phase over time. The excretion of daptomycin is primarily promoted by the kidneys, with 54–78% eliminated in urine and a plasma

elimination half-life from 7.6 to 8.8 h [3,5,6]. In addition, daptomycin is highly bound to plasma proteins, with a free fraction being around 10%. However, the pathophysiological alterations and clinical situation of patients may alter the protein binding kinetics [3,5], conditioning the time course of daptomycin in plasma and at the target site. It is essential that these elements be carefully considered to establish optimal dosage schemes based on the type of resistant microorganism, since in most cases the PK/PD efficacy indices are based on the total daptomycin values.

For all this, the development of model-informed strategies for the optimization of daptomycin treatment in patients is a fundamental tool in clinical practice. Creatinine clearance, albumin or body weight have been identified as clinically relevant covariates in the PK properties of daptomycin [17,19–22]. However, the use of therapeutic drug monitoring for the selection of an individualized daptomycin schedule could be additionally recommended [23], due to the moderate inter-individual variability on the PK parameters and the complex efficacy/safety balance. In this sense, Monte Carlo simulations represent a reliable and powerful methodology, which contributes to an optimal dose selection by integrating the variability in the PK parameters [20]. Recently, several authors have conducted a simulation-based analysis to suggest higher dose levels of daptomycin than approved to achieve bactericidal effects in adults [11,24–26] and pediatric [27,28] patients, especially on methicillin-resistant *Staphylococcus aureus* (MRSA) bacteremia. However, these studies include a narrow distribution of body weight and renal function conditions that impede a precise and specific evaluation of those effects on efficacy and safety outcomes. Therefore, the aims of the current work are (i) to develop a population pharmacokinetic/pharmacodynamic (PK/PD) model of daptomycin in patients with normal and impaired renal function, and (ii) to establish the optimal dose recommendation of daptomycin in clinical practice.

2. Materials and Methods

A prospective multi-center study in patients admitted to the General University Hospital of Castellón (Castellón, Spain) and the Arnau de Vilanova Hospital of Valencia (Valencia, Spain) was conducted. Patients who received daptomycin for any diagnosis were enrolled in the present study according to the following inclusion criteria: signed informed consent; administration daptomycin >48 h; and age >18 years. The exclusion criteria were patients with renal replacement therapy.

Daptomycin was administered by a 30-min intravenous infusion every 24 h, with daily doses ranging from 4 mg/kg up to 12 mg/kg by clinical practice.

Five serial blood samples were collected from a peripheral vein from day 4 (steady state) at the pre-dose, 0.5, 1–2, 4–10 h after the infusion ended and before the next dose. According to a TDM protocol for daptomycin adopted in these hospitals, the samples, collected in an additive-free tube, were centrifuged and the serum was stored at $-20\text{ }^{\circ}\text{C}$ until the drug concentrations were measured [29].

Demographic (age, body weight and height), treatment (dose and co-administered medications) and clinical data (microbiological data, analytical and disease covariates) were collected at patient enrolment. Serum creatinine and serum albumin were measured on the day (if not available, the nearest day) of measurement of the drug concentration. MIC were determined in all patients by e-test, but in patients where MIC were not available, the majority of the local MIC distribution to daptomycin was used (0.25 mg/L).

Creatinine clearance (CLCR) was estimated using the Cockcroft-Gault formula [30] and normal renal function (CLCR: $>90\text{ mL}/\text{min}/1.73\text{ m}^2$), mild renal impairment (CLCR: $60\text{--}90\text{ mL}/\text{min}/1.73\text{ m}^2$), moderate renal impairment (CLCR: $30\text{--}60\text{ mL}/\text{min}/1.73\text{ m}^2$), and severe renal impairment (CLCR: $<30\text{ mL}/\text{min}/1.73\text{ m}^2$) were established.

Daptomycin plasma total concentrations were determined by a validated high-performance liquid chromatography assay with UV detection, which was available at our hospital center [31]. Precision and accuracy were assessed by performing replicate analysis of quality control samples against calibration standards. Intra- and inter-assay

coefficients of the variation were always <10%. The lower limit of detection was 0.027 mg/L. Although free daptomycin fractions have been determined by ultra-performance liquid chromatography coupled to tandem mass spectrometry [32] (UPLC-MS/MS), this equipment was not available at our center and, therefore, no free daptomycin concentrations could be measured.

All data analyses were performed based on the population approach with the software NONMEM[®] (Version 7.4; Icon Inc, PA, USA). The population PK parameters were estimated using the Stochastic Approximation of the Expectation Maximization and the Importance Sampling Estimation method. The Perl-Speaks-NONMEM program (version 4.8.1, Department of Pharmaceutical Biosciences, Uppsala University, Sweden) [33] and Pirana software (version 2.9.6, Pirana Software & Consulting BV) were used for model development and evaluation. Data processing and graphical analysis were conducted using R (version 4.0.0, <http://cran.r-project.org>, accessed on 14 April 2020) and R-Studio (version v2021.09.0+354).

Structural compartmental PK models (one-, two-, and three-compartment) parametrized in apparent volume of distribution and clearances were selected to characterize the time-course of daptomycin. Due to the high plasma protein-binding capacity of daptomycin, several protein-binding relationships were evaluated to account for the unbound and total concentrations, which are described as follows [34,35]:

- Linear binding model [36,37]

$$C_{total} = C_{free} + C_{bound} = C_{free} + K_b \cdot C_{free} \quad (1)$$

where C_{total} (mg/L) is the total daptomycin concentration, C_{free} (mg/L) is the unbound daptomycin concentration, C_{bound} (mg/L) is the bound daptomycin concentration and K_b is the binding proportionality constant. This model assumes that C_{bound} is directly proportional to C_{free} with no saturable mechanism.

- Non-linear binding model with single site binding [38,39]:

$$C_{total} = C_{free} + B_{max} \cdot C_{free} / (K_D + C_{free}) \quad (2)$$

where B_{max} (mg/L) represents the maximal binding capacity and K_D (mg/L) is the equilibrium dissociation constant.

- Non-linear binding model with multiple site binding [38,40,41]

$$C_{total} = C_{free} + \sum B_{max,i} \cdot C_{free} / (K_{D,i} + C_{free}) \quad (3)$$

where $B_{max,i}$ and $K_{D,i}$ represent the binding-dissociation parameters of the i th binding site of plasma proteins.

Inter-individual variability (IIV) associated with the PK model parameters was modeled exponentially, and the residual unexplained variability (RUV) was explored using the additive error model on the logarithmic scale. The significance of the non-diagonal elements of the Ω variance-covariance matrix was evaluated.

Once the base model was selected, a graphical exploration of the correlations between the PK parameters (when eta-shrinkage was less than 35%) [42] and covariates was performed. The continuous covariate included age, height, body weight, serum creatinine, creatinine clearance, and albumin. The categorical covariates were gender and statin co-administration. A numerical evaluation of the parameter-covariate relationship was performed manually in a univariate testing.

The model selection was based on the physiological and pharmacological rationale with the principle of parsimony [43]. A decrease in 3.84 points of the objective function value (OFV) provided by NONMEM[®] between two nested models differing in one parameter was considered significant at the 5% level, together with the visual inspection of the goodness of fit (GOF) [44] and normalized prediction distribution error (NPDE) plots [45].

A model evaluation [46,47] was performed through prediction-corrected visual predictive checks (pc-VPC), [48] and a non-parametric bootstrap ($n = 1000$) was conducted to assess the statistical significance of the final parameter estimates.

Monte Carlo simulations ($n = 10,000$) were conducted assuming a log-normal distribution of PK parameters and a fixed combination of creatinine clearance (30, 60, 90 mL/min/1.73 m²) and body weight (50, 60, 70, 80, 90, and 100 kg). A 30-min intravenous infusion of daptomycin at 5–12 mg/kg every 12 (q12h) and 24 (q24h) hours or 10–17 mg/kg every 48 (q48h) hours were assumed.

The total area under the concentration curve (AUC) and unbound AUC (fAUC) were calculated after the first (AUC0-24) dose to represent the exposure at single dose conditions, respectively. Eight MIC values were considered: 0.25, 0.5, 0.75, 1, 1.5, 2, 3, and 4 mg/L.

A safety PK/PD index has been reported based on the trough levels (C_{min}) of daptomycin ≥ 24.3 mg/L, which are associated with a 50% probability of creatine phosphokinase elevation [49]. The probability of C_{min} ≥ 24.3 mg/L at 24 h was calculated for each scenario.

An optimal daptomycin dose selection was based on the probability of the target attainment (PTA) [50] $\geq 90\%$ and $\leq 20\%$ probability of achieving a C_{min} ≥ 24.3 mg/L for each scenario.

3. Results

3.1. Patient Population and Study Design

A total of 46 patients, including 157 daptomycin observations, were included in the analysis, which were logarithmically transformed for model development. Table 1 summarizes the baseline demographic, treatment and clinical data of the cohort included in the dataset.

Table 1. Summary of study design characteristics.

Characteristics		46 Patients Median (IQR)/n (%)
Demographics	Sex (Male) (n , %)	43 (93%)
	Age (years)	68 (59–81)
	Body weight (kg)	75 (65–85)
	Height (m)	1.7 (1.6–1.7)
	BMI (Kg/m ²)	25.9 (23.4–31.1)
Treatment	Dose (mg)	675 (500–765)
	Dose per kilogram (mg/kg)	9.1 (7.5–10.0)
	Treatment duration (days)	11 (7–15)
Clinical data	Serum albumin (g/dL)	2.9 (2.4–3.4)
	Serum protein (g/dL)	6.0 (5.0–6.4)
	Serum creatinine(g/dL)	0.9 (0.6–1.3)
	Creatinine clearance (mL/min/1.73 m ²)	93 (50–136)
	Renal function	
	>90 mL/min/1.73 m ²	16 (34.9%)
	60–89 mL/min/1.73 m ²	13 (28.2%)
30–59 mL/min/1.73 m ²	14 (30.4%)	
15–29 mL/min/1.73 m ²	3 (6.5%)	
<15 mL/min/1.73 m ²	0 (0%)	
Pathogenic micro-organism (36/46)	<i>S. aureus</i>	17 (47.2%)
	<i>S. epidermidis</i>	12 (33.3%)
	<i>S. hominis</i>	2 (5.6%)
	<i>E. fecalis</i>	2 (5.6%)
	<i>S. sacrohyticus</i>	2 (5.6%)
	<i>S. lugdunensis</i>	1 (2.7%)
MIC micro-organism		0.5 (0.25, 0.5)

IQR: interquartile range, n : number of individuals.

3.2. Population Pharmacokinetic Model

3.2.1. Base Population PK Model

A two-compartment model with first-order peripheral distribution and elimination kinetics using ordinary differential equations for unbound daptomycin concentrations was selected as the structural PK model (Figure 1).




Figure 1. Schematic representation of the final population PK model. IV: intravenous; CL: clearance; V1: apparent volume of distribution of the unbound central compartment; Q: intercompartmental clearance; V2: apparent volume of distribution of the unbound peripheral compartment; Bmax: maximal binding capacity; KD: equilibrium dissociation constant.

The total concentrations of daptomycin were derived by assuming non-linear protein-binding kinetics (Equation (2)). Non-linear protein binding was found to significantly improve the description of the data compared with a linear implementation ($\Delta\text{OFV} = -23$). The adequacy of the base model to capture the observed data can be found in Supplementary Material Figure S1 (GOF-NPDE) and Supplementary Material Figure S2 (pc-VPC).

3.2.2. Final Population PK Model

The covariate analysis estimated a moderate but statistically significant relationship ($\Delta\text{OFV} = -9.4$) between CLCR and CL, using a power function (0.19) centered around the median value (Equation (4)):

$$CL_i = CL \cdot \left(\frac{CLCR_i}{92.8} \right)^{0.19} \quad (4)$$

Additional continuous and categorical covariates were tested but did not result in a statistical ($\Delta\text{OFV} < 3.84$) reduction of the OFV.

The final population PK model shows a higher elimination clearance ($CL = 6.98 \text{ L/h}$) and an apparent volume of distribution of the central ($V1 = 0.95 \text{ L}$) compared to previous works using total daptomycin concentrations. This could be expected, since PK parameters are referred to unbound daptomycin concentrations. The peripheral distribution of unbound daptomycin represents a relevant PK process, based on the magnitude of the apparent volume of peripheral ($V2$) compartment and distribution (Q) clearance. A maximal binding capacity (B_{max}) and KD of 160 and 3.56 mg/L, respectively, were estimated.


Moderate inter-individual random effects were identified for clearance ($CL = 32\%$) and the apparent volume of the peripheral ($V2 = 47\%$) compartment and the RUV was 22%. Table 2 lists the parameters of the base and final population PK models, together with the bootstrap results. All PK parameters were statistically significant, as the 95% confidence intervals (95% CI) from the bootstrap analysis did not include the null value.

Table 2. Final population pharmacokinetic parameters of daptomycin in patients with normal renal function and renal impairment.

	Population PK Model Estimates			Bootstrap Results		
	Value	RSE (%)	Shrinkage (%)	Median	RSE * (%)	95%CI
<i>Fixed-Effect</i>						
CL (L/h)	6.98	14		7.01	15	[6.63–7.44]
V ₁ (L)	0.95	9		0.97	10	[0.92–1.09]
Q (L/h)	1.96	21		1.93	19	[1.43–2.48]
V ₂ (L/h)	21	19		20.5	21	[19.3–22.1]
B _{max} (mg/L)	160	26		157	24	[129–183]
K _D (mg/L)	3.56	15		3.61	12	[3.17–3.93]
CrCl on CL	0.19	12		0.19	13	[0.18–0.22]
<i>Inter-individual variability</i>						
CL (%)	32	11	12	33	10	[21–42]
V ₂ (%)	47	23	17	46	24	[52–94]
<i>Residual unexplained variability</i>						
Additive on Log-scale (%)	22	8	5	21	8	[18–24]

* RSE: relative standard error.

Due to the different doses administered across the study, a prediction-corrected VPC was conducted, showing an adequate characterization of the longitudinal profiles of the median and the dispersion of the data of the final population PK model (Figure 2). Supplementary Material Figure S3 shows the GOF and NPDE plots of the final population PK model.

**Figure 2.** Prediction-Corrected Visual Predictive check of the final population pharmacokinetic model of daptomycin. Grey lines represent the 2.5th, 50th and 97.5th experimental percentiles. Blue shaded areas represent the 95% prediction interval of the 2.5th, 50th and 97.5th percentiles. Empty grey dots represent the experimental daptomycin observations.

3.3. Population Pharmacokinetic/Pharmacodynamic Simulations and Optimal Dosage Selection

Several authors have reported the AUC of daptomycin with a bacteriostatic (465 mg·h/L) [19] or bactericidal (666 [24,51,52] and 761 mg·h/L [19]) effect for an MIC

equal to 1 mg/L. Based on the non-linear relationship (Figure 3) between total AUC and unbound AUC (fAUC), two efficacy PK/PD targets, defined as fAUC were considered: 59, and 107.5 mg·h/L.




Figure 3. Predicted relationship between total and unbound (fAUC) AUC of daptomycin. Grey and red dotted lines represent the bacteriostatic and bactericidal AUC values, respectively, for *Staphylococcus aureus* for a MIC value of 1 mg/L.

PTA simulations after daptomycin 5–12 mg/kg q24h administration assuming a bactericidal effect ($fAUC/MIC \geq 107.5$) for each combination of CLCR and body weight observed in this study, together with the probability of safety concerns ($C_{min} \geq 24.3$ mg/L), are depicted in Figure 4. PTA simulation after daptomycin 5–12 mg/kg q24h administration assuming a bacteriostatic effect ($fAUC/MIC \geq 59$) is represented in Supplementary Material Figure S4.

A dose-dependent effect is observed throughout the dose levels evaluated, indicating that lower doses may be required in patients with a high body weight and decreased renal function. A bactericidal effect ($fAUC/MIC \geq 107.5$) for Gram+ strains with $MIC \leq 0.5$ mg/L (Figure 4) could be achieved with 5–12 mg/kg daily daptomycin in patients weighing 50–100 kg, but different dose ranges would be recommended based on renal function: 5–10 mg/kg (CLCR = 30 mL/min/1.73 m²), 6–11 mg/kg (CLCR = 60 mL/min/1.73 m²), and 6–12 mg/kg (CLCR = 90 mL/min/1.73 m²). Although higher dose levels have been proposed to achieve a bactericidal effect, especially in patients at the low body weight range and with less impaired renal function, the probability of trough daptomycin levels above 24.3 mg/L was below 20%, demonstrating the adequate benefit/risk balance of the daptomycin q24h dosing regimens proposed for Gram+ strains with $MIC \leq 0.5$ mg/L. However, the results from the PTA analysis did not achieve the efficacy threshold when a bactericidal effect is expected for MRSA strains with $MIC \geq 1$ mg/L (Figure 4). High dose levels (>9 mg/kg in patients with a high body weight and/or low CLCR) would be recommended, but relevant safety concerns may appear in those scenarios (Figure 4).

Therefore, alternative dosing regimens (q12h or q48h) were evaluated to select the optimal daptomycin dosing regimens with a bactericidal effect for MRSA strains (Figures 5 and 6).




Figure 4. Probability of target attainment (PTA) after once daily (q24h) dose levels (5–12 mg/kg) of daptomycin in patients with different creatinine clearance (CLCR) and body weight (WT) for an (A) fAUC/MIC greater or equal to 107.5 (bactericidal effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L MIC: minimum inhibitory concentration; fAUC/MIC: unbound drug area under the concentration–time curve/minimum inhibitory concentration ratio.




Figure 5. Probability of target attainment (PTA) after twice daily (q12h) dose levels (5–12 mg/kg) of daptomycin in patients with different creatinine clearance (CLCR) and body weight (WT) for an

(A) $fAUC/MIC$ greater or equal to 107.5 (bactericidal effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L MIC: minimum inhibitory concentration; $fAUC/MIC$: unbound drug area under the concentration–time curve/minimum inhibitory concentration ratio.




Figure 6. Probability of target attainment (PTA) after once every two days (q48h) dose levels (5–12 mg/kg) of daptomycin in patients with different creatinine clearance (CLCR) and body weight (WT) for an (A) $fAUC/MIC$ greater or equal to 107.5 (bactericidal effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L MIC: minimum inhibitory concentration; $fAUC/MIC$: unbound drug area under the concentration–time curve/minimum inhibitory concentration ratio.

The global evaluation of the results obtained with the q12h regimen show high levels of PTA ($\geq 90\%$) in the range of doses evaluated (5–12 mg/kg), although administration every 12 h increases by well over 20% the probability of daptomycin trough levels with risk of CPK elevation. The administration of 10–17 mg/kg q48h daptomycin allows reaching PTA $\geq 90\%$ for MRSA Gram+ strains in patients with CLCR ≤ 30 mL/min/1.73 m² or body weight ≥ 70 kg. For patients with CLCR ≥ 60 mL/min/1.73 m² and body weight ≤ 60 kg, 17 mg/kg, q48h allows to achieve PTA 60–85% with a probability of trough levels less than 11%. In this sense, increasing the dose and spacing its administration at 48-h intervals would favor a satisfactory response rate in most subgroups evaluated, as well as a low probability of associated muscle toxicity. Supplementary Material Figures S5 and S6 depict the PTA simulations of daptomycin q12h and q48h assuming bacteriostatic effect.

Distribution of trough levels of daptomycin q12h, q24h or q48h obtained from the Monte Carlo simulation are represented in Supplementary Material Figures S7–S9.

4. Discussion

The selection of the optimal antibiotic treatment in patients with renal impairment represents one of the greatest challenges currently facing clinical teams. This is due to the increasingly frequent appearance of resistant strains (highly resistant *S. aureus*, methicillin-resistant *S. aureus*, and *E. faecium*, which leads to severe or life-threatening conditions [24,27,53]. The increase of PK and PD knowledge of daptomycin, together with the commitment to make a more rational and efficient decision-making process, has

allowed for the development of model-informed strategies. This has become a fundamental tool for the selection of the optimal dosing regimen for daptomycin. In this sense, a two-compartment population PK model of the unbound fraction of daptomycin, based on a non-linear binding relationship with the total (observed) daptomycin concentrations, provides new insights into the PK properties of this drug to guide dose selection of daptomycin in patients with several degrees of renal impairment and body weight ranges. In addition, the model can inform an optimal PK sampling scenario to evaluate individual safety and efficacy in daptomycin TDM.

The structural PK parameters estimated in this study are referred to the predicted unbound fraction of daptomycin. If a 10% unbound fraction of daptomycin is assumed, the CL ($CL = CL_u \times f_u$) and Q ($Q = Q_u \times f_u$) of total daptomycin concentrations are in agreement with the previously reported values [25]. Daptomycin is highly (90%) reversibly bound to albumin and α -glycoprotein [24,53]. The variation of the latter, especially in patients with renal impairment, may affect the unbound daptomycin concentration. Therefore, the characterization of the unbound fraction of daptomycin is of special relevance, since it is the active form of the drug that is able to exert the pharmacological response [25]. Thus, a PK/PD target based on unbound concentrations is more suitable for a target attainment assessment [35]. In previous works, the dose selection was based on the total daptomycin levels or assuming a linear unbound fraction of 10% for PK/PD indexes, which could lead to biased conclusions due to differences in protein binding and the apparent volume of distribution that have been reported in patients with impaired renal function [3,5]. In this work, we have developed a population PK model of daptomycin, which considers a non-linear binding kinetics to human plasma proteins, that allows to predict free daptomycin concentrations from total daptomycin values. Although albumin levels could not be related to the binding mechanism, the results show a high binding affinity of daptomycin to plasma proteins ($K_D = 1.96$ mg/L), with a maximal binding capacity reached at 160 mg/L. The maximal binding capacity reflects the concentration of the binding protein and the number of available binding sites [54]. Based on the average albumin concentration (2.95 g/L), it seems reasonable to suggest that albumin is the primary binding site of daptomycin in a 1:1 molar relationship. At the range of usual concentrations of daptomycin, saturable binding conditions appear. This leads to a more than proportional increase in fAUC as AUC increases, that, if not considered, may affect the selection of an optimal dosage regimen.

A concentration-dependent antibiotic effectiveness of daptomycin was characterized that is best correlated to AUC/MIC [52]. The selected AUC/MIC values (465 and 761) represent the most favorable and restrictive scenarios. Their corresponding fAUC/MIC were derived (Figure 3) that allows the covering of a PK/PD range in order to explore the selection of the dosage regimen according to the desired therapeutic objective. To reach a bacteriostatic effect (Table 3), a dose recommendation of 5–7 mg/kg for Gram-positive strains ($MIC \leq 0.5$ mg/L) in patients with body weight ≥ 50 kg and 5–11 mg/kg for MRSA strains once daily of daptomycin in ≥ 60 kg is in agreement with previous reports [55–63].

The schedule optimization with bactericidal effect for Gram-positive strains with $MIC \leq 0.5$ mg/L, suggested that safe and efficacious 5–12 mg/kg once daily daptomycin dosing regimens can be proposed in patients from 50 to 100 kg and different degrees of CLCR (Table 3).

Given the appearance of more resistant strains ($MIC \geq 1$ mg/L) in patients with normal renal function, PTA results (Figure 4) suggested daily regimens of daptomycin between 9 and 12 mg/kg in patients with a high body weight (≥ 80 kg) or moderate renal impairment ($CLCR = 30$ mL/min/1.73 m²) resulted in a probability of trough concentration ≥ 24.3 mg/L higher than 20%. This is correlated with CPK elevations and, therefore, muscle toxicity [1–3,5] (Figure 5). These results are in agreement with previous works, suggesting that higher dose levels than the approved ones of daptomycin may be required [19,21,49,64–66]. Therefore, it is evident for the need to explore more frequent schemes with lower dosage levels in order to guarantee optimal PTA levels and safety. According to the results from Table 3, the treatment of infections with MRSA Gram-positive strains requires higher dose levels every 48 h

in order to achieve adequate efficacy (PTA > 90%) and safety (C_{min} > 24.3 mg/L) objectives. Intensive dosing regimens (q12h) resulted in a high probability of trough concentrations of daptomycin above the safety threshold (Figure 6). Alternatively, it may be necessary to evaluate fixed-dose schemes that allow reducing the differences in the magnitude of doses associated with body weight. The risk of over-exposure in low weight patients is minor, and the under-exposure associated with the fixed dose that would lead to risks of inefficacy in heavy patients could be of little clinical relevance based on the PTA results.

Table 3. Daptomycin dosing schedules proposed based on the MIC, renal function and body weight of the patient to achieve bactericidal effect.

	Moderate Renal Impairment (CL _{CR} = 30 mL/min/1.73 m ²)	Mild Renal Impairment (CL _{CR} = 60 mL/min/1.73 m ²)	Normal Renal Function (CL _{CR} = 90 mL/min/1.73 m ²)	Body Weight
MIC ≤ 0.5 mg/L	10 mg/kg q24h	11 mg/kg q24h	12 mg/kg q24h	50 kg
	9 mg/kg q24h	10 mg/kg q24h	10 mg/kg q24h	60 kg
	8 mg/kg q24h	9 mg/kg q24h	9 mg/kg q24h	70 kg
	7 mg/kg q24h	7 mg/kg q24h	8 mg/kg q24h	80 kg
	6 mg/kg q24h	7 mg/kg q24h	7 mg/kg q24h	90 kg
	5 mg/kg q24h	6 mg/kg q24h	6 mg/kg q24h	100 kg
MIC ≤ 1 mg/L	17* mg/kg q48h	17* mg/kg q48h	17* mg/kg q48h	50 kg
	16 mg/kg q48h	17* mg/kg q48h	17* mg/kg q48h	60 kg
	14 mg/kg q48h	16 mg/kg q48h	17 mg/kg q48h	70 kg
	12 mg/kg q48h	14 mg/kg q48h	15 mg/kg q48h	80 kg
	11 mg/kg q48h	12 mg/kg q48h	13 mg/kg q48h	90 kg
	10 mg/kg q48h	11 mg/kg q48h	12 mg/kg q48h	100 kg

MIC: minimum inhibitory concentration; CL_{CR}: creatinine clearance; * PTA results <90%.

Among the main limitations of this analysis, it is important to highlight the absence of observations of free daptomycin that would allow us to identify with greater certainty the degree of binding to plasma proteins of daptomycin *in vivo*. However, similar structural PK models have been used to characterize drug binding kinetics based on total concentrations [67,68]. On the other hand, a greater number of observations per individual, as well as the availability of patients with severe renal insufficiency, would be of great relevance for the characterization of the influence of CL_{CR} on clearance and, with it, the design of dosage schemes in this subgroup of patients. In addition, the lack of women (7%) enrolled in this study due to the study design could limit the evaluation of the gender effect on the PK parameters. Therefore, additional studies, including a more balanced distribution of sub-groups of interest, are encouraged to evaluate their dosing recommendations of daptomycin.

5. Conclusions

In conclusion, the two-compartment population PK model has first allowed us to characterize the non-linear binding kinetics of daptomycin in patients with renal impairment. This served to evaluate the impact of different dosing regimens on the efficacy and safety outcomes of daptomycin treatment, based on the unbound exposure of daptomycin and individual patient characteristics.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/pharmaceutics14102226/s1>, Figure S1: Goodness-of-fit plots of the base population pharmacokinetic model of daptomycin; Figure S2: VPC of the base population pharmacokinetic model of daptomycin; Figure S3: Goodness-of-fit plots of the final population pharmacokinetic model of daptomycin; Figure S4: PTA after once daily (q24h) dose levels (5–12 mg/kg) of daptomycin in patients with different CL_{CR} and WT for an (A) fAUC/MIC greater or equal to 59 (bacteriostatic effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L; Figure S5: PTA after twice daily (q12h) dose levels (5–12 mg/kg) of daptomycin in patients with different CL_{CR} and WT for an (A) fAUC/MIC greater or equal to 59 (bacteriostatic effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L; Figure S6: PTA after once every other day (q48h) dose levels (5–12 mg/kg) of daptomycin in patients with different CL_{CR} and WT for an (A) fAUC/MIC greater

or equal to 59 (bacteriostatic effect), and (B) Predicted probability of achieving trough concentrations of total daptomycin greater or equal to 24.3 mg/L; Figure S7: Daptomycin trough concentrations (C_{min}) after single dose levels (5–12 mg/kg) of daptomycin every 24 h in patients with different CL_{CR} and WT; Figure S8: Daptomycin trough concentrations (C_{min}) after single dose levels (5–12 mg/kg) of daptomycin every 12 h in patients with different CL_{CR} and WT; Figure S9: Daptomycin trough concentrations (C_{min}) after single dose levels (10–17 mg/kg) of daptomycin every 48 h in patients with different CL_{CR} and WT.

Author Contributions: Conceptualization, T.G.-M., M.D.B.-M., R.F.-P., V.M.-S. and M.M.-S.; methodology T.G.-M., M.D.B.-M. and V.M.-S.; software, V.M.-S. and M.G.-C.; validation, M.M.-S., V.M.-S. and M.G.-C.; formal analysis, T.G.-M., M.G.-C. and V.M.-S.; investigation, T.G.-M., M.D.B.-M., V.M.-S. and M.M.-S.; resources, T.G.-M., R.F.-P. and V.M.-S.; data curation, T.G.-M. and M.D.B.-M.; writing—original draft preparation, T.G.-M. and V.M.-S.; writing—review and editing, T.G.-M. and V.M.-S.; visualization, T.G.-M. and V.M.-S.; supervision, M.D.B.-M., M.M.-S. and R.F.-P.; project administration, DB., M.M.-S. and R.F.-P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the local (General University Hospital of Castellón) and regional (Valencian Community) Ethics Committee (protocol code: FAR-DAP-2017-01, 28 May 2018). Procedures were approved by each participating investigator's Institutional Review Board for Human Investigation.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: The authors acknowledge the investigators and patients who participated in the clinical trial.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. US FDA Daptomycin (Marketed as Cubicin) Information. 2017. Available online: https://www.accessdata.fda.gov/drugsatfda_docs/Label/2017/021572s059lbl.pdf (accessed on 10 April 2022).
2. European Medicines Agency. *Cubicin: EPAR—Product Information (WC500036049.pdf)*; European Medicines Agency: London, UK, 2006.
3. European Medicines Agency. *Cubicin: EPAR—Scientific Discussion (WC500036046.pdf)*; European Medicines Agency: London, UK, 2006.
4. Vilhena, C.; Bettencourt, A. Daptomycin: A Review of Properties, Clinical Use, Drug Delivery and Resistance. *Mini.-Rev. Med. Chem.* **2012**, *12*, 202–209. [[CrossRef](#)] [[PubMed](#)]
5. European Medicines Agency. *Cubicin (Daptomycin); EMEA/H/C/000637*; European Medicines Agency: London, UK, 2006.
6. Avolio, A.D.; Pensi, D.; Baietto, L.; Pacini, G.; Di Perri, G.; Francesco, P.; de Rosa, G. Daptomycin Pharmacokinetics and Pharmacodynamics in Septic and Critically Ill Patients. *Drugs* **2016**, *76*, 1161–1174. [[CrossRef](#)]
7. Soraluca, A.; Asín-Prieto, E.; Rodríguez-Gascón, A.; Isla, A.; Barrasa, H.; Maynar, J.; Carcelero, E.; Soy, D. Population Pharmacokinetics of Daptomycin in Critically Ill Patients. *Int. J. Antimicrob. Agents* **2018**, *52*, 158–165. [[CrossRef](#)] [[PubMed](#)]
8. Nakafusa, J.; Misago, N.; Miura, Y.; Kayaba, M.; Tanaka, T.; Narisawa, Y. The Importance of Serum Creatine Phosphokinase Level in the Early Diagnosis, and as a Prognostic Factor, of *Vibrio Vulnificus* Infection. *Br. J. Dermatol.* **2001**, *145*, 280–284. [[CrossRef](#)] [[PubMed](#)]
9. DN Gilbert. *The Sanford Guide to Antimicrobial Therapy*; BI Publications Pvt Ltd.: New Delhi, India, 2006.
10. Canut, A.; Isla, A.; Betriu, C.; Gascón, A.R. Pharmacokinetic—Pharmacodynamic Evaluation of Daptomycin, Tigecycline, and Linezolid versus Vancomycin for the Treatment of MRSA Infections in Four Western European Countries. *Eur. J. Clin. Microbiol.* **2012**, *31*, 2227–2235. [[CrossRef](#)]
11. Ukimura, A.; Oda, K.; Yoshida, M.; Nishihara, M.; Kawanishi, F.; Yamada, T.; Nakano, T.; Ooi, Y.; Uchida, T.; Shibata, Y.; et al. Observational Study to Determine the Optimal Dose of Daptomycin Based on Pharmacokinetic/Pharmacodynamic Analysis. *J. Infect. Chemother.* **2019**, *26*, 379–384. [[CrossRef](#)]
12. Gray, D.A.; Wenzel, M. More Than a Pore: A Current Perspective on the In Vivo Mode of Action of the Lipopeptide Antibiotic Daptomycin. *Antibiotics* **2020**, *9*, 17. [[CrossRef](#)]
13. EUCAST Daptomycin: Rational for the EUCAST Clinical Breakpoints, Version 2. Mayo 2021. Available online: https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Rationale_documents/Daptomycin_Rationale_Document_2_0_20210512.pdf (accessed on 31 May 2021).

14. Trecarichi, E.M.; Pagano, L.; Candoni, A.; Pastore, D.; Cattaneo, C.; Fanci, R.; Nosari, A.; Caira, M.; Spadea, A.; Busca, A.; et al. Current Epidemiology and Antimicrobial Resistance Data for Bacterial Bloodstream Infections in Patients with Hematologic Malignancies: An Italian Multicentre Prospective Survey. *Clin. Microbiol. Infect.* **2015**, *21*, 337–343. [[CrossRef](#)]
15. Rolston, K.V.I.; Yadegarynia, D.; Kontoyiannis, D.P.; Raad, I.I.; Ho, D.H. The Spectrum of Gram-Positive Bloodstream Infections in Patients with Hematologic Malignancies, and the in Vitro Activity of Various Quinolones against Gram-Positive Bacteria Isolated from Cancer Patients. *Int. J. Infect. Dis.* **2006**, *10*, 223–230. [[CrossRef](#)]
16. Benvenuto, M.; Benziger, D.P.; Yankelev, S.; Vigliani, G. Pharmacokinetics and Tolerability of Daptomycin at Doses up to 12 Milligrams per Kilogram of Body Weight Once Daily in Healthy Volunteers. *Antimicrob. Agents Chemother.* **2006**, *50*, 3245–3249. [[CrossRef](#)]
17. Dvorchik, B.; Arbeit, R.D.; Chung, J.; Liu, S.; Knebel, W.; Kastrissios, H. Population Pharmacokinetics of Daptomycin. *Antimicrob. Agents Chemother.* **2008**, *48*, 2799–2807. [[CrossRef](#)] [[PubMed](#)]
18. di Paolo, A.; Tascini, C.; Polillo, M.; Gemignani, G.; Nielsen, E.I.; Bocci, G.; Karlsson, M.O.; Menichetti, F.; Danesi, R. Population Pharmacokinetics of Daptomycin in Patients Affected by Severe Gram-Positive Infections. *Int. J. Antimicrob. Agents* **2013**, *42*, 250–255. [[CrossRef](#)]
19. Chaves, R.L.; Chakraborty, A.; Benziger, D.; Tannenbaum, S. Clinical and Pharmacokinetic Considerations for the Use of Daptomycin in Patients with Staphylococcus Aureus Bacteraemia and Severe Renal Impairment. *J. Antimicrob. Chemother.* **2014**, *69*, 200–210. [[CrossRef](#)] [[PubMed](#)]
20. Soon, R.L.; Turner, S.J.; Forrest, A.; Tsuji, B.T.; Brown, J. Pharmacokinetic/Pharmacodynamic Evaluation of the Efficacy and Safety of Daptomycin against Staphylococcus Aureus. *Int. J. Antimicrob. Agents* **2013**, *42*, 53–58. [[CrossRef](#)]
21. Xu, X.; Khadzhynov, D.; Peters, H.; Chaves, R.L.; Hamed, K.; Levi, M.; Corti, N. Population Pharmacokinetics of Daptomycin in Adult Patients Undergoing Continuous Renal Replacement Therapy. *Br. J. Clin. Pharmacol.* **2017**, *83*, 498–509. [[CrossRef](#)]
22. Xie, F.; Li, S.; Cheng, Z. Population Pharmacokinetics and Dosing Considerations of Daptomycin in Critically Ill Patients Undergoing Continuous Renal Replacement Therapy. *J. Antimicrob. Chemother.* **2020**, *75*, 1559–1566. [[CrossRef](#)] [[PubMed](#)]
23. Abdul-Aziz, M.H.; Alffenaar, J.-W.C.; Bassetti, M.; Bracht, H.; Dimopoulos, G.; Marriott, D.; Neely, M.N.; Paiva, J.-A.; Pea, F.; Sjøvall, F.; et al. Antimicrobial Therapeutic Drug Monitoring in Critically Ill Adult Patients: A Position Paper. *Intensive Care Med.* **2020**, *46*, 1127–1153. [[CrossRef](#)] [[PubMed](#)]
24. Lou, Y.; Liu, Y.; Wang, J.; Cai, L.; He, L.; Yang, X.; Xu, H.; He, X.; Yang, X.; Wei, C.; et al. Population Pharmacokinetics and Individual Analysis of Daptomycin in Kidney Transplant Recipients. *Eur. J. Pharm. Sci.* **2021**, *162*, 105818. [[CrossRef](#)]
25. Grégoire, N.; Marchand, S.; Ferrandière, M.; Lasocki, S.; Seguin, P.; Vourc'h, M.; Barbaz, M.; Gaillard, T.; Launey, Y.; Asehnoune, K.; et al. Population Pharmacokinetics of Daptomycin in Critically Ill Patients with Various Degrees of Renal Impairment. *J. Antimicrob. Chemother.* **2018**, *74*, 117–125. [[CrossRef](#)]
26. Geriak, M.; Haddad, F.; Rizvi, K.; Rose, W.; Kullar, R.; LaPlante, K.; Yu, M.; Vasina, L.; Ouellette, K.; Zervos, M.; et al. Clinical Data on Daptomycin plus Ceftaroline versus Standard of Care Monotherapy in the Treatment of Methicillin-Resistant Staphylococcus Aureus Bacteremia. *Antimicrob. Agents Chemother.* **2019**, *63*, e02483-18. [[CrossRef](#)]
27. Wei, X.; Zhao, M.; Li, X.; Xiao, X. Pharmacokinetic/Pharmacodynamic Analysis of Daptomycin Against *Staphylococcus aureus* and *Enterococcus faecium* in Pediatric Patients by Monte Carlo Simulation. *J. Clin. Pharmacol.* **2020**, *60*, 768–774. [[CrossRef](#)] [[PubMed](#)]
28. Satlin, M.J.; Nicolau, D.P.; Humphries, R.M.; Kuti, J.L.; Campeau, S.A.; Lewis II, J.S.; Weinstein, M.P.; Jorgensen, J.H. Development of Daptomycin Susceptibility Breakpoints for Enterococcus Faecium and Revision of the Breakpoints for Other Enterococcal Species by the Clinical and Laboratory Standards Institute. *Clin. Infect. Dis.* **2019**, *70*, 1240–1246. [[CrossRef](#)] [[PubMed](#)]
29. Ogami, C.; Tsuji, Y.; Kasai, H.; Hiraki, Y.; Yamamoto, Y. Evaluation of Pharmacokinetics and the Stability of Daptomycin in Serum at Various Temperatures. *Int. J. Infect. Dis.* **2017**, *57*, 38–43. [[CrossRef](#)] [[PubMed](#)]
30. Cockcroft, D.W.; Gault, H. Prediction of Creatinine Clearance from Serum Creatinine. *Nephron* **1976**, *16*, 31–41. [[CrossRef](#)]
31. Tobin, C.M.; Darville, J.M.; Lovering, A.M.; MacGowan, A.P. An HPLC Assay for Daptomycin in Serum. *J. Antimicrob. Chemother.* **2008**, *62*, 1462–1463. [[CrossRef](#)]
32. Tanaka, R.; Suzuki, Y.; Goto, K.; Yasuda, N.; Koga, H.; Kai, S.; Ohchi, Y.; Sato, Y.; Kitano, T.; Itoh, H. Development and Validation of Sensitive and Selective Quantification of Total and Free Daptomycin in Human Plasma Using Ultra-Performance Liquid Chromatography Coupled to Tandem Mass Spectrometry. *J. Pharm. Biomed. Anal.* **2019**, *165*, 56–64. [[CrossRef](#)]
33. Lindbom, L.; Pihlgren, P.; Jonsson, N. PsN-Toolkit—A Collection of Computer Intensive Statistical Methods for Non-Linear Mixed Effect Modeling Using NONMEM. *Comput. Methods Programs Biomed.* **2005**, *79*, 241–257. [[CrossRef](#)]
34. Jun, H.; Rong, Y.; Yih, C.; Ho, J.; Cheng, W.; Kiang, T.K.L. Comparisons of Four Protein-Binding Models Characterizing the Pharmacokinetics of Unbound Phenytoin in Adult Patients Using Non-Linear Mixed-Effects Modeling. *Drugs R & D* **2020**, *20*, 343–358. [[CrossRef](#)]
35. Aulin, L.B.S.; de Paepe, P.; Dhont, E.; de Jaeger, A.; vande Walle, J.; Vandenberghe, W.; McWhinney, B.C.; Ungerer, J.P.J.; van Hasselt, J.G.C.; de Cock, P.A.J.G. Population Pharmacokinetics of Unbound and Total Teicoplanin in Critically Ill Pediatric Patients. *Clin. Pharmacokinet.* **2021**, *60*, 353–363. [[CrossRef](#)]
36. Charles, B.; Norris, R.; Xiao, X.; Hague, W. Population Pharmacokinetics of Metformin in Late Pregnancy. *Ther. Drug Monit.* **2006**, *28*, 67–72. [[CrossRef](#)]
37. Hennig, S.; Norris, R.; Tu, Q.; van Breda, K.; Riney, K.; Foster, K.; Lister, B.; Charles, B. Population Pharmacokinetics of Phenytoin in Critically Ill Children. *J. Clin. Pharmacol.* **2015**, *55*, 355–364. [[CrossRef](#)] [[PubMed](#)]

38. Toutain, P.L.; Bousquet-Melou, A. Free Drug Fraction vs. Free Drug Concentration: A Matter of Frequent Confusion. *J. Vet. Pharmacol. Ther.* **2002**, *25*, 460–463. [[CrossRef](#)] [[PubMed](#)]
39. Heine, R.; Kane, S.P.; Huitema, A.D.R.; Krasowski, M.D.; Maarseveen, E.M. Nonlinear Protein Binding of Phenytoin in Clinical Practice: Development and Validation of a Mechanistic Prediction Model. *Br. J. Clin. Pharmacol.* **2019**, *85*, 2360–2368. [[CrossRef](#)] [[PubMed](#)]
40. Chen, J.; Ohnmacht, C.; Hage, D.S. Studies of Phenytoin Binding to Human Serum Albumin by High-Performance Affinity Chromatography. *J. Chromatogr. B* **2004**, *809*, 137–145. [[CrossRef](#)]
41. Musteata, F.M. Calculation of Normalized Drug Concentrations in the Presence of Altered Plasma Protein Binding. *Clin. Pharmacokinet.* **2012**, *51*, 55–68. [[CrossRef](#)]
42. Savic, R.M.; Karlsson, M.O. Importance of Shrinkage in Empirical Bayes Estimates for Diagnostics: Problems and Solutions. *AAPS J.* **2009**, *11*, 558–569. [[CrossRef](#)]
43. Wade, J.R.; Beal, S.L.; Sambol, N.C. Interaction between Structural, Statistical, and Covariate Models in Population Pharmacokinetic Analysis. *J. Pharmacokinet. Biopharm.* **1994**, *22*, 165–177. [[CrossRef](#)]
44. Hooker, A.C.; Staatz, C.E.; Karlsson, M.O. Conditional Weighted Residuals (CWRES): A Model Diagnostic for the FOCE Method. *Pharm. Res.* **2007**, *24*, 2187–2197. [[CrossRef](#)]
45. Comets, E.; Brendel, K.; Mentré, F. Computing Normalised Prediction Distribution Errors to Evaluate Nonlinear Mixed-Effect Models: The Npde Add-on Package for R. *Comput. Methods Programs Biomed.* **2008**, *90*, 154–166. [[CrossRef](#)]
46. Sherwin, C.M.T.; Kiang, T.K.L.; Spigarelli, M.G.; Ensom, M.H.H. Fundamentals of Population Pharmacokinetic Modelling. *Clin. Pharmacokinet.* **2012**, *51*, 573–590. [[CrossRef](#)]
47. Kiang, T.K.L.; Sherwin, C.M.T.; Spigarelli, M.G.; Ensom, M.H.H. Fundamentals of Population Pharmacokinetic Modelling. *Clin. Pharmacokinet.* **2012**, *51*, 515–525. [[CrossRef](#)]
48. Bergstrand, M.; Hooker, A.C.; Wallin, J.E.; Karlsson, M.O. Prediction-Corrected Visual Predictive Checks for Diagnosing Nonlinear Mixed-Effects Models. *AAPS J.* **2011**, *13*, 143–151. [[CrossRef](#)] [[PubMed](#)]
49. Bhavnani, S.M.; Rubino, C.M.; Ambrose, P.G.; Drusano, G.L. Daptomycin Exposure and the Probability of Elevations in the Creatine Phosphokinase Level: Data from a Randomized Trial of Patients with Bacteremia and Endocarditis. *Clin. Infect. Dis.* **2010**, *12208*, 1568–1574. [[CrossRef](#)] [[PubMed](#)]
50. Mouton, J.W.; Dudley, M.N.; Cars, O.; Derendorf, H.; Drusano, G.L. Standardization of Pharmacokinetic/Pharmacodynamic (PK/PD) Terminology for Anti-Infective Drugs: An Update. *J. Antimicrob. Chemother.* **2005**, *55*, 601–607. [[CrossRef](#)] [[PubMed](#)]
51. Tsukada, H.; Tsuji, Y.; Yamashina, T.; Tsuruta, M.; Hiraki, Y.; Tsuruyama, M.; Ogami, C.; Kawasuji, H.; Sakamaki, I.; Yamamoto, Y. Pharmacokinetics and Pharmacodynamics of Daptomycin in a Clinical Setting. *J. Infect. Chemother.* **2020**, *26*, 230–235. [[CrossRef](#)] [[PubMed](#)]
52. Safdar, N.; Andes, D.; Craig, W.A. In Vivo Pharmacodynamic Activity of Daptomycin. *Antimicrob. Agents Chemother.* **2004**, *48*, 63–68. [[CrossRef](#)]
53. Schneider, E.K.; Huang, J.X.; Carbone, V.; Han, M.; Zhu, Y.; Nang, S.; Khoo, K.K.; Mak, J.; Cooper, M.A.; Li, J.; et al. Plasma Protein Binding Structure-Activity Relationships Related to the N-Terminus of Daptomycin. *ACS Infect. Dis.* **2017**, *3*, 249–258. [[CrossRef](#)]
54. Sellers, E.M.; Koch-Weser, J. Clinical Implications of Drug-Albumin Interaction. In *Albumin: Structure, Function and Uses*; Rosenoer, V.M., Oratz, M., Rothschild, M.A., Eds.; Pergamon Press: Oxford, UK, 1977; pp. 159–182.
55. Lai, C.-C.; Sheng, W.-H.; Wang, J.-T.; Cheng, A.; Chuang, Y.-C.; Chen, Y.-C.; Chang, S.-C. Safety and Efficacy of High-Dose Daptomycin as Salvage Therapy for Severe Gram-Positive Bacterial Sepsis in Hospitalized Adult Patients. *BMC Infect. Dis.* **2013**, *13*, 66. [[CrossRef](#)]
56. Durante-Mangoni, E.; Andini, R.; Parrella, A.; Mattucci, I.; Cavezza, G.; Senese, A.; Trojaniello, C.; Caprioli, R.; Diana, M.V.; Utili, R. Safety of Treatment with High-Dose Daptomycin in 102 Patients with Infective Endocarditis. *Int. J. Antimicrob. Agents* **2016**, *48*, 61–68. [[CrossRef](#)]
57. Casapao, A.M.; Kullar, R.; Davis, S.L.; Levine, D.P.; Zhao, J.J.; Potoski, B.A.; Goff, D.A.; Crank, C.W.; Segreti, J.; Sakoulas, G.; et al. Multicenter Study of High-Dose Daptomycin for Treatment of Enterococcal Infections. *Antimicrob. Agents Chemother.* **2013**, *57*, 4190–4196. [[CrossRef](#)]
58. Kullar, R.; Casapao, A.M.; Davis, S.L.; Levine, D.P.; Zhao, J.J.; Crank, C.W.; Segreti, J.; Sakoulas, G.; Cosgrove, S.E.; Rybak, M.J. A Multicentre Evaluation of the Effectiveness and Safety of High-Dose Daptomycin for the Treatment of Infective Endocarditis. *J. Antimicrob. Chemother.* **2013**, *68*, 2921–2926. [[CrossRef](#)] [[PubMed](#)]
59. Katz, D.E.; Lindfield, K.C.; Steenbergen, J.N.; Benziger, D.P.; Blackerby, K.J.; Knapp, A.G.; Martone, W.J. A Pilot Study of High-Dose Short Duration Daptomycin for the Treatment of Patients with Complicated Skin and Skin Structure Infections Caused by Gram-Positive Bacteria. *Int. J. Clin. Pract.* **2008**, *62*, 1455–1464. [[CrossRef](#)] [[PubMed](#)]
60. Moise, P.A.; Hershberger, E.; Amodio-Groton, M.I.; Lamp, K.C. Safety and Clinical Outcomes When Utilizing High-Dose (≥ 8 Mg/Kg) Daptomycin Therapy. *Ann. Pharmacother.* **2009**, *43*, 1211–1219. [[CrossRef](#)] [[PubMed](#)]
61. Bassetti, M.; Nicco, E.; Ginocchio, F.; Ansaldi, F.; de Florentiis, D.; Viscoli, C. High-Dose Daptomycin in Documented Staphylococcus Aureus Infections. *Int. J. Antimicrob. Agents* **2010**, *36*, 459–461. [[CrossRef](#)]
62. Parra-Ruiz, J.; Peña-Monje, A.; Tomás-Jiménez, C.; Pomares-Mora, J.; Hernández-Quero, J. Eficacia y Seguridad de Daptomicina En Dosis Elevadas (≥ 8 mg/Kg/Día). *Enferm. Infecc. Microbiol. Clin.* **2011**, *29*, 425–427. [[CrossRef](#)]

63. Kullar, R.; Davis, S.L.; Levine, D.P.; Zhao, J.J.; Crank, C.W.; Segreti, J.; Sakoulas, G.; Cosgrove, S.E.; Rybak, M.J. High-Dose Daptomycin for Treatment of Complicated Gram-Positive Infections: A Large, Multicenter, Retrospective Study. *Pharmacotherapy* **2011**, *31*, 527–536. [[CrossRef](#)]
64. Gregoire, N.; Chauzy, A.; Buyck, J.; Rammaert, B.; Couet, W.; Marchand, S. Clinical Pharmacokinetics of Daptomycin. *Clin. Pharmacokinet.* **2020**, *60*, 271–281. [[CrossRef](#)]
65. Reiber, C.; Senn, O.; Müller, D.; Kullak-ublick, G.A. Therapeutic Drug Monitoring of Daptomycin: A Retrospective Monocentric Analysis. *Ther. Drug Monit.* **2015**, *37*, 634–640. [[CrossRef](#)]
66. Cojutti, P.G.; Candoni, A.; Ramos-martin, V.; Lazzarotto, D.; Zannier, M.E.; Fanin, R.; Hope, W.; Pea, F.; Carlo, M. Population Pharmacokinetics and Dosing Considerations for the Use of Daptomycin in Adult Patients with Haematological Malignancies. *J. Antimicrob. Chemother.* **2017**, *72*, 2342–2350. [[CrossRef](#)]
67. Campos Moreno, E.; Merino Sanjuán, M.; Merino, V.; Nácher, A.; Algarra, R.V.M.; Casabó, V.G. Population Modelling to Describe Pharmacokinetics of Amiodarone in Rats: Relevance of Plasma Protein and Tissue Depot Binding. *Eur. J. Pharm. Sci.* **2007**, *30*, 190–197. [[CrossRef](#)]
68. Rodríguez-Fernández, K.; Gras-Colomer, E.; Climente-Martí, M.; Mangas-Sanjuán, V.; Merino-Sanjuán, M. Pharmacometric Characterization of Entero-Hepatic Circulation Processes of Orally Administered Formulations of Amiodarone under Complex Binding Kinetics. *Eur. J. Pharm. Sci.* **2022**, *174*, 106198. [[CrossRef](#)] [[PubMed](#)]