

RESEARCH ARTICLE

Rapid weight reduction does not impair athletic performance of Taekwondo athletes – A pilot study

Woo-Hwi Yang^{1,2*}, Oliver Heine², Marijke Grau¹

1 Department of Molecular and Cellular Sport Medicine, Institute of Cardiovascular Research and Sport Medicine, German Sport University Cologne, Cologne, Germany, **2** Olympic Training Centre Rhineland, Cologne, Germany

* ywh1235@hotmail.de



OPEN ACCESS

Citation: Yang W-H, Heine O, Grau M (2018) Rapid weight reduction does not impair athletic performance of Taekwondo athletes – A pilot study. PLoS ONE 13(4): e0196568. <https://doi.org/10.1371/journal.pone.0196568>

Editor: Clemens Fürnsinn, Medical University of Vienna, AUSTRIA

Received: September 19, 2017

Accepted: April 16, 2018

Published: April 26, 2018

Copyright: © 2018 Yang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be made publicly available due to ethical restrictions imposed by the privacy policy of German Sport University Cologne. Furthermore, participants did not provide consent for data sharing. Interested, qualified researchers may request the data by contacting our institute at sportmedizin@dshs-koeln.de.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abstract

In combat sports such as taekwondo (TKD), athletes rapidly reduce body weight to achieve a desired weight category. Competition takes place 16–24 h after weigh-in and thus, the recovery time is an important factor for competition performance. The purpose of this study was to investigate the impact of rapid weight reduction (RWR) on athletic performance and associated hemorheological properties considering relevant recovery time. Five male TKD athletes reduced body weight by 5% within 3½ days. A simulated competition day (SCD) was carried out after a 16 h recovery period. Parameters were measured before RWR, at weigh-in and before and after three TKD simulation matches (SMs) at SCD. Same set-up was conducted but without RWR as control. Basal blood parameters, red blood cells (RBC) deformability and aggregation, serum glucose and fibrinogen were determined. During SMs, heart rate (HR_{peak} , HR_{mean}), oxygen uptake (VO_{2peak} , VO_{2mean}), peak lactate (Peak La^-), difference of lactate (ΔLa^-) and energy systems (anaerobic-alactic, -lactic and aerobic) were analyzed. Basal blood parameters remained unaltered during the interventions. RBC deformability was reduced and aggregation was increased after RWR but values returned to baseline after recovery and were not affected by the SMs. Glucose level was not affected by the interventions. Kick frequency in SMs was higher after RWR which might be responsible for higher HR_{peak} , VO_{2peak} , VO_{2mean} , Peak La^- , ΔLa^- and aerobic demand. The 16 h recovery is sufficient to regenerate measured physiological and hemorheological parameters. TKD-specific performance was not negatively affected during SMs after RWR.

Introduction

The combat sport Taekwondo (TKD) is an Olympic discipline since the 2000 Olympic Games in Sydney/Australia. In TKD, all fights—from qualification to finals—take place in one day [1]. TKD athletes compete in a designated weight category which is controlled by official weigh-in. The weigh-in takes place in the afternoon prior to the competition day [1] and TKD athletes aim to fight in the next lighter weight category to compete with smaller and lighter opponents

[2–4]. In this regard, TKD athletes rapidly reduce their body weight in a range of 2–10% [2, 4–10]. Rapid weight reduction (RWR) is achieved by losing body water during a short-term period. Strategies to reduce body weight include food restriction such as fasting and dehydration, increased training intensity and volume with thermal clothing and sauna sessions [4, 5, 11]. Previous studies indicated that RWR increases oxidative stress, leads to an imbalance of electrolytes and hormones, decreases glycogen storage, changes in blood flow, and decreases plasma volume [12–18]. Further, a recent study showed that 5% RWR decreases red blood cell (RBC) deformability and increases RBC aggregation [8].

Deformability and aggregation of RBC are important factors ensuring oxygen (O₂) transport into the working muscle cell which is associated to athletes' performance [8, 19–21]. RBC must deform to pass through capillaries and to supply O₂ to the working muscle and organs. Blood fluidity, cell surface area-to-volume ratio and physiological properties of the membrane and cytoskeleton were shown to affect RBC deformability [20, 21]. RBC aggregation indicated by face-to-face rouleaux formation upon changes in plasma protein concentration [22]. Increased plasma fibrinogen concentration is believed to increase RBC aggregation. Fibrinogen affects all aspects of RBC aggregation such as aggregate size, stress, viscosity of red cell suspensions and RBC sedimentation [23–25]. It is believed that the recovery time between weigh-in and competition represents a key factor determining the performance during competition after RWR. In TKD, judo and wrestling, 16–24 h lie in-between weigh-in and competition [3, 26]. This period is used to recover body water and carbohydrate storage and to manage possible gastrointestinal distress after RWR [4, 8, 26, 27].

Analyzing recent studies implies a difference between physical performance in laboratory tests and athletic performance in real competition after RWR. Strength, power, peak oxygen uptake (VO_{2peak}), and anaerobic performance measured in a Wingate test were not affected by RWR [11, 28–31]. These appear to be controversially discussed because of different phases and percentages of RWR and diverse measurements of athletic performances [7, 11, 28, 29]. The true competitive benefit that a combat athlete may have by facing smaller opponents may overcome small harmful effects on physical capacities such as strength and aerobic or anaerobic fitness. Actually, no study assessing physical capacities in laboratory settings will ever be able to fully explain how RWR affects actual competitive performance [32].

The outcomes of aerobic and anaerobic performance tests highly depend on the time period between weigh-in and performance test [10]. Artioli et al. [10] suggest that several studies allowed no or very short recovery periods (<1h) which indicated negative effects of RWR on physical performance, but when the recovery periods are longer (>3h), RWR typically does not impair physical performance.

From aforementioned studies it is not clear if and how RWR affects TKD-specific performance but also blood flow related parameters such as deformability and aggregation after a practically relevant recovery time. Therefore, the aim of the present pilot study was to examine the changes of different energy demands, physiological and hemorheological responses after a 3½ day RWR, including 16 h recovery period and during multiple simulation matches on the simulated competition day in TKD athletes.

Materials and methods

Ethical approval

The study was approved by the ethics committee of the German Sport University Cologne (No. 53/2015). The applied protocols align with the Declaration of Helsinki. All participants signed an informed consent form.

Table 1. Nutritional- and activity status during 3½ days.

Nutritional status during 3½ days (kJ·d ⁻¹)	Activity status during 3½ days (kJ·d ⁻¹)
1806.03 ± 517.47	11461.78 ± 734.30

Data are present as mean ± SD of n = 5.

<https://doi.org/10.1371/journal.pone.0196568.t001>

Subjects and study design

Five male regional level taekwondo athletes (black or red and black striped belt; degree: at least 1st Dan) who belong to the light weight categories (range: -54 up to -68 kg) participated in this study. The anthropometric data of the participants were as follows (Mean ± SD): age: 17.40 ± 1.67 years, body weight: 57.77 ± 7.04 kg, height: 170.30 ± 4.91 cm. Average endurance training volume was between 10–12 h per week and all participants exhibited individual experiences in body weight loss. The athletes did not take any medication during all procedure and did without alcohol and nicotine consumption for at least 24 h before the study.

The study consisted of two phases, a control phase without body weight reduction and a RWR phase in which the participants were told to reduce body weight by 5% during 3½ days according to previous studies [4–8]. Due to study organization to display real weigh-in time, the duration of RWR was modified (3½ days instead of 4 days). Weigh-in was scheduled at the fourth day of RWR [4, 7, 8]. During RWR, weight loss was conducted by individual methods of athletes and included fasting, decreased fluid intake, dehydration through sauna sessions, increased training intensity/volume. Nutritional intake and activity report were shown in Table 1 [4, 8].

A two day recovery phase was scheduled between control and RWR period. Each study phase consisted of a pretest, weigh-in test after 3½ days and a competition day after a 16 h recovery time (usual method of weight recovery) (Fig 1). During 16 h recovery, athletes recovered their body water and carbohydrate storage.

On the simulated competition days (SCD), each athlete executed three TKD simulation matches (SMs; 1stSM, 2ndSM and 3rdSM) with a 1 h rest between SMs. According to the guidelines of G1 or G2 (ranking point chart by grade) World Taekwondo (WT) tournaments in Europe, the period of rest between matches should approximately last 30 min– 2 h. Thus, a mean of resting time of 1 h between matches was chosen in the study. However, these durations depend fully on number of participants (athletes) and organization of tournaments.

In each SM, one test athlete fought against a sparring partner; an opponent who was not a test athlete. The fight displayed real competition conditions: The same athletes fought against each other in both phases (control and RWR).

In accordance with the World Taekwondo rules, each SMs consisted of three rounds and 2 min each, with a resting interval of 1 min in-between. During resting intervals, subjects remained in a seated position (passive). Each match conducted within an official area (8 x 8m). A gas analyzer was placed on the back of the competing athlete and no attack to this area was allowed. The sparring partner remained in a passive position during SM such as blocking and fake motion with the intensive steps. Electronic body protectors and headgear were not used because they would interfere with the spirometry [33–35].

Anthropometry, hydration, blood sampling and processing

Anthropometric parameters were assessed and venous blood was sampled during pretest (8 am), weigh-in (3 pm) and pre (9 am) as well as post (5 pm) simulated competition day. Anthropometric parameters and hydration status were measured using an 8-electrode




Fig 1. Study design. The study consisted of a control phase without body weight reduction and a rapid weight reduction (RWR) phase. A two day recovery phase was scheduled between control and RWR period. The participants were told to reduce body weight by 5% during 3 1/2 days. Each phase consisted of a pretest, weigh-in test after 3 1/2 days and a simulated competition day after 16 h recovery time. On SCD, each athlete executed three TKD simulation matches (SMs) with a 1 h rest between SMs. Blood sampling was conducted at pretest, weigh-in test, SCD and after 3rd SM of both phases.

<https://doi.org/10.1371/journal.pone.0196568.g001>

segmental multi-frequency bioelectrical impedance analysis (BIA) for body weight (kg) and hydration (%) (seca mBCA 515, GmbH & Co. KG, Germany). mBCA 515 uses four pairs of electrodes which are positioned at each hand and foot, respectively. The 8-electrode technique enables segmental impedance measurement of the arms and legs. Impedance is measured with a current of 100 μA at frequencies between 1 and 1000 kHz [36]. A skinfold Harpenden caliper was used to measure for body fat (%). An experienced investigator measures skin fold at ten sites: cheek, chin, thorax I, triceps, subscapular, thorax II, abdomen, suprailiac, thigh and calf. The mean of duplicate or the median of triplicate measures (when the first two measures differ by more than 5%) was used [37, 38].

Venous blood was sampled from an elbow vein into sodium-heparin tubes (BD Vacutainer, USA) to measure basal blood parameters, RBC deformability and RBC aggregation. Further, blood was sampled into serum vacutainer (BD Vacutainer, USA) to measure glucose and fibrinogen. Capillary blood was additionally sampled from the earlobe into heparinized capillaries (EKF-Diagnostics GmbH, Germany) to measure RBC deformability. Basal blood parameters included RBC ($\times 10^6 \cdot \mu\text{L}^{-1}$), hemoglobin concentration (Hb) ($\text{g} \cdot \text{dL}^{-1}$), hematocrit [19] (%), mean corpuscular volume (MCV) ($\text{g} \cdot \text{dL}^{-1}$), mean cellular hemoglobin [39] (pg) and mean cellular hemoglobin concentration [39] ($\text{g} \cdot \text{dL}^{-1}$) and was measured using the Sysmex Digitana KX-21N system (Germany).

RBC deformability (venous and capillary blood) and aggregation were analyzed via the Laser-Assisted Optical Rotational Cell Analyzer (LORCA; RR Mechatronics, The Netherlands) [40]. This combined system is an ektacytometer to analyze deformability and sylectometry measurements to determine RBC aggregation. RBC deformability was immediately measured after blood sampling at various fluid shear stresses by laser diffraction analysis at 37°C. The procedure has been described in detailed elsewhere RBCs [8]. Briefly, RBC were mixed with polyvinylpyrrolidone in a 1:250 ratio [41] and sheared in a Couette system at 37°C with a laser beam directed through the sheared sample. Shear stresses between 0.3 and 50 Pa were applied and the horizontal and vertical axis of the laser diffraction pattern, a result of the deformed RBC, was used to calculate an elongation index by the LORCA software. Maximum deformability (EI max) and shear rate required for half of maximum deformability (SS1/2) were calculated using the Lineweaver-Burke equation [42].

Aggregation samples were fully oxygenated (Roller Mixer RM5, 36 rpm). Samples were then filled into the Couette system and aggregation was measured at 37°C. During aggregation

measurement RBC were first disaggregated which was achieved by applied shear rate of 500s^{-1} for 3 sec. After motor stop, RBC aggregation starts with rouleaux formation in the beginning followed by 3D aggregate formation during the following 120 sec of aggregation measurement. The laser beam transmits through the sample and the reflected light is recorded by two photodiodes and used by the LORCA software to calculate the aggregation index (AI, %). A final iteration protocol was applied to analyze the threshold shear rate balancing RBC aggregation and disaggregation. The estimated $dI_{sc\ min}$ is determined as the minimum change in backscatter intensity found during the iteration procedure enabling disaggregation with γ at $dI_{sc\ min}$ ($1/\text{sec}$) being the shear rate at which $dI_{sc\ min}$ was detected.

Glucose concentration was analyzed by a photometric system Cobas Mira Plus (Roche Diagnostic, Switzerland). Fibrinogen concentration ($\text{ng}\cdot\text{mL}^{-1}$) was measured using the Fibrinogen ELISA (Abcam, UK) according to the manufacturer's guidelines.

During all SMs, heart rate [43], blood lactate concentration ($\text{mmol}\cdot\text{L}^{-1}$) and oxygen uptake (VO_2) were additionally monitored. Capillary blood was sampled from the earlobe before and after each SM as well as from the first to seventh minute after last SM [35] to measure lactate concentration ($\text{mmol}\cdot\text{L}^{-1}$). Lactate concentration (Peak La^-) was analyzed by an enzymatic-amperometric sensor chip system (Biosen S-line, EKF diagnostic sales, GmbH, Germany).

A video camera (JVC GC-PX10 HD) was used to record the match for following analysis (i.e., to quantify the amount of time spend fighting during the SM). A Polar Pro Team dock (Polar Electro OY, Finland) was used to record heart rate. The mean heart rate (HR_{mean}) was calculated as average of the heart rate during the entire round (each SM), while peak heart rate (HR_{peak}) was estimated as the highest value during all rounds of each SM [35, 44, 45]. The oxygen uptake in SM and during rest was measured breath-by-breath using the Oxycon Mobile (CareFusion, Germany). The portable spirometric device measures the oxygen concentration through an electrochemical sensor and data are telemetrically transmitted and recorded on a personal computer [46]. The gas analyzer was calibrated using calibration gas (16% O_2 , 5% CO_2 ; RiessnerGase, Germany), and the turbine volume transducer was calibrated using a 3 L syringe (Hans Rudolph Inc., USA).

Data were recorded until 6 min after the end of the SM. The mean oxygen uptake ($\text{VO}_{2\text{mean}}$) was calculated as the average VO_2 of each SM and the peak oxygen uptake ($\text{VO}_{2\text{peak}}$) was determined as highest value of each SM.

Calculation of energy contribution and video analysis

Evaluations of aerobic, anaerobic-lactic and anaerobic-alactic system were conducted by the measurement of oxygen uptake during activity, peak blood lactate concentration, and the fast phase of excess oxygen uptake after exercise ($\text{EPOC}_{\text{FAST}}$), respectively [35].

Aerobic energy (W_{AER}) was calculated by subtracting rest oxygen uptake ($\text{VO}_{2\text{REST}}$) from VO_2 during a round by the trapezoidal method in which the area under the curve is divided into pieces and then the sum of each trapezoid is calculated to estimate the integral [35, 47, 48]. $\text{VO}_{2\text{REST}}$ was determined in the standing position, with the last 30 s of a 5 min period used as a reference. The anaerobic-alactic system (W_{PCR}) was calculated considering the oxygen uptake during the interval between rounds, and the fast component of excess post-exercise ($\text{EPOC}_{\text{FAST}}$) after the third round [33, 47–49]. In this study, the kinetics of post-match oxygen consumption was fitted by mono- and bi-exponential models. The slow component of the bi-exponential model was negligible. Hence, the post-match VO_2 data was fitted to a mono-exponential model and W_{PCR} was obtained by calculating the integral of the exponential part [33, 47–49].

The contribution of the anaerobic-lactic system (W_{La^-}) was estimated as the lactate concentration after combat, assuming that the accumulation of $1\ \text{mmol}\cdot\text{L}^{-1}$ (La^-) is equivalent to $3\text{ml}\ \text{O}_2\ \text{kg}^{-1}$

of body mass [50]. The caloric quotient of 20.92 kJ [51] was used in all three different energy systems. The difference of lactate (ΔLa^-) was calculated as the lactate concentration after the round, minus the lactate concentration at the beginning of the round. Total energy demand (W_{TOTAL}) was calculated as the sum of the three energy systems ($W_{\text{AER}} + W_{\text{PCR}} + W_{\text{La}^-}$) in kJ [35].

According to the criteria of Santos et al. [52], the recorded matches were analyzed using the Dartfish Version 7 software. Time-motion analyzes were conducted in an attempt to calculate work/rest ratios for each match. The events were measured in seconds and tenths of seconds, using the marking tool in the analysis software. An attack was considered the time from the beginning of a foot or hand movement in the direction of the opponent up to the athlete stopped his attack movement or unable to continue attack (because of a fall or referee pause). Step movement was also determined as the entire phase between attacks when there was no break in the match. Pause was determined as the stops in SM ordered by the referee. Step time was added to the referee pause-times to determine the phase without attack (PWA), together with the stopwatch stop-time called out by the referee. Attack was considered the total time during which one of the athletes tried to attack or succeeded in attacking his opponent [35]. For the ratio between High Intensity Movement (HIM) and Low Intensity Movement (LIM), sum of time without attacks was divided by the number of attacks ratio [35]. The analyses showed a high intra-class correlation [53] of 0.93 which corresponds to other studies [35, 54, 55].

Statistical analysis

Statistical software GraphPad Prism 6 (LaJolla, USA) was used for the statistical analyses of the data and graphical representation of data. The data are presented as means \pm SD. The mean value of SM for performance parameters and energy demands was calculated by all 3 rounds of each SM and statistically compared. The variables of each phase (control and RWR) were compared using a Friedman repeated-measures rank test with post-hoc Dunn's. The same measurement time e.g., 1stSM (control) vs. 1stSM (RWR) was analyzed using a Wilcoxon signed ranks test. Statistical difference was considered to be significant for $P < 0.05$ and $P < 0.01$. The effect size ($r = Z/\sqrt{n}$) was calculated for the non-parametric tests [56]. Whiskers box plot present minimum to maximum and median values. Thresholds for small, medium, and large effects were 0.10, 0.30, and 0.50, respectively [57]. Additionally, a Pearson's two-tailed correlation and linear regression analyses were performed between kick frequency (KF) and $\text{VO}_{2\text{peak}}$, $\text{VO}_{2\text{mean}}$, peak La^- and ΔLa^- .

Results

Anthropometric data, hydration and glucose

In RWR, athletes significantly reduced body weight between 2ndpretest und 2ndweigh-in ($P = 0.003$; $r = -0.63$). Hydration status was significantly decreased in 2ndweigh-in compared to 2ndpretest and again increased on 2ndSCD compared to 2ndweigh-in ($P = 0.021$; $r = -0.63$, $P = 0.035$; $r = -0.63$, respectively). Body weight was unaltered in the control phase. Body fat was not affected by the interventions. Serum glucose concentration showed a significant increase after 3rdSM compared to SCD of control phase ($P = 0.003$; $r = -0.63$) and another significant difference on SCD between control and RWR ($P = 0.031$; $r = -0.63$) (Fig 2A–2C).

Basal blood parameters

Basal blood parameters are presented in Table 2. The hematocrit concentration was significantly lower at 2ndpretest compared to 1stpretest ($P = 0.032$; $r = -0.64$). Other basal blood parameters were not affected by the interventions.




Fig 2. Anthropometric data, hydration and glucose. (A) Body weight significantly decreased at weigh-in of RWR phase ($P = 0.003$). (B) Hydration status was significantly decreased during weigh-in compared to pretest and increased on SCD ($P = 0.021$, $P = 0.035$, respectively) of RWR phase. (C) Body fat remained unaffected by the interventions. (D) Serum glucose concentration was significantly increased after 3rdSM of control phase ($P = 0.003$) and a significant difference was detected between SCD of control and RWR phase ($P = 0.031$). All data are presented as box plot ($n = 5$).

<https://doi.org/10.1371/journal.pone.0196568.g002>

Hemorheological properties

Calculated EI max of venous blood showed no differences between the two set-ups (control/RWR) and no alteration during the interventions. In RWR, SS1/2 (Pa) significantly increased from pretest to weigh-in. Values were also significantly higher at weigh-in compared to weigh-in of control phase ($P = 0.019$; $r: -0.63$, $P = 0.031$; $r: -0.63$, respectively). EI max values of capillary blood were higher in control phase at weigh-in and SCD compared to RWR ($P = 0.031$; $r: -0.63$, respectively). In RWR, SS1/2 of capillary blood was significantly higher at pretest compared to control pretest ($P = 0.031$; $r: -0.63$) (Fig 3A–3D).

Table 2. Basal blood parameters of all measurements.

	1 st Pretest	1 st Weigh-in	1 st SCD	1 st After 3 rd SM	2 nd Pretest	2 nd Weigh-in	2 nd SCD	2 nd After 3 rd SM
RBC (*10 ⁶ ·μL ⁻¹)	5.89 ± 0.26	5.28 ± 0.14	5.39 ± 0.26	5.28 ± 0.28	5.08 ± 0.20	5.68 ± 0.44	5.03 ± 0.16	5.07 ± 0.14
Hb (g·dL ⁻¹)	17.96 ± 0.75	15.62 ± 0.91	15.72 ± 0.46	15.32 ± 1.48	14.76 ± 0.50	16.56 ± 2.11	14.96 ± 1.15	15.30 ± 0.97
Hct (%)	50.30 ± 1.83	44.94 ± 2.30	46.00 ± 0.74	45.06 ± 2.83	43.58 ± 1.60 ^{a*}	46.50 ± 6.96	43.12 ± 2.85	43.40 ± 2.18
MCV (fl)	85.52 ± 3.03	85.02 ± 3.27	85.40 ± 3.12	85.38 ± 3.40	85.78 ± 3.02	84.78 ± 3.37	85.70 ± 3.14	85.66 ± 3.29
MCH (pg)	30.58 ± 1.83	29.56 ± 1.25	29.22 ± 1.86	28.98 ± 1.97	29.08 ± 1.63	30.42 ± 1.76	29.72 ± 1.58	30.20 ± 2.02
MCHC (g·dL ⁻¹)	35.72 ± 1.12	34.76 ± 0.99	34.16 ± 1.38	33.94 ± 1.36	33.88 ± 0.99	35.76 ± 1.75	34.68 ± 1.49	35.28 ± 1.89

Data are present as mean ± SD of n = 5.

RBC: red blood cell; PLT: platelet; Hb: hemoglobin; Hct: hematocrit; MCV: mean corpuscular volume; MCH: mean corpuscular hemoglobin; MCHC: mean corpuscular hemoglobin concentration; SCD: simulated competition day; SM: simulation match.

a: 1stPretest

*P<0.05

<https://doi.org/10.1371/journal.pone.0196568.t002>

AI (%) was significantly higher at pretest in control phase compared to RWR (P = 0.031; r:-0.63). AI was significantly higher at weigh-in of RWR compared to control (P = 0.031; r = 0.63). The disaggregation threshold showed no significant difference during the interventions and between set-ups. Serum fibrinogen concentration was not affected by the interventions (Fig 4A–4C).

Physiological parameters

HR_{mean} significantly increased during 3rdSM of RWR compared to 3rdSM control (P = 0.031; r:-0.63). HR_{peak} was significantly higher in 3rdSM of RWR compared to 2ndSM of RWR (P = 0.012; r:-0.64). VO_{2mean} was significantly higher at 3rdSM of RWR compared to control (P = 0.047; r:-0.63). VO_{2peak} and absolute VO_{2peak} were significantly higher at 2ndSM and 3rdSM of RWR compared to control (P = 0.031; r:-0.64, respectively). Peak La⁻ and ΔLa⁻ were significantly decreased in 3rdSM compared to 1stSM of control (P = 0.006; r:-0.64, P = 0.002; r:-0.63, respectively) and significantly lower compared to corresponding values of RWR (P = 0.031; r:-0.63, P = 0.021; r:-0.63, respectively) (Table 3). Kick frequency (KF) was significant different between control and RWR in 1stSM and 3rdSM (P = 0.031; r:-0.64, P = 0.031; r:-0.63, respectively) (Table 4).

Moderate positive correlations and linear regressions were observed between VO_{2peak}, VO_{2mean} and KF (r = 0.64; 95%CI: 0.36–0.81; R² = 0.41; F = 19.44; P<0.001, r = 0.56; 95%CI: 0.25–0.77; R² = 0.32; F = 12.92; P = 0.0012, respectively). Moderate negative correlations and linear regressions were calculated for peak La⁻, ΔLa⁻ and KF (r = 0.54; 95%CI: -0.75 - -0.22; R² = 0.29; F = 11.57; P = 0.002, r = 0.54; 95%CI: -0.75 - -0.23; R² = 0.29; F = 11.62; P = 0.002, respectively). (Fig 5A–5D).

Energy demands during SM

W_{TOTAL} and W_{PCR} showed no significant difference during interventions and no difference between set-ups. W_{La⁻} was significantly lower during 3rdSM compared to 1stSM of control (P = 0.002; r:-0.63, P = 0.005; r:-0.64, respectively), and significantly higher values were measured during 3rdSM of RWR compared to control (P = 0.021; r:-0.63, P = 0.031; r:-0.63, respectively). W_{AER} was significantly higher during 2ndSM of RWR compared to 2ndSM of control (P = 0.031; r:-0.63) (Table 3).




Fig 3. RBC maximum deformability (EI max) and shear rate for half of maximum deformability SS1/2 (Pa). (A) EI max of venous blood showed no significant difference during interventions nor between set-ups. (B) During RWR, SS1/2 of venous blood significantly increased at weigh-in with significantly higher values compared to pretest of control phase ($P = 0.019$; $P = 0.031$, respectively). (C) EI max of capillary blood was significantly higher at weigh-in and SM in RWR compared to control phase ($P = 0.031$). (D) SS1/2 of capillary blood was significantly lower at pretest in control phase compared to RWR ($P = 0.031$). All data are presented as box plot ($n = 5$).

<https://doi.org/10.1371/journal.pone.0196568.g003>

The technical actions and the time spent on each SM are described in Table 4. Kick frequency, attack time, sum of attack time, sum without attack time, attacks/without attacks ratio and sum of time without attacks/number of attacks ratio were identified.




Fig 4. Aggregation index (AI, %), disaggregation threshold (γ at disc min, $1 \cdot s^{-1}$) and fibrinogen concentration. (A) AI showed significant differences in pretest and weigh-in between control and RWR ($P = 0.003$, $P = 0.031$, respectively). (B) Disaggregation threshold and (C) fibrinogen concentration were not affected by the interventions. All data are presented as box plot ($n = 5$).

<https://doi.org/10.1371/journal.pone.0196568.g004>

Discussion

The effect of rapid weight reduction on TKD-specific performance parameters after a practically relevant recovery time has not been investigated so far. Thus, to our knowledge this is the

Table 3. Physiological parameters and calculated energy demands of Taekwondo simulation match during control and rapid weight reduction phases.

	1 st SM (control)	2 nd SM (control)	3 rd SM (control)	1 st SM (RWR)	2 nd SM (RWR)	3 rd SM (RWR)
HR _{mean} (beats·min ⁻¹)	181.8 ± 8.2	182.8 ± 6.5	178.8 ± 8.2	179.6 ± 10.9	176.8 ± 8.7	185.8 ± 3.2 ^c
HR _{peak} (beats·min ⁻¹)	192.4 ± 5.8	190.6 ± 4.4	193.6 ± 8.3	193.8 ± 8.8	188.6 ± 5.4	196.2 ± 5.0 ^e
VO _{2mean} (mL·kg ⁻¹ ·min ⁻¹)	34.2 ± 2.9	35.5 ± 1.7	33.2 ± 2.0	34.9 ± 2.2	37.3 ± 1.9	38.7 ± 3.0 ^c
VO _{2peak} (mL·kg ⁻¹ ·min ⁻¹)	40.0 ± 2.9	40.9 ± 2.9	40.8 ± 2.0	40.8 ± 3.3	44.0 ± 2.7 ^b	46.7 ± 6.1 ^c
VO _{2peak} (mL·min ⁻¹)	2259.0 ± 252.3	2315.3 ± 219.0	2327.1 ± 295.9	2304.3 ± 199.3	2487.0 ± 221.9 ^b	2632.1 ± 223.7 ^c
Peak La ⁻ (mmol·L ⁻¹)	9.2 ± 3.9	7.2 ± 4.2	5.6 ± 4.2 ^{a**}	8.2 ± 4.4	7.4 ± 3.7	8.1 ± 2.9 ^c
ΔLa ⁻ (mmol·L ⁻¹)	2.7 ± 1.1	2.0 ± 1.2	1.5 ± 1.0 ^{a**}	2.2 ± 1.5	2.0 ± 0.8	2.3 ± 0.9 ^c
W _{PCR} (kJ)	18.8 ± 3.5	19.9 ± 3.3	19.5 ± 2.3	21.3 ± 5.5	21.0 ± 5.5	22.2 ± 4.9
W _{PCR} (%)	19 ± 3	20 ± 4	21 ± 3	21 ± 6	20 ± 6	20 ± 4
W _{La⁻} (kJ)	10.1 ± 5.5	7.7 ± 5.9	5.6 ± 4.7 ^{a**}	8.3 ± 6.6	7.2 ± 3.9	8.6 ± 4.2 ^c
W _{La⁻} (%)	9 ± 4	7 ± 4	5 ± 4 ^{a*}	8 ± 5	6 ± 3	8 ± 3 ^c
W _{AER} (kJ)	72.8 ± 10.4	75.3 ± 7.5	71.9 ± 12.5	73.9 ± 9.1	79.6 ± 8.5 ^b	82.4 ± 7.2
W _{AER} (%)	72 ± 3	73 ± 2	74 ± 2	71 ± 1	74 ± 4	72 ± 2
W _{TOTAL} (kJ)	101.7 ± 16.1	102.5 ± 11.6	97.0 ± 17.9	103.5 ± 12.9	107.8 ± 8.5	113.2 ± 8.4

Data are present as mean ± SD of $n = 5$.

RWR: phase of rapid weight reduction, SM: simulation match, HR_{mean}: mean heart rate, HR_{peak}: highest heart rate, VO_{2mean}: mean oxygen uptake, VO_{2peak}: highest oxygen uptake, Peak La⁻: highest lactate concentration, ΔLa⁻: difference of lactate concentration after the round minus at the beginning of the round, W_{PCR}: anaerobic-alactic demand, W_{La⁻}: anaerobic-lactic demand, W_{AER}: aerobic demand.

a: first simulation match of control

b: second simulation match of control

c: third simulation match of control

e: second simulation match of rapid weight reduction phase

* $P < 0.05$

** $P < 0.01$

<https://doi.org/10.1371/journal.pone.0196568.t003>

Table 4. Kick frequency, attack time, total time of attacks, total time without attacks, ratio between attacks and time without attacks, and ratio between total time without attacks and number of attacks (HIM vs. LIM).

	1 st SM (control)	2 nd SM (control)	3 rd SM (control)	1 st SM (RWR)	2 nd SM (RWR)	3 rd SM (RWR)
Kick frequency (number of attacks)	15.73 ± 5.83	16.87 ± 4.16	16.53 ± 3.44	20.73 ± 5.10 ^a	20.11 ± 6.56	21.73 ± 6.31 ^c
Attack time (s)	0.87 ± 0.04	0.86 ± 0.05	0.86 ± 0.06	0.84 ± 0.05	0.85 ± 0.06	0.86 ± 0.07
Sum of attacks time (s)	14.07 ± 5.99	15.40 ± 5.68	13.80 ± 3.63	17.60 ± 5.18	18.00 ± 7.97	18.80 ± 6.53
Sum of time without attacks (s)	106.13 ± 5.95	104.67 ± 5.54	106.27 ± 3.61	102.53 ± 5.28	101.93 ± 7.91	101.20 ± 6.50
Attacks/without attacks ratio	0.27 ± 0.37	0.15 ± 0.06	0.13 ± 0.04	0.17 ± 0.07	0.18 ± 0.10	0.19 ± 0.08
Sum of time without attacks/number of attacks ratio	8 ± 3	7 ± 2	7 ± 2	5 ± 2	6 ± 2	5 ± 2
HIM vs. LIM	1:8	1:7	1:7	1:5	1:6	1:5

Data are present as mean ± S.D. of n = 5.

RWR: phase of rapid weight reduction, SM: simulation match, HIM: high intensity movement; LIM: low intensity movement.

a: first simulation match of control

c: third simulation match of control

*P<0.05

<https://doi.org/10.1371/journal.pone.0196568.t004>

first study investigating the effects of a 16 h recovery time after weight reduction by 5% within 3½ days on physiological and hemorheological responses and energy demands with special regards to multiple TKD SMs. The major findings indicate that RWR impairs hemorheological but not hematological parameters. The values returned to pretest level after the 16 h recovery period. Performance related parameters such as VO_{2peak} were higher during SMs after RWR, most likely because of higher kick frequency during this phase. Hematological parameters were monitored during RWR and SMs and compared to a control phase with no weight reduction. Basal blood parameters included RBC count, hematocrit, hemoglobin concentration, MCV, MCH and MCHC. Blood parameters give some indication of hydration status and blood viscosity. MCHC for example was described to sensitively reflect cell viscosity within RBC [39] and increased hematocrit was shown to negatively influence blood viscosity. Alterations of these parameters were associated with a reduction in performance capacity because increased viscosity reduces peripheral blood flow and cardiac output [21, 58, 59]. Increases in athletes' hematocrit may be caused by extensive loss of body water during training or fasting [60]. Here, MCH(C) and hematocrit were not affected by the interventions, although hydration status was significantly decreased after RWR. Several studies indicate that hematocrit does not necessarily increase loss of body water. Loss of body water < 3% of body mass does not affect the plasma volume [7, 8, 61–63]. However, hematocrit was significantly higher at 1stpretest compared to 2ndpretest. Hematocrit is affected by a variety of factors including hydration status and abovementioned mechanisms. This might have influenced the values. Inaccuracies of measurement might also explain the high hematocrit results and might thus represent a possible limitation of the study.

Besides MCH(C) and hematocrit, other blood values were also not affected by the intervention. In regard of RBC deformability, MCV is of special interest. Both, increased and decreased MCV values were described to alter RBC deformability [64]. The values described for this study were not affected by RWR or altered upon SMs and were all within normal range (80–98 fl) [65]. To evaluate the effects of RWR, recovery phase and SMs on RBC deformability, EI max was measured in capillary (arterial-venous mix blood) and venous blood, respectively. EI max of capillary and of venous blood remained unaltered during RWR, respectively, and results thus in part reflect our recent findings [8]. Values of capillary blood were in general




Fig 5. Pearson's two-tailed correlation and linear regression between kick frequency (KF) and peak oxygen uptake (VO_{2peak}), mean oxygen uptake (VO_{2mean}), peak lactate (peak La^-) and difference of lactate ΔLa^- . (A) Moderate positive correlation and linear regression between KF and VO_{2peak} (B) Moderate positive correlation and linear regression between KF and VO_{2mean} (C) Moderate negative correlation and linear regression between KF and peak ΔLa^- (D) Moderate negative correlation and linear regression between KF and peak La^- .

<https://doi.org/10.1371/journal.pone.0196568.g005>

higher in control compared to RWR with significant differences at weigh-in before SMs but this difference might not be related to the interventions.

EI max represents a theoretical values and might not be sufficient to state the effect of RWR on deformability. Thus, shear rate necessary for one half of maximum deformability (SS1/2) was calculated. In venous blood, SS1/2 significantly increased after RWR indicating that more shear stress is necessary to deform RBC and thus indicates reduced deformability. In capillary blood, no such behavior was measured. These differences between capillary and venous blood might be caused by differing flow resistance in vascular systems [66]. A decrease in RBC deformability might also be caused by food restriction during RWR. Several studies associated a reduction in RBC deformability to decrease in blood glucose and thus reduced ATP levels [8, 67, 68]. Glucose levels were reduced on weigh-in of RWR and might in part relate to the reduced RBC deformability. A recent study indicated that food restriction and reduced glucose levels reduce nitric oxide production in RBC which in turn reduces RBC deformability [8]. It

remains to be investigated whether RWR facilitates oxidative/nitrosative stress which was shown to reduce RBC deformability [69].

RBC aggregation just like RBC deformability define blood flow in the microcirculation and might thus be of importance in terms of performance capacity. The results presented herein showed higher RBC aggregation value at weigh-in of RWR compared to the control situation. Further, shear rate balancing aggregation and disaggregation was higher at weigh-in after RWR. Both results are in accordance to recent findings by Yang et al. [8]. Dehydration as monitored for RWR might have caused an increase in blood viscosity thus leading to hyper-aggregation [8, 39, 70].

RBC aggregation is a physiological phenomenon that occurs at the access of the capillaries. RBC aggregation is influenced by the plasma protein fibrinogen with increased aggregation behavior at higher fibrinogen levels [71]. RBC aggregation can influence hemodynamics, RBC distribution, and flow dynamics in the microcirculation. Low protein intake through caloric restriction appears to relate to the increase of fibrinogen and aggregation [72]. Hence, serum fibrinogen levels were measured but results suggest that fibrinogen levels were not affected by RWR. Solá et al. [73] and Poggi et al. [74] suggested that other, yet unknown factors, are responsible for an increase in RBC aggregation without a parallel change in fibrinogen levels.

Both, RBC deformability and aggregation values returned to pretest level after a 16 h recovery period and no changes for the parameters were found upon SMs. The data thus suggest that, although RWR might impair hemorheological parameters, recovery time between weigh-in and competition is sufficient to recover blood parameters. In case of reduced performance capacity on competition day, it seems unlikely that this might be associated to impaired hemorheological responses. Nevertheless, it remains to be investigated whether repeated RWR applications might have a bigger effect.

TKD-specific performance parameters were monitored during SMs. $VO_{2\text{mean}}$, $VO_{2\text{peak}}$, absolute $VO_{2\text{peak}}$ and aerobic demand were higher in 2ndSM and 3rdSM of RWR compared to control. Accordingly, HR_{mean} and HR_{peak} were increased in 3rdSM of RWR compared to 3rdSM of control. This might be related to increased kick frequency measured during SMs of the RWR phase as also indicated by lactate measurements [35, 45]. Other studies confirmed that RWR does not impact performance (determined in wrestlers and boxer) after a recovery period of 3–20 h [3, 75].

Metabolic systems were measured during SMs. The aerobic system was found to dominate during all simulated matches. The results might be explained by frequent low intensity movement which includes the pauses and step movements [35, 52, 76, 77]. Furthermore, the ratio of high intensity to low intensity movement (HIM to LIM) was from 1:5 to 1:8.

Anaerobic-lactic system showed no alteration in both phases. Recovery interval during pauses and rounds with LIM might be sufficient for the recovery of phosphocreatine, which indicated no alteration between control and RWR phases [35, 78, 79]. La^- , ΔLa^- and anaerobic-lactic demand were also higher in RWR than those in control because more HIM was done in RWR. Indeed, our regression analyses showed moderate positive and negative relationships between these parameters. The results are thus in accordance to a computer simulation program which indicates that the aerobic system is used during moderate and increasing duration of exercise while anaerobic-lactic system is less required (Fig 6) [80, 81]. Similar results were reported by Campos et al. [35] showing that the aerobic system (66%) was predominantly used during TKD simulation matches.

To avoid an impact of RWR on performance parameters, TKD athletes delimitate body weight reduction to 5% in praxis. And to protect athletes health situation, World Taekwondo should consider a random control of body weight in the morning of the competition day to ensure no discrepancy of body weight between weigh-in and competition day.




Fig 6. Averages of each energy system during all TKD simulation matches (30 matches) in kJ (A) and % (B). Black: An example of VO_2 ($mL \cdot kg^{-1} \cdot min^{-1}$) during a simulation match, red: Anaerobic-alactic demand, violet: Anaerobic-lactic demand, blue: aerobic demand (kJ), yellow: lactate, green: difference of lactate ($mmol \cdot L^{-1}$). All illustration of energy systems and lactate values based on mean values of each parameters.

<https://doi.org/10.1371/journal.pone.0196568.g006>

Conclusion

The findings of the present pilot study indicated that RWR by 5% body weight does not impair physiological and hemorheological parameters on the simulated competition when a 16 h recovery time was scheduled between weigh-in and competition day. However, it remains to be investigated whether repeated RWRs lead to different results in the season.

Supporting information

S1 Checklist. TREND statement checklist.
(PDF)

Acknowledgments

We would like to thank all of the subjects for participating in this study, and Anke Schmitz and Thomas Dietz for excellent technical assistance.

Author Contributions

Conceptualization: Woo-Hwi Yang, Oliver Heine, Marijke Grau.

Data curation: Woo-Hwi Yang, Marijke Grau.

Formal analysis: Woo-Hwi Yang.

Funding acquisition: Oliver Heine, Marijke Grau.

Investigation: Woo-Hwi Yang, Oliver Heine, Marijke Grau.

Methodology: Woo-Hwi Yang, Oliver Heine, Marijke Grau.

Project administration: Woo-Hwi Yang.

Resources: Oliver Heine, Marijke Grau.

Supervision: Oliver Heine, Marijke Grau.

Visualization: Woo-Hwi Yang.

Writing – original draft: Woo-Hwi Yang.

Writing – review & editing: Woo-Hwi Yang, Marijke Grau.

References

1. World Taekwondo. Competition Rules & Interpretation. 2016. Available from: <http://www.worldtaekwondofederation.net/wp-content/uploads/2017/01/WTF-Competition-Rules-Interpretation-Nov-15-2016-Burnaby-Canada.pdf>.
2. Artioli GG, Gualano B, Franchini E, Scagliusi FB, Takesian M, Fuchs M, et al. Prevalence, magnitude, and methods of rapid weight loss among judo competitors. *Med Sci Sports Exerc.* 2010; 42(3):436–442. <https://doi.org/10.1249/MSS.0b013e3181ba8055> PMID: 19952804
3. Reale R, Cox GR, Slater G, Burke LM. Weight Re-Gain is Not Linked to Success in a Real Life Multi-Day Boxing Tournament. *Int J Sports Physiol Perform.* 2016:1–26. <https://doi.org/10.1123/IJSP.2015-0781> PMID: 26752203
4. Yang WH, Grau M, Kim P, Schmitz A, Heine O, Bloch W, et al. Physiological and psychological performance of taekwondo athletes is more affected by rapid than by gradual weight reduction. *Arch Budo.* 2014; 10:169–177.
5. Brito CJ, Roas AF, Brito IS, Marins JC, Cordova C, Franchini E. Methods of body mass reduction by combat sport athletes. *Int J Sport Nutr Exerc Metab.* 2012; 22(2):89–97. Epub 2012/02/22. PMID: 22349031.
6. Petterson S, Ekström MP, Berg CM. Practices of weight regulation among elite athletes in combat sports: a matter of mental advantage? *J Athl Train.* 2013; 48(1):99–108. <https://doi.org/10.4085/1062-6050-48.1.04> PMID: 23672331
7. Yang WH, Heine O, Mester J, Grau M. Impact of rapid weight reduction on health and performance related indicators of athletes representing the Olympic combat sports. *Arch Budo.* 2017; 13:147–160.
8. Yang WH, Heine O, Pauly S, Kim P, Bloch W, Mester J, et al. Rapid Rather than Gradual Weight Reduction Impairs Hemorheological Parameters of Taekwondo Athletes through Reduction in RBC-NOS Activation. *PLoS One.* 2015; 10(4):e0123767. Epub 2015/04/16. <https://doi.org/10.1371/journal.pone.0123767> PMID: 25875585.

9. Franchini E, Brito CJ, Artioli GG. Weight loss in combat sports: physiological, psychological and performance effects. *J Int Soc Sports Nutr.* 2012; 9(1):52. <https://doi.org/10.1186/1550-2783-9-52> PMID: [23237303](https://pubmed.ncbi.nlm.nih.gov/23237303/)
10. Artioli GG, Saunders B, Iglesias RT, Franchini E. It is time to ban rapid weight loss from combat sports. *Sports Med.* 2016; 46(11):1579–1584. <https://doi.org/10.1007/s40279-016-0541-x> PMID: [27102173](https://pubmed.ncbi.nlm.nih.gov/27102173/)
11. Koral J, Dosseville F. Combination of gradual and rapid weight loss: effects on physical performance and psychological state of elite judo athletes. *J Sports Sci.* 2009; 27(2):115–120. Epub 2008/12/06. <https://doi.org/10.1080/02640410802413214> PMID: [19058087](https://pubmed.ncbi.nlm.nih.gov/19058087/).
12. Oppliger RA, Case HS, Horswill CA, Landry GL, Shelter AC. American College of Sports Medicine position stand. Weight loss in wrestlers. *Med Sci Sports Exerc.* 1996; 28(6):9–12. Epub 1996/06/01. PMID: [8926865](https://pubmed.ncbi.nlm.nih.gov/8926865/).
13. Costill DL, Cote R, Fink W. Muscle water and electrolytes following varied levels of dehydration in man. *J Appl Physiol.* 1976; 40(1):6–11. Epub 1976/01/01. <https://doi.org/10.1152/jappl.1976.40.1.6> PMID: [1248983](https://pubmed.ncbi.nlm.nih.gov/1248983/).
14. Vaccaro P, Zauner C, Cade J. Changes in body weight, hematocrit and plasma protein concentration due to dehydration and rehydration in wrestlers. *J Sports Med Phys Fitness.* 1976; 16(1):45. PMID: [1263474](https://pubmed.ncbi.nlm.nih.gov/1263474/)
15. Sawka MN, Pandolf KB. Effects of body water loss on physiological function and exercise performance. In: Gisolfi CV, Lamb DR. *Perspectives in exercise science and sports medicine.* Vol 3. Carmel, IN: Benchmark Press; 1990. pp. 1–38.
16. Houston M, Marrin D, Green H, Thomson J. The effect of rapid weight loss on physiological functions in wrestlers. *Physician Sportsmed.* 1981; 9(11):73–78.
17. Weissinger E, Housh TJ, Johnson GO, Evans SA. Weight loss behavior in high school wrestling: wrestler and parent perceptions. *Pediatric Exercise Science.* 1991; 3(1):64–73.
18. Woods ER, Wilson CD, Masland RP. Weight control methods in high school wrestlers. *J Adolesc Health Care.* 1988; 9(5):394–397. PMID: [3170306](https://pubmed.ncbi.nlm.nih.gov/3170306/)
19. Ahmadizad S, El-Sayed MS, MacLaren DP. Effects of water intake on the responses of haemorheological variables to resistance exercise. *Clin Hemorheol Microcirc.* 2006; 35(1–2):317–327. Epub 2006/08/11. PMID: [16899951](https://pubmed.ncbi.nlm.nih.gov/16899951/).
20. Tripette J, Loko G, Samb A, Gogh BD, Sewade E, Seck D, et al. Effects of hydration and dehydration on blood rheology in sickle cell trait carriers during exercise. *Am J Physiol Heart Circ Physiol.* 2010; 299(3):H908–914. Epub 2010/06/29. <https://doi.org/10.1152/ajpheart.00298.2010> PMID: [20581085](https://pubmed.ncbi.nlm.nih.gov/20581085/).
21. Connes P, Simmonds MJ, Brun JF, Baskurt OK. Exercise hemorheology: classical data, recent findings and unresolved issues. *Clin Hemorheol Microcirc.* 2013; 53(1–2):187–199. Epub 2012/10/09. <https://doi.org/10.3233/CH-2012-1643> PMID: [23042105](https://pubmed.ncbi.nlm.nih.gov/23042105/).
22. Rampling MW, Meiselman HJ, Neu B, Baskurt OK. Influence of cell-specific factors on red blood cell aggregation. *Biorheology.* 2004; 41(2):91–112. Epub 2004/04/20. PMID: [15090679](https://pubmed.ncbi.nlm.nih.gov/15090679/).
23. Maeda N, Seike M, Nakajima T, Izumida Y, Sekiya M, Shiga T. Contribution of glycoproteins to fibrinogen-induced aggregation of erythrocytes. *Biochim Biophys Acta.* 1990; 1022(1):72–78. PMID: [2302404](https://pubmed.ncbi.nlm.nih.gov/2302404/)
24. Baskurt O NB, Meiselman HJ. *Red blood cell aggregation:* CRC Press; 2011.
25. Maeda N, Shiga T. Opposite effect of albumin on the erythrocyte aggregation induced by immunoglobulin G and fibrinogen. *Biochim Biophys Acta.* 1986; 855(1):127–135. PMID: [3942735](https://pubmed.ncbi.nlm.nih.gov/3942735/)
26. Davis P. Comment on: "It is time to ban rapid weight loss from combat sports". *Sports Med.* 2017:1–3. <https://doi.org/10.1007/s40279-016-0562-5> PMID: [27251334](https://pubmed.ncbi.nlm.nih.gov/27251334/)
27. Reale R, Slater G, Burke LM. Individualised dietary strategies for Olympic combat sports: Acute weight loss, recovery and competition nutrition. *Eur J Sport Sci.* 2017:1–14. <https://doi.org/10.1080/17461391.2016.1218939> PMID: [27800713](https://pubmed.ncbi.nlm.nih.gov/27800713/)
28. Serfass RC, Stull GA, Alexander JF, Ewing JL Jr. The effects of rapid weight loss and attempted rehydration on strength and endurance of the handgripping muscles in college wrestlers. *Res Q Exerc Sport.* 1984; 55(1):46–52.
29. Webster S, Rutt R, Weltman A. Physiological effects of a weight loss regimen practiced by college wrestlers. *Med Sci Sports Exerc.* 1990; 22(2):229–234. PMID: [2355820](https://pubmed.ncbi.nlm.nih.gov/2355820/)
30. Reljic D, Feist J, Jost J, Kieser M, Friedmann-Bette B. Rapid body mass loss affects erythropoiesis and hemolysis but does not impair aerobic performance in combat athletes. *Scand J Med Sci Sports.* 2016; 26(5):507–517. Epub 2015/04/29. <https://doi.org/10.1111/sms.12485> PMID: [25916419](https://pubmed.ncbi.nlm.nih.gov/25916419/).
31. Fogelholm GM, Koskinen R, Laakso J, Rankinen T, Ruokonen I. Gradual and rapid weight loss: effects on nutrition and performance in male athletes. *Med Sci Sports Exerc.* 1993; 25(3):371–377. Epub 1993/03/01. PMID: [8455453](https://pubmed.ncbi.nlm.nih.gov/8455453/).

32. Artioli GG, Saunders B, Iglesias RT, Franchini E. Authors' Reply to Davis: "It is Time to Ban Rapid Weight Loss from Combat Sports". *Sports Med.* 2017;1–5. <https://doi.org/10.1007/s40279-016-0562-5> PMID: 27251334
33. Beneke R, Beyer T, Jachner C, Erasmus J, Hütler M. Energetics of karate kumite. *Eur J Appl Physiol.* 2004; 92(4–5):518–523. <https://doi.org/10.1007/s00421-004-1073-x> PMID: 15138826
34. Crisafulli A, Vitelli S, Cappai I, Milia R, Tocco F, Melis F, et al. Physiological responses and energy cost during a simulation of a Muay Thai boxing match. *Appl Physiol Nutr Metab.* 2009; 34(2):143–150. <https://doi.org/10.1139/H09-002> PMID: 19370044
35. Campos FAD, Bertuzzi R, Dourado AC, Santos VGF, Franchini E. Energy demands in taekwondo athletes during combat simulation. *Eur J Appl Physiol.* 2012; 112(4):1221–1228. <https://doi.org/10.1007/s00421-011-2071-4> PMID: 21769736
36. Peine S, Knabe S, Carrero I, Brundert M, Wilhelm J, Ewert A, et al. Generation of normal ranges for measures of body composition in adults based on bioelectrical impedance analysis using the seca mBCA. *Int J Body Compos Res.* 2013; 11: 67–76.
37. Eston R, Reilly T. *Kinanthropometry and exercise physiology laboratory manual: tests, procedures and data: volume two: physiology*: Routledge; 2013.
38. Pařízková J. *Lean body mass and depot fat during ontogenesis in humans. Body fat and physical fitness*: Springer; 1977. pp. 24–51.
39. Bull BS, Aller R, Houwen B. MCHC-Red Cell Index or Quality Control Parameter. *Haematology.* 1996:43–43.
40. Hardeman MR, Dobbe JG, Ince C. The Laser-assisted Optical Rotational Cell Analyzer (LORCA) as red blood cell aggregometer. *Clin Hemorheol Microcirc.* 2001; 25(1):1–11. Epub 2002/01/16. PMID: 11790865.
41. Hardeman MR, Goedhart P, Breederveld D. Laser diffraction ellipsometry of erythrocytes under controlled shear stress using a rotational viscosimeter. *Clin Chim Acta.* 1987; 165(2–3):227–234. Epub 1987/06/15. PMID: 3652448.
42. Baskurt OK, Boynard M, Cokelet GC, Connes P, Cooke BM, Forconi S, et al. New guidelines for hemorheological laboratory techniques. *Clin Hemorheol Microcirc.* 2009; 42(2):75–97. Epub 2009/05/13. <https://doi.org/10.3233/CH-2009-1202> PMID: 19433882.
43. Ingebrigtsen J, Bendiksen M, Randers MB, Castagna C, Krstrup P, Holtermann A. Yo-Yo IR2 testing of elite and sub-elite soccer players: performance, heart rate response and correlations to other interval tests. *J Sports Sci.* 2012; 30(13):1337–1345. Epub 2012/08/08. <https://doi.org/10.1080/02640414.2012.711484> PMID: 22867048.
44. Capranica L, Lupo C, Cortis C, Chiodo S, Cibelli G, Tessitore A. Salivary cortisol and alpha-amylase reactivity to taekwondo competition in children. *Eur J Appl Physiol.* 2012; 112(2):647–652. <https://doi.org/10.1007/s00421-011-2023-z> PMID: 21643917
45. Chiodo S, Tessitore A, Cortis C, Cibelli G, Lupo C, Ammendolia A, et al. Stress-related hormonal and psychological changes to official youth Taekwondo competitions. *Scand J Med Sci Sports.* 2011; 21(1):111–119. <https://doi.org/10.1111/j.1600-0838.2009.01046.x> PMID: 20030779
46. Perret C, Mueller G. Validation of a new portable ergospirometric device (Oxycon Mobile®) during exercise. *Int J Sports Med.* 2006; 27(05):363–367.
47. de Moraes Bertuzzi RC, Franchini E, Kokubun E, Kiss MAPDM. Energy system contributions in indoor rock climbing. *Eur J Appl Physiol.* 2007; 101(3):293–300. <https://doi.org/10.1007/s00421-007-0501-0> PMID: 17602238
48. de Campos Mello F, de Moraes Bertuzzi RC, Grangeiro PM, Franchini E. Energy systems contributions in 2,000 m race simulation: a comparison among rowing ergometers and water. *Eur J Appl Physiol.* 2009; 107(5):615. <https://doi.org/10.1007/s00421-009-1172-9> PMID: 19707782
49. Lopes-Silva JP, da Silva Santos JF, Branco BHM, Abad CCC, de Oliveira LF, Loturco I, et al. Caffeine ingestion increases estimated glycolytic metabolism during taekwondo combat simulation but does not improve performance or parasympathetic reactivation. *PLoS one.* 2015; 10(11):e0142078. <https://doi.org/10.1371/journal.pone.0142078> PMID: 26539982
50. di Prampero PE, Ferretti G. The energetics of anaerobic muscle metabolism: a reappraisal of older and recent concepts. *Respir Physiol.* 1999; 118(2):103–115.
51. Gastin PB. Energy system interaction and relative contribution during maximal exercise. *Sports Med.* 2001; 31(10):725–741. PMID: 11547894
52. Santos VG, Franchini E, Lima-Silva AE. Relationship between attack and skipping in taekwondo contests. *J Strength Cond Res.* 2011; 25(6):1743–1751. <https://doi.org/10.1519/JSC.0b013e3181ddfb0f> PMID: 21512402

53. Araujo MP, Nobrega AC, Espinosa G, Hausen M, Castro RR, Soares PP, et al. Proposal of a New Specific Cardiopulmonary Exercise Test for Taekwondo Athletes. *J Strength Cond Res.* 2017; 31(6):1525–1535. Epub 2015/12/24. <https://doi.org/10.1519/JSC.0000000000001312> PMID: 28538301.
54. Marcon G, Franchini E, Jardim JR, Barros Neto TL. Structural analysis of action and time in sports: judo. *J Quant Anal Sports.* 2010; 6(4):1–15.
55. Salvador A, Suay F, Martínez–Sanchis S, Simon VM, Brain PF. Correlating testosterone and fighting in male participants in judo contests. *Physiol Behav.* 1999; 68(1):205–209.
56. Rosenthal R, Cooper H, Hedges L. Parametric measures of effect size. *The handbook of research synthesis.* 1994:231–244.
57. Cohen J. A power primer. *Psychol Bull.* 1992; 112(1):155–159. PMID: 19565683
58. Brun JF, Varlet-Marie E, Raynaud de Mauverger E. Relationships between insulin sensitivity measured with the oral minimal model and blood rheology. *Clin Hemorheol Microcirc.* 2012; 51(1):29–34. Epub 2012/01/14. <https://doi.org/10.3233/CH-2011-1426> PMID: 22240352.
59. McArdle WD, Katch FI, Katch VL. *Exercise physiology: nutrition, energy, and human performance.* Lippincott Williams & Wilkins; 2010.
60. Brun JF, Bouchahda C, Chaze D, Benhaddad AA, Micallef JP, Mercier J. The paradox of hematocrit in exercise physiology: which is the "normal" range from an hemorheologist's viewpoint? *Clin Hemorheol Microcirc.* 2000; 22(4):287–303. Epub 2000/11/18. PMID: 11081466.
61. Francesconi RP, Hubbard RW, Szlyk PC, Schnakenberg D, Carlson D, Leva N, et al. Urinary and hematologic indexes of hypohydration. *J Appl Physiol.* 1987; 62(3):1271–1276. Epub 1987/03/01. <https://doi.org/10.1152/jappl.1987.62.3.1271> PMID: 3571082.
62. Armstrong LE, Maresh CM, Castellani JW, Bergeron MF, Kenefick RW, LaGasse KE, et al. Urinary indices of hydration status. *Int J Sport Nutr.* 1994; 4(3):265–279. Epub 1994/09/01. PMID: 7987361.
63. Armstrong LE, Soto JA, Hacker FT Jr., Casa DJ, Kavouras SA, Maresh CM. Urinary indices during dehydration, exercise, and rehydration. *Int J Sport Nutr.* 1998; 8(4):345–355. Epub 1998/12/08. PMID: 9841955.
64. Bizjak DA, Brinkmann C, Bloch W, Grau M. Increase in red blood cell-nitric oxide synthase dependent nitric oxide production during red blood cell aging in health and disease: a study on age dependent changes of rheologic and enzymatic properties in red blood cells. *PLoS one.* 2015; 10(4):e0125206. <https://doi.org/10.1371/journal.pone.0125206> PMID: 25902315
65. Spell DW, Jones DV, Harper WF, Bessman JD. The value of a complete blood count in predicting cancer of the colon. *Cancer Detect Prev.* 2004; 28(1):37–42. <https://doi.org/10.1016/j.cdp.2003.10.002> PMID: 15041076
66. Baskurt OK, Meiselman KM. *Blood rheology and hemodynamics.* Thieme Medical Publishers: NY 10001; 2003. pp. 435–450.
67. Henkelman S, Dijkstra-Tiekstra MJ, de Wildt-Eggen J, Graaff R, Rakhorst G, van Oeveren W. Is red blood cell rheology preserved during routine blood bank storage? *Transfusion.* 2010; 50(4):941–948. Epub 2009/12/17. <https://doi.org/10.1111/j.1537-2995.2009.02521.x> PMID: 20003060.
68. Grau M, Pauly S, Ali J, Walpurgis K, Thevis M, Bloch W, et al. RBC-NOS-dependent S-nitrosylation of cytoskeletal proteins improves RBC deformability. *PLoS One.* 2013; 8(2):e56759. Epub 2013/02/21. <https://doi.org/10.1371/journal.pone.0056759> PMID: 23424675; PubMed Central PMCID: PMC3570529.
69. Mohanty J, Nagababu E, Rifkind JM. Red blood cell oxidative stress impairs oxygen delivery and induces red blood cell aging. *Front Physiol.* 2014; 5:84. <https://doi.org/10.3389/fphys.2014.00084> PMID: 24616707
70. Tikhomirova IA. The effect of dehydration on macro- and microrheological blood properties. *Clin Hemorheol Microcirc.* 2002; 26(2):85–90. Epub 2002/06/26. PMID: 12082256.
71. Falcó C, Vayá A, Simó M, Contreras T, Santaolara M, Aznar J. Influence of fibrinogen levels on erythrocyte aggregation determined with the Myrenne aggregometer and the Sefam erythro-aggregometer. *Clin Hemorheol Microcirc.* 2005; 33(2):145–151. PMID: 16151262
72. Gaudard A, Varlet-Marie E, Bressolle F, Mercier J, Brun JF. Nutrition as a determinant of blood rheology and fibrinogen in athletes. *Clin Hemorheol Microcirc.* 2004; 30(1):1–8. Epub 2004/02/18. PMID: 14967877.
73. Solá E, Vayá A, Corella D, Santaolara ML, Espana F, Estellés A, et al. Erythrocyte hyperaggregation in obesity: determining factors and weight loss influence. *Obesity.* 2007; 15(8):2128–2134. <https://doi.org/10.1038/oby.2007.253> PMID: 17712132
74. Poggi M, Palareti G, Biagi R, Legnani C, Parenti M, Babini A, et al. Prolonged very low calorie diet in highly obese subjects reduces plasma viscosity and red cell aggregation but not fibrinogen. *Int J Obes Relat Metab Disord.* 1994; 18(7):490–496. PMID: 7920875

75. Horswill CA, Scott JR, Dick RW, Hayes J. Influence of rapid weight gain after the weigh-in on success in collegiate wrestlers. *Med Sci Sports Exerc.* 1994; 26(10):1290–1294. PMID: [7799774](#)
76. Bridge CA, McNaughton LR, Close GL, Drust B. Taekwondo exercise protocols do not recreate the physiological responses of championship combat. *Int J Sports Med.* 2013; 34(7):573–581. Epub 2013/01/09. <https://doi.org/10.1055/s-0032-1327578> PMID: [23296399](#).
77. Hausen M, Soares PP, Araújo MP, Porto F, Franchini E, Bridge CA, et al. Physiological responses and external validity of a new setting for taekwondo combat simulation. *PLoS One.* 2017; 12(2):e0171553. <https://doi.org/10.1371/journal.pone.0171553> PMID: [28158252](#)
78. Matsushige KA, Hartmann K, Franchini E. Taekwondo: Physiological responses and match analysis. *J Strength Cond Res.* 2009; 23(4):1112–1117. <https://doi.org/10.1519/JSC.0b013e3181a3c597> PMID: [19528839](#)
79. Gaitanos GC, Williams C, Boobis LH, Brooks S. Human muscle metabolism during intermittent maximal exercise. *J Appl Physiol.* 1993; 75(2):712–719. <https://doi.org/10.1152/jappl.1993.75.2.712> PMID: [8226473](#)
80. Rodríguez FA, Mader A. World Book of Swimming From Science to Performance. In: Seifert L, Chollet D, Mujika I, editors. *Energy systems in swimming.* New York: Nova; 2011. pp. 225–240.
81. Mader A. Glycolysis and oxidative phosphorylation as a function of cytosolic phosphorylation state and power output of the muscle cell. *Eur J Appl Physiol.* 2003; 88(4–5):317–338. <https://doi.org/10.1007/s00421-002-0676-3> PMID: [12527960](#)