

CASE REPORT

Metastatic yolk sac tumor masquerading as multifocal hepatocellular carcinoma in a young adult: A case report

Kamran Mushtaq^{1,2}  | Muhammad Umair Khan^{1,2}  | Deema Al Soub³ |
Sheija Mary Koshy⁴ | Maher Petkar⁴ | Khalid Mohsin Al Ejji¹

¹Department of Gastroenterology, Hamad Medical Corporation, Doha, Qatar

²Executive and Continuing Professional Education, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA

³Department of Palliative Medicine, National Center for Cancer Care and Research (NCCCR), Doha, Qatar

⁴Department of Laboratory Medicine and Pathology, Hamad General Hospital, Doha, Qatar

Correspondence

Kamran Mushtaq, Department of Gastroenterology, Hamad Medical Corporation, Doha, Qatar.
Email: kmushtaq@hamad.qa and bkamrans@hotmail.com

Funding information

Qatar National Library

Abstract

Primary yolk sac tumor of the liver is very rare and can present as multifocal liver lesions. Multifocal nature may mimic other diagnoses such as hepatocellular carcinoma. Early recognition and therapeutic intervention are important as the prognosis of metastatic yolk sac tumors is poor. We present a case of a young adolescent who presented with bleeding per rectum abdominal pain and multiple liver lesions.

KEYWORDS

germ cell tumor, metastasis, multifocal hepatocellular carcinoma, tumor in young adult, yolk sac tumor

1 | INTRODUCTION

Extragenital germ cell tumors occur much more commonly in males than in females and are usually seen in young adults.¹ These are aggressive neoplasms and can arise virtually anywhere, but typically, the site of origin is in the midline (mediastinum, retroperitoneum, or pineal gland). The diagnosis should be considered in any patient with a poorly defined epithelial malignancy, particularly young individuals. They are also associated with a high tumor burden on the diagnosis.^{2,3} Yolk sac tumors of the liver are extremely rare and may not be considered until

the biopsy is performed. In young patients with multiple liver lesions, one of the differential diagnoses to consider is the yolk sac tumor. We present one patient with multiple liver lesions, which mimicked multifocal liver lesions.

2 | CASE PRESENTATION

A 19-year-old boy with no chronic medical illness presented with abdominal pain, weight loss, and bleeding per rectum for the last 2 months. His abdominal pain was vague, colicky in nature, and associated with reduced

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

appetite and food aversion. He also developed altered bowel habits mainly diarrhea and occasionally blood mixed in stools. No extraintestinal symptoms were reported. No family history of inflammatory bowel disease or any malignancy was present. No history of any prior surgery. He smoked cigarettes occasionally for <1 month. The clinical exam was remarkable for mild right upper quadrant tenderness. No palpable lymph nodes. The digital rectal exam was negative for fresh blood or melena. The scrotal exam was normal. Initial labs revealed mild anemia and slightly elevated transaminases (Table 1).

US abdomen revealed an enlarged liver measuring 19.7 cm in length. Multiple heterogeneous hyperechoic lesions were noted in the liver, the largest measuring 9.9 × 9.7 in the right lobe and 10 × 8.3 in the left lobe. The main portal vein measured 15 mm in diameter. There was a filling defect in the intrahepatic main portal vein extending into the main right and left branches, suggestive of thrombosis. A few periportal lymph nodes were noted, the largest measuring 17 × 14 mm. Mild ascites.

Based on the initial ultrasound malignancy was suspected and an MRI of the abdomen was performed, which showed enlarged liver with multiple focal lesions 11 cm in maximal dimension, involving all segments with the largest two at the left lobe and the subdiaphragmatic part of the right lobe. Tumor thrombus of the left and

right portal branches as well as some of the segmental branches. The lesions had more radiological features of multi-centric primary malignant liver neoplasm particularly hepatocellular carcinoma than metastases. Dilated intrahepatic bile ducts particularly at the left lobe were seen. Enlarged lymph nodes at the retrocaval, porta hepatis, upper paraaortic, and mesenteric regions 23 mm in maximal short axis, impressive of metastatic lymph nodes (Figure 1A,B). Alpha-fetoprotein (AFP) was requested and came very high (Table 1).

The patient was discussed in the tumor board although the radiologic impression was multifocal hepatocellular carcinoma plan was to biopsy the lesion and perform a colonoscopy and pan CT scan to rule out other primary lesions. A colonoscopy was performed as well to rule out primary colonic malignancy given a history of bleeding per rectum.

Contrast-enhanced CT scan revealed multiple large necrotic liver masses in all segments, showing heterogeneous enhancement, the largest at segment VII, measuring 13.0 × 9.6 cm. large filling defect was seen at the distal part of the portal vein, likely representing tumor thrombus. A heterogenous soft tissue mass was seen at the sigmoid mesentery infiltrating the sigmoid colon wall. Multiple enhancing peritoneal nodules were seen at the abdomen and pelvis, the largest in the pelvic region measuring 3.8 × 2.8 cm suggesting peritoneal metastasis. Multiple enlarged lymph nodes at mesenteric, retrocaval, and paraaortic regions. Other structures were unremarkable. Moderate free fluid is seen in the abdomen and pelvis. No lung or bony lesions were seen (See Figure 2).

Colonoscopic examination revealed a circumferential ulcerative and infiltrative mass lesion of around 6–7 cm, extending from 19 to 26 cm from the anal verge (Figure 3). The rest of the colonic mucosa appeared normal, and biopsies were taken; suspicion of the primary colonic tumor was raised. Gastroscopy was also performed to rule out the GI origin of the tumor and was normal. An ultrasound-guided liver biopsy was performed on one of the larger liver lesions.

The liver biopsy consisted of multiple fragments of tumor tissue, composed of predominantly clear cells arranged in reticular, solid, and focal hepatoid and microcystic growth patterns. In places, the tumor cells exhibited marked nuclear pleomorphism with a high nuclear-cytoplasmic ratio. Mitotic figures were easily identifiable. Frequent hyaline globules were noted, and foci of necrosis were also seen (Figure 4A,B).

A wide panel of immunostains was performed, which revealed that the tumor was positive for Cytokeratin AE1/AE3, Glypican 3, CDX2 (strong patchy positivity), and SALL4 (Figure 4C,D). The hyaline globules stained weakly for AFP, but the tumor cells were AFP-negative.

TABLE 1 Laboratory results

Laboratory parameters	Initial lab	Reference
Hemoglobin	11.9	13–17 g/dl
White blood cells	12.2	4–10 ³ /μl
Platelets	375	150–400 ³ /μl
MCV	57.8	80–100 fl
Urea	4.4	2–7 mmol/L
Creatinine	80	62–106 μmol/L
ALT	61	5–50 U/L
AST	44	5–35 U/L
ALP	120	20–150 U/L
Bilirubin	12	3–21 μmol/L
INR	1.4	202–415 μmol/L
Alpha-fetoprotein	2492	0–6 IU/ml
Uric acid	532	200–430 μmol/L
Beta-HCG	4	<5 mIU/ml
HIV	Negative	
HBsAg	Negative	
Anti-HCV Ab	Negative	

Abbreviations: ALT, alanine transaminase; AST, aspartate transaminase; HBV, hepatitis B virus; hCG, human choriongonadotropin; HCV, hepatitis C infection; HIV, human immunodeficiency virus; INR, International normalized ratio; MCV, mean corpuscular volume.




FIGURE 1 (A) MRI abdomen T2: Showing multiple liver lesions with necrosis and heterogenous enhancement. (B) MRI abdomen T1: Showing multiple liver lesions with necrosis and heterogenous enhancement

Immunostains for arginase, Hep par 1, CD30, Oct3/4, and S100 were negative.

The tumor morphology, along with a combination of positive staining for cytokeratin, Glypican 3, SALL4, CDX2, and high serum AFP levels were highly suggestive of a yolk sac tumor. In addition, in this biopsy, no other germ cell tumor morphology was noted. The appearances in the separately sent sigmoid colon biopsy showed similar tumor morphology and were likely representing metastasis of the sigmoid colon.

After the diagnosis was confirmed and revealed to the family discussion was made regarding further treatment. It was discussed with the patient that the likelihood of cure is very low and there is a high risk of treatment with chemotherapy. The patient was seen by the palliative care team as part of multidisciplinary management. The patient and family opted for high-risk chemotherapy. Given the high tumor burden, there was a high risk of tumor lysis syndrome as well as the dose was adjusted for liver and renal parameters. The patient was started on Bleomycin, Etoposide 50% dose reduction, Cisplatin 25% dose reduction, and dexamethasone 5 mg IV. For tumor



FIGURE 2 Venous phase CT scan of abdomen showing multiple large heterogenous liver lesions with thrombosis in the portal vein




FIGURE 3 Colonoscopy showing a circumferential mass lesion in the sigmoid colon

lysis syndrome Rasburicase IV pretreatment, with allopurinol was started. After the first cycle, the patient became severely ill with respiratory distress and was admitted to ICU requiring noninvasive ventilation. After his condition stabilized, he received a second cycle of chemotherapy on day 5 of the first cycle. After the second cycle, his clinical condition deteriorated with episodes of vomiting resulting in aspiration pneumonia, and the patient was intubated and started on mechanical ventilation. The patient was neutropenic and was started on broad-spectrum antibiotics including Meropenem IV and Vancomycin IV. On day 6 of the first cycle, the patient had a cardiac arrest and could not be revived.




FIGURE 4 (A) Low power view of the liver biopsy showing fragments of a tumor exhibiting variable architectural patterns (H and E $\times 2$). (B) High power view of the tumor, which exhibits marked cytological atypia with occasional intracellular hyaline globules (H and E $\times 20$). (C) The tumor cells are strongly positive for SALL4. (D) The tumor cells displaying strong Glypican 3 positivity

3 | DISCUSSION

Yolk sac tumors are germ cell tumors that most commonly arise from the gonads.⁴ Extragonadal germ cell tumors are rare and primary yolk sac tumors of the liver are extremely rare with only few case reports being described.⁵ Primary yolk sac tumors of the liver arise in young adults and the prognosis is variable with several case reports showing successful treatment with chemotherapy.

These tumors can pose a significant diagnostic challenge as multiple liver lesions in young adults have a broad differential diagnosis. Multifocal liver lesions in young adults and children can be of benign or malignant origin. Common benign causes include focal nodular hyperplasia, liver abscesses, and hepatic adenomas. While causes of malignant multifocal liver lesions in adults include Infantile hepatoblastoma, lymphoma, hepatocellular carcinoma, hepatic epithelioid hemangioendotheliomas, and metastases.^{6,7} MRI abdomen with a histologic diagnosis is key to establishing a definite diagnosis. In our case, there was a high probability of a primary liver tumor with a second likely differential of liver metastases from the colon until the biopsy results revealed the histology of a yolk sac tumor from the liver lesion as well as from the sigmoid colon. Correct diagnosis is essential for early recognition and early initiation of appropriate therapy.

Histologically, yolk sac tumors display marked heterogeneity, with numerous architectural patterns described. A combination of patterns is common, in varying proportions. The various histological patterns include microcystic/reticular, macrocystic, papillary, solid, sarcomatoid, glandular, hepatoid, myxomatous, perivascular, and polyvesicular vitelline patterns. Schillar-Duval bodies are another characteristic finding but are not required for diagnosis. Hyaline globules are usually noted in the hepatoid pattern, which are usually AFP-positive.

Primary hepatic yolk sac tumors are rare and to date, several case reports have been reported.^{8–14} There are no specific radiologic diagnostic features of yolk sac tumors and hence their ability to masquerade other liver lesions.¹⁰ Several treatment modalities have been reported. The treatment depends on whether the tumor is resectable or not, the presence of metastasis, and the initial response to chemotherapy. Successful treatment with surgical resection has also been achieved.⁵ Treatment by embolization for liver yolk sac tumors has been reported.¹⁴ While Liver transplant as a treatment option has also been reported successfully.^{11,12} Despite these options, the mainstay treatment modality in the extragonadal yolk sac tumors is chemotherapy as these tumors are highly sensitive to chemotherapy. The main chemotherapy is bleomycin etoposide- and cisplatin-based. Other regimens have also been used. It is important to consider palliative care in patients

with metastasis and high tumor burden. Discussing end-of-life issues is challenging especially with young patients and their families as the diagnosis itself had a deep psychological impact hence early multidisciplinary psychological and palliative care is crucial.¹⁵

Yolk sac histology and the presence of metastasis are predictors of poor survival in germ cell tumors.³ Other established predictors of poor prognosis include mediastinal primary origin, nonpulmonary metastases, and elevated markers (AFP > 10,000 ng/ml, LDH > 10× upper limit of normal, beta-HCG > 50,000 IU/L).¹⁶ Our patient was classified as a poor prognosis group due to the presence of poor prognostic markers including yolk sac histology and colonic metastasis.

4 | CONCLUSION

Yolk sac tumors are rare germ cell tumors that are extragonadal and rarely present in the Liver. The yolk sac tumor of the liver can mimic other liver lesions such as hepatocellular carcinoma and metastatic lesions. Imaging and tumor markers are helpful, but the ultimate diagnosis of a yolk sac tumor is made by histology. Metastatic yolk sac tumor of the liver has a poor prognosis. Physicians evaluating multiple liver lesions with elevated AFP, especially in young adults should consider germ cell tumors such as yolk sac tumors as a differential diagnosis.

AUTHOR CONTRIBUTIONS

KM, MUK, and DS were involved in writing the case, reviewing the literature, and clinical care of the patient. SMK and MP reviewed the histopathologic specimens. KA reviewed the literature and was involved in the clinical care of the patient. All authors reviewed the manuscript.

ACKNOWLEDGMENT

None.

FUNDING INFORMATION

The funding was provided by the Qatar National Library, Doha Qatar.

CONFLICT OF INTEREST

All authors declare no relevant financial conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

ETHICS STATEMENT

The case report was approved by the Medical Research Center (MRC) of Hamad Medical Corporation (MRC-04-21-904), Doha, Qatar.

CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

ORCID

Kamran Mushtaq  <https://orcid.org/0000-0002-1787-640X>

Muhammad Umair Khan  <https://orcid.org/0000-0003-3013-4352>

REFERENCES

1. Stang A, Trabert B, Wentzensen N, et al. Gonadal and extragonadal germ cell tumours in the United States, 1973-2007. *Int J Androl.* 2012;35(4):616-625.
2. Hainsworth JD, Greco FA. Extragonadal germ cell tumors and unrecognized germ cell tumors. *Semin Oncol.* 1992;19(2):119-127.
3. Mortazavi N, Mahzooni P, Taheri D, Jalilian M, Novin K. Germ cell tumor's survival rate in young patients. *Iran J Cancer Prev.* 2015;8(4):e3440.
4. Pinto MT, Cárcano FM, Vieira AGS, Cabral ERM, Lopes LF. Molecular biology of pediatric and adult male germ cell tumors. *Cancer.* 2021;13(10):2349.
5. Gunawardena SA, Siriwardana HPP, Wickramasinghe SYLS, Madurawe MN, Fernando R. Primary endodermal sinus (yolk sac) tumour of the liver. *Eur J Surg Oncol.* 2002;28(1):90-91.
6. Hegde SV, Dillman JR, Lopez MJ, Strouse PJ. Imaging of multifocal liver lesions in children and adolescents. *Cancer Imaging.* 2013;12(3):516-529.
7. Karakuchi N, Toyota K, Kajikawa R, et al. Alpha-fetoprotein-producing colon carcinoma with rapidly increasing liver metastases that resulted in death: a case report. *J Anus Rectum Colon.* 2021;5(1):100-106.
8. Narita T, Moriyama Y, Ito Y. Endodermal sinus (yolk sac) tumour of the liver. A case report and review of the literature. *J Pathol.* 1988;155(1):41-47.
9. Villaschi S, Balistreri P. Endodermal sinus tumour of the liver. *Histopathology.* 1991;18(1):86-88.
10. Szucs RA, Narla LD. Primary endodermal sinus tumor of the liver detected by CT. *Pediatr Radiol.* 1992;22(6):449-450.
11. Lenci I, Tariciotti L, Baiocchi L, et al. Primary yolk sac tumor of the liver: incidental finding in a patient transplanted for hepatocellular carcinoma. *Transpl Int.* 2008;21(6):598-601.
12. Abramson LP, Pillai S, Acton R, Melin-Aldana H, Superina R. Successful orthotopic liver transplantation for treatment of a hepatic yolk sac tumor. *J Pediatr Surg.* 2005;40(7):1185-1187.
13. Whelan JS, Stebbings W, Owen RA, Calne R, Clark PI. Successful treatment of a primary endodermal sinus tumor of the liver. *Cancer.* 1992;70(9):2260-2262.

14. Higuchi T, Kikuchi M. Yolk sac tumor of the liver treated with transcatheter arterial embolization. *Am J Gastroenterol.* 1993;88(7):1125-1126.
15. Oing C, Giannatempo P, Honecker F, Oechsle K, Bokemeyer C, Beyer J. Palliative treatment of germ cell cancer. *Cancer Treat Rev.* 2018;71:102-107.
16. International Germ Cell Consensus Classification: a prognostic factor-based staging system for metastatic germ cell cancers. International Germ Cell Cancer Collaborative Group. *J Clin Oncol.* 2016;15(2):594-603. Sep 21 [cited 2021 Aug 20]. doi:[10.1200/JCO.1997.15.2.594](https://doi.org/10.1200/JCO.1997.15.2.594)

How to cite this article: Mushtaq K, Khan MU, Al Soub D, Koshy SM, Petkar M, Al Ejji KM. Metastatic yolk sac tumor masquerading as multifocal hepatocellular carcinoma in a young adult: A case report. *Clin Case Rep.* 2023;11:e6861. doi:[10.1002/ccr3.6861](https://doi.org/10.1002/ccr3.6861)