



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review

Molecular pathogenesis of secondary bacterial infection associated to viral infections including SARS-CoV-2

Sounik Manna^a, Piyush Baidara^b, Santi M. Mandal^{c,*}^a Department of Microbiology, Midnapore College (Autonomous), Paschim Medinipur, India^b Department of Molecular Microbiology and Immunology, University of Missouri, Columbia, USA^c Central Research Facility, Indian Institute of Technology Kharagpur, Kharagpur 721302, India

ARTICLE INFO

Article history:

Received 11 April 2020

Received in revised form 11 June 2020

Accepted 6 July 2020

Keywords:

Viral infection

Secondary bacterial infection

Immune response

SARS-CoV-2

ABSTRACT

Secondary bacterial infections are commonly associated with prior or concomitant respiratory viral infections. Viral infections damage respiratory airways and simultaneously defects both innate and acquired immune response that provides a favorable environment for bacterial growth, adherence, and facilitates invasion into healthy sites of the respiratory tract. Understanding the molecular mechanism of viral-induced secondary bacterial infections will provide us a chance to develop novel and effective therapeutic approaches for disease prevention. The present study describes details about the secondary bacterial infection during viral infections and their immunological changes. The outcome of discussion avails an opportunity to understand possible secondary bacterial infections associated with novel SARS-CoV-2, presently causing pandemic outbreak COVID-19.

© 2020 Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

Introduction	1398
Mechanisms of viral predisposition to secondary bacterial infections	1398
Virus facilitates bacteria adherence and colonization	1398
Impaired innate and adaptive immune response	1398
Virus-induced immune suppression	1399
Exploitation of bacteria through viral infection	1399
Secondary infection of bacteria for viral infection	1400
Relationship between pathological changes and host responses	1400
Viral factors implicated in severity of infection	1400
Role of inflammatory responses in disease severity	1401
SARS-CoV-2 and secondary bacterial infection	1401
Conclusion	1402
Funding	1402
Conflict of interests	1402
Ethical approval	1402
Acknowledgement	1402
References	1402

* Corresponding author.

E-mail address: mandalsm@gmail.com (S.M. Mandal).

Introduction

Bacteria and viruses often occupy the same niches, however, interest in their potential collaboration in promoting wellness or disease development has only recently gained attention. The interaction of some bacteria and viruses are well characterized and researchers are typically more interested in the location of the infection than the manner of cooperation [1]. There are two overarching types of bacteria-virus disease causing interactions by direct interactions that in some way aid the viruses, and indirect interactions of aiding bacteria. The virus-promoting direct interactions occur when the virus exploits a bacterial component to facilitate penetration into the host cell [2]. On the other hand, indirect interactions result in increased bacterial pathogenesis as a consequence of viral infection. Enteric viruses mainly utilize the direct pathway, while respiratory viruses indirectly affect bacteria [3]. The causative agent of the current pandemic outbreak, SARS-CoV-2 is also reported to associate with secondary bacterial and fungal coinfections as patients showed the symptoms of atypical bacterial pneumonia [1–3]. It has been known that SARS-CoV is the nearest virus to SARS-CoV-2 and their systemic comparison may further help in understanding the disease pathogenesis during COVID-19 and secondary bacterial infections [4]. However, SARS-CoV-2 is reported to evolve in the human body that might affect its virulence, infectivity, and transmissibility [5], thus the biology of SARS-CoV-2 further needs to be understood in detail, especially in case of secondary bacterial infections. This review focuses on some key examples of how virus-bacteria interactions impact the infection process across the two organ systems and provide evidence supporting this as an emerging premise in infectious disease pathogenesis.

Human body is inhabited by a diverse microbial community that is collectively coined as commensal microbiota. Recent research has greatly advanced our understanding of how commensal microbiota affects host health. Among the various kinds of pathogenic infections of the host, viral infections constitute one of the most serious public health problems worldwide. During the infection process, viruses may have substantial and intimate interactions with the commensal microbiota [4]. A plethora of evidence suggests that the commensal microbiota regulated by invading viruses through diverse mechanisms, thereby having stimulatory or suppressive roles in viral infections. Furthermore, the integrity of the commensal microbiota can be troubled by invading viruses, causing dysbiosis in the host, and further influencing virus infectivity. In the present article, we discuss current insights into the regulation of viral infection by the commensal microbiota. We also draw attention to the disruption of microbiota homeostasis by several viruses [5].

Viral infections predispose patients to secondary bacterial infections, which often have a more severe clinical course. The mechanisms underlying post-viral bacterial infections are complex and include multifactorial processes mediated by interactions between viruses, bacteria, and the host immune system [6]. Studies over the past 15 years have demonstrated that unique microbial communities reside on the mucosal surfaces of the gastrointestinal tract and the respiratory tract, which has both direct and indirect effects on host defense against viral infections [7]. The antiviral immune responses induced by acute respiratory infections such as influenza are associated with changes in microbial composition and function (“dysbiosis”) in the respiratory and gastrointestinal tract, which in turn may alter subsequent immune function against secondary bacterial infection or alter the dynamics of inter-microbial interactions, thereby enhancing the proliferation of potentially pathogenic bacterial species [4]. Here, we summarize the literature on the interactions between host microbial communities and host defense, including how influenza and other acute respiratory

viral infections disrupt these interactions, thereby contributing to the pathogenesis of secondary bacterial infections.

Mechanisms of viral predisposition to secondary bacterial infections

Respiratory viruses are often encouraged, secondary bacterial infections by supplementing the outgrowth opportunistic bacterial pathogens. Viral infections damage the respiratory airway by both histologically and functionally [8]. Cell loss, goblet cell hyperplasia, altered mucus secretion, reduced ciliary beat frequency, dis-coordinated mucociliary clearance function along with reduced oxygen exchange are characteristics of viral infection [9]. These effects are associated with different molecular mechanisms by which the predisposition of the virus occurs in the respiratory tract that facilitates the bacterial infection, simultaneously (Fig. 1).

Virus facilitates bacteria adherence and colonization

Epithelial cells of the respiratory tract facilitate bacterial adherence using different mechanisms during viral infection, while the disease severity varies upon virus, bacterial strain, and experimental model used. Respiratory syncytial virus (RSV) reported to bind directly with *Haemophilus influenzae* and *Staphylococcus pneumoniae*, so enhancing bacterial proximity to the epithelial monolayer and supplementing attachment to the host cell receptors [10]. Interestingly, RSV glycoprotein expressed and localized on the host cell membrane upon infection that is further act as bacterial receptors for pneumococcal binding [11,12]. Respiratory viruses are also able to up-regulate the expression of host cell membrane protein to facilitate their binding [9,10]. Influenza virus, known to induce the production of type I interferons (IFNs) that results in reduced production of CCL2 (chemokine (C-C motif) ligand 2) and so decreased recruitment of macrophages thus results in enhanced colonization of *S. pneumoniae* in mice [11]. Influenza virus also makes mice susceptible to pneumonia caused by *Staphylococcus aureus* where both virus and bacterial load increased during co-infection [12,13]. RSV, parainfluenzavirus-3, and influenza viruses reported increasing the bacterial adherence upon infection, in both primary and immortalized epithelial cells with different differences [9]. Recently, it has been reported that surface glycoprotein adhesion molecule-1 (ICAM-1) presents up-regulated expression during RSV and adenovirus infection. ICAM-1 presented as a cognate ligand for the Type IV pilus of non-typeable *H. influenzae* induced secondary bacterial infection via adherence promotion [14]. RSV infections also confirmed to increase adherence for *S. pneumoniae* in human nasopharyngeal cells (HEp-2) and pneumocyte type II cells (A549). The enhanced pneumococcal adherence in epithelial cells results in bacterial accumulation and thus secondary bacterial infection may occur [15,16].

Impaired innate and adaptive immune response

Secondary bacterial infection with viral infections can drastically change the immunity level by the uncontrolled bacterial growth [17,18]. Viral infection impaired the phagocyte functions and induced alveolar macrophages (AM) depletion results in the facilitation of bacterial infection via apoptosis [19]. Infection in AM caused by the influenza virus reported to result in low-level production of cytokines and chemokines which is also required for recruitment and activation of neutrophils [18]. Influenza virus infection can diminish NADPH oxidase-dependent phagocytic bacterial clearance and so increase the chances of secondary bacterial infections [20]. Virus-induced dysregulation of proinflammatory cytokine response played a major role in promoting secondary bacterial infections [21,22]. While type I IFNs are well reported for




Fig. 1. Proposed model for viral-induced secondary bacterial infections.

antiviral and immuno-stimulatory properties, their, inappropriate or excessive production may have harmful effects. Results suggested that IFNs have an important role in the production of specific cytokines like immunosuppressive IL-10 and pro-inflammatory IL-6, suppression of which is key to the linkage between innate and adaptive immune response, involving IL-7 and IL-23, dendritic cells, macrophages, natural killer cells, CD4⁺ and CD8⁺, T cells, all these results in the reduced ability to efficiently eliminate bacterial co-infections and thus promote secondary bacterial infections. Dysregulation of antimicrobial peptides is another way used by the upper respiratory tract (URT) viruses to facilitate secondary bacterial infections. Expression of antimicrobial peptides lipocalin2, CAMP, REG3B, S100A8, and S100A9 found down-regulated during URT infections [23]. In another study, RSV infection reported to reduce the gene expression of chinchilla beta-defensin 1 (an orthologue of human beta defensin3, hBD-3), and thus RSV infection found to induce 10–100 increase in nasopharynx (NP) load of *H. influenza* [24].

Virus-induced immune suppression

Bacterial superinfections are associated with viral immune suppression condition which is not only limited to the respiratory tract but also related to several organ failures. These immunosuppressive viruses include especially HIV, cytomegalovirus (CMV), measles virus (MeV), and others. The prime characteristic of HIV infection is the severe reduction of CD4⁺ T cells that make the patient much more susceptible to develop bacterial superinfections like tuberculosis [25]. It has been reported that HIV infection ease the *Mycobacterium* infection along with various characteristics to macrophages that constrains bacterial clearance which promotes bacterial colonization with disease development. These HIV associated features to macrophages comprise the up-regulation of cell surface receptors for *Mycobacterium* entry, manipulation of bactericidal pathways used by macrophages, transformed chemotaxis, induction of immune response causing Th1/Th2 imbalance and apoptotic response that mediated by diminished tumor necrosis factor [26]. CMV infection is also connected with immune system paralysis that is characterized by enhanced IL-10 and NFκB production, lymphopenia, reduction of IFN-γ producing T cells,

and increase tissue damage by dysregulated cytokine production. Immune response during CMV infection is generated a favorable environment for secondary bacterial infections [27–29]. MeV infection is another example of immunosuppressive characteristics that are associated with a secondary bacterial infection. MeV reported to have associated with several weeks of immune suppression, loss of delayed-type hypersensitivity responses, and thus increased susceptibility to secondary bacterial infections [30]. It has been reported that MeV infection destroys pre-existing antibodies that increase the vulnerability to pathogens that result in secondary bacterial infections [31].

Exploitation of bacteria through viral infection

The majority of reported direct bacteria-virus interactions are associated with infection in the gastrointestinal tract caused by viruses. In body system, commensal bacteria are considered the first line of defense against invading pathogens by out-competing their disease-promoting counterparts and limiting tissue accessibility. Undoubtedly, enteric viruses encounter these large numbers of diverse commensal bacteria, but rather than always preventing infection, some viruses evolved to exploit this contact facilitating the disease developmental process [32].

Under in vitro conditions, viruses may able to directly bind to their target cells and undergo replication. However, this strategy may prove problematic in the gastrointestinal tract whereas a large number of bacteria occupy tissue surfaces, directly competing for receptor binding sites, and reducing the likelihood of pathogenic bacterial proliferation or virus attachment. Other components, like mucus or enzymatic secretions, may also interfere or assist the infection process. To circumvent this, rather than compete for host cell-binding sites, some viruses can utilize bacterial ligands to enhance their association with more accessible host cells, initiating infection. This same strategy may be employed by some viruses that may not exclusively target the host's epithelial cells rather uses bacteria to assist infection of other cell types in addition to or exclusive of epithelial cells [33,34].

Although the nature of the interaction remains the same, there may be additional benefits to viruses with bacteria interactions other than direct disease progression. Studies have also shown

that association with fecal microbiota increased poliovirus environmental fitness and stability, as exposure to bacteria or their polysaccharides decreased the efficacy of virus inactivation by heat and bleach, potentially aiding viral survival in the environment [35]. This observation was further supported by the higher susceptibility to inactivation with heat observed in a poliovirus mutant that did not bind LPS as efficiently. Furthermore, the introduction of these bacterial polysaccharide components has been found to enhance wild type poliovirus binding to its host cells expressing its receptor [35]. In short, gastrointestinal microbiota not only increase poliovirus infectivity but may also promote virus transfer to the next host. Thus, there are numerous ways that direct viral interaction with bacteria aid viral pathogenesis, and this topic is an emerging area of study in microbiology.

Bacterial species often benefit from viral infections, the virus-induced disease state can allow normally harmless bacteria to become opportunistically pathogenic. Under normal, healthy circumstances, direct competition between microbes limits pathogen invasion by saturating colonization sites, priming barrier immunity to produce antimicrobials, and increasing the immune response to invading microorganisms [36]. When microbial populations are disrupted, niches previously inaccessible to invading pathogens become available, and the microenvironment where native microbiota previously was outcompeted their disease-causing counterparts are compromised. The overall ways viruses aid bacteria pathogenesis include a complex combination of cellular receptor up-regulation, disruption of the epithelial layers, displacement of commensal bacteria, and immune system suppression [37]. Arguably, the interactions between influenza viruses and pathogenic bacteria (i.e., *S. pneumoniae*, *S. aureus*, and *H. influenzae*) remain the best-studied within the human body, and they exemplify some of these mechanisms with both viruses and bacteria benefitting from the relationship. Influenza viruses not only damage the host epithelium (e.g., apoptosis), they also provide potential binding sites for bacteria through three mechanisms such as neuraminidase cleavage of sialic acid from host cells, bacterial host receptor up-regulation, and host regeneration of the common bacterial receptors fibrin and fibrinogen [38,39]. This pattern of host damage is common amongst upper respiratory tract viruses and bacteria [40,41], and these interactions are summarized in Table 1. Thus, these numerous dynamics that exist between the influenza virus and bacteria exemplify the multiple complex ways of viral infection that can indirectly assist bacterial infection [42,43].

Secondary infection of bacteria for viral infection

Epidemiological and clinical research clearly says that bacterial secondary infection can significantly increase after viral infections [44]. As an example, Up to 75% of that influenza-infected patient that goes on to acquire pneumonia, and they are established that have bacterial co-infection [45]. Bacterial secondary infection after influenza infection looks to occur normally. Data revealed that up to 65% of the laboratory established that cases of influenza infection showed bacterial secondary infection. However, Klein et al. [46] established through a meta-analysis that this figure ranged between 11 and 35%. In the site of influenza epidemic or pandemic bacterial secondary infection can have an alarming situation that is predominantly risk to immune-compromised or immune-suppressed patients. Immunosuppression is related to more severe illness and much of a higher risk of mortality from a secondary bacterial infection [47]. During pandemic Swine influenza in 2009, there was an arising patient in hospital pneumonia cases as a consequence of secondary bacterial pneumonia, which was recognized in 29–55% of mortalities [48–50].

Relationship between pathological changes and host responses

Viral infections are different with specific characteristics those are developed in the infected area and for that reason, bacterial colonization occurs. Bacteria can easily adhere and penetrate the infected region of the host cell [51]. Influenza virus produces neuraminidase, which also helps to increase the adhesion of some bacterial species by eliminating sialic acid which helps to depict the host cell receptors [39,52]. Some other bacteria such as *Streptococci* have sialic acid which allows for direct attachment with the viral hemagglutinin (HA) and this is expressed by influenza-infected host cells [52,53]. When the viral infection occurs they damaged the host cell by directly or inflammation of immune cell responses. The wound or damaged tissue welcomes the bacterial cells which can easily adhere and this viral infection increased by bacterial adhesion. For example, the contacts of apical receptors like integrins help the adhesion of bacteria such as *S. aureus* and *Pseudomonas aeruginosa* [54–57]. At the onset of viral infection, the host inflammatory responses are the reason for the up-regulation in the expression of host receptor molecules and other molecules that bacteria can use as a receptor [52,58]. As an example, Cundell et al. [59] reported that increased performance of the G-protein-coupled platelet-activating factor (PAF) is utilized by some bacteria such as *S. pneumoniae*, for their attachment and colonization in the endothelial cells [59,60]. In difference, it has been recommended that the platelet-activating factor (PAF) receptor does not disturb the preliminary bacterial adherence and colonization but is more involved with secondary bacterial transition or spread into the blood and facilitates the growth of invasive disease [61].

The respiratory viruses enter through the upper respiratory tract and these virus-infected in the acute infection of the respiratory tract. When the influenza virus or other viral infections are beginning in the respiratory tract than they might influence the secondary bacterial infection such as pneumonia by changing the microbial composition in the upper respiratory tract. For that secondary infection, there is developed the growth of pathogens and large bacterial loads has occurred in the lower respiratory tract [61]. Acute viral infection and secondary bacterial colonization have a bidirectional relationship.

Viral factors implicated in severity of infection

Humans are constantly exposed to viral agents with endogenous or exogenous in origin, but few of them are the cause of significant pathogenicity or severity of the diseases. Viruses are require host cells to replicate and assemble infectious offspring. The character of the viral factors are more important which imparted in the disease progression and tissue damage [62]. Recently, Miller et al. [63], demonstrated the role of viral factors showed their severity in infant respiratory tract illness. Both, host and viral risk factors are necessary to understand which may help to develop the disease prevention strategies. When a viral strain move between hosts is accountable to new symptoms appearance or stably adapted to new hosts. Viruses are able to initiate infection, and start their replication by the help of specific virulence factors [64]. It is well documented that different viral strains possess variable viral factors and therefore different degree of pathogenicity observed with different virulence [65]. One of the most important viral factor is located in their genome to defeat the host barriers, such as a point mutation in 5' untranslated region (5' UTR) of polio vaccine strain but the strain without mutation shows the greater pathogenicity in hosts [64]. Thus, viral genome is important factor to control the tropism, virus entry to host, shedding and transmission.

Table 1
List of secondary bacterial infections and their immune responses during viral infection.

Viral infection	Secondary bacterial disease	Bacterial name	Immune response
Influenza	Pneumonia, Otitis media, Sinusitis, meningitis	<i>S. pneumoniae</i> , <i>S. aureus</i> , <i>S. pyrogenes</i> , <i>M. catarrhalis</i> , <i>N. meningitidis</i>	loss of ciliary function, produce INF, Mucus response. Destroy phagocytic cell, produce cytokines including TNF- α , IL-6, and pro-IL-1 β .
Coronavirus	Pneumonia	<i>H. influenzae</i>	activate the innate immune response or induce costimulatory signals for adaptive immunity
Adenovirus	Pneumonia	<i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>S. aureus</i> , <i>M. catarrhalis</i>	Produce cytokines, loss of ciliary function, produce INF, Mucus response.
Measles virus	Otitis media, Pneumonia, Tracheobronchitis.	<i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>S. aureus</i> ,	activate the innate immune response, produce cytokines including TNF- α , IL-6, and pro-IL-1 β , Mucus response.
Human rhinovirus	Pneumonia, Otitis media, Sinusitis,	<i>S. pneumoniae</i> , <i>H. influenzae</i> , <i>S. aureus</i>	Alters in pulmonary immune response, produce cytokines, inflammatory cell recruitment.
Parainfluenza virus	Pneumonia	<i>S. pneumoniae</i>	Destroy phagocytic cell, produce cytokines including TNF- α , IL-6, and pro-IL-1 β .
Respiratory tract viral infection	Pneumonia, lungs problem	<i>S. pneumoniae</i> , <i>S. aureus</i> , <i>H. influenzae</i> , <i>pseudomonas</i> .	Mucus response, IFN secretion, epithelial cell death, cytokine release, loss of ciliary function.
UTI viral infection BK Virus, Herpes virus	UTI infection is occur	<i>E. Coli</i> , <i>K. pneumoniae</i> , <i>P. aeruginosa</i>	chemokines and cytokines such as CXCL8, CCL2, interleukins (IL-6, IL-8, IL-10, IL-17A), and granulocyte colony stimulating factor (G-CSF) are release.

Viruses are able to developed immunomodulation pathway to undermine the host immune response. The virus-encoded decoy receptors are targeted to cytokines and chemokines of host cells [66]. Usually, primary bacterial clearance needs innate immunity with bacterial phagocytosis by local AM. The phagocytic activity of CD11c + alveolar macrophages in broncho alveolar lavage fluid was lower. To define the effect of interferon c (IFNc) were immunized with exogenous IFNc which inhibited alveolar macrophage-mediated phagocytosis of pneumococci both in vitro and in vivo. There was also repressed the external expression of the scavenger receptor which is responsible for phagocytosis of pneumococci by alveolar macrophages. Thirdly, interferon improved the surface expression of MHC class II antigen for that region interferon c inhibits bacterial phagocytosis and destroys innate defense against pneumococcal infection. in vivo treatment with the interferon c specific antibody had little consequence on the progress of viral infection but prohibited increased susceptibility to pneumococcal infection by developed macrophage expression of inflammatory cytokines and inhibiting pre-regulation of MHC class II expression [67]. The post-viral bacterial infection mechanism is very complex because different multifunctional interaction occurs between the virus, bacteria, and also host immune system. The pathogenesis of the respiratory super-infection has been endorsed by the direct mucosal or epithelial damage by the influenza virus. This viral infection increased the bacterial colonization of the upper and lower respiratory tracts. The host immune responses are the main cause of increased susceptibility to secondary bacterial infections. Another way, modern research suggests that our microbial populations be present in on our mucosal surfaces and probably affect our immune responses which create the characters of ecological relations among hosts and pathogens. The past 10 years result revealed the complex relationship of secondary microbial infections in the host body during viral infection [68]. In the time of secondary infection, different bacteria affect the physical substance in the host body, number of genes, and metabolic activities in different organ systems. Actually, respiratory viral infections can decrease the host immune responses, for that reason microbial growth conditions in the URT, LRT is high. When the influenza virus attacks host, body induced antiviral interferon pathways but this immune system is inadequate during host defense against secondary bacterial infections, therefore the pathogenic bacteria can easily proliferate and infected the host cell. Thus, molecular basis of virulence is polygenic in nature by consideration of viral factors. Therefore, viruses are efficiently altering the host cell response and exhibit greater pathogenicity [69].

Role of inflammatory responses in disease severity

The epithelial cells and mucosal immune cells in lung airway are the hotspot of respiratory viruses. These primary cells are getting first infected with viral agents and immediately strat to produce a range of mediators such as type I interferon (IFN), proinflammatory cytokines, and chemokines, which are the major molecules for the development of inflammation process in infected area. These are key factors in virus control molecules until vaccines are not available. It is well documented that infections induce the excessive production of reactive oxidative species (ROS), which are one of the important inflammatory mediators [70], and consequences of several adverse physiological responses [71]. The expression of genes involved in respiratory process in host, their biogenesis signalling and antioxidant enzymes are regulated at the onset of viral infection. Overproduction of ROS along with mitochondrial damage increase the inflammation process in infected zone [72]. The entry of viral agents into lung and their colonization trigger for acute exacerbations of chronic obstructive pulmonary disease (AECOPD). A details study on patients with AECOPD revealed the elevation level of IL-6, TNF- α , and MCP-1 which are also increased several folds during coinfection [73].

Co-infections of viral and bacterial pathogens increase the disease susceptibility and outcome is more severe with higher mortality. In healthy lung tissue, neutrophils, monocytes, macrophages, dendritic cells, natural killer (NK), and other innate lymphocyte (ILC), B and T cells are found in abundance where AM are mostly abundant. After establishment of infection both pathogen-associated molecular pattern (PAMPs) and damage-associated molecular patterns (DAMPs) can engage in pattern recognition receptors (PRRs) to active the cellular signalling process for the production of soluble interferons (IFNs) to limit the replication process of invading pathogens. Interestingly, this PRR ligation [e.g., toll-like receptors (TLRs) 2 and 4] start to induce the overproduction of cytokines and chemokines which regulate the activation and recruitment of inflammatory molecules in the lung [74]. Recognition of PAMPs by specific PRRs have substantial impact on disease susceptibility and increased pathogen transmission. Therefore, it is important to accurately tune the immune and inflammatory mechanisms to restore the lung tissues from maximum damage [75].

SARS-CoV-2 and secondary bacterial infection

Like other well studied respiratory viral infections including the 1918 influenza outbreak [76] and 2009 H1N1 pandemic [77], the current pandemic outbreak of SARS-CoV-2 is also reported to

associate with secondary bacterial infections, thus poor outcomes and fatalities [2,78]. Recent studies reported a low rate of associated secondary bacterial and fungal infections with COVID-19 [3], however, SARS-CoV-2 is a novel coronavirus that suggested as proximally originated [79] and thus we can't ignore the fact that our current knowledge SARS-CoV-2 is very limited. Very recently, cohort studies of tuberculosis, sequelae, and COVID-19 co-infection revealed SARS-CoV-2 associated infection in patients with TB, however, the number of patients were less and no associated disease pathogenesis need to be determined [80,81]. Further investigation and detailed studies are needed to explore the role of latent or active TB associated with COVID-19 disease severity and progression [82]. Another cohort study of 11 patients suggested that the cases with pre-existing chronic obstructive pulmonary disease or complicated with secondary bacterial pneumonia as the most severe outcome cases of COVID-19 [83].

SARS-CoV-2 is very much similar to the SARS-CoV, though it caused ongoing pandemic outbreak COVID-2. SARS-CoV has been already reported to regulate immune function-related gene expression in human monocytes. Immune-related gene expression suggested that SARS-CoV infection downregulates IFN- α/β -inducible and cathepsin/proteasome genes while differential regulated genes include, TLR/TLR-signaling, cytokine/cytokine receptor-related, chemokine/chemokine receptor-related, lysosome-related, MHC/chaperon-related, and fibrosis-related genes [84]. In a different study, SARS-CoV also reported suppressing type 1 IFN production [85]. Downregulation and differential regulation of immune genes suggested a possible encouraging environment for secondary bacterial infections. Additionally, SARS patients has been reported to have secondary bacterial infections and confirmed the presence of different bacteria in various human samples, including *Escherichia coli*, *K. pneumoniae*, *P. aeruginosa*, Methicillin-Resistant *S. aureus* (MRSA), etc. [86]. During the SARS outbreak, increased rates of MRSA infections have been reported in SARS patients that further suggested the role of secondary bacterial infections in viral disease pathogenesis [87].

A recent study, confirmed SARS-CoV-2 infection in pregnant women and studied the clinical features and neonatal outcomes. The study suggested timely use of antibiotics might prevent COVID-19 associated secondary bacterial infections and can reduce mortality and complications during pregnancy [88].

SARS-CoV-2 RNA has also been found in stool samples of COVID-19 patients [89]. This raises the question of gastrointestinal infection of SARS-CoV-2 and also suggested a possible fecal-oral route of disease transmission. Further, high expression levels of ACE2 mRNA in the gastrointestinal system, avail indirect evidence of SARS-CoV-2 infection [90,91]. Preliminary studies suggested gastrointestinal infection [92,93] and strong interaction of SARS-CoV-2 with the gastrointestinal system that has high microbiome diversity and possible chances of immune suppression and secondary bacterial infections. Additionally, a recent study reviewed the secondary bacterial and fungal infection in coronavirus patients and suggested guidelines for the use and prescription of antimicrobial in SARS-CoV, SARS-CoV-2, and related viral infections [3]. However, further, clinical studies may provide future directions for SARS-CoV-2 and related secondary bacterial infections.

Conclusion

Viral infections avail bacterial infections by using multiple strategies, including providing a more susceptible site for adhesion, altering immune response, and invasive infection by cell and tissue damage. Secondary bacterial infections then make the clinical outcomes more severe that is an alarming situation in concern of public health. However, antibiotics can diminish the effect

of secondary bacterial infections, in-depth understanding of the viral-host, bacterial-host, and viral-bacterial interaction is important to fight against these infections. Especially in the focus of rapid increasing antibiotic resistance and adaptation of microbes may lead to escape from vaccine-induced immunity. The eventual goal of understanding the molecular pathogenesis of viral infection associated secondary bacterial infections is to development of more efficient diagnostics and therapeutic strategies. Further, it becomes an important fight against the current or future potential of pandemics and the emergence of novel viruses like SARS-CoV-2.

Funding

No funding sources.

Conflict of interests

None declared.

Ethical approval

Not required.

Acknowledgement

SMM is grateful to IIT-Kharagpur for providing necessary facilities and support.

References

- [1] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020;382:727–33.
- [2] Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395(10229):1054–62.
- [3] Rawson TM, Moore LSP, Zhu N, Ranganathan N, Skolimowska K, Gilchrist M, et al. Bacterial and fungal co-infection in individuals with coronavirus: a rapid review to support COVID-19 antimicrobial prescribing. *Clin Infect Dis* 2020, <http://dx.doi.org/10.1093/cid/ciaa530>.
- [4] Xu J, Zhao S, Teng T, Abdalla AE, Zhu W, Xie L, et al. Systematic comparison of two animal-to-human transmitted human coronaviruses: SARS-CoV-2 and SARS-CoV. *Viruses* 2020;12(2):244.
- [5] Shen Z, Xiao Y, Kang L, Ma W, Shi L, Zhang L, et al. Genomic diversity of severe acute respiratory syndrome-coronavirus 2 in patients with coronavirus disease 2019. *Clin Infect Dis* 2020, <http://dx.doi.org/10.1093/cid/ciaa203>.
- [6] McCullers JA. The co-pathogenesis of influenza viruses with bacteria in the lung. *Nat Rev Microbiol* 2014;12(4):252–62.
- [7] Hament JM, Aerts PC, Flerer A, Van Dijk H, Harmsen T, Kimpen JLL, et al. Direct binding of respiratory syncytial virus to pneumococci: a phenomenon that enhances both pneumococcal adherence to human epithelial cells and pneumococcal invasiveness in a murine model. *Pediatr Res* 2005;58(6):1198–203.
- [8] Avadhanula V, Wang Y, Portner A, Adderson E. Nontypeable *Haemophilus influenzae* and *Streptococcus pneumoniae* bind respiratory syncytial virus glycoprotein. *J Med Microbiol* 2007;56(Pt. 9):1133–7.
- [9] Avadhanula V, Rodriguez CA, DeVincenzo JP, Wang Y, Webby RJ, Ulett GC, et al. Respiratory viruses augment the adhesion of bacterial pathogens to respiratory epithelium in a viral species- and cell type-dependent manner. *J Virol* 2006;80(4):1629–36.
- [10] Smith CM, Sandrini S, Datta S, Freestone P, Shafeeq S, Radhakrishnan P, et al. Respiratory syncytial virus increases the virulence of *Streptococcus pneumoniae* by binding to penicillin binding protein 1a a new paradigm in respiratory infection. *Am J Respir Crit Care Med* 2014;190(2):196–207.
- [11] Nakamura S, Davis KM, Weiser JN. Synergistic stimulation of type I interferons during influenza virus coinfection promotes *Streptococcus pneumoniae* colonization in mice. *J Clin Invest* 2011;121(9):3657–65.
- [12] Iverson AR, Boyd KL, McAuley JL, Plano LR, Hart ME, McCullers JA. Influenza virus primes mice for pneumonia from *Staphylococcus aureus*. *J Infect Dis* 2011;203(6):880–8.
- [13] Smith AM, Adler FR, Ribeiro RM, Gutenkunst RN, McAuley JL, McCullers JA, et al. Kinetics of coinfection with influenza a virus and *Streptococcus pneumoniae*. *PLoS Pathog* 2013;9(3):e1003238.
- [14] Novotny LA, Bakaletz LO. Intercellular adhesion molecule 1 serves as a primary cognate receptor for the type IV pilus of nontypeable *Haemophilus influenzae*. *Cell Microbiol* 2016;18(8):1043–55.

- [15] Nguyen DT, Louwen R, Elberse K, Van Amerongen G, Yüksel S, Luijckx A, et al. *Streptococcus pneumoniae* enhances human respiratory syncytial virus infection in vitro and in vivo. *PLoS One* 2015;10(5):e0127098.
- [16] Hament JM, Aerts PC, Flerer A, Van Dijk H, Harmsen T, Kimpfen JLL, et al. Enhanced adherence of *Streptococcus pneumoniae* to human epithelial cells infected with respiratory syncytial virus. *Pediatr Res* 2004;55(6):972–8.
- [17] Lijek RS, Weiser JN. Co-infection subverts mucosal immunity in the upper respiratory tract. *Curr Opin Immunol* 2012;24(4):417–23.
- [18] Braciale TJ, Sun J, Kim TS. Regulating the adaptive immune response to respiratory virus infection. *Nat Rev Immunol* 2012;12(4):295–305.
- [19] Ghoneim HE, Thomas PG, McCullers JA. Depletion of alveolar macrophages during influenza infection facilitates bacterial superinfections. *J Immunol* 2013;191(3):1250–9.
- [20] Sun K, Metzger DW. Influenza infection suppresses NADPH oxidase-dependent phagocytic bacterial clearance and enhances susceptibility to secondary methicillin-resistant *Staphylococcus aureus* infection. *J Immunol* 2014;192(7):3301–7.
- [21] Mehta D, Petes C, Gee K, Basta S. The role of virus infection in deregulating the cytokine response to secondary bacterial infection. *J Interferon Cytokine Res* 2015;35(12):925–34.
- [22] Davidson S, Maini MK, Wack A. Disease-promoting effects of type I interferons in viral, bacterial, and coinfections. *J Interferon Cytokine Res* 2015;35(4):252–64.
- [23] Robinson KM, Kolls JK, Alcorn JF. The immunology of influenza virus-associated bacterial pneumonia. *Curr Opin Immunol* 2015;34:59–67.
- [24] McGillivray G, Mason KM, Jurcisek JA, Peeples ME, Bakaletz LO. Respiratory syncytial virus-induced dysregulation of expression of a mucosal B-defensin augments colonization of the upper airway by non-typeable *Haemophilus influenzae*. *Cell Microbiol* 2009;11(9):1399–408.
- [25] Vittor AY, Garland JM, Gilman RH. Molecular diagnosis of TB in the HIV positive population. *Ann Glob Health* 2014;80(6):476–85.
- [26] Bell LCK, Noursadeghi M. Pathogenesis of HIV-1 and mycobacterium tuberculosis co-infection. *Nat Rev Microbiol* 2018;16(2):80–90.
- [27] Limaye AP, Kirby KA, Rubenfeld GD, Leisenring WM, Bulger EM, Neff MJ, et al. Cytomegalovirus reactivation in critically ill immunocompetent patients. *JAMA* 2008;300(4):413–22.
- [28] Al-Omari A, Aljamaan F, Alhazzani W, Salih S, Arabi Y. Cytomegalovirus infection in immunocompetent critically ill adults: literature review. *Ann Intensive Care* 2016;6(1):110.
- [29] Alyazidi R, Murthy S, Slyker JA, Gantt S. The potential harm of cytomegalovirus infection in immunocompetent critically ill children. *Front Pediatr* 2018;6:96.
- [30] Griffin DE. Measles virus-induced suppression of immune responses. *Immunol Rev* 2010;236:176–89.
- [31] Mina MJ, Kula T, Leng Y, Li M, De Vries RD, Knip M, et al. Measles virus infection diminishes preexisting antibodies that offer protection from other pathogens. *Science* 2019;366(6465):599–606.
- [32] Berkhout B. With a little help from my enteric microbial friends. *Front Med* 2015;2:30.
- [33] Karst SM, Wobus CE. A working model of how noroviruses infect the intestine. *PLoS Pathog* 2015;11(2):e1004626.
- [34] Racaniello VR. One hundred years of poliovirus pathogenesis. *Virology* 2006;344(1):9–16.
- [35] Robinson CM, Jesudhasan PR, Pfeiffer JK. Bacterial lipopolysaccharide binding enhances virion stability and promotes environmental fitness of an enteric virus. *Cell Host Microbe* 2014;15(1):36–46.
- [36] Khosravi A, Mazmanian SK. Disruption of the gut microbiome as a risk factor for microbial infections. *Curr Opin Microbiol* 2013;16(2):221–7.
- [37] McCullers JA. Insights into the interaction between influenza virus and pneumococcus. *Clin Microbiol Rev* 2006;19(3):571–82.
- [38] Lillebaek T, Dirksen A, Vynnycky E, Baess I, Thomsen VØ, Andersen ÅB. Stability of DNA patterns and evidence of *Mycobacterium tuberculosis* reactivation occurring decades after the initial infection. *J Infect Dis* 2003;188(7):1032–9.
- [39] McCullers JA, Bartmess KC. Role of neuraminidase in lethal synergism between influenza virus and *Streptococcus pneumoniae*. *J Infect Dis* 2003;187(6):1000–9.
- [40] Murphy TF, Bakaletz LO, Smeesters PR. Microbial interactions in the respiratory tract. *Pediatr Infect Dis J* 2009;28(10 Suppl):S121–6.
- [41] Bosch AATM, Biesbroek G, Trzcinski K, Sanders EAM, Bogaert D. Viral and bacterial interactions in the upper respiratory tract. *PLoS Pathog* 2013;9(1):e1003057.
- [42] Hendaus MA, Jomha FA, Alhammadi AH. Virus-induced secondary bacterial infection: a concise review. *Ther Clin Risk Manag* 2015;11:1265–71.
- [43] Hament J-M, Kimpfen JL, Flerer A, Wolfs TF. Respiratory viral infection predisposing for bacterial disease: a concise review. *FEMS Immunol Med Microbiol* 1999;26(3–4):189–95.
- [44] Gupta RK, George R, Nguyen-Van-Tam JS. Bacterial pneumonia and pandemic influenza planning. *Emerg Infect Dis* 2008;14(8):1187–92.
- [45] Zambon MC. The pathogenesis of influenza in humans. *Rev Med Virol* 2001;11(4):227–41.
- [46] Klein EY, Monteforte B, Gupta A, Jiang W, May L, Hsieh YH, et al. The frequency of influenza and bacterial coinfection: a systematic review and meta-analysis. *Influenza Other Respi Viruses* 2016;10(5):394–403.
- [47] Rice TW, Rubinson L, Uyeki TM, Vaughn FL, John BB, Miller RR, et al. Critical illness from 2009 pandemic influenza A virus and bacterial coinfection in the United States. *Crit Care Med* 2012;40(5):1487–98.
- [48] Bacterial coinfections in lung tissue specimens from fatal cases of 2009 pandemic influenza A (H1N1)—United States, May–August 2009. *Morb Mortal Wkly Rep* 2009;58(38):1071–4.
- [49] Gill JR, Sheng ZM, Ely SF, Guinee DG, Beasley MB, Suh J, et al. Pulmonary pathologic findings of fatal 2009 pandemic influenza A/H1N1 viral infections. *Arch Pathol Lab Med* 2010;134(2):235–43.
- [50] Weinberger DM, Simonsen L, Jordan R, Steiner C, Miller M, Viboud C. Impact of the 2009 influenza pandemic on pneumococcal pneumonia hospitalizations in the United States. *J Infect Dis* 2012;205(3):458–65.
- [51] Selinger DS, Reed WP, McLaren LC. Model for studying bacterial adherence to epithelial cells infected with viruses. *Infect Immun* 1981;32(2):941–4.
- [52] Peltola VT, McCullers JA, Fink RJ, Fedson DS. Respiratory viruses predisposing to bacterial infections: role of neuraminidase. *Pediatr Infect Dis J* 2004;23(1 suppl):s87–97.
- [53] Okamoto S, Kawabata S, Nakagawa I, Okuno Y, Goto T, Sano K, et al. Influenza A virus-infected hosts boost an invasive type of *Streptococcus pyogenes* infection in mice. *J Virol* 2003;77(7):4104–12.
- [54] Sanford BA, Shelokov A, Ramsay MA. Bacterial adherence to virus-infected cells: a cell culture model of bacterial superinfection. *J Infect Dis* 1978;137(2):176–81.
- [55] Davison VE, Sanford BA. *Staphylococcus aureus* Adherence of to influenza A virus-infected Madin-Darby canine kidney cell cultures. *Infect Immun* 1981;32(1):118–26.
- [56] Bucior I, Pielage JF, Engel JN. *Pseudomonas aeruginosa* Pili and Flagella mediate distinct binding and signaling events at the apical and basolateral surface of airway epithelium. *PLoS Pathog* 2012;8(4):e1002616.
- [57] Smith AM, McCullers JA. Secondary bacterial infections in influenza virus infection pathogenesis. *Curr Top Microbiol Immunol* 2014;385:327–56.
- [58] Hakansson A, Kidd A, Wadell G, Sabharwal H, Svanborg C. Adenovirus infection enhances in vitro adherence of *Streptococcus pneumoniae*. *Infect Immun* 1994;62(7):2707–14.
- [59] Cundell DR, Gerard NP, Gerard C, Idanpaan-Heikkilä I, Tuomanen EI. *Streptococcus pneumoniae* anchor to activated human cells by the receptor for platelet-activating factor. *Nature* 1995;377(6548):435–8.
- [60] van der Sluijs KF, van der Poll T, Lutter R, Juffermans NP, Schultz MJ. Bench-to-bedside review: bacterial pneumonia with influenza – pathogenesis and clinical implications. *Crit Care* 2010;14(2):219.
- [61] McCullers JA, Iverson AR, Murray PJ. The platelet activating factor receptor is not required for exacerbation of bacterial pneumonia following influenza. *Scand J Infect Dis* 2008;40(1):11–7.
- [62] MacLachlan NJ, Dubovy EJ. Pathogenesis of viral infections and diseases. *Fenner's Vet Virol* 2011:43–74.
- [63] Miller AL, Gerard C, Schaller M, Gruber AD, Humbles AA, Lukacs MW. Deletion of CCR1 attenuates pathophysiologic responses during respiratory syncytial virus infection. *J Immunol* 2006;176(4):2562–7.
- [64] Albrecht T, Fons M, Boldogh IR, Alan S. In: Samuel Baron, editor. *Medical Microbiology*. 4th ed. Galveston (TX): University of Texas Medical Branch at Galveston; 1996. ISBN 0963117211. PMID 21413282.
- [65] Fuentes-González AM, Contreras-Paredes A, Manzo-Merino J, Lizano M. The modulation of apoptosis by oncogenic viruses. *Virology* 2013;10:182.
- [66] Felix J, Savvides S. Mechanisms of immunomodulation by mammalian and viral decoy receptors: insights from structures. *Nat Rev Immunol* 2017;17:112–29.
- [67] Akiko I, Ellen FF, Ryan DM. Early local immune defenses in the respiratory tract. *Nat Rev Immunol* 2017;17(1):7–20.
- [68] Denise EM, David WC, Stuart C. Secondary bacterial infections associated with influenza pandemics. *Front Microbiol* 2017;8:1041.
- [69] John CK, Jeffery KT. The role of viral, host, and secondary bacterial factors in influenza pathogenesis. *Am J Pathol* 2015;185(6):1528–36.
- [70] Yilong F, Andy Y, Peck GS, Francesca B, Pei-yong S, Maxime H. Modulation of inflammation and pathology during dengue virus infection by p38 MAPK inhibitor SB203580. *Antiviral Res* 2014;110:151–7.
- [71] Banerjee D, Mandal SM, Das A, Hegde ML, Das S, Bhakat KK, et al. Preferential repair of oxidized base damage in the transcribed genes of mammalian cells. *J Biol Chem* 2011;286(8):6006–16.
- [72] Shitao L, Bishi F, Meshram Chetan D. Innate immune and inflammatory responses to respiratory viruses. *Mediators Inflamm* 2019. Article ID 3146065, 2 pages.
- [73] Jingdong Z, Yue S, Lingxin X, Weijie Z, Ying L, Peter GG, et al. The expression of IL-6, TNF- α , and MCP-1 in respiratory viral infection in acute exacerbations of chronic obstructive pulmonary disease. *J Immunol Res* 2017. Article ID 8539294, 9 pages.
- [74] Iwasaki A, Pillai PS. Innate immunity to influenza virus infection. *Nat Rev Immunol* 2014;14:315–28, <http://dx.doi.org/10.1038/nri3665>.
- [75] Wenting Z, Qinghua H, Weiwei X, Yue Z, Jiang P, Huan X, et al. Advances in anti-tumor treatments targeting the CD47/SIRP α Axis. *Front Immunol* 2020;11:18.
- [76] Morens DM, Taubenberger JK, Fauci AS. Predominant role of bacterial pneumonia as a cause of death in pandemic influenza: implications for pandemic influenza preparedness. *J Infect Dis* 2008;198(7):962–70.
- [77] MacIntyre CR, Chughtai AA, Barnes M, Ridda I, Seale H, Toms R, et al. The role of pneumonia and secondary bacterial infection in fatal and serious outcomes of pandemic influenza A(H1N1)pdm09. *BMC Infect Dis* 2018;18(1):637.
- [78] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395(10223):507–13.

- [79] Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2. *Virological* 2020;26(4):450–2.
- [80] Motta I, Centis R, D'Ambrosio L, García-García J-M, Goletti D, Gualano G, et al. Tuberculosis, COVID-19 and migrants: preliminary analysis of deaths occurring in 69 patients from two cohorts. *Pulmonology* 2020;26(4):233–40.
- [81] Tadolini M, Codecasa LR, Jé-Mía García-García, Blanc F-X, Borisov S, Alffenaar J-W, et al. Active tuberculosis, sequelae and COVID-19 co-infection: first cohort of 49 cases. *Eur Respir J* 2020;56(1):2001398.
- [82] Liu Y, Bi L, Chen Y, Wang Y, Fleming J, Yu Y, et al. Active or latent tuberculosis increases susceptibility to COVID-19 and disease severity. *MedRxiv* 2020, <http://dx.doi.org/10.1101/2020.03.10.20033795>.
- [83] Dong X, Cao Y yuan, Lu X xia, Zhang J jin, Du H, Yan Y qin, et al. Eleven faces of coronavirus disease 2019. *Allergy Eur J Allergy Clin Immunol* 2020;75(7):1699–709.
- [84] Hu W, Yen YT, Singh S, Kao CL, Wu-Hsieh BA. SARS-CoV regulates immune function-related gene expression in human monocytic cells. *Viral Immunol* 2012;25(4):277–888.
- [85] Siu KL, Chan CP, Kok KH, Chiu-Yat Woo P, Jin DY. Suppression of innate antiviral response by severe acute respiratory syndrome coronavirus M protein is mediated through the first transmembrane domain. *Cell Mol Immunol* 2014;11(2):141–9.
- [86] Tan FLS, Loo WL, Tan SG, Wong CY, Tan Y-M. Severe acute respiratory syndrome in surgical patients: a diagnostic dilemma. *ANZ J Surg* 2005;75:21–6.
- [87] Yap FHY, Gomersall CD, Fung KSC, Ho P-L, Ho O-M, Lam PKN, et al. Increase in methicillin-resistant *Staphylococcus aureus* acquisition rate and change in pathogen pattern associated with an outbreak of severe acute respiratory syndrome. *Clin Infect Dis* 2004;39(4):511–6.
- [88] Yu N, Li W, Kang Q, Xiong Z, Wang S, Lin X, et al. Clinical features and obstetric and neonatal outcomes of pregnant patients with COVID-19 in Wuhan, China: a retrospective, single-centre, descriptive study. *Lancet Infect Dis* 2020;20(5):559–64.
- [89] Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, et al. First case of 2019 novel coronavirus in the United States. *N Engl J Med* 2020;382(10):929–36.
- [90] Harmer D, Gilbert M, Borman R, Clark KL. Quantitative mRNA expression profiling of ACE 2, a novel homologue of angiotensin converting enzyme. *FEBS Lett* 2002;532(1–2):107–10.
- [91] Yan R, Zhang Y, Guo Y, Xia L, Zhou Q. Structural basis for the recognition of the 2019-nCoV by human ACE2. *BioRxiv* 2020;367(6485):1444–8.
- [92] Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology* 2020;158(6):1831–3.e3.
- [93] Gao QY, Chen YX, Fang JY. Novel coronavirus infection and gastrointestinal tract. *J Dig Dis* 2019;21(3):125–6.