

# Thermoregulatory, metabolic, and cardiovascular responses during 88 min of full-body ice immersion – A case study

Coen C. W. G. Bongers<sup>1</sup>  | Thijs M. H. Eijsvogels<sup>1</sup>  | Dick H. J. Thijssen<sup>1,2</sup> | Maria T. E. Hopman<sup>1</sup> 

<sup>1</sup>Radboud Institute for Health Sciences, Department of Physiology, Radboud University Medical Center, Nijmegen, The Netherlands

<sup>2</sup>Research Institute for Sport and Exercise Science, Liverpool John Moores University, Liverpool, United Kingdom

## Correspondence

Maria T. E. Hopman, Department of Physiology 392, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands.  
Email: maria.hopman@radboudumc.nl

## Abstract

Exposure to extreme cold environments is potentially life-threatening. However, the world record holder of full-body ice immersion has repeatedly demonstrated an extraordinary tolerance to extreme cold. We aimed to explore thermoregulatory, metabolic, and cardiovascular responses during 88 min of full-body ice immersion. We continuously measured gastrointestinal temperature ( $T_{gi}$ ), skin temperature ( $T_{skin}$ ), blood pressure, and heart rate (HR). Oxygen consumption ( $VO_2$ ) was measured at rest, and after 45 and 88 min of ice immersion, in order to calculate the metabolic heat production.  $T_{skin}$  dropped significantly ( $28\text{--}34^\circ\text{C}$  to  $4\text{--}15^\circ\text{C}$ ) and  $VO_2$  doubled ( $5.7\text{--}11.3\text{ ml kg}^{-1}\text{ min}^{-1}$ ), whereas  $T_{gi}$  ( $37.6^\circ\text{C}$ ), HR (72 bpm), and mean arterial pressure (106 mmHg) remained stable during the first 30 min of cold exposure. During the remaining of the trial,  $T_{skin}$  and  $VO_2$  remained stable, while  $T_{gi}$  gradually declined to  $37.0^\circ\text{C}$  and HR and mean arterial blood pressure increased to maximum values of 101 bpm and 115 mmHg, respectively. Metabolic heat production in rest was 169 W and increased to 321 W and 314 W after 45 and 80 min of ice immersion. Eighty-eight minutes of full-body ice immersion resulted in minor changes of  $T_{gi}$  and cardiovascular responses, while  $T_{skin}$  and  $VO_2$  changed markedly. These findings may suggest that our participant can optimize his thermoregulatory, metabolic, and cardiovascular responses to challenge extreme cold exposure.

## KEYWORDS

cold exposure, G-Tummo meditation, hypothermia, thermoregulation

## 1 | INTRODUCTION

Hypothermia, defined as a core body temperature below  $35^\circ\text{C}$ , is associated with life-threatening complications, as has been dramatically illustrated by polar expeditions (i.e., Scott's Terra Nova Expedition), ship wrecks (i.e.,

Titanic), and military operations under extreme cold (i.e., World War II Eastern Front) (Shetty, 2003; Wissler, 2003). Furthermore, athletes competing in colder climates are at risk to develop hypothermia as well (Melau, Mathiassen, Stensrud, Tipton, & Hisdal, 2019). Even nowadays, hypothermia is still estimated to result in  $\sim 1,500$  deaths annually

Coen C. W. G. Bongers and Thijs M. H. Eijsvogels are contributed equally.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Physiological Reports* published by Wiley Periodicals, Inc. on behalf of The Physiological Society and the American Physiological Society.

in the USA (Brown, Brugger, Boyd, and Paal (2012)). In marked contrast, the world record holder of full-body ice immersion has repeatedly demonstrated an extraordinary tolerance to extreme cold when exposed to full-body ice immersion for more than 1 hr. His physiological responses during these cold challenges are unknown, but may reveal new information about regulatory mechanisms that allow for the extraordinary ability to sustain prolonged exposure to extreme cold.

The purpose of this case study was to examine the thermoregulatory, cardiovascular, and metabolic responses of the world record holder of full-body ice immersion during prolonged full-body contact with crushed ice. We expected to observe a drop in  $T_{gi}$  and concomitant drop in skin temperature (to minimize heat loss) and an elevation of the metabolic rate (e.g., shivering) to increase heat production (Sessler, 2009).

## 2 | CASE PRESENTATION AND METHODS

The world record holder of full-body ice immersion is a Caucasian male (50 year), with a height of 183 cm, and a body mass of 85 kg ( $BMI = 25.4 \text{ kg/m}^2$  and  $BSA = 2.07 \text{ m}^2$ ). He was positioned in an open-top container filled with crushed ice to the shoulder level (2.45 p.m.). The participant was wearing shorts only, whereas his body was in direct contact with the crushed ice. Core temperature (gastrointestinal temperature;  $T_{gi}$ ), skin temperature ( $T_{skin}$ ), blood pressure, and heart rate (HR) were measured continuously before, during and after ice exposure. Oxygen consumption was used to determine metabolic activity at rest and after 45 and 88 min of crushed ice immersion. According to prior mutual agreement, ice immersion was finished after 88 min. Signed written informed consent was obtained prior to the study. Ethical approval was waived as ice exposure was part of a demonstration session of the participant, so no additional burden was present. All measurements were conducted in line with the ethical standards of the Declaration of Helsinki.

The participant performed Asian G-Tummo meditation, 1 hr before and during the experiment. G-Tummo meditation consists of both somatic and neurocognitive components. The somatic component involves specialized breathing techniques, defined as “the vase,” that are accompanied with isometric muscle contractions. Moreover after inhalation that practitioners are holding their breath during which they contract both the abdominal and pelvic muscles (lower belly is shaping like a vase) (Evans-Wentz, 1958; Kozhevnikov, Elliott, Shephard, & Gramann, 2013). The neurocognitive component consists

of meditative visualization that requires the generation and maintenance of mental images of flames at specific locations in the body accompanied by intense sensations of bodily heat in the spine (Evans-Wentz, 1958; Kozhevnikov et al., 2013).

### 2.1 | Thermoregulatory responses

$T_{gi}$  was measured continuously (every 20 s and averaged per minute) using a telemetric temperature capsule (CorTemp system, HQ Inc) and an external recorder, which has been demonstrated to be safe and reliable (Byrne & Lim, 2007; Gant, Atkinson, & Williams, 2006). The participant ingested an individually calibrated capsule 5 hr prior to the experiment, to ensure passage through the stomach (Wilkinson, Carter, Richmond, Blacker, & Rayson, 2008).

$T_{skin}$  was measured continuously (every 20 s and averaged per minute) using wireless temperature recorders (iButton DS1922L, Dallas Semiconductor Corp) (Marken Lichtenbelt et al., 2006; Smith et al., 2009). The temperature recorders were attached to the skin using Tegaderm Film (Tegaderm).  $T_{skin}$  was measured at eight distinct locations (forehead, chest, scapula, shoulder, elbow, hand, thigh, and calf), and a weighted average  $T_{skin}$  per minute was calculated according to the ISO-9886 norm (2004).

### 2.2 | Cardiovascular responses

Responses in HR and arterial blood pressure were continuously measured during the first 80 min of ice immersion using an automated blood pressure monitoring device attached to the third digit of the left hand at a sampling rate of 100 Hz (Nexfin, BMEYE). A pulse oximeter (Nellcor N-200 E, Nellcor) and a manual sphygmomanometer were used every 10 min as a control for the continuous Nexfin measurements.

### 2.3 | Metabolic responses

Oxygen consumption ( $VO_2$ ) and carbon dioxide ( $VCO_2$ ) production were measured by indirect calorimetry using a calibrated gas exchange analyzer (Quark CPET, Cosmed) with a breathing mask. The mask was carefully placed over the mouth and nose to avoid of air leakage. Metabolic heat production ( $M$  in Watt) was subsequently calculated using the following formula:

$$M = VO_2 \frac{\left(\frac{RER-0.7}{0.3}\right) * E_c + \left(\frac{1.0-RER}{0.3}\right) * E_f}{60} * 1000 \quad (1)$$

where  $\text{VO}_2$  is the oxygen consumption in L/min, RER is the respiratory exchange ratio ( $\text{VCO}_2/\text{VO}_2$ ), and  $E_c$  and  $E_f$  are the caloric equivalents per liter of oxygen for the oxidation of carbohydrates (21.13 kJ) and fats (19.62 kJ), respectively (Murgatroyd, Shetty, & Prentice, 1993). During the test,  $\text{VO}_2$  and metabolic heat production were measured at baseline, after 45 min and after 80 min of ice immersion, and averaged over a 3-min recording.

### 3 | RESULTS


#### 3.1 | Thermoregulatory responses

Temperature of the crushed ice was  $0.14 \pm 0.06^\circ\text{C}$ .  $T_{\text{gi}}$  during the 10 min before ice immersion was stable at  $37.7 \pm 0.1^\circ\text{C}$ .  $T_{\text{gi}}$  did not change during the first 32 min of full-body contact with ice (Figure 1a). During the remaining 56 min,  $T_{\text{gi}}$  gradually decreased to  $37.0^\circ\text{C}$ .

$T_{\text{skin}}$  of the forehead (not in contact with ice) demonstrated a gradual decrease from  $34^\circ\text{C}$  (pre-immersion) to  $32^\circ\text{C}$  at the end of ice immersion (Figure 1b).  $T_{\text{skin}}$  of the upper chest (measured just above ice level) started at  $32^\circ\text{C}$  (pre-immersion) and reached a plateau at  $26^\circ\text{C}$  after 10 min of ice immersion (Figure 1b).  $T_{\text{skin}}$  of the shoulder, elbow, hand, and scapula (i.e., body areas which are in direct contact with ice) rapidly decreased from  $28\text{--}34^\circ\text{C}$  (pre-immersion) to temperatures as low as  $4\text{--}15^\circ\text{C}$  within 20 min of ice immersion, and remained stable at this level until the end of immersion. The thigh and calf, also areas in direct contact with ice, demonstrated a drop in  $T_{\text{skin}}$  from  $33\text{--}35^\circ\text{C}$  (pre-immersion) to  $18\text{--}20^\circ\text{C}$  during ice immersion. The decrease observed in the thigh and calf was slightly less than that of parts of the upper body that were in contact with the ice.

#### 3.2 | Cardiovascular responses


Resting HR before ice immersion was 77 bpm (range: 69–88 bpm) and HR remained relatively stable at 72 bpm (range: 66–85 bpm) during the first 40 min of ice immersion (Figure 2a). HR started to increase gradually during the last 40 min of ice immersion, reaching a maximum of 101 bpm. Mean arterial blood pressure at rest was 106 mmHg (94–123 mmHg) and remained stable during the first 60 min at 106 mmHg (87–123 mmHg) (Figure 2b). Mean arterial blood pressure increased to 115 mmHg (107–123 mmHg) during the last 10 min of ice immersion (70–80 min). The changes in HR and blood pressure across the ice immersion were confirmed by data obtained from a pulse oximeter and manual sphygmomanometer every 10 min, respectively, which suggested that our continuous heart rate and mean arterial blood pressure measurements were reliable.



**FIGURE 1** (a)  $T_{\text{gi}}$  before, during, and after ice immersion. Vertical dotted lines delineate the start ( $t = 0$  min) and end ( $t = 88$  min) of ice immersion. (b)  $T_{\text{skin}}$  before, during, and after ice immersion. Vertical dotted lines delineate the start ( $t = 0$  min) and end ( $t = 88$  min) of ice immersion. (c) Metabolic heat production before (baseline) and during (45 and 88 min) ice immersion

#### 3.3 | Metabolic responses

Oxygen consumption increased significantly from  $0.49 \pm 0.08$  L/min at rest to  $0.96 \pm 0.05$  L/min and  $0.93 \pm 0.07$  L/min after 45 and 80 min of ice immersion, respectively. Accordingly, metabolic heat production at baseline was 169 W and increased to 321 W and 312 W after 45 and 80 min of ice immersion (Figure 1c). No shivering or bodily movement was observed by the researchers or reported by the participant.



**FIGURE 2** (a) HR before and during ice immersion. A vertical dotted line delineates the start ( $t = 0$  min) of ice immersion, while measurements were terminated after 80 min of recording. Bpm, beats per minute. (b) Blood pressure responses before and during ice immersion. Systolic, diastolic and mean arterial blood pressure are presented. A vertical dotted line delineates the start ( $t = 0$  min) of ice immersion, while measurements were terminated after 80 min of recording. The transient (5 min) drop in blood pressure after 60 min is likely due to technical problems related to the start of a new recording

## 4 | DISCUSSION

In this study, we report that our participant maintained a stable  $T_{gi}$ , HR, and blood pressure during the first part of crushed ice immersion (i.e., 30-min). In the second part of crushed ice immersion, our participant was still able to maintain a  $T_{gi}$  above  $37^{\circ}\text{C}$ , whereas HR and blood pressure were only modestly increased. There was, however, a drop in  $T_{skin}$  immediately upon full-body ice immersion. Despite 88 min of full-body contact with ice, clinically relevant complications or signs of (thermal) discomfort were not observed. Our study describes the physiological responses which counteract an extreme cold exposure in humans.

$T_{gi}$  dropped from  $37.7$  to  $37.0^{\circ}\text{C}$  after 88 min of crushed ice immersion. The decrease of  $0.7^{\circ}\text{C}$  is modest compared to previous studies that reported the onset of hypothermia after  $\sim 30$  min of cold water immersion (Giesbrecht, 2000; Goheen et al., 1997; Holmer, 1995; Tikuisis, 1997). Furthermore, swimming exercise on a self-selected pace for 80 min in water at  $10^{\circ}\text{C}$  results in decrease in rectal temperature of

$2.6^{\circ}\text{C}/\text{h}$ , in which five out of 10 swimmers did not complete the 90 min (Tipton, Eglin, Gennser, & Golden, 1999). Moreover metabolic heat production in the swimmers was more than twofold higher than the heat production in our participant. Additionally, a recent study demonstrated a drop in gastrointestinal temperature of  $1.4^{\circ}\text{C}/\text{h}$  in triathletes swimming in  $10^{\circ}\text{C}$  water while wearing a wetsuit (Melau et al., 2019), which was in line with the study by Tipton and colleagues demonstrating that rectal temperature decreased with  $1.6^{\circ}\text{C}/\text{h}$  and  $1.1^{\circ}\text{C}/\text{h}$  while swimming without wetsuit in  $16^{\circ}\text{C}$  and  $18^{\circ}\text{C}$  water, respectively (Tipton et al., 1999). A study by Wakabayashi, Hanai, Yokoyama, & Nomura (2006) demonstrated that the  $T_{gi}$  of healthy Japanese male participants ( $\text{BSA} = 1.79 \text{ m}^2$ ) decreased with  $1.1 \pm 0.1^{\circ}\text{C}$  after 60 min of water immersion at  $26^{\circ}\text{C}$  in normal swim suit. Although clear differences exist between ice immersion and cold-water immersion (e.g., conductive capacity of water vs. ice), the ability to sustain the cold tolerance and maintain a relatively stable  $T_{gi}$  is remarkable. One explanation for this finding may be the  $\sim 2$ -fold increase in metabolic heat production in the absence of shivering or any other voluntary bodily movement. The level of oxygen consumption achieved during ice immersion is, in daily living and under normal conditions, comparable to light intensity exercise such as walking (Cramer & Jay, 2016). The increased heat production and oxygen consumption during ice immersion might partly be due to the activity of brown adipose tissue, which has been suggested to be responsible for non-shivering thermogenesis (Marken Lichtenbelt et al., 2009; Nedergaard & Cannon, 2010). However, in a previous study using the same participant and his monozygotic twin brother (who was not cold acclimatized), cold air exposure ( $13^{\circ}\text{C}$ ) did not elicit differences in brown fat activity (Vosselman, Vijgen, Kingma, Brans, & Marken Lichtenbelt, 2014). Therefore, it is not likely that the increased heat production ( $\sim 150 \text{ W}$ ) during crushed ice immersion can be explained by brown fat activity in our participant.

Before and during ice immersion, our participant performed Asian G-Tummo meditation to enhance metabolism and maintain  $T_{gi}$ . Previous studies indicated that this meditation technique is associated with an increased metabolism and core temperature (Benson et al., 1982; Kozhevnikov et al., 2013). As our participant performed Asian G-Tummo meditation prior to ice immersion as well, his metabolic heat production in rest was higher, which could explain the relatively high baseline  $T_{gi}$ . However,  $T_{gi}$  was still within reference values for resting core body temperature (Wunderlich, 1868). We were unable to perform a control experiment in which he was exposed to the same stimulus without meditation. Therefore, it remains speculative whether such meditation represents the principle explanation for our results.

Repeated exposure to extreme cold, that is, cold training, may have activated and/or enhanced metabolic-insulative

body responses. One such mechanisms relate to an enhanced vasoconstriction capacity (Makinen, 2010). Peripheral vasoconstriction is a powerful mechanism to reduce heat loss, but may also cause cold injury due to prolonged low skin temperatures. Previous studies demonstrated that blood vessels typically dilate after 5 to 10 min of cold exposure, resulting in an increase in  $T_{skin}$  (Cheung & Daanen, 2012). This cold induced vasodilation is followed by a new phase of vasoconstriction, which is known as the “hunting reaction” of Lewis (1941), and is observed in the hand, feet and face (Brajkovic & Ducharme, 2006; Daanen, 2003; Greenfield, Kernohan, Marshall, Shepherd, & Whelan, 1951). Previous studies indicated that these local responses are more pronounced in individuals regularly exposed to cold (e.g., individuals from polar tribes, fisherman, fish filleters, etc.) (Daanen, 2003) and with cold acclimation (Livingstone, Nolan, & Keefe, 1993; Purkayastha, Ilavazhagan, Ray, & Selvamurthy, 1993). Whether the cyclic process of alternating cutaneous vasoconstriction and vasodilation “trains” the human body to prevent tissue damage is currently unknown. Interestingly, after cessation of the experiment, the skin displayed a generalized characteristic pattern of the “hunting reaction” of Lewis. Additionally, we were not able to draw a venous blood sample at the final stage of ice immersion, suggesting severe vasoconstriction. Therefore, our results suggest that our participant may have optimized his physiological response to cold stress to effectively counteract the cold stimulus through profound (peripheral) vasoconstriction and a higher metabolism that could have contributed to the cold tolerance.

Our study has some limitations. First, we describe the physiological responses of 88 min of ice immersion in a single participant, making it impossible to extrapolate our findings to the general population. Nevertheless, case reports with remarkable findings can yield valuable information and are important in hypothesis generation for further research. This study is further limited by the absence of a control condition without G-Tummo meditation. A randomised controlled intervention study should be performed to examine whether G-Tummo medication is able to alter these responses in participants who are a priori not used to ice immersion. Another limitation of our study, is that we visually checked whether our participant demonstrated shivering thermogenesis, whereas it would have been better to quantify shivering thermogenesis by performing EMG measurements in order to examine the presence of low-level shivering or an increased muscle tone. Furthermore, we did not measure the percentage of body fat of our participant, which could have influenced our results, as a higher body fatness is associated with a higher insulative capacity and thereby a lower heat loss in cold environments. Additionally, it has been demonstrated that heat production increases during cold exposure, due to the development of electrical activity in the

“resting muscles,” defined as thermoregulatory muscle tone, that occurs before the onset of shivering (Burton, 1937). Unfortunately, we did not measure muscle tone using EMG signals in our experiment, and we cannot establish whether a higher muscle tone can explain the increased heat production in our participant. Finally, our results were not corrected for peripheral vasoconstriction. However, as the hands were not in direct contact with ice and the fact that we did not experience problems with taking these measurements, it is likely that peripheral vasoconstriction did not importantly affect our outcomes and main conclusions.

In conclusion, our participant tolerated 88 min of full body crushed ice immersion, during which  $T_{gi}$  did not drop more than  $0.7^{\circ}\text{C}$ , while no shivering was observed. Potentially, he optimized his physiological responses to effectively counteract the cold stimulus through profound (peripheral) vasoconstriction and marked elevation of metabolism.

## ACKNOWLEDGMENTS

The authors thank the participant for cooperation and willingness to share the data collected in this case report.

## CONFLICT OF INTEREST

The authors report no conflict of interest.

## ORCID

Coen C. W. G. Bongers  <https://orcid.org/0000-0003-0055-5308>

Thijs M. H. Eijsvogels  <https://orcid.org/0000-0003-0747-4471>

Maria T. E. Hopman  <https://orcid.org/0000-0001-9504-5452>

## REFERENCES

- Benson, H., Lehmann, J. W., Malhotra, M. S., Goldman, R. F., Hopkins, J., & Epstein, M. D. (1982). Body temperature changes during the practice of g Tum-mo yoga. *Nature*, *295*, 234–236.
- Brajkovic, D., & Ducharme, M. B. (2006). Facial cold-induced vasodilation and skin temperature during exposure to cold wind. *European Journal of Applied Physiology*, *96*, 711–721.
- Brown, D. J., Brugger, H., Boyd, J., & Paal, P. (2012). Accidental hypothermia. *The New England Journal of Medicine*, *367*, 1930–1938.
- Burton, A. B. (1937). The motor mechanism of shivering and thermal muscular tone. *American Journal of Physiology*, *119*, 284–285.
- Byrne, C., & Lim, C. L. (2007). The ingestible telemetric body core temperature sensor: A review of validity and exercise applications. *British Journal of Sports Medicine*, *41*, 126–133.
- Cheung, S. S., & Daanen, H. A. (2012). Dynamic adaptation of the peripheral circulation to cold exposure. *Microcirculation*, *19*, 65–77.
- Cramer, M. N., & Jay, O. (2016). Biophysical aspects of human thermoregulation during heat stress. *Autonomic Neuroscience*, *196*, 3–13.
- Daanen, H. A. (2003). Finger cold-induced vasodilation: A review. *European Journal of Applied Physiology*, *89*, 411–426.
- Evans-Wentz, W. Y. (1958). *Tibetan yoga and secret doctrines*. London: Oxford University Press.
- Gant, N., Atkinson, G., & Williams, C. (2006). The validity and reliability of intestinal temperature during intermittent running. *Medicine and Science in Sports and Exercise*, *38*, 1926–1931.

- Giesbrecht, G. G. (2000). Cold stress, near drowning and accidental hypothermia: A review. *Aviation Space and Environmental Medicine*, *71*, 733–752.
- Goheen, M. S., Ducharme, M. B., Kenny, G. P., Johnston, C. E., Frim, J., Bristow, G. K., & Giesbrecht, G. G. (1997). Efficacy of forced-air and inhalation rewarming by using a human model for severe hypothermia. *Journal of Applied Physiology*, *83*, 1635–1640.
- Greenfield, A. D., Kernohan, G. A., Marshall, R. J., Shepherd, J. T., & Whelan, R. F. (1951). Heat loss from toes and fore-feet during immersion in cold water. *Journal of Applied Physiology*, *4*, 37–45.
- Holmer, I. (1995). Prediction of responses to cold. *Arctic Medical Research*, *54*(Suppl 2), 48–54.
- ISO-9886. (2004). *Ergonomics - Evaluation of thermal strain by physiological measurements*. p. 15–16.
- Kozhevnikov, M., Elliott, J., Shephard, J., & Gramann, K. (2013). Neurocognitive and somatic components of temperature increases during g-Tummo meditation: Legend and reality. *PLoS ONE*, *8*, e58244.
- Lewis, T. (1941). Observations on some normal and injurious effects of cold upon the skin and underlying tissues: I. Reactions to cold, and injury of normal skin. *British Medical Journal*, *2*, 795–797.
- Livingstone, S. D. N., Nolan, R. W., & Keefe, A. A. (1993). *Changes in cold tolerance during a 100-day polar ski expedition*. Amsterdam: Elsevier.
- Makinen, T. M. (2010). Different types of cold adaptation in humans. *Frontiers in Bioscience*, *2*, 1047–1067.
- Melau, J., Mathiassen, M., Stensrud, T., Tipton, M., & Hisdal, J. (2019). Core temperature in triathletes during swimming with wetsuit in 10°C cold water. *Sports*, *7*, 130.
- Murgatroyd, P. R., Shetty, P. S., & Prentice, A. M. (1993). Techniques for the measurement of human energy expenditure: A practical guide. *International Journal of Obesity*, *17*, 549–568.
- Nedergaard, J., & Cannon, B. (2010). The changed metabolic world with human brown adipose tissue: Therapeutic visions. *Cell Metabolism*, *11*, 268–272.
- Purkayastha, S. S., Ilavazhagan, G., Ray, U. S., & Selvamurthy, W. (1993). Responses of Arctic and tropical men to a standard cold test and peripheral vascular responses to local cold stress in the Arctic. *Aviation, Space, and Environmental Medicine*, *64*, 1113–1119.
- Sessler, D. I. (2009). Thermoregulatory defense mechanisms. *Critical Care Medicine*, *37*, S203–S210.
- Shetty, M. R. (2003). Cause of death among passengers on the Titanic. *Lancet*, *361*, 438.
- Smith, A. D., Crabtree, D. R., Bilzon, J. L., & Walsh, N. P. (2009). The validity of wireless iButtons and thermistors for human skin temperature measurement. *Physiological Measurement*, *31*, 95–114.
- Tikusis, P. (1997). Prediction of survival time at sea based on observed body cooling rates. *Aviation, Space, and Environmental Medicine*, *68*, 441–448.
- Tipton, M., Eglin, C., Gennser, M., & Golden, F. (1999). Immersion deaths and deterioration in swimming performance in cold water. *Lancet*, *354*, 626–629.
- van Marken Lichtenbelt, W. D., Daanen, H. A., Wouters, L., Fronczek, R., Raymann, R. J., Severens, N. M., & Van Someren, E. J. (2006). Evaluation of wireless determination of skin temperature using iButtons. *Physiology & Behavior*, *88*, 489–497.
- van Marken Lichtenbelt, W. D., Vanhommel, J. W., Smulders, N. M., Drossaerts, J. M., Kemerink, G. J., Bouvy, N. D., ... Teule, G. J. (2009). Cold-activated brown adipose tissue in healthy men. *The New England Journal of Medicine*, *360*, 1500–1508.
- Vosselman, M. J., Vijgen, G. H., Kingma, B. R., Brans, B., & van Marken Lichtenbelt, W. D. (2014). Frequent extreme cold exposure and brown fat and cold-induced thermogenesis: A study in a monozygotic twin. *PLoS ONE*, *9*, e101653.
- Wakabayashi, H., Hanai, A., Yokoyama, S., & Nomura, T. (2006). Thermal insulation and body temperature wearing a thermal swimsuit during water immersion. *Journal of Physiological Anthropology*, *25*, 331–338. <https://doi.org/10.2114/jpa2.25.331>
- Wilkinson, D. M., Carter, J. M., Richmond, V. L., Blacker, S. D., & Rayson, M. P. (2008). The effect of cool water ingestion on gastrointestinal pill temperature. *Medicine and Science in Sports and Exercise*, *40*, 523–528. <https://doi.org/10.1249/MSS.0b013e31815cc43e>
- Wissler, E. H. (2003). Probability of survival during accidental immersion in cold water. *Aviation, Space, and Environmental Medicine*, *74*, 47–55.
- Wunderlich, C. A. (1868). *Temperature in diseases - a manual of medical thermometry*. Leipzig: The New Sydenham society.

**How to cite this article:** Bongers CCWG, Eijsvogels TMH, Thijssen DHJ, Hopman MTE. Thermoregulatory, metabolic, and cardiovascular responses during 88 min of full-body ice immersion – A case study. *Physiol Rep*. 2019;7:e14304. <https://doi.org/10.14814/phy2.14304>