



Article

PM_{2.5}, Population Exposure and Economic Effects in Urban Agglomerations of China Using Ground-Based Monitoring Data

Yonglin Shen ¹ and Ling Yao ^{2,3,*}

¹ College of Information Engineering, China University of Geosciences, Wuhan 430074, China; shenyl@cug.edu.cn

² State Key Laboratory of Resources and Environmental Information System, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

³ Jiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing Normal University, Nanjing 210023, China

* Correspondence: yaoling@reis.ac.cn

Received: 2 May 2017; Accepted: 29 June 2017; Published: 3 July 2017

Abstract: This paper adopts the PM_{2.5} concentration data obtained from 1497 station-based monitoring sites, population and gross domestic product (GDP) census data, revealing population exposure and economic effects of PM_{2.5} in four typical urban agglomerations of China, i.e., Beijing-Tianjin-Hebei (BTH), the Yangtze River delta (YRD), the Pearl River delta (PRD), and Chengdu-Chongqing (CC). The Cokriging interpolation method was used to estimate the PM_{2.5} concentration from station-level to grid-level. Next, an evaluation was conducted mainly at the grid-level with a cell size of 1 × 1 km, assisted by the urban agglomeration scale. Criteria including the population-weighted mean, the cumulative percent distribution and the correlation coefficient were applied in our evaluation. The results showed that the spatial pattern of population exposure in BTH was consistent with that of PM_{2.5} concentration, as well as changes in elevation. The topography was also an important factor in the accumulation of PM_{2.5} in CC. Moreover, the most polluted urban agglomeration based on the population-weighted mean was BTH, while the least was PRD. In terms of the cumulative percent distribution, only 0.51% of the population who lived in the four urban agglomerations, and 2.33% of the GDP that was produced in the four urban agglomerations, were associated with an annual PM_{2.5} concentration smaller than the Chinese National Ambient Air Quality Standard of 35 µg/m³. This indicates that the majority of people live in the high air polluted areas, and economic development contributes to air pollution. Our results are supported by the high correlation between population exposure and the corresponding GDP in each urban agglomeration.

Keywords: fine particulate matter; population exposure; population-weighted mean; urban agglomeration

1. Introduction

Fine particulate matter (also known as PM_{2.5} or respirable particles), which is suspended in the atmosphere with an aerodynamic diameter no more than 2.5 microns, has been recognized as a serious environmental concern due to its significant adverse effects on human health, climate biogeochemical cycle, and atmospheric chemistry [1,2]. Negative health effects are important consequences of population exposure to PM_{2.5}, which makes exposure information essential for policy-makers to develop appropriate risk control policies [3].

In previous studies worldwide, the characteristics, chemical compositions, sources, and formation mechanism of PM_{2.5} have been researched [4,5], and conclusions were made that pollution arose

at several levels of scale, e.g., a sampling site scale for traffic non-exhaust and re-suspended dust sources, an urban scale for combustion and industrial sources, and a regional scale [6]. Moreover, terrain conditions are one of the most important factors that affect PM_{2.5} concentration distribution [7]. Yao and Lu [8] found that the spatial distribution of annual average PM_{2.5} concentration coincides with China's three gradient terrains. In China, many mega-cities have been investigated, e.g., Beijing [9,10], Shanghai [11], Guangzhou [12], Nanjing [13,14], Chengdu [15], Shenyang [16], and Changchun [17]. Furthermore, studies conducted in recent years have also focused on a regional scale, such as urban agglomeration. Urban agglomeration is an extended city or town area comprising the built-up area of one mega-city or more, with surrounding suburbs linked by continuous urban areas. Shimou [18] reviewed the differences among urban agglomerations in America, Europe, and China, and concluded that strict administration systems and regulations made the development much more difficult if a city tried to expand beyond their political borders in China. Zhao et al. [19] analyzed the chemical compositions, seasonal variations, and regional pollution events of PM_{2.5} in the Beijing-Tianjin-Hebei (hereinafter, BTH, also known as Jing-Jin-Ji), China. Moreover, other urban agglomerations of China were also examined, including the Yangtze River delta (YRD) [20], Pearl River delta (PRD) [21,22], and Chang-Zhu-Tan [23]. A comparison between cities and urban agglomerations has been conducted; Chan and Yao [24] presented an overview of air pollution in China and provided a detailed comparison between Beijing, Shanghai and the PRD.

To better quantify the adverse health impacts of PM_{2.5}, population exposure estimates that can better incorporate the spatial variability of PM_{2.5} concentrations are required [1]. Aleksandropoulou and Lazaridis [25] evaluated the population exposure to PM_{2.5} in outdoor environments in the metropolitan areas of Greece from 2001 to 2010. Pant et al. [26] estimated the population exposure to PM_{2.5}, and assessed the representability of ambient air quality monitoring stations to serve as surrogates for population exposure in New Delhi, India. Requia et al. [3] evaluated the spatial patterns of link-based PM_{2.5} emissions and subsequent human exposure in a large Canadian metropolitan area. Furthermore, evaluations have also been conducted in mega-cities of China, like Beijing [27,28]. Population of exposure to pollutant can be estimated either with satellite images or with ground-based monitoring data, where the former use satellite-derived aerosol optical depth (AOD) as a proxy for PM_{2.5} [1]. The variations of population exposure to PM_{2.5} across the PRD [29] and mainland China [30] have been stated by using the 0.1° × 0.1° moderate resolution imaging spectroradiometer (MODIS) satellite data. However, the accuracy of AOD-based PM_{2.5} retrieval methods may vary significantly due to seasons and locations [31,32].

In this study, we focused on the assessment of population exposure and economic effects on PM_{2.5} in typical urban agglomerations of China, i.e., BTH, YRD, PRD and Chengdu-Chongqing (hereinafter, CC, also known as Cheng-Yu). Station-based PM_{2.5} datasets across the four urban agglomerations were used, and criteria including population-weighted mean, cumulative percent distribution, and correlation coefficient were employed in our evaluation. This provided a unique opportunity for us to study the impact of grid-level PM_{2.5} data in exposure estimates for regional populations, as well as to further validate the economic effects on PM_{2.5} concentrations.

2. Materials and Methods

2.1. Study Area

The spatial domain of this study included the four most typical urban agglomerations of China, BTH, YRD, PRD and CC. In 2014, the population of the study areas accounted for 26% of the total in China, and the gross domestic product (GDP) of these regions accounted for 43.8% of the national GDP. The locations of the study areas are presented in Figure 1.

The BTH is the biggest urbanized region in northern China, which was born out of political pressure rather than economic prosperity. Ten cities including Beijing, Tianjin, Baoding, Langfang,

Tangshan, Qinhuangdao, Shijiazhuang, Zhangjiakou, Chengde and Cangzhou in the BTH urban agglomeration are officially recognized in the government publications.




Figure 1. The location of the study areas and stations for PM_{2.5} monitoring. The color scale represents the elevation variability across the urban agglomeration.

The YRD region is the most economically developed and rapidly urbanizing city cluster in East China. It is located on the east coast of China bordering the East China Sea. The newest YRD urban agglomeration consists of Shanghai, Nanjing, Wuxi, Changzhou, Suzhou, Nantong, Yancheng, Yangzhou, Zhenjiang, Taizhou, Hangzhou, Ningbo, Jiaxing, Huzhou, Shaoxing, Jinhua, Zhoushan, Taizhou, Hefei, Wuhu, Ma'an Shan, Tongling, Anqing, Chuzhou, Chizhou, and Xuancheng.

The PRD, which is currently known as the Guangdong-Hong Kong-Macau Greater Bay area, is situated in South China adjacent to the South China Sea, which is considered as one of the country's chief economic regions and manufacturing centers. The PRD urban agglomeration generally comprises Guangzhou, Shenzhen, Zhuhai, Foshan, Dongguan, Zhongshan, Jiangmen, Zhaoqing, Huizhou, Hong Kong and Macau. As there is no ground-based monitoring data in Hong Kong and Macau, they were not considered in this study.

The CC is a national level urban agglomeration that leads the development of West China. It consists of Chongqing, Chengdu, Zigong, Luzhou, Deyang, Mianyang, Suining, Neijiang, Leshan, Nanchong, Meishan, Yibin, Guang'an, Dazhou, Ya'an and Ziyang, among which Chongqing and Chengdu are the core cities.

2.2. Datasets

The primary data used in this study were station-based PM_{2.5} concentrations, population census data, GDP census data and a 30 × 30 m digital elevation model (DEM). Annual PM_{2.5} concentration data were obtained from the China National Environmental Monitoring Center (CNEMC) website [33]. By the end of 2014, approximately 1497 monitoring sites had been established to report the overall air

quality in China. The population and GDP census data were derived from the National Earth System Science Data Sharing Infrastructure [34] with 1×1 km sized grids. The census data showed a high density of people living in and around the major urban areas.

The original $PM_{2.5}$ concentration data were at discrete points, i.e., at fixed monitoring stations. To present a localized representation of population exposure to $PM_{2.5}$ and economic effects on it, we used a geographic information system based grid mapping where grids of 1×1 km were applied to $PM_{2.5}$ data. Thus, the spatial interpolation method was applied to estimate the concentration of pollutants from station-level to grid-level to characterize concentrations across the entire area. In this study, we used Cokriging method incorporated with DEM data to calculate the pollutant concentration at each grid from the measured data at 1497 monitoring stations. Cokriging is a multivariate extension of kriging where the auxiliary information (in this case, DEM data) is incorporated in the estimation at unsampled locations by accounting for spatial correlations of the primary variable with secondary attributes [35]. The mapping processes divided regions of the BTH, YRD, PRD and CC into 146,598, 166,151, 36,103 and 208,879 available grid cells, respectively. Next, the processes of population exposure and the economic effects of urban agglomerations on those grid cells were analyzed by mathematical methods.

2.3. Estimation of Population Exposure

The population exposure was estimated both at the grid-level and region-level (i.e., at the urban agglomeration level). Population exposure at the grid-level was proposed as an important indicator to measure the population exposure to $PM_{2.5}$ air pollution, which requires both population distribution data and $PM_{2.5}$ concentration distribution data. It can be calculated by,

$$\psi_i = P_i C_i \quad (1)$$

where ψ_i represents the i th grid of population exposure; i designates each computational cell in the domain; P_i is the population at a given cell location; and C_i is the particulate concentration in the same cell location.

At the urban agglomeration level, the population exposure was measured by the population-weighted mean $PM_{2.5}$ [36]. Calculating the population weighted mean $PM_{2.5}$ has implications for estimating the impact of air pollution on public health and providing insights into pollution mitigation policies for individual urban agglomeration and cross-regional collaboration. If the distribution of $PM_{2.5}$ within an urban agglomeration concurs with the spatial patterns of population, the resulting population-weighted mean $PM_{2.5}$ tends to be larger than its original $PM_{2.5}$ value, and vice versa [37]. The population weighted mean $PM_{2.5}$ (denoted by ϵ) is calculated as

$$\epsilon = \frac{\sum_{i=1}^n \psi_i}{P_o} \quad (2)$$

where $P_o = \sum_{i=1}^n P_i$ is the total population in the domain of interest (in this case, an urban agglomeration); and n represents the total number of grids in the urban agglomeration.

2.4. Estimation of Cumulative Percentage Distribution

Meanwhile, this paper also calculated the cumulative $PM_{2.5}$ concentrations in each study area, as cumulative percentage is a way of expressing frequency distribution. It calculates the percentage of the cumulative frequency within each interval, as much as relative frequency distribution calculates the percentage of frequency [27]. The cumulative percent distribution of population or GDP by annual $PM_{2.5}$ concentrations is calculated as

$$\gamma_\tau = \frac{\int_0^\tau P_{C<\tau}}{P_o} \times 100\% \quad (3)$$

where γ_τ represents the cumulative percent distribution of population or GDP by the PM_{2.5} concentration τ ; τ designates a certain concentration of PM_{2.5}; $\int_0^\tau P_{C<\tau}$ is the cumulative population or GDP under the condition that the PM_{2.5} concentration is less than τ , i.e., $C < \tau$.

2.5. Correlations among PM_{2.5}, Population, GDP and Population Exposure

To measure the effects of population growth and economic development on PM_{2.5} concentrations, correlation coefficients were used to quantify the correlation and dependence among PM_{2.5}, population, GDP, and population exposure. The correlation coefficient (r) measures the linear correlation between two variables x and y , and has a value between +1 and −1, where 1 is total positive linear correlation, 0 is no linear correlation, and −1 is total negative linear correlation. For each urban agglomeration, r was counted at both grid-level and city-level.

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2}} \quad (4)$$

where n represents the number to variable; x_i and y_i are the i th of variables x and y , respectively; and \bar{x} and \bar{y} are the mean of variables x and y , respectively. If r is counted at the grid-level, then x and y represent the values of each grid of 1×1 km, and n represents the number of cells; and if r is counted at the city-level, then x and y represent the values of each city, and n represents the number of cities.

3. Results and Discussion

This study was performed in three parts: (1) analysis of the spatial patterns of PM_{2.5} concentrations in four major urban agglomerations of China; (2) evaluation of population exposure to PM_{2.5}; and (3) evaluation of economic impacts on PM_{2.5}.

3.1. Spatial Patterns of PM_{2.5} Concentration

Urban PM_{2.5} originates mainly from sources such as coal combustion, traffic-related emissions, fugitive dust, biomass burning, agricultural activities and regional transported aerosols [38,39]. The World Health Organization (WHO) put forward an annual limit of 10 $\mu\text{g}/\text{m}^3$ to mitigate its impact on human health [40]. The Chinese government adopted the interim Target 1 (IT1) of the air quality guidelines (AQG) established by the WHO. In 2012, the new Chinese National Ambient Air Quality Standard (NAAQS) was amended and issued by the Ministry of Environmental Protection of China (MEP). In the NAAQS, the annual PM_{2.5} concentration Grade II limit is 35 $\mu\text{g}/\text{m}^3$ [41]. As shown in Table 1, the overall minimum and maximum of annual PM_{2.5} concentration in four urban agglomerations were 32.91 $\mu\text{g}/\text{m}^3$ and 120.11 $\mu\text{g}/\text{m}^3$, respectively, and were 3–12 times the standard set by the WHO of 10 $\mu\text{g}/\text{m}^3$, and 1–3 times that of the NAAQS of 35 $\mu\text{g}/\text{m}^3$. Additionally, the overall mean and standard deviation of PM_{2.5} concentration were 63.4 $\mu\text{g}/\text{m}^3$ and 14.03 $\mu\text{g}/\text{m}^3$, respectively. Figure 2 shows the spatial patterns of PM_{2.5} in the urban agglomerations. Since the four urban agglomerations had different maximum and the minimum concentrations (Table 1), the annual PM_{2.5} concentration of each urban agglomeration in Figure 2 was plotted in a subfigure with a different color scale to highlight the spatial variability across the corresponding urban agglomeration.

3.1.1. Beijing-Tianjin-Hebei Urban Agglomeration

The PM_{2.5} concentration of the BTH urban agglomeration ranged from 37.21 $\mu\text{g}/\text{m}^3$ to 120.11 $\mu\text{g}/\text{m}^3$. The mean and standard deviation of PM_{2.5} concentration in BTH were 74.97 $\mu\text{g}/\text{m}^3$ and 19.46 $\mu\text{g}/\text{m}^3$, respectively, which were the highest in the four urban agglomerations (Table 1). As shown in Figure 2a, the higher PM_{2.5} concentrations were distributed in the southern areas of the BTH, and they decreased gradually from south to north. In particular, north Beijing and Tangshan were superior to their respective area in the south. The trends were consistent with the elevation changes

shown in Figure 1. The air quality was relatively good in Zhangjiakou, Chengde, Qinhuangdao, and the corresponding PM_{2.5} concentration in these cities were 47.52 µg/m³, 59.45 µg/m³ and 66.06 µg/m³, respectively. The worst was Baoding with a concentration of 116.21 µg/m³, followed by Shijiazhuang of 108.51 µg/m³.

Table 1. Statistical information of the urban agglomerations.

Urban Agglomeration	C_{min} (µg/m ³)	C_{max} (µg/m ³)	C_{mean} (µg/m ³)	C_{std} (µg/m ³)	C_{pwm} (µg/m ³)
BTH	37.21	120.11	74.97	19.46	86.11
YRD	38.51	75.43	59.24	7.68	58.32
PRD	32.91	52.96	41.34	4.54	41.12
CC	51.15	73.97	63.32	4.1	63.63
Overall	32.91	120.11	63.74	14.02	64.62

Notes: C_{min} , C_{max} , C_{mean} and C_{std} represent the minimum, maximum, mean and standard deviation of PM_{2.5} concentration in the domain of interest; and C_{pwm} is the population-weighted mean of PM_{2.5} concentration.




Figure 2. The spatial patterns of PM_{2.5} concentrations in the four urban agglomerations. (a) Beijing-Tianjin-Hebei; (b) Yangtze River delta; (c) Pearl River delta; and (d) Chengdu-Chongqing. Each subfigure has a different color scale to highlight the spatial variability across the corresponding urban agglomeration.

Several previous studies have revealed the sources of PM_{2.5} pollutants in the BTH [19]. Zhao et al. [19] pointed out that BTH suffers from serious secondary pollution, and motor vehicle exhaust highly contribute to the PM_{2.5} concentration. From the perspective of transportation process, pollution sources can be placed into three categories: local emissions, local transportation and regional transportation. Furthermore, local emissions, local transportation, and regional transportation have all significantly contributed to high fine particle loadings in these areas [38].

The local emission sources for air pollution include power plants, domestic heating, and industrial, vehicular, and biogenic sources [24]. In Beijing and Tianjing, motor vehicle exhausted highly contributes to the annual PM_{2.5} concentrations. In addition to transportation, coal combustion was a vital factor affecting PM_{2.5} concentration, especially in Shijiazhuang and Chengde, where the PM_{2.5} pollution was dominated by coal combustion [19]. During winter, domestic heating emissions, mainly from coal combustion, contribute highly to the air quality degradation observed in these cities [42]. For example, domestic heating in Beijing usually starts in mid-November and ends in the following March, which is the major source of SO₂ in winter [24].

Pollution source of local transportation is defined as the one that comes from other cities of the same urban agglomeration. Recent studies have shown that changes in atmospheric circulation (e.g., the weakened northerly winds and the development of inversion anomalies in the lower troposphere) may be an important reason for the increased haze pollution in northern China [43]. The prevailing wind in Beijing is from the north and the northwest, particularly in winter and spring, respectively, and the concentrations of air pollutants generally decrease with increasing wind speed [24].

Pollution source of regional transportation is the one that comes from areas beyond the urban agglomeration. Several studies have shown that the enhanced PM_{2.5} pollutants in BTH is not only due to the primary emissions from local sources (such as industrial, domestic and agricultural sources), but also the regional transportation contribution (e.g., from nearby Shandong and Henan Provinces) and secondary production. Furthermore, the climate of the BTH is characterized by stagnant weather with weak wind and relatively low boundary layer height, which is a favorable atmospheric condition for the accumulation, formation and processing of aerosols [39].

3.1.2. Yangtze River Delta Urban Agglomeration

The PM_{2.5} concentration of the YRD urban agglomeration ranged from 38.51 µg/m³ to 75.43 µg/m³. The mean and standard deviation of PM_{2.5} concentration of the YRD were 59.24 µg/m³ and 7.68 µg/m³, respectively (Table 1). The mean concentration of PM_{2.5} in the YRD region was below average level (63.4 µg/m³), and lower than that of the BTH (74.97 µg/m³) and CC (63.32 µg/m³), but higher than that of the PRD region (43.39 µg/m³).

Industrial sources including power plants, other fuel combustion facilities, and non-combustion processes were the major contributors to the PM_{2.5} pollution in the YRD [44]. As shown in Figure 2b, the PM_{2.5} concentrations distributed in the northwest areas of the YRD were high, and decreased gradually from north and northwest to southeast. There are three capital cities (i.e., Hangzhou, Nanjing and Hefei) and a municipality (i.e., Shanghai) in the YRD. Hefei had the worst air quality (76.6 µg/m³), followed by Nanjing (68.23 µg/m³), Hangzhou (53.63 µg/m³), and Shanghai (49.59 µg/m³). Shanghai is located at downstream of the YRD, and the prevailing winds from the east carry clean air into the region [45,46]. The major abundant metal elements of the mass concentrations of PM_{2.5} in Shanghai, Nanjing, Hangzhou and Ningbo were crustal elements (e.g., Al, Ca, Fe and Mg) and trace metals (e.g., Zn, Pb, Cu, Cr, V and Ni) [20].

3.1.3. Pearl River Delta Urban Agglomeration

The PM_{2.5} concentration of the PRD urban agglomeration ranged from 32.91 µg/m³ to 52.96 µg/m³. The mean and standard deviation of PM_{2.5} concentration of the PRD were respectively 41.34 µg/m³ and 4.54 µg/m³ (Table 1). The mean concentration of PM_{2.5} in the PRD was the lowest in

the four urban agglomerations mainly because of fewer coal-based industries and good dispersion weather conditions [39]. Nevertheless, it was still four times the standard set by the WHO of $10 \mu\text{g}/\text{m}^3$, and higher than NAAQS at $6.34 \mu\text{g}/\text{m}^3$.

In the PRD, the level of secondary pollutants was higher than that of dusts and primary pollutants [47]. $\text{PM}_{2.5}$ emissions mainly concentrated in Guangzhou, Foshan, Shenzhen and Dongguan, which possess advanced industry, high energy consumption, and dense population. However, the climate of the PRD is affected by monsoons. The prevailing winds are from the north in winter and the south in summer [1]. As shown in Figure 1, PRD is a low-lying area surrounding the Pearl River estuary, and the elevations of north PRD and parts of Jiangmen are relatively higher than that of other regions. Affected by the topography and climate characteristics, $\text{PM}_{2.5}$ concentrations easily gather in the north of PRD in winter. As shown in Figure 2c, $\text{PM}_{2.5}$ concentrations were lower near the coastline, and vice versa. Much higher $\text{PM}_{2.5}$ concentrations were observed in the north of PRD, with orange and red colors showing annual $\text{PM}_{2.5}$ concentrations higher than that of southern coastal areas.

3.1.4. Chengdu-Chongqing Urban Agglomeration

The $\text{PM}_{2.5}$ concentration of the CC urban agglomeration ranged from $51.15 \mu\text{g}/\text{m}^3$ to $73.97 \mu\text{g}/\text{m}^3$. The mean and standard deviation of $\text{PM}_{2.5}$ concentration were $63.32 \mu\text{g}/\text{m}^3$ and $4.1 \mu\text{g}/\text{m}^3$, respectively (Table 1). Although its mean $\text{PM}_{2.5}$ concentrations out of the four urban agglomerations was not the lowest, the standard deviation was. This suggested that concentration change in the whole urban agglomeration was relatively smooth and steady.

The maximum $\text{PM}_{2.5}$ concentration in the CC was nearly two times the NAAQS of $35 \mu\text{g}/\text{m}^3$, which was partly due to its unique topographic condition [48]. Local $\text{PM}_{2.5}$ emissions accompanied by low wind speed and high relative humidity conditions are major causes of visibility impairment in the Sichuan Basin [49]. As shown in Figure 2d, there are two hotspots in the CC: one is centered at the Sichuan Basin, where the capital city of Sichuan Province (i.e., Chengdu) and the downtown area of Chongqing are located, and another is centered at Dazhou.

Regarding the first hotspot (Figure 1), the elevation of cities there was lower than that of adjacent areas. Chengdu has typical basin climate characteristics, i.e., a precipitation period is basically from July to September with relatively high humidity, and static wind frequency, and there is atmospheric stability subject to neutral weather in winter. These special topographic and climatic conditions are important causes of the accumulation of $\text{PM}_{2.5}$. Static wind and temperature inversion weather occur frequently, especially in winter and autumn, resulting in continuous heavy pollution [50]. Similarly, local contributors are likely to play a predominant role in the downtown of Chongqing. Influenced by its specific topographic conditions, Chongqing is located in the region of the lowest wind speed over China. For example, the annual average wind speed was between 0.9 m/s and 1.6 m/s from 1979 to 2007 [38]. The specific geographic and meteorological conditions favor the accumulation of regional and local pollutants, and stable weather conditions with low wind speed, low mixing heights, and high relative humidity can greatly enhance pollution levels [43].

In the second hotspot where the city of Dazhou is located, a variety of factors contributed to the higher concentration. The air pollution of Dazhou was mainly caused by the discharge of industrial dust and soot emissions from heavy industries (e.g., iron, steel, and cement), vehicular emissions, and urban fugitive dust. In addition, the topography of Dazhou is an objective factor, as it is surrounded by mountains on three sides in a narrow area. These anthropic and natural factors made Dazhou unusually prominent in the air pollution of the CC, thus, the eventual formation of one of the two hotspots.

3.2. Population Exposure to $\text{PM}_{2.5}$

The total population of the four urban agglomerations was estimated at about 355 million, of which about 77.8 million reside in BTH, and 129.9, 31.4, and 115.9 respectively in YRD, PRD, and CC. A comparison of the spatial distributions of population exposure among the four urban agglomerations

in Figure 3 showed that areas in the mega-cities were often associated with higher population exposure to PM_{2.5}. For example, as shown in Figure 3a, a larger proportion of the population was exposed to high PM_{2.5} concentrations in Beijing, Tianjin, and Shijiazhuang, which are capital cities or municipalities in the BTH. Nevertheless, the spatial pattern of population exposure in the BTH was consistent with that of PM_{2.5} concentration, and both of them decreased gradually from south to north. On 1 April 2017, the Xiongan new area (also known as the Hung an District) was set up in Baoding, Hebei Province and was a key component of a massive mega region developing around Beijing, Tianjin, and Hebei. With further economic development of the region, it is expected that the population will rise sharply. As shown in Figure 2a, PM_{2.5} concentration of this area was quite high. As the population increasing, the population exposure to PM_{2.5} should be very serious. Thus, it is suggested that during the process of the district construction, much attention should be paid to environmental pollution control.




Figure 3. The population exposure to PM_{2.5} in the four urban agglomerations. (a) Beijing-Tianjin-Hebei; (b) Yangtze River delta; (c) Pearl River delta; and (d) Chengdu-Chongqing.

In the YRD as shown in Figure 3b, central area had the highest PM_{2.5} concentration, followed by the northern area, and the air quality in the south was the best. In the PRD, population density had a strong effect on population exposure. Therefore, the spatial pattern of the PM_{2.5} concentrations was opposite to that of the corresponding population exposure (Figure 3c). As mentioned above, the PM_{2.5} concentrations of PRD decreased from the north to the southern coastal areas; whereas the population exposure to PM_{2.5} was more serious in the southern coastal areas such as Guangzhou and Shenzhen,

which are two major mega-cities situated in the PRD region. The population exposure distribution also showed that these two cities have the most important position in this urban agglomeration. Furthermore, the PRD region—especially the areas between the two mega-cities—has undergone rapid urbanization and many smaller towns like Dongguan with a population of about one million have been established in the last two decades. The air quality in those cities and towns has been deteriorating over the last decade [24], and the corresponding population exposure is very grim.

As shown in Figure 3d, the outside of the CC had a relatively low population exposure, and the middle part had a level of moderate severity except for the three hotspots, i.e., Chengdu, Chongqing, and Dazhou.

Based on the grid-level annual $PM_{2.5}$ concentration and population census data in each urban agglomeration, the population-weighted mean of $PM_{2.5}$ in the BTH, YRD, PRD, and CC urban agglomerations were $86.11 \mu\text{g}/\text{m}^3$, $58.32 \mu\text{g}/\text{m}^3$, $41.12 \mu\text{g}/\text{m}^3$ and $63.63 \mu\text{g}/\text{m}^3$, respectively (Table 1), which were approximately 2 to 4 times as high as the global population-weighted mean ($20 \mu\text{g}/\text{m}^3$) [51]. The population-weighted mean of $PM_{2.5}$ across all four urban agglomerations was $64.62 \mu\text{g}/\text{m}^3$. We found that the population-weighted mean of $PM_{2.5}$ in the YRD, PRD, and CC were under $64.62 \mu\text{g}/\text{m}^3$, while that of BTH was far more than that.

In terms of the population-weighted mean of $PM_{2.5}$ concentrations, the most polluted urban agglomeration was BTH and the least was PRD. For the BTH, the spatial average of $PM_{2.5}$ concentration in 2014 was $74.97 \mu\text{g}/\text{m}^3$, while the corresponding population-weighted mean of $PM_{2.5}$ was much higher at $86.11 \mu\text{g}/\text{m}^3$, which indicated that more people were living in highly polluted areas. Hence, the spatial average of the $PM_{2.5}$ concentration would underestimate the mean pollution exposure, and the population weighted average would be a better indicator of the public exposure to $PM_{2.5}$ pollution [1]. The only urban agglomeration with the population-weighted mean less than that of the spatial average was CC, which was indicative that more people were living in low pollution areas.

Using the 1×1 km grid-level $PM_{2.5}$ concentration, the cumulative percentage of the population (0–100%) in the four typical urban agglomerations of China are shown in Figure 4a. The results showed that the WHO AQG ($10 \mu\text{g}/\text{m}^3$) for $PM_{2.5}$ was exceeded by 100% of the population in the study area, while China's NAAQS was exceeded by 99.49% of the population. Figure 4a also shows the cumulative percent distribution of population by annual $PM_{2.5}$ concentration in BTH, YRD, PRD and CC urban agglomerations. Overall, only 0.51% of the population lived in the four urban agglomerations with an annual average $PM_{2.5}$ concentration smaller than China's NAAQS Grade II of $35 \mu\text{g}/\text{m}^3$, while the quantity was 5.8% for the PRD, and no one in the BTH, YRD and CC urban agglomerations lived under this guideline. It should be noted that all populations lived in the four urban agglomerations where the WHO AQG was exceeded.




Figure 4. Cumulative percent distribution by annual average $PM_{2.5}$ concentrations which is estimated from station-based monitoring data in 2014 (a) Cumulative percent distribution of population; (b) Cumulative percent distribution of GDP. The dashed line represents the computing on the overall four urban agglomerations.

However, there were some special cases including areas with higher PM_{2.5} concentration, but lower population density (e.g., industrial parks). This can be divided into two scenarios: (1) The industrial park is seriously polluted, but has fewer employees (the levels of population exposure can be directly indicated by Figure 3); and (2) the industrial park is seriously polluted and has many employees, but the vast majority of employees are migrants or live outside the same 1 × 1 km grid. The population exposure of this scenario can be measured by the population weighted mean of PM_{2.5}.

3.3. Economic Effects to PM_{2.5}

Fast-economic development and high energy consumption have led to record heavy haze pollution days in many regions of China [8,16]. The total GDP of the four urban agglomerations had been estimated at approximately 29.4 trillion, of which about 6.1, 13.3, 5.8, and 4.2 trillion were produced by BTH, YRD, PRD and CC, respectively.

Based on the grid-level annual PM_{2.5} concentration and GDP census data in each urban agglomeration, the cumulative percentage distribution of the GDP (0–100%) was estimated (Figure 4b). Overall, only 2.33% of the GDP was produced in the four urban agglomerations with annual average PM_{2.5} concentration lower than the China's NAAQS Grade II of 35 µg/m³. And the quantity was 11.8% for PRD and none of the GDP in the BTH, YRD, and CC urban agglomerations was produced under this guideline. It should be noted that all GDP of the four urban agglomerations exceeded the WHO AQG (10 µg/m³).

We calculated the correlation coefficient between PM_{2.5} concentration and the corresponding population in each urban agglomeration with counting at the grid-level (denoted by R_{pp} , as shown in Table 2). R_{pp} in the BTH, YRD, PRD and CC were 0.26, −0.08, −0.04, and 0.04 respectively, and overall for the four urban agglomerations was 0.03. All studied regions showed that PM_{2.5} concentration and population were independent. The correlation coefficient with counting at the city-level (denoted by R'_{pp}), where the R'_{pp} in the BTH, YRD, PRD, and CC respectively were 0.13, −0.19, 0.08 and −0.02, respectively, and across the overall four urban agglomerations was −0.05. The results showed that population was not a direct factor influencing PM_{2.5} concentrations at local scale, i.e., at a 1 × 1 km grid-level or city-level. The uncorrelated relationship of PM_{2.5} concentrations with population was consistent with previous findings conducted in other places around the world [39,51,52].

Table 2. Correlation coefficient of PM_{2.5} concentrations, population, GDP, population exposure in urban agglomeration.

Urban Agglomeration	Grid-Level			City-Level		
	R_{pp}	R_{pg}	R_{eg}	R'_{pp}	R'_{pg}	R'_{eg}
BTH	0.26	0.11	<u>0.81</u>	0.13	<u>0.69</u>	<u>0.76</u>
YRD	−0.08	−0.07	<u>0.91</u>	−0.19	−0.17	<u>0.79</u>
PRD	−0.04	−0.10	0.38	0.08	<u>0.6</u>	<u>0.73</u>
CC	0.04	0.03	<u>0.88</u>	−0.02	−0.08	<u>0.92</u>
Overall	0.03	−0.06	<u>0.72</u>	−0.05	0.28	<u>0.64</u>

Notes: R_{pp} , R_{pg} , and R_{eg} respectively are the correlation coefficient between PM_{2.5} concentration and population; PM_{2.5} concentration and GDP; and population exposure and GDP, which are counted in the grid-level. While, R'_{pp} , R'_{pg} , and R'_{eg} respectively are the correlation coefficient between PM_{2.5} concentration and population; PM_{2.5} concentration and GDP; and population exposure and GDP, which are counted in the city-level. The underscored values are highlighted to indicate the high dependence.

The correlation coefficient between PM_{2.5} concentration and GDP at the grid-level (denoted by R_{pg}) also showed that there was no connection between the two variables. The weak correlation was partly due to transporting the pollution, and the impact of economic development on PM_{2.5} concentrations was revealed in larger spatial scales, i.e., bigger than a 1 × 1 km cell scale. At the city-level (denoted by R'_{pg}), a good correlation coefficient of 0.69 and 0.6 were found in the BTH and PRD, respectively, and this indicated that the GDP of the internal city was a vital factor for the increase

of PM_{2.5} concentrations in these two urban agglomerations. For example, previous studies had shown that the growth in GDP had increased the demand for vehicular transport, and exacerbated Beijing's air pollution problems [24].

We also calculated the correlation coefficient between population exposure and the corresponding GDP in each urban agglomeration. Most values at both grid-level and city-level showed good correlation coefficients, except for the PRD at grid-level, which was only 0.38. The correlation coefficient of population and corresponding GDP at the grid-level also showed a weak dependence, and the value was 0.41. The spatial trends of population exposure and population were related closely as economic growth attracted a large migrant population into the PRD, and provided cheap labor for economic development. There were more than 23 million migrants living in the PRD in 2009, accounting for more than 40% of the total resident population in the region [29]. Most migrants were concentrated in Shenzhen, Guangzhou, Dongguan and Foshan; however, the population census data only related to the resident population, which decreased the correlation coefficient between population exposure and the corresponding GDP in the PRD.

4. Conclusions

In summary, the population exposure and economic effects on PM_{2.5} in BTH, YRD, PRD and CC urban agglomerations were evaluated. The station-based PM_{2.5} concentrations, population and GDP census data were used, as well as the population exposure estimated for each grid sized 1 × 1 km and for each urban agglomeration. The latter was measured by population-weighted mean PM_{2.5} and cumulative percent distribution. The economic effects on PM_{2.5} were also measured by cumulative percent distribution, as well as correlation coefficient.

PM_{2.5} concentrations were generally higher in the urban agglomeration of North China (e.g., the BTH) than those observed in the south (e.g., the YRD, PRD, and CC). Domestic heating and unfavorable meteorological conditions for pollution dispersion significantly contributed to high fine particle loadings in the BTH. PM_{2.5} concentrations in the BTH increased gradually from north to south, which were consistent with trends in elevation changes. PM_{2.5} concentrations in YRD and PRD also tended to be lower in the coastal regions than in the inland. There were two concentration hotspots in the CC, one centered at the Sichuan Basin and another at Dazhou. Furthermore, areas in mega-cities were often associated with higher population exposure to PM_{2.5}, and the spatial pattern of population exposure in the BTH was consistent with the spatial pattern of PM_{2.5} concentrations. The highest PM_{2.5} concentration was in central YRD. The spatial distribution of PM_{2.5} concentrations was the opposite to that of the corresponding population exposure in the PRD. The population exposure outside of the CC was lower than that of the middle part. In terms of population-weighted mean of PM_{2.5} concentrations, the most polluted urban agglomeration was BTH, and the least was PRD. In terms of the cumulative percent distribution of population, only 0.51% of the population in the four urban agglomerations lived with an annual average PM_{2.5} concentrations smaller than the NAAQS standard. Only 2.33% of the GDP was produced in the four urban agglomerations with annual average PM_{2.5} concentration smaller than NAAQS. Generally, fast-economic development has led to serious population exposure to PM_{2.5} in most urban agglomerations.

In the future, multi-year series of PM_{2.5} concentrations will be used, and further studies will focus on inter-annual variations, seasonal changes, as well as the long-range spatial transport effects related to population exposure of PM_{2.5} concentrations, and the economic effects upon this.

Acknowledgments: This work is partially supported by the National Natural Science Foundation of China under Grant No. 41501459, by the National Marine Hazard Mitigation Service of China under Grant No. 2016AA053, and by the Young Talents Training Fund of State Key Laboratory of Resources and Environment Information System of China under Grant No. O8R8B6E0YA. The authors would like to thank Prof. Xiaohui Yuan from the Department of Computer Science and Engineering at University of North Texas, who provided general advice on the review and helped with editorial contributions. Acknowledgement for the data support from "National Earth System Science Data Sharing Infrastructure, National Science & Technology Infrastructure of

China (<http://geo.ckcest.cn>)” and “National Earth System Science Data Sharing Infrastructure, National Science & Technology Infrastructure of China (<http://www.geodata.cn>)”.

Author Contributions: Ling Yao principally conceived the idea for the study and was responsible for the design of the study, and Yonglin Shen was responsible for setting up experiments, completing the experiments and retrieving data and also wrote the initial draft of the manuscript. Both authors participated in some form in the concept, experimentation, writing and/or editing of this manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lin, C.; Li, Y.; Lau, A.K.H.; Deng, X.; Tse, T.K.T.; Fung, J.C.H.; Li, C.; Li, Z.; Lu, X.; Zhang, X.; et al. Estimation of long-term population exposure to PM_{2.5} for dense urban areas using 1-km modis data. *Remote Sens. Environ.* **2016**, *179*, 13–22. [[CrossRef](#)]
2. Yeh, H.L.; Hsu, S.W.; Chang, Y.C.; Chan, T.C.; Tsou, H.C.; Chang, Y.C.; Chiang, P.H. Spatial analysis of ambient PM_{2.5} exposure and bladder cancer mortality in Taiwan. *Int. J. Environ. Res. Public Health* **2017**, *14*, 508. [[CrossRef](#)] [[PubMed](#)]
3. Requia, W.J.; Dalumpines, R.; Adams, M.D.; Arain, A.; Ferguson, M.R.; Koutrakis, P. Modeling spatial patterns of link-based PM_{2.5} emissions and subsequent human exposure in a large Canadian metropolitan area. *Atmos. Environ.* **2017**, *158*, 172–180. [[CrossRef](#)]
4. Ledoux, F.; Kfoury, A.; Delmaire, G.; Roussel, G.; El Zein, A.; Courcot, D. Contributions of local and regional anthropogenic sources of metals in PM_{2.5} at an urban site in northern France. *Chemosphere* **2017**, *181*, 713–724. [[CrossRef](#)] [[PubMed](#)]
5. Murillo, J.H.; Rojas Marín, J.F.; Mugica Álvarez, V.; Solórzano Arias, D.; Guerrero, V.H.B. Chemical characterization of filterable PM_{2.5} emissions generated from regulated stationary sources in the metropolitan area of Costa Rica. *Atmos. Pollut. Res.* **2017**, *8*, 709–717. [[CrossRef](#)]
6. Squizzato, S.; Cazzaro, M.; Innocente, E.; Visin, F.; Hopke, P.K.; Rampazzo, G. Urban air quality in a mid-size city—PM_{2.5} composition, sources and identification of impact areas: From local to long range contributions. *Atmos. Res.* **2017**, *186*, 51–62. [[CrossRef](#)]
7. Jazcilevich, A.D.; García, A.R.; Caetano, E. Locally induced surface air confluence by complex terrain and its effects on air pollution in the valley of Mexico. *Atmos. Environ.* **2005**, *39*, 5481–5489. [[CrossRef](#)]
8. Yao, L.; Lu, N. Spatiotemporal distribution and short-term trends of particulate matter concentration over China, 2006–2010. *Environ. Sci. Pollut. Res.* **2014**, *21*, 9665–9675. [[CrossRef](#)] [[PubMed](#)]
9. Chen, Y.; Schleicher, N.; Chen, Y.; Chai, F.; Norra, S. The influence of governmental mitigation measures on contamination characteristics of PM_{2.5} in Beijing. *Sci. Total Environ.* **2014**, *490*, 647–658. [[CrossRef](#)] [[PubMed](#)]
10. Lang, J.; Zhang, Y.; Zhou, Y.; Cheng, S.; Chen, D.; Guo, X.; Chen, S.; Li, X.; Xing, X.; Wang, H. Trends of PM_{2.5} and chemical composition in Beijing, 2000–2015. *Aerosol Air Qual. Res.* **2017**, *17*, 412–425. [[CrossRef](#)]
11. Ye, B.; Ji, X.; Yang, H.; Yao, X.; Chan, C.K.; Cadle, S.H.; Chan, T.; Mulawa, P.A. Concentration and chemical composition of PM_{2.5} in Shanghai for a 1-year period. *Atmos. Environ.* **2003**, *37*, 499–510. [[CrossRef](#)]
12. Wang, X.; Bi, X.; Sheng, G.; Fu, J. Chemical composition and sources of PM₁₀ and PM_{2.5} aerosols in Guangzhou, China. *Environ. Monit. Assess.* **2006**, *119*, 425–439. [[CrossRef](#)] [[PubMed](#)]
13. Yang, H.; Yu, J.Z.; Ho, S.S.H.; Xu, J.; Wu, W.; Wan, C.H.; Wang, X.; Wang, X.; Wang, L. The chemical composition of inorganic and carbonaceous materials in PM_{2.5} in Nanjing, China. *Atmos. Environ.* **2005**, *39*, 3735–3749. [[CrossRef](#)]
14. Shen, G.F.; Yuan, S.Y.; Xie, Y.N.; Xia, S.J.; Li, L.; Yao, Y.K.; Qiao, Y.Z.; Zhang, J.; Zhao, Q.Y.; Ding, A.J.; et al. Ambient levels and temporal variations of PM_{2.5} and PM₁₀ at a residential site in the mega-city, Nanjing, in the western Yangtze River delta, China. *J. Environ. Sci. Health Part A* **2014**, *49*, 171–178. [[CrossRef](#)] [[PubMed](#)]
15. Tao, J.; Zhang, L.; Engling, G.; Zhang, R.; Yang, Y.; Cao, J.; Zhu, C.; Wang, Q.; Luo, L. Chemical composition of PM_{2.5} in an urban environment in Chengdu, China: Importance of springtime dust storms and biomass burning. *Atmos. Res.* **2013**, *122*, 270–283. [[CrossRef](#)]
16. Fontes, T.; Li, P.; Barros, N.; Zhao, P. Trends of PM_{2.5} concentrations in China: A long term approach. *J. Environ. Manag.* **2017**, *196*, 719–732. [[CrossRef](#)] [[PubMed](#)]
17. Fang, C.; Zhang, Z.; Jin, M.; Zou, P.; Wang, J. Pollution characteristics of PM_{2.5} aerosol during haze periods in Changchun, China. *Aerosol Atmos. Chem.* **2017**, *17*, 888–895. [[CrossRef](#)]

18. Shimou, Y. *Spatial Expansion of the Metropolitans in China*; Chinese Technology Press: Beijing, China, 1998.
19. Zhao, P.S.; Dong, F.; He, D.; Zhao, X.J.; Zhang, X.L.; Zhang, W.Z.; Yao, Q.; Liu, H.Y. Characteristics of concentrations and chemical compositions for PM_{2.5} in the region of Beijing, Tianjin, and Hebei, China. *Atmos. Chem. Phys.* **2013**, *13*, 4631–4644. [CrossRef]
20. Ming, L.; Jin, L.; Li, J.; Fu, P.; Yang, W.; Liu, D.; Zhang, G.; Wang, Z.; Li, X. PM_{2.5} in the Yangtze River delta, China: Chemical compositions, seasonal variations, and regional pollution events. *Environ. Pollut.* **2017**, *223*, 200–212. [CrossRef] [PubMed]
21. Hagler, G.S.W.; Bergin, M.H.; Salmon, L.G.; Yu, J.Z.; Wan, E.C.H.; Zheng, M.; Zeng, L.M.; Kiang, C.S.; Zhang, Y.H.; Lau, A.K.H.; et al. Source areas and chemical composition of fine particulate matter in the Pearl River delta region of China. *Atmos. Environ.* **2006**, *40*, 3802–3815. [CrossRef]
22. Wang, X.; Chen, W.; Chen, D.; Wu, Z.; Fan, Q. Long-term trends of fine particulate matter and chemical composition in the Pearl River delta economic zone (prdez), China. *Front. Environ. Sci. Eng.* **2016**, *10*, 53–62. [CrossRef]
23. Tang, X.; Chen, X.; Tian, Y. Chemical composition and source apportionment of PM_{2.5}—A case study from one year continuous sampling in the Chang-Zhu-Tan urban agglomeration. *Atmos. Pollut. Res.* **2017**, *8*, 885–899. [CrossRef]
24. Chan, C.K.; Yao, X. Air pollution in mega cities in China. *Atmos. Environ.* **2008**, *42*, 1–42. [CrossRef]
25. Aleksandropoulou, V.; Lazaridis, M. Trends in population exposure to particulate matter in urban areas of Greece during the last decade. *Sci. Total Environ.* **2017**, *581–582*, 399–412. [CrossRef] [PubMed]
26. Pant, P.; Habib, G.; Marshall, J.D.; Peltier, R.E. PM_{2.5} exposure in highly polluted cities: A case study from New Delhi, India. *Environ. Res.* **2017**, *156*, 167–174. [CrossRef] [PubMed]
27. Zhang, A.; Qi, Q.; Jiang, L.; Zhou, F.; Wang, J. Population exposure to PM_{2.5} in the urban area of Beijing. *PLoS ONE* **2013**, *8*, e63486. [CrossRef] [PubMed]
28. Wang, G.; Gu, S.; Chen, J.; Wu, X.; Yu, J. Assessment of health and economic effects by PM_{2.5} pollution in Beijing: A combined exposure-response and computable general equilibrium analysis. *Environ. Technol.* **2016**, *37*, 3131–3138. [CrossRef] [PubMed]
29. Liu, Z.; Huang, H.; Werners, S.E.; Yan, D. Construction area expansion in relation to economic-demographic development and land resource in the Pearl River delta of China. *J. Geogr. Sci.* **2016**, *26*, 188–202. [CrossRef]
30. Yao, L.; Lu, N. Particulate matter pollution and population exposure assessment over mainland China in 2010 with remote sensing. *Int. J. Environ. Res. Public Health* **2014**, *11*, 5241–5250. [CrossRef] [PubMed]
31. Wu, J.; Yao, F.; Li, W.; Si, M. Viirs-based remote sensing estimation of ground-level PM_{2.5} concentrations in Beijing–Tianjin–Hebei: A spatiotemporal statistical model. *Remote Sens. Environ.* **2016**, *184*, 316–328. [CrossRef]
32. Wang, S.; Zhou, C.; Wang, Z.; Feng, K.; Hubacek, K. The characteristics and drivers of fine particulate matter (PM_{2.5}) distribution in China. *J. Clean. Prod.* **2017**, *142 Pt 4*, 1800–1809. [CrossRef]
33. CNEMC. China national Environmental Monitoring Center (CNEMC). Available online: <http://www.cnemc.cn> (accessed on 18 April 2017).
34. NESSDSI. National Earth System Science Data Sharing Infrastructure. Available online: www.geodata.cn/ (accessed on 18 April 2017).
35. Goovaerts, P. Ordinary cokriging revisited. *Math. Geosci.* **1998**, *30*, 21–42.
36. Mahmud, A.; Hixson, M.; Kleeman, M.J. Quantifying population exposure to airborne particulate matter during extreme events in California due to climate change. *Atmos. Chem. Phys.* **2012**, *12*, 7453–7463. [CrossRef]
37. Zhang, H.; Wang, Z.; Zhang, W. Exploring spatiotemporal patterns of PM_{2.5} in China based on ground-level observations for 190 cities. *Environ. Pollut.* **2016**, *216*, 559–567. [CrossRef] [PubMed]
38. Yang, F.; Tan, J.; Zhao, Q.; Du, Z.; He, K.; Ma, Y.; Duan, F.; Chen, G.; Zhao, Q. Characteristics of PM_{2.5} speciation in representative megacities and across China. *Atmos. Chem. Phys.* **2011**, *11*, 5207–5219. [CrossRef]
39. Zhang, Y.L.; Cao, F. Fine particulate matter (PM_{2.5}) in China at a city level. *Sci. Rep.* **2015**, *5*, 14884. [CrossRef] [PubMed]
40. World Health Organization (WHO). *WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide*; WHO/SDE/PHE/OEH/06.02 Vols.; World Health Organization: Geneva, Switzerland, 2005.

41. MEP. Ambient Air Quality Standards (gb3095-2012). Ministry of Environmental Protection of the People's Republic of China (MEP). Available online: http://kjs.mep.gov.cn/hjbhbz/bzwb/dqhjbh/dqhjzlbz/201203/t20120302_224165.shtml (accessed on 18 April 2017).
42. Pui, D.Y.H.; Chen, S.C.; Zuo, Z. PM_{2.5} in China: Measurements, sources, visibility and health effects, and mitigation. *Particuology* **2014**, *13*, 1–26. [[CrossRef](#)]
43. Zhang, Y.; Cai, J.; Wang, S.; He, K.; Zheng, M. Review of receptor-based source apportionment research of fine particulate matter and its challenges in China. *Sci. Total Environ.* **2017**, *586*, 917–929. [[CrossRef](#)] [[PubMed](#)]
44. Huang, C.; Chen, C.H.; Li, L.; Cheng, Z.; Wang, H.L.; Huang, H.Y.; Streets, D.G.; Wang, Y.J.; Zhang, G.F.; Chen, Y.R. Emission inventory of anthropogenic air pollutants and VOC species in the Yangtze River delta region, China. *Atmos. Chem. Phys.* **2011**, *11*, 4105–4120. [[CrossRef](#)]
45. Hua, Y.; Cheng, Z.; Wang, S.; Jiang, J.; Chen, D.; Cai, S.; Fu, X.; Fu, Q.; Chen, C.; Xu, B.; et al. Characteristics and source apportionment of PM_{2.5} during a fall heavy haze episode in the Yangtze River delta of China. *Atmos. Environ.* **2015**, *123*, 380–391. [[CrossRef](#)]
46. Xu, J.; Chang, L.; Qu, Y.; Yan, F.; Wang, F.; Fu, Q. The meteorological modulation on PM_{2.5} interannual oscillation during 2013 to 2015 in Shanghai, China. *Sci. Total Environ.* **2016**, *572*, 1138–1149. [[CrossRef](#)] [[PubMed](#)]
47. Lin, Z.J.; Tao, J.; Chai, F.H.; Fan, S.J.; Yue, J.H.; Zhu, L.H.; Ho, K.F.; Zhang, R.J. Impact of relative humidity and particles number size distribution on aerosol light extinction in the urban area of Guangzhou. *Atmos. Chem. Phys.* **2013**, *13*, 1115–1128. [[CrossRef](#)]
48. Tian, M.; Wang, H.; Chen, Y.; Zhang, L.; Shi, G.; Liu, Y.; Yu, J.; Zhai, C.; Wang, J.; Yang, F. Highly time-resolved characterization of water-soluble inorganic ions in PM_{2.5} in a humid and acidic mega city in Sichuan Basin, China. *Sci. Total Environ.* **2017**, *580*, 224–234. [[CrossRef](#)] [[PubMed](#)]
49. Wang, H.; Shi, G.; Tian, M.; Zhang, L.; Chen, Y.; Yang, F.; Cao, X. Aerosol optical properties and chemical composition apportionment in Sichuan Basin, China. *Sci. Total Environ.* **2017**, *577*, 245–257. [[CrossRef](#)] [[PubMed](#)]
50. Liao, T.; Wang, S.; Ai, J.; Gui, K.; Duan, B.; Zhao, Q.; Zhang, X.; Jiang, W.; Sun, Y. Heavy pollution episodes, transport pathways and potential sources of PM_{2.5} during the winter of 2013 in Chengdu (China). *Sci. Total Environ.* **2017**, *584–585*, 1056–1065. [[CrossRef](#)] [[PubMed](#)]
51. Van Donkelaar, A.; Martin, R.V.; Brauer, M.; Kahn, R.; Levy, R.; Verduzco, C.; Villeneuve, P.J. Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: Development and application. *Environ. Health Perspect.* **2010**, *118*, 847–855. [[CrossRef](#)] [[PubMed](#)]
52. Ivy, D.; Mulholland, J.A.; Russell, A.G. Development of ambient air quality population-weighted metrics for use in time-series health studies. *J. Air Waste Manag. Assoc.* **2008**, *58*, 711–720. [[CrossRef](#)] [[PubMed](#)]

