



Melanocortin 3 receptor-expressing neurons in the ventromedial hypothalamus promote glucose disposal

Amy K. Sutton^{a,1}, Paulette B. Goforth^{b,1}, Ian E. Gonzalez^a, James Dell'Orco^c, Hongjuan Pei^d, Martin G. Myers Jr^{a,c}, and David P. Olson^{a,d,2}

^aDepartment of Molecular and Integrative Physiology, University of Michigan Medical School, Ann Arbor, MI 48109; ^bDepartment of Pharmacology, University of Michigan Medical School, Ann Arbor, MI 48109; ^cDepartment of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI 48109; and ^dDivision of Endocrinology, Department of Pediatrics, University of Michigan Medical School, Ann Arbor, MI 48109

Edited by Susan G. Amara, National Institutes of Health, Bethesda, MD, and approved March 9, 2021 (received for review February 20, 2021)

The ventromedial hypothalamus (VMH) is a critical neural node that senses blood glucose and promotes glucose utilization or mobilization during hypoglycemia. The VMH neurons that control these distinct physiologic processes are largely unknown. Here, we show that melanocortin 3 receptor (MC3R)-expressing VMH neurons (VMH^{MC3R}) sense glucose changes both directly and indirectly via altered excitatory input. We identify presynaptic nodes that potentially regulate VMH^{MC3R} neuronal activity, including inputs from proopiomelanocortin (POMC)-producing neurons in the arcuate nucleus. We find that VMH^{MC3R} neuron activation blunts, and their silencing enhances glucose excursion following a glucose load. Overall, these findings demonstrate that VMH^{MC3R} neurons are a glucose-responsive hypothalamic subpopulation that promotes glucose disposal upon activation; this highlights a potential site for targeting dysregulated glycemia.

hypothalamus | ventromedial nucleus | hyperglycemia

The central nervous system contributes to systemic glucose homeostasis. Glucose sensing neurons are found within the ventromedial hypothalamus (VMH), a brain structure that plays a prominent role in both the counterregulatory response to hypoglycemia (1–3), and glucose disposal (4). Direct VMH infusion of melanocortin agonists, endogenously produced in the arcuate nucleus (ARC), promotes glucose uptake into skeletal muscle independent of insulin release (5, 6). Furthermore, VMH-specific reexpression of MC3Rs in an MC3R-null background increases glucose utilization without altering energy balance (7), suggesting that VMH^{MC3R} neurons may regulate glucose disposal.

Using genetic tools, we find that VMH^{MC3R} neurons respond directly and indirectly to altered glucose levels. Potential presynaptic regulators of VMH^{MC3R} neurons include melanocortin-producing ARC neurons and other brain regions important for glycemic control. We demonstrate that VMH^{MC3R} neuron activity is both necessary and sufficient for glucose disposal, and that activation of this population has the potential to attenuate diet-induced glucose intolerance.

Results

VMH^{MC3R} Neurons Sense Glucose Changes Both Directly and Indirectly and Receive Input from Glucose-Sensing Regions. *Mc3R-2a-Cre* mice (8) were crossed to the *Rosa26^{Ai14idTomato}* reporter for electrophysiological analysis of VMH^{MC3R} neurons (Fig. 1 A–C). Application of the MC3R-selective agonist d-trp8-γMSH (10 nM) depolarized VMH^{MC3R} neurons (Fig. 1 D and E) (9). Decreasing glucose from 2 to 0.1 mM suppressed the activity of VMH^{MC3R} neurons; returning glucose to 2 mM depolarized VMH^{MC3R} neurons and restored their firing (Fig. 1 F, G, and I). Increasing glucose to 5 mM did not significantly affect VMH^{MC3R} membrane potential or firing (Fig. 1 F, H, and I). The hyperpolarization of VMH^{MC3R} neurons in low glucose persisted during blockade of action potential (AP) firing and synaptic input in four of seven neurons, indicating direct glucose regulation (Fig. 1J).

Likewise, low glucose activated an outward current in five of nine cells (Fig. 1 K and L).

Indirect regulation of VMH^{MC3R} neurons is suggested by the observation that hypoglycemia inhibits nearly all VMH^{MC3R} neurons, yet only a subset sensed glucose directly. Indeed, lowering glucose without synaptic blockade decreased the frequency and amplitude of spontaneous excitatory postsynaptic currents (sEPSCs) on VMH^{MC3R} neurons, thus demonstrating indirect glucose-dependent regulation via presynaptic input (Fig. 1 M–O). There was no change in spontaneous inhibitory postsynaptic currents (sIPSCs).

Given these glucose-dependent alterations in excitatory input, we used monosynaptic retrograde tracing to inputs to VMH^{MC3R} neurons. Projection-specific modified rabies virus tracing was performed in the bed nucleus of the stria terminalis (BNST), a site targeted by MC3R axons relevant for VMH-mediated blood glucose regulation (Fig. 1P) (2). BNST-projecting VMH^{MC3R} neurons receive inputs from several brain areas relevant to glucose regulation including POMC cells of the ARC (Fig. 1S), and cells in the lateral parabrachial nucleus (IPBN) (Fig. 1W).

VMH^{MC3R} Neurons Regulate Glucose Disposal. To directly test the capacity of VMH^{MC3R} neurons to promote glucose disposal, we chemogenetically activated VMH^{MC3R} neurons using the Cre-dependent modified human muscarinic receptor hM3Dq (AAV8-hSyn-Flex-hM3Dq-mCherry; Fig. 2 A–D) 15 min prior to a glucose tolerance test (GTT) (Fig. 2E). Preactivation of VMH^{MC3R} neurons blunts glucose excursion in comparison to controls (Fig. 2 F and G). This effect appears to be largely independent of insulin secretion, as insulin is unchanged 5 min after glucose administration (Fig. 2 H and I).

To test the potential benefit of activating VMH^{MC3R} neurons in a glucose-intolerant state, we chemogenetically activated VMH^{MC3R} neurons prior to a GTT in mice fed a 60% high-fat diet (HFD) (Fig. 2J). Preactivation of VMH^{MC3R} neurons promotes glucose disposal in HFD conditions (Fig. 2 K and L), suggesting that driving VMH^{MC3R} neuron activity may be sufficient to promote glucose disposal in glucose-intolerant states. Since VMH^{MC3R} activation without a glucose load does not significantly alter peripheral glucose levels in mice on either chow or HFD conditions (Fig. 2 M and N), the effects of VMH^{MC3R} neuronal activation may be most impactful during a glucose load.

To examine the contribution of VMH^{MC3R} neurons to glucose homeostasis, we silenced these neurons using a Cre-dependent

Author contributions: A.K.S., P.B.G., and D.P.O. designed research; A.K.S., P.B.G., I.E.G., J.D., and H.P. performed research; M.G.M. and D.P.O. contributed new reagents/analytic tools; A.K.S. and P.B.G. analyzed data; and A.K.S., P.B.G., and D.P.O. wrote the paper.

Competing interest statement: NovoNordisk provides research grant support to D.P.O. and M.G.M. Ionis Pharmaceuticals provides grant support to M.G.M.

This open access article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹A.K.S. and P.B.G. contributed equally to this work.

²To whom correspondence may be addressed. Email: dpolson@med.umich.edu.

Published April 7, 2021.




Fig. 1. VMH^{MC3R} neurons are regulated by melanocortins and glucose, with inputs from local brain regions. (A–C) VMH^{MC3R} neurons visualized in *Mc3R-2a-Cre, Td* mice. Current-clamp recordings show depolarization of VMH^{MC3R} neurons by d-trp8-γMSH with synaptic blockade (D and E; 4/5 neurons), decreased AP firing (F and G; 10/10 neurons), and hyperpolarization in low glucose (F and I; 11/13 neurons). Approximately 50% (4/7) of VMH^{MC3R} neurons directly respond to low glucose with synaptic blockers present (J), likely due to a net outward current (K and L). Indirect regulation of VMH^{MC3R} neurons in low glucose was measured by simultaneously recording EPSC and IPSCs (M). Low glucose decreased amplitude and frequency of sEPSCs, but not sIPSCs (N and O). Identification of presynaptic inputs to VMH^{MC3R} neurons projecting to the BNST using modified rabies virus tracing (P–R) identifies ARC inputs (S, purple) containing αMSH (S, green). Additional inputs include the PVH (U), MeA (V), IPBN (W), and BMP (X). Data are represented as mean ± SEM (G, H, and O), or box plots ± maximum/minimum (E, I, and J). Significance was determined using a repeated-measures one-way ANOVA (E, G, H, J, K, and O), or mixed-effects analysis followed by Tukey's post hoc if applicable (I) with **P* < 0.05, ***P* < 0.01, and ****P* < 0.001.

tetanus toxin (AAV8-hSyn-Flex-TetTox-GFP; Fig. 2 O and P). Loss of VMH^{MC3R} neuronal activity does not alter blood glucose levels in the basal state (Fig. 2Q) but does increase blood glucose during a GTT (Fig. 2R and S). VMH^{MC3R} neuron silencing did not affect body weight or food intake (body weight at *t* = 45 d post injection [mean ± SEM]: TetTox, 30.4 ± 1.9 g; control, 29.6 ± 1.0 g; two-way ANOVA), suggesting that the impaired glucose handling was not secondary to obesity alone.

Discussion

The VMH regulates both glucose mobilization during the counterregulatory response and glucose disposal in response to hormonal factors (1–4). However, the cellular mechanisms coordinating these seemingly opposing physiologic functions are largely unknown. Here, we show that VMH^{MC3R} neurons respond to melanocortin agonists and their activation promotes glucose disposal in both chow and HFD conditions. Silencing of VMH^{MC3R} neurons disrupts normal glucose handling, underscoring an important role for these neurons in glucose homeostasis. These neurons sense glucose directly and indirectly via excitatory synaptic inputs, which potentially arise from ARC POMC neurons known to modulate blood glucose (10) and PBN neurons. VMH^{MC3R} neurons do not directly respond to high glucose but appear situated to integrate input from upstream glucose sensing neurons and/or other stimuli to promote glucose uptake.

Activation of SF1-Cre VMH neurons (which represent most VMH neurons) mimics the counterregulatory response (CRR), suggesting that the autonomic and neuroendocrine effects on glucose mobilization during the CRR supersede the ability of peripheral tissues to dispose of glucose (2, 3). We show that VMH^{MC3R} neurons are a subset of VMH cells that promote glucose disposal, a role opposing pan-VMH activation. Our results highlight a unique circuit node capable of lowering peripheral blood glucose in both normal and HFD conditions and demonstrate the importance of fine-tuned analysis of neural circuits in the assessment of discrete physiologic outputs. Taken together, VMH^{MC3R} neurons may

be a potential site for targeted therapies of dysregulated glycemia, including glucose intolerance and type II diabetes.

Methods

Animals. Animals were bred and housed according to guidelines approved by the University of Michigan Committee on the Care and Use of Animals. Male mice (8–23 wk) were used for all behavioral studies. Mice were provided ad libitum food (standard chow, Purina Laboratory Diet 5001; 60% HFD, Research Diets D12492) and water, unless otherwise noted, and acclimated to intraperitoneal (i.p.) injections 3 d prior to experimentation. Following experimentation, mice were perfused, immunohistochemistry (IHC) performed as previously described (11), and data points excluded if extra-VMH viral transduction occurred. IHC was performed for dsRed (Clontech), GFP (Invitrogen), Fos (Cell Signaling 9F6), and αMSH (Phoenix Pharmaceuticals).

Stereotaxic Injections. Surgeries were performed as previously described (11). VMH-directed injections (25 nL/site) were performed relative to bregma (anteroposterior [A/P]: −1.0; mediolateral [M/L]: ± 0.25; dorsoventral [D/V]: −5.5). Mice recovered for at least 2 wk before testing. For rabies tracing, helper virus (11) was injected followed by EnvA-dB19G-rabies-mCh injections (BNST) (A/P: +0.65; M/L: +0.4; D/V: −3.8) 3 wk later. Mice were perfused and IHC performed 5 d after virus injection.

Electrophysiology. Coronal slices (250 μm) were prepared from 4- to 12-wk-old *Mc3R-2a-Cre, Td* mice of either sex in oxygenated ice-cold artificial cerebrospinal fluid (ACSF) solution containing the following (in mM): 125 NaCl, 3 KCl, 1.25 NaH₂PO₄, NaHCO₃, 2 glucose, 1 MgCl₂, and 2.5 CaCl₂, plus 0.2 mM ascorbic acid and 1 mM kynurenic acid, pH 7.4, 300–305 mOsm, and allowed to recover ≥1 h (kynurenic acid) at 33–34 °C. Patch-clamp recordings from VMH^{MC3R} neurons were made using borosilicate electrodes (3–7 MΩ) filled with either perforated patch internal (in mM): 130 K gluconate, 10 KCl, 1 EGTA, 10 Hepes, 2 MgCl₂, amphotericin B (0.15 mg/mL) for current-clamp recordings of Vm/AP firing in response to glucose (Fig. 1 F–J) or whole-cell internal solution (in mM): 130 K gluconate, 10 KCl, 1 EGTA, Hepes, 0.6 NaGTP, 2 MgATP, and 8 phosphocreatine, pH 7.2, 285–295 mOsm, for all other experiments. Slices were perfused with ACSF solution (33–34 °C, 1.5–2 mL/min) and bubbled with 5% CO₂/95% O₂. When stated, synaptic input and AP firing were blocked by inclusion of 1 μM tetrodotoxin, 20 μM D-APV, 10 μM CNQX, and 50 μM picrotoxin.




Fig. 2. VMH^{MC3R} neuron activity alters glucose disposal. DREADD(hM3Dq)-mediated activation of VMH^{MC3R} neurons (A and B) using i.p. CNO increases activation-induced Fos in the VMH (C and D). Preactivation of VMH^{MC3R} neurons using hM3Dq blunts glucose excursion during a GTT (E–G; $n = 7$) without altering plasma insulin levels 5 min after glucose injection (H and I; $n = 4$). (J–L) Activation of VMH^{MC3R} neurons promotes glucose disposal during HFD conditions. (M and N) While HFD-fed mice differ in baseline glucose levels compared to chow conditions (main effects of food and time without interaction, three-way ANOVA), activation of VMH^{MC3R} neurons does not affect glucose disposal in chow or HFD mice ($n = 4$) absent a glucose load. Tetanus toxin silencing of VMH^{MC3R} neurons (O and P) does not alter basal blood glucose (Q) and increases glucose excursion upon a glucose load (R and S) (TetTox, $n = 5$; control, $n = 7$). Significance was determined using a repeated-measures two-way ANOVA (F, G, K, and L), paired *t* test (F, G, K, and L; AUC, J), two-way ANOVA (R and S), three-way ANOVA (M and N), or an unpaired *t* test (Q, R, and S, AUC), all with Tukey's post hoc as applicable, with * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$.

Spontaneous EPSCs and IPSCs were measured from neurons voltage clamped to -50 mV, producing inward sEPSCs and outward sIPSCs, and analyzed using Synaptosoft software. Data were not corrected for a junction potential of ~ 15 mV. Analyzed neurons had uncompensated stable series resistances (< 30 M Ω).

GTT. Following a 4-h fast, mice were injected with glucose (2 g/kg, i.p.), and blood glucose monitored by tail vein (OneTouch Ultra 2 glucometer). For hM3Dq experiments, mice received vehicle (10% β -cyclodextran) or CNO (0.3 mg/kg) 15 min prior to glucose administration counterbalanced across at least 1 wk. Insulin analysis was performed in a separate experiment using blood collected via tail vein; insulin levels were measured by enzyme-linked immunosorbent assay (Crystal Chem). Separate experiments were performed with 60% HFD (Research Diets).

VMH^{MC3R} Silencing. *Mc3R-2a-Cre* mice were injected with AAV8-hSyn-Flex-TetTox and compared to two control groups: *Mc3R-2a-Cre* mice + AAV8-hSyn-Flex-GFP, or WT mice + AAV8-hSyn-Flex-TetTox, which showed no statistical difference on any measured variable and were pooled into one control group. GTT was performed 3 wk postsurgery.

Statistical Analyses. Paired and unpaired *t* tests, one-way, two-way, or three-way ANOVAs followed by Tukey's multiple comparisons (if applicable), or mixed model analyses were calculated using GraphPad Prism 8 as appropriate, with significance denoted at $P < 0.05$.

Data Availability. All study data are included in the article and/or supporting information.

- O. Chan, R. Sherwin, Influence of VMH fuel sensing on hypoglycemic responses. *Trends Endocrinol. Metab.* **24**, 616–624 (2013).
- T. H. Meek *et al.*, Functional identification of a neurocircuit regulating blood glucose. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E2073–E2082 (2016).
- J. N. Flak *et al.*, Ventromedial hypothalamic nucleus neuronal subset regulates blood glucose independently of insulin. *J. Clin. Invest.* **130**, 2943–2952 (2020).
- Z. Song, B. E. Levin, J. J. McArdle, N. Bakhos, V. H. Routh, Convergence of pre- and postsynaptic influences on glucosensing neurons in the ventromedial hypothalamic nucleus. *Diabetes* **50**, 2673–2681 (2001).
- C. Toda *et al.*, Distinct effects of leptin and a melanocortin receptor agonist injected into medial hypothalamic nuclei on glucose uptake in peripheral tissues. *Diabetes* **58**, 2757–2765 (2009).
- C. K. Gavini, W. C. Jones 2nd, C. M. Novak, Ventromedial hypothalamic melanocortin receptor activation: Regulation of activity energy expenditure and skeletal muscle thermogenesis. *J. Physiol.* **594**, 5285–5301 (2016).
- K. Begriche *et al.*, Genetic dissection of the functions of the melanocortin-3 receptor, a seven-transmembrane G-protein-coupled receptor, suggests roles for central and peripheral receptors in energy homeostasis. *J. Biol. Chem.* **286**, 40771–40781 (2011).
- H. Pei *et al.*, Lateral hypothalamic MC3R-expressing neurons modulate locomotor activity, energy expenditure, and adiposity in male mice. *Endocrinology* **160**, 343–358 (2019).
- P. Grieco, P. M. Balse, D. Weinberg, T. MacNeil, V. J. Hruby, *D*-Amino acid scan of γ -melanocyte-stimulating hormone: Importance of Trp(8) on human MC3 receptor selectivity. *J. Med. Chem.* **43**, 4998–5002 (2000).
- J. W. Hill *et al.*, Direct insulin and leptin action on pro-opiomelanocortin neurons is required for normal glucose homeostasis and fertility. *Cell Metab.* **11**, 286–297 (2010).
- A. K. Sutton *et al.*, Paraventricular, subparaventricular and periventricular hypothalamic IRS4-expressing neurons are required for normal energy balance. *Sci. Rep.* **10**, 5546 (2020).