

# ‘Heart development and morphogenesis’ is a novel pathway for human ovarian granulosa cell differentiation during long-term *in vitro* cultivation-a microarray approach

WIESŁAWA KRANC<sup>1</sup>, MACIEJ BRAŻERT<sup>2</sup>, PIOTR CELICHOWSKI<sup>3</sup>, ARTUR BRYJA<sup>1</sup>, MARIUSZ J. NAWROCKI<sup>1</sup>, KATARZYNA OŻEGOWSKA<sup>2</sup>, MAURZYCY JANKOWSKI<sup>1</sup>, MICHAŁ JESETA<sup>4</sup>, LESZEK PAWELCZYK<sup>2</sup>, ANDRZEJ BRĘBOROWICZ<sup>5</sup>, DOMINIK RACHOŃ<sup>6</sup>, MARIUSZ T. SKOWROŃSKI<sup>7</sup>, MAŁGORZATA BRUSKA<sup>1</sup>, MACIEJ ZABEL<sup>8,9</sup>, MICHAŁ NOWICKI<sup>3</sup> and BARTOSZ KEMPISTY<sup>1,3,4</sup>

<sup>1</sup>Department of Anatomy, Poznan University of Medical Sciences, 60-781 Poznań; <sup>2</sup>Division of Infertility and Reproductive Endocrinology, Department of Gynecology, Obstetrics and Gynecological Oncology, Poznan University of Medical Sciences, 60-535 Poznań; <sup>3</sup>Department of Histology and Embryology, Poznan University of Medical Sciences, 60-781 Poznań, Poland; <sup>4</sup>Department of Obstetrics and Gynecology, University Hospital and Masaryk University, 62500 Brno, Czech Republic; <sup>5</sup>Department of Pathophysiology, Poznań University of Medical Sciences, 60-806 Poznań; <sup>6</sup>Department of Clinical and Experimental Endocrinology, Medical University of Gdańsk, 80-211 Gdańsk; <sup>7</sup>Faculty of Biology and Biotechnology, University of Warmia and Mazury in Olsztyn, 10-719 Olsztyn; <sup>8</sup>Division of Histology and Embryology, Department of Human Morphology and Embryology, Wrocław Medical University, 50-368 Wrocław; <sup>9</sup>Division of Anatomy and Histology, University of Zielona Góra, 65-046 Zielona Góra, Poland

Received July 13, 2018; Accepted December 6, 2018

DOI: 10.3892/mmr.2019.9837

**Abstract.** Granulosa cells (GCs) have many functions in the endocrine system. Most notably, they produce progesterone following ovulation. However, it has recently been proven that GCs can change their properties when subjected to long-term culture. In the present study, GCs were collected from hyper-stimulated ovarian follicles during *in vitro* fertilization procedures. They were grown *in vitro*, in a long-term manner. RNA was collected following 1, 7, 15 and 30 days of culture. Expression microarrays were used for analysis, which allowed to identify groups of genes characteristic for particular cellular processes. In addition, reverse transcription-quantitative polymerase chain reaction (RT-qPCR) was performed to validate the obtained results. Two ontological groups characteristic for processes associated with the development and morphogenesis of the heart were identified during the analyses: ‘Heart development’ and ‘heart morphogenesis’. The results of the microarrays revealed that the highest change in expression was demonstrated by the *lysyl Oxidase*, *oxytocin receptor*,

*nexilin F-actin binding protein*, and *cysteine-rich protein 3* genes. The lowest change was exhibited by *odd-skipped related transcription factor 1*, *plakophilin 2*, *transcription growth factor-β receptor 1*, and *kinesin family member 3A*. The direction of changes was confirmed by RT-qPCR results. In the present study, it was suggested that GCs may have the potential to differentiate towards other cell types under long-term *in vitro* culture conditions. Thus, genes belonging to the presented ontological groups can be considered as novel markers of proliferation and differentiation of GCs towards the heart muscle cells.

## Introduction

The folliculogenesis is a compound process that leads to the formation of the fully grown ovarian follicle (1,2). The developed follicle is composed of several cell layers, including theca cells (TCs) and granulosa cells (GCs). Our recent study has shown that porcine GCs undergo permanent morphological modification when kept in the primary *in vitro* culture system. After long-term culture, GCs underwent transformation into spindle-like structures, which was accompanied by changes in their gene expression profile. Moreover, the GCs display structural and functional role in the active pre-ovulatory and ovulatory follicle, forming follicular architecture and expressing a number of hormone receptors. Some of that properties seem to prevail *ex vivo*, as we have observed a time-dependent expression of *LHR*, *FSHR*, *CYP19*, and progesterone receptor genes, in porcine GCs during long-term *in vitro* culture.

*Correspondence to:* Dr Bartosz Kempisty, Department of Anatomy, Poznań University of Medical Sciences, 6 Święcickiego Street, 60-781 Poznań, Poland  
E-mail: bkempisty@ump.edu.pl

**Key words:** human granulosa cells, *in vitro* culture, proliferation, differentiation, heart development

Using a porcine model, we have recently shown that GCs, isolated from pre-ovulatory ovarian follicles, can be successfully cultured *in vitro* for short and/or long time periods. Moreover, we found that these GCs are characterized by increased *in vitro* proliferation capability. Although the increase in cells' proliferation is accompanied by cellular ageing and subsequently decreased by a number of passages, the GCs have also displayed significant differentiability capacity. It was also presented that these cells may differentiate even into cell types unassociated with the reproductive process. Kossowska-Tomaszczuk recently determined that after long-term cultivation and under influence of several substances, including LIF and dexamethasone, GCs may differentiate into chondroblasts, osteoblasts and other adult precursor cells (3,4). These experiments have also shown how the huge differentiability plasticity of human GCs during primary *in vitro* culture can be used in clinical situations. This may open a new gate in the potential application of human granulosa cells, which are nowadays mostly treated as remnant IVF material, mostly in autologous and donor-host grafts, as they exhibit increased stem-like specificity.

The experiments by Kossowska-Tomaszczuk open the possibility of finding new stemness markers in human ovarian granulosa cells. Microarray assays allow for robust and complex transcriptomic analysis, leading to the discovery of new molecular markers of cells function. In this work, we have used the microarrays to describe the transcriptomic profile of human GCs during their long-term primary *in vitro* culture. We have detected the expression of genes belonging to more than five hundred gene ontology groups, associated with the 'biology of human GCs' during their *in vitro* proliferation and differentiation. In this study, we have investigated the 'heart development and morphogenesis' ontological group, as the potential source of new markers of GCs plasticity.

## Materials and methods

A large part of the materials and methods section is based on other, published work of the author, presenting results from the same cycle of studies describing human granulosa cells (5).

*Patients and collection of granulosa cells.* The GCs were derived from patients undergoing *in vitro* fertilization (IVF) procedures, who had given their informed, written consent to be included in this protocol. The study group consisted of 8 patients, aged 18-40 years, with diagnosed infertility, referred to the Division of Infertility and Reproductive Endocrinology, Poznan University of Medical Sciences, Poznan, Poland. Patients underwent IVF procedure, based on controlled ovarian hyperstimulation protocol, adjusted to the patient's initial infertility workup and ovarian response. Stimulation was performed with human recombinant FSH (Gonal-F, Merck Serono) and highly purified hMG-HP (Menopur, Ferring). The injections with cetrorelix acetate (Cetrotide, Merck Serono) were administered in an adequate dose, to suppress pituitary function. Ovulation triggering was based on the subcutaneous injection of 6,500 U of hCG (Ovitrelle, Merck-Serono). GC containing follicular fluid was collected during transvaginal, ultrasound-guided oocyte pick-up, performed 36 h after human chorionic gonadotropin administration. The content of follicles

that were over 16 mm in diameter was immediately passed to an embryologist, who isolated the oocyte, and pooled the follicular fluid from each ovary. The fresh pooled samples were centrifuged for 10 min at 200 g, to separate and collect GCs. Patients with polycystic ovary syndrome (PCOS), endometriosis, and diminished ovarian reserve (serum antimüllerian hormone (AMH) less than 0.7 ng/ml, and/or day 2-3 FSH serum level higher than 15 mU/ml, and/or antral follicle count less than 9) were excluded from the study. This research has been approved by Poznan University of Medical Sciences Bioethical Committee with resolution 558/17.

*Primary cell culture.* The collected cells were washed twice by centrifugation at 200 x g for 10 min at RT in the culture medium. Medium consisted of Dulbecco's modified Eagle's medium (DMEM, Sigma; Merck KGaA, Darmstadt, Germany), 2% foetal bovine serum FBS (FBS; Sigma; Merck KGaA), 4 mM L-glutamine (stock 200 mM, Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA), 10 mg/ml gentamicin (Invitrogen; Thermo Fisher Scientific, Inc.), 10,000 U/ml penicillin, and 10,000 µg/ml streptomycin (Invitrogen; Thermo Fisher Scientific, Inc.). Cells were cultivated at 37°C under aerobic conditions (5% CO<sub>2</sub>). Once adherent cells were more than 90% confluent, they were detached with 0.05% trypsin-EDTA (Invitrogen; Thermo Fisher Scientific, Inc.) for 1-2 min and counted using a 'Neubauer improved' counting chamber (ISO LAB Laborgerate GmbH, DIN EN ISO Certified 9001). GCs were cultivated for 30 days. The medium was changed twice a week. Finally, total RNA was isolated from GCs after 1, 7, 15 and 30 days.

*Total RNA isolation.* Total RNA was isolated at 4 time periods, after 1, 7, 15 and 30 days of cultivation. The improved Chomczyński-Sacchi method was used for the RNA isolation (6). The GCs were suspended in 1 ml of a monophasic solution of guanidine thiocyanate and phenol (TRI Reagent®, Sigma; Merck KGaA). The chloroform was then added, with the samples centrifuged to obtain three separate phases. RNA was located in the upper, aqueous phase. RNA was then stripped with 2-propanol (Sigma; Merck KGaA, catalogue number I9516), added in an amount adequate for 1 ml of TRI-reagent, and washed with 75% ethanol. Samples prepared in that way were used for further analysis.

*Microarray expression analysis and statistics.* Total RNA (100 ng) from each pooled sample was subjected to two rounds of sense cDNA amplification (Ambion® WT Expression kit). The obtained cDNA was used for biotin labelling and fragmentation, with the use of GeneChip® WT Terminal Labeling and Hybridization (Affymetrix). Biotin-labelled fragments of cDNA (5.5 µg) were hybridized to the Affymetrix® Human Genome U219 Array (48°C/20 h). Microarrays were then washed and stained according to the technical protocol using the Affymetrix GeneAtlas Fluidics Station. The array strips were scanned employing Imaging Station of the GeneAtlas System. Preliminary analysis of the scanned chips was performed using Affymetrix GeneAtlas™ Operating Software. The quality of gene expression data was confirmed according to the quality control criteria provided by the software. The obtained CEL files were imported into downstream data analysis software.

All of the presented analyses and graphs were performed using Bioconductor and R programming languages. Each CEL file was merged with a description file. The Robust Multiarray Averaging (RMA) algorithm was used to correct background, normalize, and summarize the results. To determine the statistical significance of the analyzed genes, moderated t-statistics from the empirical Bayes method were performed. The obtained P-value was corrected for multiple comparisons using Benjamini and Hochberg's false discovery rate. The selection of significantly altered genes was based on a P-value beneath 0.05 and expression higher than two-fold.

Differentially expressed genes were subjected to selection by examination of genes involved in heart development and morphogenesis. The differentially expressed gene list (separated for up- and down-regulated genes) was uploaded to the DAVID software (Database for Annotation, Visualization and Integrated Discovery) (7).

Subsequently, the relationship between the genes belonging to chosen GO terms was analysed with the use of GOplot package (8). The GoPlot package had calculated the Z-score: The number of upregulated genes minus the number of down-regulated genes divided by the square root of the count. This information allowed to estimate the direction of changes in each gene-ontology term.

Moreover, interactions between differentially expressed genes/proteins belonging to the chosen GO terms were investigated by STRING10 software (Search Tool for the Retrieval of Interacting Genes) (9). The list of gene names was used as a query for the interaction prediction. The search criteria were based on co-occurrences of genes/proteins in scientific texts (text mining), co-expression, and experimentally observed interactions. The results of these analyses generated a gene/protein interaction network where the intensity of the edges reflected the strength of the interaction score.

Finally, the functional interactions between the genes belonging to the chosen GO BP terms were investigated using REACTOME FIViz application from the Cytoscape 3.6.0 software. This app accesses the pathways stored in the Reactome database, allowing for pathway enrichment analysis of a set of genes, visualization of hit pathways using manually laid-out pathway diagrams directly in Cytoscape, and investigation of functional relationships among genes in hit pathways. The app can also access the Reactome Functional Interaction (FI) network, a highly reliable, manually curated pathway-based protein functional interaction network covering over 60% of human proteins.

**Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) analysis.** The RT-qPCR method was performed to confirm the results obtained in the analysis of expression microarrays. Three genes were selected from each heatmap: The ones showing highest, lowest, and intermediate-level of expression. Changes in the level of expression of those genes were then examined. Three biological samples of each gene were used for the analysis. Each test was performed in 3 replicates. Reverse transcription was based on the protocols and reagents of SABiosciences (RT<sup>2</sup> First Stand kit-330401), using a Veritimer 96-well Thermal Cycler. One microgram of each gene's RNA transcript was used for reverse transcription.

Table I. Oligonucleotide sequences of primers used for reverse transcription-quantitative polymerase chain reaction analysis.

Gene	Primer sequences (5'-3')
PCNA	F: GGCGTGAACCTCACCAGTAT R: TCTCGGCATATACGTGCAA
LOX	F: CAGAGGAGAGTGGCTGAAGG R: CCAGGTAGCTGGGGTTTACA
OXTR	F: TTCTTCGTGCAGATGTGGAG R: GGACGAGTTGCTCTTTTTGC
BMP4	F: AAGCGTAGCCCTAAGCATCA R: TGGTTGAGTTGAGGTGGTCA
TPM1	F: CTCTCAAAGATGCCCAGGAG R: TCTCATCTGCTGCCTTCTCA
TNNC1	F: ACCTCTCCGCATGTTTGAC R: CCACACCCTTCATGAACTCC
TTN	F: GGCATCCCCAAACCTAAAAT R: TTTTGCCACTGCTGATTCTG
TGFBR1	F: TGTTGGTACCCAAGGAAAGC R: CACTCTGTGGTTTGGAGCAA
SMARCD3	F: CTCTGAAGAGGCCCATGAAG R: GAACTTCCGCTTCTGTTTGC
NDST1	F: TCACCTTCAACCTGGGCTAC R: ACGGACTGGTTGTGGAAAAG
GAPDH	F: TCAGCCGCATCTTCTTTTGC R: ACGACCAAATCCGTTGACTC
ACTB	F: AAAGACCTGTACGCCAACAC R: CTCAGGAGGAGCAATGATCTTG
HPRT	F: TGGCGTTCGTGATTAGTGATG R: ACATCTCGAGCAAGACGTT

F, forward; R, reverse; PCNA, proliferating cell nuclear antigen; LOX, Lysyl Oxidase; OXTR, oxytocin receptor; BMP4, bone morphogenetic protein 4; TPM1, tropomyosin 1; TNNC1, troponin 1; TTN, titin; TGFBR1, transcription growth factor- $\beta$  receptor 1; SMARCD3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily D member 3; NDST1, N-deacetylase and N-sulfotransferase 1; ACTB,  $\beta$ -actin; HPRT, hypoxanthine phosphoribosyltransferase 1.

Real-time PCR was performed using the 7900HT Fast Real-Time PCR System (Applied Biosystems; Thermo Fisher Scientific, Inc.), RT<sup>2</sup> SYBR<sup>®</sup> Green ROX<sup>™</sup> qPCR Master Mix (Qiagen Sciences, Gaithersburg, MD, USA) and sequence-specific primers (Table I). *Glyceraldehyde-3-phosphate dehydrogenase (GADPH)*,  $\beta$ -actin (*ACTB*), *hypoxanthine phosphoribosyltransferase 1 (HRPT1)* were used as reference genes. Gene expression was analyzed using the relative quantification (RQ) method. The q-PCR starters were designed using Primer3Plus software ([primer3plus.com/cgi-bin/dev/primer3plus.cgi](http://primer3plus.com/cgi-bin/dev/primer3plus.cgi)).

**Statistical analysis.** The analysis of the presented results was conducted using Bioconductor ([www.bioconductor.org](http://www.bioconductor.org)) and R programming languages (R version 3.5.1; [www.r-project.org](http://www.r-project.org)). To determine the statistical significance of the analysed genes, moderated t-statistics from the empirical Bayes method were performed. The P-value was corrected for multiple




Figure 1. Heat map representation of differentially expressed genes belonging to the 'heart morphogenesis' and 'heart development' Gene Ontology Biological Process terms. Arbitrary signal intensity acquired from microarray analysis is represented by colours (green indicates higher and red indicates lower expression). Log<sub>2</sub> signal intensity values for any single gene were resized to Row Z-Score scale (from -2, the lowest expression to +2, the highest expression for a single gene).

comparisons using Benjamini and Hochberg's false discovery rate.  $P < 0.05$  was considered to indicate a statistically significant difference. The statistical significance of enriched GO terms and KEGG pathways was performed by DAVID database software (v.6.8; david.ncifcrf.gov). Statistical significance was calculated using Benjamini method. Each GO term and KEGG pathway were considered significantly enriched if they contained at least 5 differentially expressed genes and showed  $P < 0.05$ . Statistical analysis of RT-qPCR results was conducted using Real Statistics Resource Pack add-on for MS Excel 2016 (Microsoft Corporation, Redmond, WA, USA).

## Results

Whole transcriptome profiling by Affymetrix microarray allowed for the analysis of expression of gene transcripts in human ovarian granulosa cells during long-term *in vitro* culture. Using Affymetrix® Human HgU 219 Array we have examined the expression of 22480 transcripts. Genes with a fold change higher than abs (2) and with a corrected P-value lower than 0.05 were considered differentially expressed. This set of genes consisted of 2278 different transcripts.

DAVID (Database for Annotation, Visualization and Integrated Discovery) software was used for extraction of gene ontology biological process terms (GO BP). Up- and down-regulated gene sets were subjected to DAVID search separately, and only gene sets where adj. P-value was lower than 0.05 were selected. The DAVID software analysis showed that differentially expressed genes belong to 582 Gene ontology groups and 45 KEGG pathways. In this study, we focused on 'heart morphogenesis' and 'heart development' GO BP terms. These sets of genes were subjected to hierarchical clusterization procedure and presented as heat-maps (Fig. 1). The gene symbols, fold changes in expression, Entrez gene IDs and corrected P-values of those genes were shown in Table II.

Moreover, genes that formed one particular GO group can also belong to other different GO term categories. Because of that, we explored the gene intersections between selected GO BP terms. The relation between those GO BP terms was presented as circle plot (Fig. 2).

STRING interaction network was generated among differentially expressed genes belonging to each of selected GO BP terms. Using this prediction method yielded a molecular interaction network formed between protein products of studied genes (Fig. 3).

RT-qPCR was conducted to validate the results obtained during microarray analysis. The outcomes were presented and compared in a form of a bar graph (Fig. 4).

As can be seen, the direction of changes in expression was confirmed in all examples. However, the scale of differences in transcript levels varied between both of the methods analysed.

The GCs have undergone extensive morphological modifications through the time of the culture. They changed their shape from star-like structures to broader, fibroblast-like form. These observable changes were presented as photographs, taken under an inverted microscope, using a relief contrast (Fig. 5).

## Discussion

Ovarian follicles are formed during the process of folliculogenesis and oogenesis. The follicle consists of the oocyte, follicular fluid and oocyte-associated cells-granulosa cells (GCs) (10). In physiological conditions, GCs largely influence the growth and development of the ovarian follicle. In addition, they produce gap junctions that allow for their bi-directional communication with the oocyte (11,12). During this dialogue, nutrients and metabolites are delivered to the oocyte (13). In addition, the key role of GCs in physiological conditions is the production of estradiol during follicular growth and the secretion of progesterone after ovulation (14).

Table II. Gene symbols, fold changes in expression, Entrez gene IDs and corrected P-values of studied genes.

Nazwa	foldD7_D1	foldD15_D1	foldD30_D1	adj.P.Val.D7_D1	adj.P.Val.D15_D1	adj.P.Val.D30_D1	Entrez.Gene ID
LOX	53,155,736	80,241,544	102,191,488	0.0008508	0.0005524	0.0004038	4,015
OXTR	38,386,169	2,260,881	54,854,523	0.0006796	0.001012	0.0003993	5,021
NEXN	35,206,319	19,134,152	29,664,545	0.0012665	0.0019093	0.0010376	91,624
CSRP3	27,412,789	1,585,293	1,057,341	0.0191031	0.6877273	0.9672829	8,048
TPMI	17,883,412	16,942,974	18,049,628	0.0004891	0.0004529	0.0003078	7,168
ITGA3	16,927,809	18,842,178	22,470,094	0.0277094	0.0222552	0.016508	3,675
CDK1	13,880,751	18,568,254	12,196,334	0.002233	0.0014905	0.0018488	983
TWIST1	13,410,656	16,061,334	20,741,443	0.0013489	0.0010638	0.000668	7,291
ANKRD1	11,887,373	2,508,097	2,617,912	0.0247916	0.2787596	0.2411828	27,063
MICAL2	1,054,092	9,421,967	12,712,885	0.0068803	0.0072788	0.0042113	9,645
BMP4	10,489,539	17,221,545	21,243,468	0.0204669	0.0100715	0.0070848	652
DKK1	8,954,961	9,635,247	10,055,325	0.0024355	0.0020428	0.0015034	22,943
COL5A1	8,845,619	15,695,652	18,51,511	0.016744	0.0071873	0.005229	1,289
PRICKLE1	7,301,909	4,726,273	9,353,689	0.0106664	0.0202006	0.0056925	144,165
GLI2	6,93,552	7,103,782	14,112,758	0.0191991	0.0163042	0.0056787	2,736
CCNB1	6,641,253	8,124,624	7,003,387	0.0318394	0.0211764	0.0237061	891
STIL	56,288	540,042	4,900,424	0.0141252	0.0134497	0.0143661	6,491
TTN	4,908,741	1,271,044	1,144,088	0.0047259	0.4647411	0.6871576	7,273
NDRG4	4,817,497	7,098,036	3,324,764	0.0375828	0.0171188	0.0672424	65,009
CPE	4,531,794	6,031,618	8,102,031	0.0083776	0.0043779	0.0024281	1,363
CYR61	4,318,043	424,915	4,154,475	0.0011941	0.0011841	0.0009317	3,491
HEG1	4,138,551	4,604,733	4,311,688	0.0083402	0.0059934	0.0059049	57,493
FZD2	3,767,679	3,278,544	13,843,888	0.0057158	0.0072683	0.0006712	2,535
TNNC1	3,349,751	1,217,197	1,066,087	0.0053652	0.4460103	0.8215354	7,134
FHL2	3,073,142	3,898,276	2,849,228	0.0282662	0.0139549	0.0283799	2,274
MEF2A	3,019,241	3,647,594	2,798,678	0.0041095	0.002502	0.0039542	4,205
WHSC1	2,810,461	3,935,288	3,454,839	0.017237	0.0063714	0.0075821	7,468
TMEM65	2,790,505	19,071	3,094,836	0.0273694	0.0922242	0.0162561	157,378
SMARCD3	2,528,987	2,267,934	175,636	0.0106455	0.0135429	0.0390055	6,604
NDS1	2,511,612	1,661,929	2,158,918	0.0456627	0.0897835	0.0622311	3,340
OXCT1	2,382,383	180,451	1,980,136	0.0045742	0.0135215	0.0075821	5,019
PARVA	2,313,442	2,094,473	3,017,466	0.0060141	0.0080188	0.0021427	55,742
PCNA	2,300,166	2,407,186	2,057,527	0.0170402	0.0125668	0.0210382	5,111
CAMK2D	2,259,943	2,712,418	2,455,167	0.0133284	0.0063681	0.0076605	817
SPARC	2,230,562	2,335,873	2,990,594	0.0204267	0.0152481	0.0060567	6,678
KIF3A	219,478	1,850,005	2,377,716	0.0242452	0.04543	0.0142647	11,127
TGFBF1	2,188,996	1,585,418	146,801	0.026412	0.0065456	0.0556766	7,046
PKP2	2,108,275	183,479	2,092,256	0.0277094	0.0459483	0.0231586	5,318
OSR1	2,071,946	4,175,767	2,370,344	0.0060753	0.0009141	0.0029158	130,497




Figure 2. Representation of the mutual association between 'heart morphogenesis' and 'heart development' GO Biological Process terms. The ribbons indicate which gene belongs to which categories. The genes were sorted by logFC from most to least changed gene. FC, fold change; GO, Gene Ontology.

The process of folliculogenesis begins with the change of GCs' shape from flattened to cuboidal through numerous mitotic divisions of cells. A characteristic feature is the production of follicle stimulating hormone receptor (FSHR) by the GCs. In preovulatory follicle, four types of GCs can be distinguished: The outermost layer is the granulosa membrane, the periantral granulosa is located inside the follicle, the intermediate layer forms the *cumulus oophorus*, with *corona radiata* being the final layer. Granulosa cells forming the *corona radiata* directly surround the oocyte (15). The individual types of GCs forming the four layers are functionally different, secreting different molecules and expressing different receptors (16).

Recent studies indicate that there is a pool of stem cells in mature mouse and human ovaries that support oogenesis and folliculogenesis. Thanks to these stem cells, the primary follicular resource can be produced during the reproductive period. Previously, it was thought that the number of primary follicles is determined and established shortly before the birth of the female (17,18). In recent years, however, the mentioned stem cells have become the major focus of scientists. It is proven that every tissue or organ (including the ovary) contains a population of stem cells capable of differentiating (19-22).

Many authors point to new possibilities of using GCs in regenerative medicine and broadly understood transplantology. GC containing follicular fluid, transferred for utilization during routine *in vitro* fertilization procedures, may be the source of stem cells. Several authors clearly indicate these previously unnoticed properties of GCs. Many articles follow a different lead, suggesting that GCs are subject to the transdifferentiation process. Transdifferentiation is the transformation of already differentiated cells into another cell type, which consists of two stages: cells dedifferentiation, followed by the natural developmental path towards a specific cell lineage. During this process, the mature cell is reprogrammed and changes into another cell type under the influence of the appropriate factors (23-25). Kossowska-To maszczuk *et al* (3,4) conducted experiments in which they indicated that GCs have stem-like properties. Luteinized GC subpopulations express the *OCT-4* gene, escalation of which is characteristic for mesenchymal stem cells. In addition, the research team carried out the differentiation of GCs cells towards osteoblasts, neuroblasts and chondroblasts. They succeeded to obtain populations of differentiated cells (3,4). The results of our research indicate that GCs have




Figure 3. STRING-generated interaction network among differentially expressed genes belonging to the 'heart morphogenesis' and 'heart development' Gene Ontology Biological Process terms. The intensity of the edges reflects the strength of the interaction score.

stem cell properties and can differentiate, in the long-term primary *in vitro* culture, towards e.g.: osteoblasts or muscle cells (unpublished results). The results of our research have been confirmed by the literature. Oki *et al* (26) stated that pig-derived GCs may transdifferentiate towards osteoblasts both *in vitro* and *in vivo*. They conducted a microarray analysis clearly indicating that GCs show the GC-specific marker genes, together with osteogenic markers, during transdifferentiation (26). Brevini *et al* (27) proved that GCs, under the influence of external factors (Vascular endothelial growth factor-VEGF and 5-azacytidine-5-aza-CR), can differentiate towards muscle cells. Their research confirms the change in the phenotype of GCs. In addition, the differentiated GCs expressed the genes characteristic for muscle cells (including Desmin and Myosin heavy chain).

Following up on Brevini *et al* (27), we have analyzed the results of microarrays in terms of the possibility of GC differentiation into cardiomyocytes. It turned out that, in long-term

*in vitro* culture, GCs express some genes involved in the process of heart morphogenesis and development.

In our research, we identified 39 genes responsible for heart development and morphogenesis processes. All of these genes are presented on heatmaps (Fig. 1). Among all genes, 22 are involved in both: Heart development and heart morphogenesis ontology groups. The remaining 17 genes are characteristic for the heart development process. The expression of these genes could confirm the theory that GCs can differentiate into cells proprietary to the heart tissue, during *in vitro* cell culture.

It is known that the heart is formed from the mesoderm that arises during the process of embryo gastrulation. In humans, the mesoderm begins to form at the third week of embryo development (28). At this stage, the mesodermal cells are still precardiocytes. Mesodermal cells differentiate into cardiac cells in response to endodermal induction signals—primarily bone morphogenetic proteins (BMPs) (29). Five transcription factors that play a primary role in cardiac development




Figure 4. RT-qPCR validation of microarray results, presented in a form of a bar graph. FC was presented in its logarithmic form to provide clear comparability of the results. PCNA, Proliferating cell nuclear antigen; LOX, Lysyl Oxidase; OXTR, oxytocin receptor; BMP4, Bone morphogenetic protein 4; TPM1, Tropomyosin 1; TNNC1, Troponin 1; TTN, Titin; TGFBR1, Transcription growth factor-β receptor 1; SMARCD3, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily D member 3; NDST1, N-deacetylase and N-sulfotransferase 1; RT-qPCR, reverse transcription-quantitative polymerase chain reaction; FC, fold change.




Figure 5. Pictures of human ovarian granulosa cell cultures used for the experiment, presenting changes in their morphology following 24 h, and 7, 15 and 30 days of *in vitro* culture. All of the photographs were taken using an inverted microscope, under relief phase contrast (magnification, x10, 20 and 40, as indicated).



in various organisms have been identified as NKX2.5, Mef2, GATA, Tbx and Hand (30). Precardiac cells are multipotent and differentiate in myocardial, endothelial and smooth muscle cells through a process called progressive lineage restriction (31). The last stage of cardiac development is the differentiation of the atrial and ventricular specific myocytes and heart conduction cells (32). In our studies, we showed the expression of the *Mef2A* gene, and therefore suggest that GCs, in long-term *in vitro* culture, show features of mesenchymal stem cells that can differentiate into heart cells.

In addition, this is confirmed by the number of studies that describe the transition of multipotent mesodermal precursor cells into myoblasts, and then into mature myocytes. The *myocyte-specific enhancer factor 2 (MEF2)* (33) is responsible for the above process. Xu *et al* (34), in their studies, proved that overexpression of *Mef2A* causes cardiomyopathy and hypertension. Liu *et al* (35) demonstrated that *Mef2* becomes active during heart formation in the early stages of embryogenesis and is responsible for the differentiation of atria and ventricles of the heart. Other studies demonstrate the key role of *Mef2A* in maintaining the appropriate number of mitochondria and the proper architecture of the heart's cells. *Mef2a*-deficient mice showed high mortality within one week after birth and marked dilation of the ventricle. These mice have been proven to be susceptible to sudden death (36). The above studies indicate the key role played by *Mef2a* in the formation of the heart. This, together with the findings of our research that shows that GCs can give rise to heart cells, opens up a way for further research into the application of human granulosa in heart regeneration medicine. As it has already been mentioned, inducers of the endoderm, such as BMPs, are an important group of factors involved in the development of the heart. In long-term *in vitro* culture, GCs express the *BMP4* gene. According to the literature, this protein is involved in the development of endocardial epithelium and the correct formation of particular chambers of the heart. Deficiency of *BMP4* in mice causes an abnormal number of cardiac cells, defective valve structure, defects in the septum of the heart, and abnormal heart valve formation (37,38).

Another gene characteristic for both morphogenesis and heart development is *LOX (Lysyl oxidase)*. This gene shows the biggest change in expression in relation to the initial condition during long-term *in vitro* culture. Li *et al* (39) demonstrated that *lysyl oxidase* is present in the smooth muscle cells of the blood vessels of rats. In addition, Kaneda *et al* (40) noticed that *LOX* may be a suppressor of ovarian or stomach tumours. Our research indicates that this gene is expressed in granulosa cells and thus they could possibly differentiate into the blood vessels of the heart and be a suppressor of pathological changes in the ovary.

The gene responsible for the formation of the oxytocin receptor (*oxytocin receptor-OXTR*) showed prominently increased expression. To date, the expression of this gene has been associated with fertility, lactation and reproductive behaviour. Recently, mice research has suggested that *OXTR* plays a large role in the regulation and development of social behaviour (41). In addition, in recent years, the role of the above gene in the formation of the cardiovascular system has been indicated (42).

Another gene characteristic for the development and morphogenesis of the heart is *NEXN (Nexilin F-actin binding protein)* and *Cysteine-rich protein 3 (CSR3)* also known as *Muscle LIM protein (MLP)*. These genes show activity in the fetal and adult heart, as well as skeletal muscle adhesion and migration during embryogenesis, and are responsible for hyperthyroid cardiomyopathy (43-45).

During the long-term *in vitro* cultivation of GCs, it was noted that these cells show expression of cyclin-dependent kinases (CDKs), specifically *CDK1*. These compounds are responsible for the regulation of the cell cycle of cardiomyocytes (46). This serves as further evidence that *in vitro*, GCs may be differentiated into cardiomyocytes. The above genes showed clearly increased expression during long-term *in vitro* GC culture.

Among the genes characteristic for the development and morphogenesis of the heart, none showed a drastic decline during long-term *in vitro* culture. Some of these genes showed only a smaller change in expression. These include *OSRI*, *PKP2*, *TGFBR1*, *KIF3A*.

The *OSRI* gene is expressed in GCs under physiological conditions but also participates in the formation of cardiac septa (47,48). We can assume that even a small change in the expression of this gene may indicate the possibility of GC differentiation. Similar observations were found in the case of the *PKP2* gene (*plakophilin*). Plakophilin is a component of desmosomes and is necessary for the proper functioning and distribution of connexins, a type gap junction protein. It was noted that, during the silencing of the *PKP2* gene in rat cardiac myocytes, there was a reduction in the number of gap junctions between the heart cells. This led to changes within the right ventricle (49). Other studies indicate that connexin is also present in GCs and is necessary for bi-directional communication between the oocyte and granulosa cells (50).

Another gene that seems to be characteristic for the processes related to the development of heart is *TNNC1 (troponin c)*. This gene is responsible for the proper functioning of the muscle fibres of the heart, with all of its known mutations leading to cardiomyopathy (51). Another very important gene, demonstrating that GCs in long-term *in vitro* culture can undergo differentiation/transdifferentiation into cardiac muscle, is *SMARCD3 (actin-dependent regulator of chromatin, subfamily D)*. Lickert *et al* (52) noticed that this gene is expressed in the hearts of early mouse embryos. Deficiency of this gene caused defects in heart morphogenesis, as well as abnormal differentiation of myocardium and skeletal muscles. In humans, the impairment of this gene is associated with the occurrence of many congenital heart defects (52). Other studies show that, in general, mesodermal cells may undergo transdifferentiation towards cardiac myocytes. *SMARCD3*, among others, is deeply involved in this process. It is noted that the presence of this gene can cause the transdifferentiation of mesoderm cells into cardiomyocytes (53). It can, therefore, be suggested that GCs, in long-term *in vitro* culture, show stem-like potential and may undergo differentiation or transdifferentiation towards cardiomyocytes.

It may be concluded that GCs have many characteristics of stem cells in long-term *in vitro* culture. The presented results

suggest that it is possible to differentiate GCs into cells that build the heart and, at the same time, it may become a new possibility in the treatment of heart diseases and regenerative medicine of blood vessels. However, the conclusions were based only on large-scale transcriptomic analysis. While validated by more focused RT-qPCR tests in terms of the direction of changes, the scale of differences in transcript levels varied between the methods. This may be caused by many factors, mainly associated with the huge array of processes occurring from the time of transcription to expression of protein products. Therefore, the conclusions of this study require further analyses on the protein level, followed by practical trials of obtaining a stable, differentiated population of cardiac cells coming from GC cultures, which could later be applied in possible clinical situations.

### Acknowledgements

Not applicable.

### Funding

The present study was supported by the Polish National Science Centre (grant nos. 2014/15/B/NZ7/00999 and UMO-2016/21/B/NZ9/03535) and from Poznan University of Medical Sciences (grant no. 502-01-02227367-08414).

### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### Authors' contributions

WK conducted the experiments, chose the models and wrote the manuscript. MBrą designed the experiments and wrote the manuscript. PC performed data analysis and figure preparation, and wrote the manuscript. ArB conducted the experiments, performed data analysis and prepared the figures. MJN wrote the manuscript and performed data analysis. KO conducted the experiments, revised the medical methodology, and performed material extraction and preparation. MaJ conducted the experiments, and performed data analysis and language corrections. MiJ designed the experiments and performed models analysis. LP designed the study and revised the medical methodology (particularly that of *in vitro* fertilization procedures). AnB designed the study, and revised the methodology and the manuscript. DR designed the study. MTS designed the study, revised the methodology and critically revised the manuscript for important intellectual content. MBru designed the experiments, supervised the project, was involved in the medical procedure design and approved the final draft of the manuscript. MZ revised the methodology, performed data analysis, provided writing assistance and approved the final draft of the manuscript. MN supervised the project, designed the experiments (particularly the primary cell culture procedure), provided editorial supervision and approved the final draft of the manuscript. BK supervised the project, designed the study, revised the methodology, provided

editorial supervision, and wrote the manuscript. All authors read and approved the final manuscript

### Ethics approval and consent to participate

The present study has been approved by Poznan University of Medical Sciences Bioethical Committee with resolution 558/17. All patients provided written informed consent.

### Patient consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### References

- Rybska M, Knap S, Jankowski M, Jeseta M, Bukowska D, Antosik P, Nowicki M, Zabel M, Kempisty B and Jaśkowski JM: Cytoplasmic and nuclear maturation of oocytes in mammals-living in the shadow of cells developmental capability. *Med J Cell Biol* 6: 13-17, 2018.
- Rybska M, Knap S, Jankowski M, Jeseta M, Bukowska D, Antosik P, Nowicki M, Zabel M, Kempisty B and Jaśkowski JM: Characteristic of factors influencing the proper course of folliculogenesis in mammals. *Med J Cell Biol* 6: 33-38, 2018.
- Kossowska-Tomaszczuk K, De Geyter C, De Geyter M, Martin I, Holzgreve W, Scherberich A and Zhang H: The multipotency of luteinizing granulosa cells collected from mature ovarian follicles. *Stem Cells* 27: 210-219, 2009.
- Kossowska-Tomaszczuk K and De Geyter C: Cells with stem cell characteristics in somatic compartments of the ovary. *Biomed Res Int* 2013: 310859, 2013.
- Kranc W, Brązert M, Ożęgowska K, Nawrocki MJ, Budna J, Celichowski P, Dyszkiewicz-Konwińska M, Jankowski M, Jeseta M, Pawelczyk L, *et al*: Expression profile of genes regulating steroid biosynthesis and metabolism in human ovarian granulosa cells-a primary culture approach. *Int J Mol Sci* 18: pii: E2673, 2017.
- Chomczynski P and Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156-159, 1987.
- Huang DW, Sherman BT, Tan Q, Kir J, Liu D, Bryant D, Guo Y, Stephens R, Baseler MW, Lane HC and Lempicki RA: DAVID bioinformatics resources: Expanded annotation database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Res* 35: W169-W175, 2007.
- Walter W, Sánchez-Cabo F and Ricote M: GPlot: An R package for visually combining expression data with functional analysis. *Bioinformatics* 31: 2912-2914, 2015.
- von Mering C, Jensen LJ, Snel B, Hooper SD, Krupp M, Foglierini M, Jouffre N, Huynen MA and Bork P: STRING: Known and predicted protein-protein associations, integrated and transferred across organisms. *Nucleic Acids Res* 33: D433-D437, 2004.
- Kranc W, Budna J, Kahan R, Chachuła A, Bryja A, Ciesiółka S, Borys S, Antosik MP, Bukowska D, Brussov KP, *et al*: Molecular basis of growth, proliferation, and differentiation of mammalian follicular granulosa cells. *J Biol Regul Homeost Agents* 31: 1-8, 2017.
- Mora JM, Fenwick MA, Castle L, Baithun M, Ryder TA, Moberley M, Carzaniga R, Franks S and Hardy K: Characterization and significance of adhesion and junction-related proteins in mouse ovarian follicles. *Biol Reprod* 86: 153, 1-14, 2012.
- Kempisty B, Ziółkowska A, Piotrowska H, Zawierucha P, Antosik P, Bukowska D, Ciesiółka S, Jaśkowski JM, Brussov KP, Nowicki M and Zabel M: Real-time proliferation of porcine cumulus cells is related to the protein levels and cellular distribution of Cdk4 and Cx43. *Theriogenology* 80: 411-420, 2013.
- Su YQ, Sugiura K and Eppig JJ: Mouse oocyte control of granulosa cell development and function: Paracrine regulation of cumulus cell metabolism. *Semin Reprod Med* 27: 32-42, 2009.

14. Jeppesen JV, Kristensen SG, Nielsen ME, Humaidan P, Dal Canto M, Fadini R, Schmidt KT, Ernst E and Yding Andersen C: LH-receptor gene expression in human granulosa and cumulus cells from antral and preovulatory follicles. *J Clin Endocrinol Metab* 97: E1524-E1531, 2012.
15. Findlay JK, Kerr JB, Britt K, Liew SH, Simpson ER, Rosairo D and Drummond AE: Ovarian physiology: Follicle development, oocyte and hormone relationships. *Anim Reprod* 6: 16-19, 2009.
16. Nguyen T, Lee S, Hatzirodos N, Hummizsch K, Sullivan TR, Rodgers RJ and Irving-Rodgers HF: Spatial differences within the membrana granulosa in the expression of focimatrix and steroidogenic capacity. *Mol Cell Endocrinol* 363: 62-73, 2012.
17. Guignon CJ and Magre S: Contribution of germ cells to the differentiation and maturation of the ovary: Insights from models of germ cell depletion. *Biol Reprod* 74: 450-458, 2006.
18. Johnson J, Canning J, Kaneko T, Pru JK and Tilly JL: Germline stem cells and follicular renewal in the postnatal mammalian ovary. *Nature* 428: 145-150, 2004.
19. Parte S, Bhartiya D, Telang J, Daithankar V, Salvi V, Zaveri K and Hinduja I: Detection, characterization, and spontaneous differentiation in vitro of very small embryonic-like putative stem cells in adult mammalian ovary. *Stem Cells Dev* 20: 1451-1464, 2011.
20. Virant-Klun I, Rozman P, Cvjeticanin B, Vrtacnik-Bokal E, Novakovic S, Rulicke T, Dovc P and Meden-Vrtovec H: Parthenogenetic embryo-like structures in the human ovarian surface epithelium cell culture in postmenopausal women with no naturally present follicles and oocytes. *Stem Cells Dev* 18: 137-149, 2009.
21. Virant-Klun I, Zech N, Rozman P, Vogler A, Cvjeticanin B, Klemenc P, Malicev E and Meden-Vrtovec H: Putative stem cells with an embryonic character isolated from the ovarian surface epithelium of women with no naturally present follicles and oocytes. *Differentiation* 76: 843-856, 2008.
22. Virant-Klun I: Postnatal oogenesis in humans: A review of recent findings. *Stem Cells Cloning* 8: 49-60, 2015.
23. Buganim Y and Jaenisch R: Transdifferentiation by defined factors as a powerful research tool to address basic biological questions. *Cell Cycle* 11: 4485-4486, 2012.
24. Shen CN, Burke ZD and Tosh D: Transdifferentiation, metaplasia and tissue regeneration. *Organogenesis* 1: 36-44, 2004.
25. Dzafic E, Stimpfel M and Virant-Klun I: Plasticity of granulosa cells: On the crossroad of stemness and transdifferentiation potential. *J Assist Reprod Genet* 30: 1255-1261, 2013.
26. Oki Y, Ono H, Motohashi T, Sugiura N, Nobusue H and Kano K: Dedifferentiated follicular granulosa cells derived from pig ovary can transdifferentiate into osteoblasts. *Biochem J* 447: 239-248, 2012.
27. Brevini TAL, Pennarossa G, Rahman MM, Paffoni A, Antonini S, Ragni G, deEguileor M, Tettamanti G and Gandolfi F: Morphological and molecular changes of human granulosa cells exposed to 5-azacytidine and addressed toward muscular differentiation. *Stem Cell Rev* 10: 633-642, 2014.
28. Moorman A, Webb S, Brown NA, Lamers W and Anderson RH: Development of the heart: (1) formation of the cardiac chambers and arterial trunks. *Heart* 89: 806-814, 2003.
29. Meilhac SM, Lescroart F, Blanpain C and Buckingham ME: Cardiac cell lineages that form the heart. *Cold Spring Harb Perspect Med* 4: a013888, 2014.
30. Olson EN: Gene regulatory networks in the evolution and development of the heart. *Science* 313: 1922-1927, 2006.
31. Epstein JA: Franklin H. Epstein Lecture. Cardiac development and implications for heart disease. *N Engl J Med* 363: 1638-1647, 2010.
32. Wu SM, Fujiwara Y, Cibulsky SM, Clapham DE, Lien CL, Schultheiss TM and Orkin SH: Developmental origin of a bipotential myocardial and smooth muscle cell precursor in the mammalian heart. *Cell* 127: 1137-1150, 2006.
33. Breitbart RE, Liang CS, Smoot LB, Laheru DA, Mahdavi V and Nadal-Ginard B: A fourth human MEF2 transcription factor, hMEF2D, is an early marker of the myogenic lineage. *Development* 118: 1095-1106, 1993.
34. Xu J, Gong NL, Bodi I, Aronow BJ, Backx PH and Molkenin JD: Myocyte enhancer factors 2A and 2C induce dilated cardiomyopathy in transgenic mice. *J Biol Chem* 281: 9152-9162, 2006.
35. Liu N, Williams AH, Kim Y, McAnally J, Bezprozvannaya S, Sutherland LB, Richardson JA, Bassel-Duby R and Olson EN: An intragenic MEF2-dependent enhancer directs muscle-specific expression of microRNAs 1 and 133. *Proc Natl Acad Sci USA* 104: 20844-20849, 2007.
36. Naya FJ, Black BL, Wu H, Bassel-Duby R, Richardson JA, Hill JA and Olson EN: Mitochondrial deficiency and cardiac sudden death in mice lacking the MEF2A transcription factor. *Nat Med* 8: 1303-1309, 2002.
37. McCulley DJ, Kang JO, Martin JF and Black BL: BMP4 is required in the anterior heart field and its derivatives for endocardial cushion remodeling, outflow tract septation, and semilunar valve development. *Dev Dyn* 237: 3200-3209, 2008.
38. Abdelwahid E, Rice D, Pelliniemi LJ and Jokinen E: Overlapping and differential localization of Bmp-2, Bmp-4, Msx-2 and apoptosis in the endocardial cushion and adjacent tissues of the developing mouse heart. *Cell Tissue Res* 305: 67-78, 2001.
39. Li W, Nellaiappan K, Strassmaier T, Graham L, Thomas KM and Kagan HM: Localization and activity of lysyl oxidase within nuclei of fibrogenic cells. *Proc Natl Acad Sci USA* 94: 12817-12822, 1997.
40. Kaneda A, Wakazono K, Tsukamoto T, Watanabe N, Yagi Y, Tatematsu M, Kaminishi M, Sugimura T and Ushijima T: Lysyl oxidase is a tumor suppressor gene inactivated by methylation and loss of heterozygosity in human gastric cancers. *Cancer Res* 64: 6410-6415, 2004.
41. Takayanagi Y, Yoshida M, Bielsky IF, Ross HE, Kawamata M, Onaka T, Yanagisawa T, Kimura T, Matzuk MM, Young LJ and Nishimori K: Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient mice. *Proc Natl Acad Sci USA* 102: 16096-16101, 2005.
42. Gutkowska J and Jankowski M: Oxytocin revisited: Its role in cardiovascular regulation. *J Neuroendocrinol* 24: 599-608, 2012.
43. Hassel D, Dahme T, Erdmann J, Meder B, Hugel A, Stoll M, Just S, Hess A, Ehlermann P, Weichenhan D, *et al*: Nexilin mutations destabilize cardiac Z-disks and lead to dilated cardiomyopathy. *Nat Med* 15: 1281-1288, 2009.
44. Zhao Y, Wei YJ, Cao HQ and Ding JF: Molecular cloning of NELIN, a putative human cytoskeleton regulation gene. *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao (Shanghai)* 33: 19-24, 2001.
45. Geier C, Gehmlich K, Ehler E, Hassfeld S, Perrot A, Hayess K, Cardim N, Wenzel K, Erdmann B, Krackhardt F, *et al*: Beyond the sarcomere: CSRP3 mutations cause hypertrophic cardiomyopathy. *Hum Mol Genet* 17: 2753-2765, 2008.
46. Ikenishi A, Okayama H, Iwamoto N, Yoshitome S, Tane S, Nakamura K, Obayashi T, Hayashi T and Takeuchi T: Cell cycle regulation in mouse heart during embryonic and postnatal stages. *Dev Growth Differ* 54: 731-738, 2012.
47. Lan CW, Chen MJ, Jan PS, Chen HF and Ho HN: Differentiation of human embryonic stem cells into functional ovarian granulosa-like cells. *J Clin Endocrinol Metab* 98: 3713-3723, 2013.
48. Zhou L, Liu J, Olson P, Zhang K, Wynne J and Xie L: Tbx5 and Ors1 interact to regulate posterior second heart field cell cycle progression for cardiac septation. *J Mol Cell Cardiol* 85: 1-12, 2015.
49. Oxford EM, Musa H, Maass K, Coombs W, Taffet SM and Delmar M: Connexin43 remodeling caused by inhibition of plakophilin-2 expression in cardiac cells. *Circ Res* 101: 703-711, 2007.
50. Kempisty B, Ziólkowska A, Ciesiółka S, Piotrowska H, Antosik P, Bukowska D, Nowicki M, Brüssow KP and Zabel M: Study on connexin gene and protein expression and cellular distribution in relation to real-time proliferation of porcine granulosa cells. *J Biol Regul Homeost Agents* 28: 625-635, 2014.
51. Parvatiyar MS, Landstrom AP, Figueiredo-Freitas C, Potter JD, Ackerman MJ and Pinto JR: A mutation in TNNC1-encoded cardiac troponin C, TNNC1-A31S, predisposes to hypertrophic cardiomyopathy and ventricular fibrillation. *J Biol Chem* 287: 31845-31855, 2012.
52. Lickert H, Takeuchi JK, Von Both I, Walls JR, McAuliffe F, Adamson SL, Henkelman RM, Wrana JL, Rossant J and Bruneau BG: Baf60c is essential for function of BAF chromatin remodelling complexes in heart development. *Nature* 432: 107-112, 2004.
53. Takeuchi JK and Bruneau BG: Directed transdifferentiation of mouse mesoderm to heart tissue by defined factors. *Nature* 459: 708-711, 2009.

