

Chapter 11

Cell Defence and Survival

Abstract Central to immune defence mechanisms is the role of transcription factor nuclear factor kappa B (NF- κ B). This is a complex biochemical topic with ever more controls revealed. NF- κ B determines the production of proinflammatory cytokines and chemokines. Pharmacologists step in with possible means of control. Other systems involved in defence include the cyclooxygenase 2 (Cox-2) enzyme and peroxisome proliferator-activated receptors. Insulin receptor activation needs to be seen in context. The mTOR system directs uptake of nutrients by cells. mTOR is suppressed by rapamycin, whose usage is now quite considerable in the control of transplant rejection.

11.1 Transcription Factor NF-kappa

Nuclear factor kappa B (NF- κ B) was identified in 1986 as a transcription factor (TF) that binds to the enhancer of the B cell immunoglobulin- κ light chain gene. NF- κ B proteins are cell homologues of viral v-Rel. NF- κ B functions as homo- and heterodimers (p50/p65) (Fig. 11.1a) that share a 300-amino acid DNA binding domain at the amino terminal called the Rel homology domain (RHD) (Fig. 11.1b), which allows nuclear localization and dimerization [1]. These units are held in the cell cytoplasm bound to an inhibitory unit I-kappa B (I κ B). The I κ B proteins have ankyrin repeats that attach to the RHDs. When the I κ B units are phosphorylated on two serine residues, they are ubiquitinated and degraded by the 26S proteasome. Thus, the I κ Bs are removed, and the RelA (p65) RHDs then move rapidly into nuclei to cause activation of target genes [2–4]. In the nuclei, the dimeric Rel homology domains bind to a decameric DNA sequence 5'GGGRNYYCC-3', where R indicates A or G, Y is C or T, and N is any base. The ability of the dimers to recognise slightly differing DNA targets means that a whole range of genes is regulated differentially.

Accordingly NF- κ B is a family of TFs that play a crucial role in the immune and inflammatory responses of cells and in cell survival [5–7]. Various stimuli activate NF- κ B [8, 9]: (a) proinflammatory cytokines interleukin 1 (IL-1) and tumour necrosis factor- α (TNF α), (b) oxidants like H₂O₂, (c) ceramide, (d) protein kinase




Fig. 11.1a Activation of nuclear factor kappa B (*NF-κB*) by inflammatory stimuli. Microbial products or cytokines that induce *NF-κB* trigger phosphorylation of unit I-kappa B (*IκB*) kinase (*IKK*) -α and -β. The heterodimers *IKK*α or -β then bind to the regulatory *IKK*γ, Nemo, and each heterodimer [1] phosphorylates the *IκB* inhibitory proteins, so that they undergo proteasomal degradation. *NF-κB* p50, p65 is then freed and will move into the nuclei. With regard to the role of ubiquitination [2, 3], it regulates three steps: (a) degradation of the inhibitor *IκB*, (b) processing of the *NF-κB* precursors and (c) *Traf* 6 ubiquitination, which activates the *IKK*. Proteasomes are implicated in *NF-κB* activation: (a) in generation of p50 from the precursor p105 and (b) in the degradation of *IκB*α. Types I and II interferons induce immunoproteasomes in which the reactions are much quicker. Calpain activates *NF-κB* since it degrades the PEST (proline, glutamate, serine, threonine) sequence in the inhibitor *IκB*α. **b** Structures of nuclear factor kappa B (*NF-κB*) family members. All five *NF-κB* members are related through the conserved Rel homology domain (*RHD*), which contains sequences for DNA binding, dimerization, inhibitor association and nuclear localization. The members c-Rel, RelA and RelB contain a C-terminal transactivation domain (*TAD*). The function of the *TAD*s is enhanced by phosphorylation and acetylation. The second group (the p105/p50 and p100/p52) arises as precursors p100 and p105 that are partially proteolysed by the 26S proteasome to yield the mature p52 and p50 units. They contain C-terminal sequences with inhibitory ankyrin (*ANK*) repeats, and this means that they can also function as unit I-kappa B (*IκB*s) and retain Rel proteins in the cytoplasm. The five to seven ankyrin repeats of inhibitory *IκB* proteins interact with *NF-κB* dimers and mask their nuclear localization signals (*NLS*s)

B (Akt), (e) protein kinase C (PKC) [10] and (f) viruses. Intracellular oxidants [11, 12] help the I κ B inhibitors to be dissociated from the NF- κ B units. Hence, intracellular reductants like glutathione and thioredoxin have differential roles in the redox regulation of target proteins, and S-nitrosylation can inhibit NF- κ B DNA binding [13]. Yet, within nuclei it is the more reducing environment there that enables NF- κ B RHDs to bind onto the DNA.

As you will now expect, zymosan proinflammatory agent stimulates NF- κ B, causing release of IL-8 so that neutrophils are attracted [14]. That process can be inhibited by nitric oxide (NO) or by IL-10 inhibition of NF- κ B [15]. Examination of synovial tissue in rheumatoid arthritis shows marked expression of NF- κ B1 at sites adjacent to the cartilage-pannus junction. NF- κ B activation upregulates chemokines and adhesion molecules and neutrophils and macrophages are then recruited.

Extra comment about Nemo (I κ B kinase- γ , IKK γ) is appropriate as this is a developing field [16]. Nemo has two coiled domains, CC1 and CC2, and a leucine zipper. Nemo interacts with many proteins involved in NF- κ B activation. It is polyubiquitinated but on differing lysine residues according to the stimulus [17]. Its potential is also influenced by sumoylation and phosphorylation.

11.1.1 The Classical Canonical NF- κ B Pathway

The IL-1 β and TNF α cytokine receptors connect via upstream kinases to the activation of NF- κ B [17, 18]. The two IKK kinases α and β have an activation loop like that of the mitogen-activated protein kinase kinase (MAPKK) proteins, which bears two serines that are phosphorylated in response to IL-1 or TNF α . Cytokine IL-1 can activate either IKK β or the IKK α .

As shown in Fig. 11.1a, serine-threonine kinases IKK α and IKK β form homo- or heterodimers in conjunction with stabilising protein Nemo (the NF- κ B essential modulator), also termed IKK γ [4, 19]. Binding of Nemo to Lys63-polyUb-RIP1 forms the active kinase complex IKK γ , IKK β -P, IKK α -P that phosphorylates the inhibitor unit I κ B α [20], which then dissociates and is destroyed by ubiquitination. Accordingly, NF- κ B as p50/RelA is free to enter nuclei. Oscillations in intranuclear NF- κ B coupled to cycles of phosphorylation of RelA and I κ B are required for NF- κ B-dependent gene expression [21]. As revealed in Table 11.1, NF- κ B activates a range of genes relevant to inflammation and innate immunity. There are genes for the cytokines and chemokines and for regulation of the cell cycle, survival, adhesion and invasiveness. Both apoptotic and antiapoptotic genes are induced. Genes with a role in feedback regulation of the NF- κ B, such as I κ B α and A20, are induced early. *What does A20 protein do?* It inhibits the IKK complex, it inhibits RIP (Fig. 12.5), and it inhibits TAK1 kinase. By inhibiting RIP, A20 stops TNF α -induced NF- κ B activation, a step that requires Richard TAXIBI [21b].

NF- κ B is a key participant in multiple processes affecting the immune system and inflammation. It can interact with other TFs like Interferon Regulatory Factors (IRFs), signal transducers and activators of transcription (STATs) and the p53. Such promiscuity might explain the many roles of NF- κ B in cell functions and fate [22].

Table 11.1 Common genes activated by nuclear factor kappa B (NF- κ B)

Class	Target gene
Immunoreceptors	Immunoglobulin κ light chain
	Interleukin 2 receptor α -chain
	Major histocompatibility complex class I/II
	Platelet-activating factor receptor
	Tissue factor
Cell adhesion molecules (CAMs)	Endothelial leucocyte adhesion molecule 1 (ELAM-1)
	Vascular cell adhesion molecule 1 (VCAM1)
	Intercellular cell adhesion molecule (ICAM1)
Cytokines and growth factors	Tumour necrosis factor- α (TNF α)
	Interleukin 1 β (IL-1 β), IL-2, IL-6, IL-8, IL-12
	Transforming growth factor- β
	Granulocyte colony-stimulating factor (G-CSF)
Chemokines	Monocyte chemotactic peptide 1 (MCP-1)
	Macrophage inflammatory proteins (MIPs) 1 α , 1 β
Acute-phase proteins	C-reactive protein (CRP)
	Lipopolysaccharide-binding protein (LBP)
Others	Nitric oxide synthases (NOSs)
	Cyclooxygenase 2 (Cox-2)
	Phospholipase A2 (PLA2)
	Metalloproteinases

Hydrogen peroxide (from ROS) prolongs nuclear localization of NF- κ B in activated cells [22b]. Fortunately, there is control by heat shock proteins HSP70 and HSP90 which regulate the IKK complex [22c].

NF- κ B plays a crucial role in inflammatory processes. Activation of p38 mitogen-activated protein kinase (MAP) kinase is well recognised to activate NF- κ B [23]. If a single injection of lipopolysaccharide (LPS) is given intraperitoneally to mice, there is transient NF- κ B activation in lung cell types. When the LPS is infused by a pump, there is widespread NF- κ B activation and neutrophilic infiltration that begets lung injury following release of cytokine IL-8 ([MIP2] of mice) (Table 11.1). Since the search is on for means of controlling ARDS (adult respiratory distress syndrome), an inhibitor of NF- κ B could be found to reduce the lung inflammation. As for natural restraints PIAS [24] or MK2 [25] or Cezzare [22b] or CYLD deubiquitinating enzyme [26] block NF- κ B activation. In resting neutrophils, RhoA GTPase inhibits the NF- κ B to prevent TNF α production, and yet in LPS-stimulated neutrophils RhoA activates NF- κ B and supports TNF α formation [27].

11.1.1.1 The Roles of Subunits: subtle feedback control mechanisms

The various homodimers and heterodimers of NF- κ B are kept inactive by association with cytoplasmic inhibitory proteins [28]. They are I κ B α , I κ B β and I κ B ϵ and also the p105 and p100 precursors of p50 and p52. Are there roles for the α -, β -, γ -subunits of the IKK complex? [29]. IKK α is a histone H3 kinase that moves into nuclei to associate with the promoters of NF- κ B-responsive genes [30, 31]. IKK α

contributes to NF- κ B DNA binding on gene promoters; thus, it derepresses IL-8, cIAP2 and more. Yet, it creates a negative feed-back loop for NF- κ B and macrophage activation [32, 32b, 33]. In fact, IKK α resolves inflammation by switching off the canonical pathway. By accelerating turnover of cRel and RelA, IKK α attenuates IKK β driven NF- κ B activation and so inflammation. It can also promote adaptive immunity via the non-canonical pathway [33].

When Nemo/IKK γ shuttles into nuclei, it creates feedback by repression of NF- κ B gene expression. Nemo binds to other proteins like RIP and A20, and it regulates the IKK-related kinases IKK ϵ and TBK1 (TANK binding kinase 1), which regulate interferon regulatory factors IRF3 and IRF7.

As for IKK β , it can increase the activity of hypoxia-inducible factor 2 α (HIF2 α) [33b], so linking hypoxia to innate immunity and inflammation, and it activates a subset of (IFN γ)-stimulated genes. For sure IKK β promotes the canonical pathway of NF- κ B activation (Fig. 11.1a) [29]. IKK β promotes the MAP kinases and cell proliferation (e.g. in skin keratinocytes) [29]. All the actions are tissue specific. Yet, IKK β exerts negative control over the production of IL-1 β [34]. Furthermore in some situations IKK β can suppress M1 host defence macrophages, so that it is anti-inflammatory [34b], but it promotes M2 immune regulation macrophages [34c].

11.1.2 *The Alternative Noncanonical Pathway*

There is an alternative means of NF- κ B activation [35]. Lymphotoxin, BAFF (B cell-activating factor), CD40L or RANKL (receptor activator of NF- κ B ligand) activate NIK (NF- κ B-inducing kinase), and it then acts on IKK α homodimer, whose target is NF- κ B2, p100. Phosphorylation-dependent ubiquitination of the p100 is only partial. The N-terminal p52 is released and joins with RelB to enter the nuclei (Fig. 11.2). This alternative path via p52/RelB enables BAFF (BLys) to activate B lymphocytes. The alternative path essentially provides organogenic chemokines [4] for the development of lymphoid tissue [36] and adaptive immunity. The NF- κ B2 is also required for establishment of thymic central tolerance through an Aire-dependent pathway [37]. AIRE is the autoimmune regulator gene, and NF- κ B2 is required in medullary thymic epithelial cells.

Whenever Rankl stimulates osteoclastogenesis, the NF- κ B2 p100 path is implicated [38]. Acting on receptor rank on bone stromal cells, Rankl stimulates their NFATc1 and c-fos so that osteoclasts are formed. This activation is reinforced by Ca²⁺-calmodulin (CaM) stimulation of NFATc and of CaM kinase IV, which stimulates cAMP response element-binding protein (CREB) on the promoter for osteoclast marker genes. NIK has been shown by the study of mouse knockouts to control the activities of lymphocytes and osteoclasts in inflammatory arthritis [39]. NIK^{-/-} mice have no peripheral lymph nodes, defective T and B cells and impaired Rankl-induced osteoclastogenesis. One might also be interested that when ultraviolet light acts on skin to cause immunosuppression by induction of Treg cells, Rankl on keratinocytes first stimulates Rank on Langerhans cells. Vitamin D₃, which is immunosuppressive, also induces Rankl [40].




Fig. 11.2 Comparison of the canonical and alternative paths of nuclear factor kappa B (*NF- κ B*) activation. Nemo also connects IKK complexes with IKK ϵ and TBK1 kinases. Monarch-1 suppresses the noncanonical NF- κ B2 activation by causing proteasomal degradation of NF- κ B-inducing kinase (*NIK*). In T lymphocytes, the NF- κ B2/p100 is a late-acting negative-feedback molecule affecting T cell receptor activation of the NF- κ B [41]

11.1.3 More about NF- κ B Control

All NF- κ B activation pathways share the proteasome-mediated step (Fig. 11.1) that causes the degradation of the inhibitory proteins. Posttranslational modifications like phosphorylation [42], acetylation [43] or prolyl isomerization modulate the activity of the p65 (or p50) subunits. Thus, acetylation of multiple lysine residues by p300/CBF acetyltransferases (Fig. 4.8) modifies the behaviour of NF- κ B components. Acetylation works as an intranuclear molecular switch [44]. Acetylation of p65 stops inhibition by I κ B α and controls nuclear NF- κ B. Oxidative stress, via its formation of H₂O₂, which enhances degradation of I κ B α , enhances NF- κ B activity [22b]; thus, macrophages have increased production of cyclooxygenase 2 (Cox-2), prostaglandin E2 (PGE2), and matrix metalloproteinase 1 (MMP-1) [45].

Activation of NF- κ B-inducible genes (Table 11.1) usually takes place by means of the p65/p50 DNA binding heterodimers of the classical pathway. Significant fluctuation in p50 levels can alter the abundance and composition of NF- κ B complexes within a cell. When animals or cells are exposed to repeated small doses of LPS, the output of cytokines via NF- κ B activation is thwarted, and the animals or cells show *endotoxin tolerance*. The explanation is that p50/p50 homodimers are formed instead of the p65/p50, and the consequence is suppression of NF- κ B-inducible genes [46]. p65 has a transactivation domain, whereas p50 does not. p50/p50 homodimers (as produced by IL-10 or by protein kinase A) are inhibitory. Cells that are tolerant to endotoxin still respond to LPS with the degradation of I κ B and nuclear translocation of p50 and p65. Yet, they cannot form IL-1 β . In such cells, it was found that RelB

represses inflammatory gene expression [47]. Alternatively, in endotoxin tolerance, mobilization of Toll-like receptor 4 (TLR4) into lipid rafts is curtailed [48].

p53 turns out to be a buffer of the innate immune system because p53 works as an antagonist of NF- κ B to dampen chronic inflammation. p53^{null} macrophages show very high release of cytokines IL-1, IL-6 and IL-12 in response to LPS or IFN γ . In fact, p53 downregulates Stat1 and proinflammatory cytokines [49]. In some cells, like leucocytes, I κ B ζ negatively regulates intranuclear NF- κ B activity [50]. We have noted how LPS exposure can attenuate TNF α gene transcription because NF- κ B p50 homodimers bind to the TNF α promoter. In addition, LPS induces Bcl-3 (B cell lymphoma 3), which is an anti-inflammatory regulator [32b,51]. Bcl-3 is only located in cell nuclei, and it associates with p50/p50 homodimers and acts as a negative regulator of TLR signalling [52]. Bcl-3 reduces transcription of proinflammatory cytokines. It mediates some of the anti-inflammatory activities of IL-10 (Sect. 12.16) Bcl-6 is another transcriptional repressor protein. Repression by Bcl-6 is cell-type specific. Bcl-6 negatively regulates the expression of the NF- κ B1 p105/p50 subunit. In macrophages, it negatively regulates the chemokines monocyte chemotactic peptide 1 (MCP-1), MCP-3 and MIP-1 [53]. Indeed, in Bcl-6^{-/-} mice there is florid T helper lymphocyte Th2-mediated inflammation. Bcl-6 is essential for formation of germinal centres and for a normal antibody response [54]. Bcl-6 suppresses p53 expression in germinal centre B lymphocytes. Bcl-6 interacts with TF Miz-1 to suppress p21 and thus to allow proliferation of germinal centre B cells [55]. Activating mutations or chromosomal translocations that affect Bcl-6 are relevant to B cell lymphomas [56, 57]. The t (3;14) translocation affecting Bcl-6 is part of diffuse large B cell lymphoma (DLBCL).

One should consider the pharmacological means of control of NF- κ B (Table 11.2). Commensal gut bacteria like *Bacteroides* reduce the activity of NF- κ B in gut

Table 11.2 Mediators and drugs that inhibit nuclear factor kappa B (NF- κ B)

Physiological

Interleukins 10 and 13 transforming growth factor- β
 I κ BNS inhibitor, heat shock proteins HSP70 and HSP90, Nemo-binding domain peptide, CYLD
 Cyclic adenosine monophosphate (cAMP), glucocorticoids, oestrogen, melanocortins
 Nitric oxide late in macrophage activation^a S-nitrosylation [9]
 T suppressor lymphocytes (Treg) tumor suppressor p53

Pharmacological

Cyclopentenone prostaglandin 15-deoxy-PGJ2, and the EETs, prostaglandin E2 (PGE2)
 Salicylates, acetylsalicylic acid, sulindac (nonsteroidal anti-inflammatory, NSAID), sulfasalazine, mesalamine
 Antioxidants: vitamin E, N-acetylcysteine, taurine-chloramine, lipoic acid
 Lipoxin A4, pyrrolidinedithiocarbamate, conjugated linoleic acid (CLA)
 PPAR γ agonists like the thiazolidinediones (TZDs) and PPAR α
 Immunosuppressant cyclosporin 1,25(OH)₂ vitamin D₃ alcohol
 ACE inhibitors, statins-HMG-CoA reductase inhibitors
 Flavonoids, silymarin, green tea polyphenols, curcumin, capsaicin, flavopiridol
 Deoxyspergualin, geldanamycin, retinoids, vitamin D analogues
 Proteasome inhibitors: velcade, lactacystine, epoxomicin, peptide aldehydes

^aEarly in macrophages, nitric oxide induces NF- κ B, but later it inhibits it

mucosal epithelial cells. Actually, *Bacteroides* species activate peroxisome proliferator-activated receptor- γ (PPAR γ) receptors, which take RelA out of cell nuclei of the gastrointestinal (GI) epithelium. Hence, cytokine production via NF- κ B is suppressed [58]. Clearly, inflammatory reactions at the gut mucosa can cause much trouble. Hence, note that local GI macrophages show downregulation of their TREM1 (triggering receptors expressed on myeloid cell 1)-activating receptors. TREM would amplify cytokine-induced responses. This is explained by the local environment, in which there is plentiful IL-10 (suppressive cytokine) and transforming growth factor- β (TGF β) [59].

Enhanced NF- κ B activity occurs in the gut mucosa in inflammatory bowel diseases. In Crohn's disease, there is high expression of the immunoproteasome so that there is enhanced processing of the p105 \rightarrow p50 and enhanced degradation of inhibitor I κ B α . Then p50/c-Rel is important for IFN γ induction of proteasomes via IL-12-driven Th1 cell responses [60].

Phosphorylation of cRel is needed for antiviral responses. Accordingly, TBK1 (Tank binding kinase-1) and IKK ϵ phosphorylate the C-terminal of cRel so that p50/cRel can bind at NF- κ B sites [61]. TBK also phosphorylates IRF3. This means that interferons IFN α / β are induced (Chap. 14) [62].

11.2 How to Inhibit NF- κ B

Anti-inflammatory cytokines IL-10 and IL-13 suppress the activation of NF- κ B. Unlike many cytokines, IL-10 does not activate the NF- κ B. Heat shock proteins inhibit NF- κ B by stopping the degradation of I κ B [22c]. There is accumulating information on drugs and antioxidants like N-acetylcysteine [63], resveratrol of red wine or the polyunsaturated fatty acid (PUFA) conjugated lineoleic acid (CLA) [64] that inhibit NF- κ B activation and thus have anti-inflammatory properties.

Those with therapeutic inclinations will be interested in physiological mediators and drugs that can be used to inhibit NF- κ B (Table 11.2). The immunosuppressant cyclosporin stops RelA activation and so reduces IL-2 formation by T lymphocytes [65]. Decoy deoxynucleotides or negative mutants of NF- κ B are used by the biochemist. A recent review of pharmacological aspects of NF- κ B [66] listed inhibitory peptides for NF- κ B that have been used in anti-inflammatory exploration. The cyclopentenone 15-deoxyPGJ2 is anti-inflammatory by virtue of its ability to modulate TFs [67]. It also acts to potentiate nitric oxide (NO), stopping neutrophil ingress.

In this amazing list of diverse agents, note that T regulatory (suppressor) lymphocytes act, in part, by inhibition of NF- κ B. Thus, Li et al. [68] demonstrated how suppressor lymphocytes stop the presentation of CD86 on antigen-presenting cells (APCs). I κ BNS is an inhibitor that binds to nuclear Rel/p65. It accounts for negative selection of thymocytes [69] and T lymphocytes or macrophages [32b].

Dendritic cells (DCs) use NF- κ B for their maturation, and they use NF- κ B to help form Th1 lymphocytes and CD8 cytotoxic T cells. NF- κ B p50-RelA determines the differentiation of DCs and their survival, and their formation of IL-12 depends

on NF- κ B p50-cRel [70]. Not surprisingly, we know of numerous microorganisms that are able to subvert NF- κ B activation to gain entry into the body [71]. Conversely, if there is p38 MAP kinase activation, that will lead to NF- κ B induction of inflammatory genes [72]. NF- κ B supports adaptive immunity and the elimination of microbial invaders.

Glucocorticoids, on that list, are immunosuppressive agents. Their glucocorticoid receptors (GRs) suppress the activity of CREB-binding protein (CBP)/p300 induce corepressors like histone deacetylase 2 (HDAC2) [73]. Similarly, PGE2 is anti-inflammatory by virtue of the fact that it enhances the expression of I κ B α , and ineffectual p50/p50 homodimers form [74]. It turns out that 1,25(OH)₂ vitamin D₃ and 1,24(OH)₂D₂ inhibit TNF α expression by macrophages. What is more, these forms of vitamin D increase the activity of I κ B α ; thus, they inhibit NF- κ B activity [75].

Obviously, using PPAR γ and - α agonists (Fig 4.14) is a way of dealing with inflammation. PPAR α knockout mice show augmented inflammation. In them, there is marked expression of TNF α and IL-1 β and expression of Fas ligand in the course of carrageenan-induced paw oedema or carrageenan-induced pleurisy [76].

11.3 Gene Knockout or Transgenic Experimental Animals

The involvement of effector molecules in physiological or pathological processes can now be elucidated or confirmed by the use of gene knockout (ko) or transgenic (tg) mice, into which an extra gene is inserted. Already, examples have been mentioned. Mouse models that lack almost all the known genes for G protein α -subunits have been generated to give insight into G protein signal pathways. G α q-deficient platelets fail to aggregate in response to thrombin, adenosine diphosphate (ADP) or thromboxane A₂ (TxA₂) due to lack of agonist-induced phospholipase C activation. Mice lacking G α q have increased bleeding times and are protected against thromboembolism. Mice with knockout of gene granulocyte-macrophage colony-stimulating factor (GM-CSF) develop silting of the lungs by “alveolar proteinosis”, which is a clear indication of the cause of this rare condition. Transgenic mice that overexpress phospholipase A₂, which generates leukotrienes and platelet-activating factor (PAF) (Fig. 6.2b), develop exuberant atherosclerosis. They show enhanced oxidation of their plasma low-density lipoprotein (LDL). Mice that lack neutrophil elastase (NE) cannot clear infections with intracellular bacteria [77]. Mice deficient in cathepsin G succumb to *Staphylococcus aureus* [78]. The subtleties of NF- κ B controls are gradually being elucidated by techniques like this (cf. [79]). B lymphocytes with RelA^{-/-} deficiency were killed easily by exposure to TNF α since they had reduced expression of Flice (i.e. caspase 8) and of Bcl-2 antiapoptotic protein. Clearly, NF- κ B enables B cell survival by eliminating the effects of TNF α toxicity. This may remind one that c-Flip, inhibitor of the extrinsic apoptotic pathway, is known to promote NF- κ B signalling and cell survival [80]. RNA interference (RNAi) is a mechanism for sequence-specific posttranscriptional inhibition of gene expression acting on double-stranded RNA molecules. Small interfering RNAs can be applied to mammalian cells. Thus,

Table 11.3 How nuclear factor kappa B (NF- κ B) supports antiapoptosis/cell survival

 NF κ B supports:

Inhibitor of apoptosis proteins 1 and 2 and c-FLIP, which inhibit caspase 8

Protein A1 and Bcl-XL, which stop mitochondrial release of cytochrome C

A20 protein, which thwarts (a) IKK, (b) RIP and (c) TAK1 (Fig. 11.2)

XIAP

 GADD45 β , which inhibits JNK

 FHC, which stops ROS stimulation of JNK

FHC ferritin heavy chain, *GADD45* growth arrest and DNA damage 45, *IKK* unit I-kappa B kinase, *JNK* Jun N-terminal kinase, *ROS* reactive oxygen species, *XIAP* X chromosome inhibitor of apoptosis TAK1 = transforming growth factor β -activated kinase 1.

RNAi was used to show how phospholipase D (PLD-1) colocalises with actin filaments at the adhesive surfaces of macrophages [81]. RNAi has been used to assess the role of p50 NF- κ B1 protein in the maturation of monocyte-derived DCs. siRNA for p50 reduces the transcription of IL-12 p40. Absence of NF- κ B1 p50 or of c-Rel leads to impaired survival of DCs and lack of IL-12 production. IL-12 production in macrophages is blocked when the IFN γ receptor or the TNFR1 are removed by gene deletions in mice. NF- κ B2, p52, derived from precursor p100, when eliminated in knockout mice, results in DCs with enhanced RelB activity. Enhanced RelB is accompanied by enhanced MHC class II molecule expression on APCs and an increased ability to induce CD4 lymphocytes (Table 11.3).

11.4 The Control of Cyclooxygenase 2

Various pathways help to stimulate the expression of Cox-2, such as ROS, tumour promoters, growth factors, oncogenes, and cytokines. Her2/Neu related to the epidermal growth factor receptor (EGFR) does this in breast cancer. Also, in breast and colon cancers there is activation of NFAT (nuclear factor of activated T cells), which leads to Cox-2 induction and so formation of PGE2. Tumours with elevated Cox-2 will be invasive [82]. Indeed, typical metastatic gene products are Cox-2, intercellular cell adhesion molecule 1 (ICAM1), vascular endothelial growth factor (VEGF), and metalloproteinase 9 (see Chap. 18).

Generally, transcription of Cox-2 gene is promoted by PKC, Ras signalling, and especially MAP kinase induction of NF- κ B and C/EBP and factors that stimulate activator protein 1 (AP-1) [83, 84] (Fig. 11.3). This is similar to induction of cytokines. Consider the situation when *Helicobacter pylori* causes chronic gastritis. Study of TFs in gastric epithelial cells showed Cox-2 promoter induction by NF- κ B, AP-1 and C/EBP (NF-IL-6) [85]. In Fig. 11.3, note the canonical TATA motif and several important enhancer elements affected by transactivators. Proinflammatory mediators can activate ribosomal S6 kinase (RSK), which enhances C/EBP β binding to the Cox-2 promoter. Sphingosine-1-P, by stimulation of NF- κ B, also stimulates Cox-2 activity [86].

Cox-2 expression in neutrophils is promoted by ROS induction of NF- κ B in response to cell membrane events [87]. Likewise, when monocytes interact with




Fig 11.3 Transcription factors affecting the cyclooxygenase 2 (*Cox-2*) promoter. (as in airway smooth muscle cells) C/EBP = CCAAT/enhancer binding protein

platelets or P-selectin-coated surfaces, there is activation of NF- κ B and so Cox-2. In a second phase, IL-1 stabilizes the Cox-2 messenger RNA (mRNA), which it does by silencing of the binding protein ARE; thus, there is enhanced translation [88]. Induction of Cox-2 in macrophages is also biphasic and requires C/EBP β and C/EBP δ TFs [89]. Conserved elements in the macrophage Cox-2 gene promoter as revealed by analysis after exposure to LPS are -CRE2/NF- κ B-C/EBP/AP-1/CRE1/E-box-TATA. The E-box mediates transcriptional repression, whilst the other *cis*-elements are activating. At 6 h after LPS administration, Cox-2 gene transcription was completely dependent on phospho-c-Jun along with the p50 of NF- κ B [90]. One can compare this information with the details of the promoter for Cox-2 in airway smooth muscle cells.

With regard to the discussion in Chap. 4 (Sect. 4.3.1), one can expect that there will be histone acetylation on H3/H4 at the Cox-2 promoter and phosphorylation of H3 histones.

There are different patterns of histone H4 acetylation induced by IL-1 β or by bradykinin. Whereas IL-1-induced transcription utilises NF-IL-6, NF- κ B and CRE, bradykinin uses CRE [91]. Induction of Cox-2 in macrophages is complicated. Normally, any response to NF- κ B activation is rapid. Full Cox-2 expression in macrophages is delayed since there first has to be induction of TNF α by IFN γ [92]. In mesangial cells, TGF β is able to stimulate Cox-2 [93]. When mouse macrophages respond to virus or double-stranded RNA (dsRNA), there is selective induction of Jun N-terminal kinase (JNK) and p38 and therefore activation of NF- κ B that activates Cox-2 [94].

As a lesson, consider how the peptidoglycan of bacterial cell walls stimulates TLR2 receptors; hence, there is NF- κ B activation that activates Cox-2, and PGE2 is produced. This PGE2 then acts on EP2/EP4 surface receptors of the macrophages, so that cAMP is produced and protein kinase A is activated, which then accounts for Ser276 phosphorylation on the p65 of NF- κ B. The consequence is that peptidoglycan elicits the secretion of IL-6 [95]. PGE2 induction of Stat3 is operative in this process [96].

11.5 PPAR γ and PPAR α in Inflammation and Immunity

Transactivation of a particular gene requires a large complex of proteins (Figs. 4.8, 4.19). In their inactive state, PPARs are complexes in the cell cytoplasm along with corepressor proteins. On activation, they dissociate and move into the nuclei, where PPARs form heterodimers with retinoid X receptors (Fig. 4.14). PPAR γ receptors are highly expressed in macrophage-derived foam cells in atheroma. It was found that PPAR γ receptors are involved in the differentiation of monocyte macrophages, and that PPAR γ receptors help to regulate their inflammatory activities [97]. In macrophages, PPAR γ serves as a stimulus for the anti-inflammatory action of carbon monoxide.

PPAR γ will promote the survival of lymphocytes since they suppress ROS by boosting antioxidants, and they attenuate any decline in adenosine triphosphate (ATP) [98]. Yet, fatty acid ligands for PPAR γ can inhibit proliferation of activated T lymphocytes through inhibition of IL-2 formation and induced T cell apoptosis. The PPAR γ associates with NFAT, so the DNA binding and transcriptional activation of the IL-2 promoter is blocked. Also, PPAR γ receptors are expressed on DCs, and PPAR γ ligands inhibit the release of IL-12 [99]. DCs treated with synthetic PPAR γ ligands have reduced ability to stimulate lymphocytes. People are getting excited about the potential use of PPAR γ ligands in conditions like rheumatoid arthritis [100]. PPAR γ agents will decrease synovial production of TNF α , IL-6, IL-8, metalloproteinases [101] and no doubt IL-12 (Fig. 12.12).

Both PPARs and liver X receptors of the nuclear-receptor superfamily function in combination with the GRs to repress inflammatory response genes [102]. NCoR (nuclear receptor corepressor) and SMRT corepressor complexes, which are able to interact with NF- κ B and AP-1, maintain the repression of inflammatory response genes (cf. Fig. 11.4). PPAR α receptors respond to a different set of ligands/drugs. PPAR α receptors are also expressed on monocyte macrophages, and ligands can induce apoptosis of those cells. PPAR α receptors dampen endothelial cell inflammatory responses. In aged mice, oxidative stress-induced NF- κ B is apparent in many tissues. Fortunately, PPAR α agonists restore the cellular redox balance, so there is amelioration of the production of inflammatory cytokines [103] since NF- κ B is inhibited (cf. Fig 11.4). Fenofibrate has been shown to repress IL-17 expression in cultured splenocytes and by Th-17 cells exposed to caecal bacterial antigens.




Fig 11.4 Peroxisome proliferator-activated receptor- γ (PPAR γ) with glucocorticoids suppresses inflammatory genes. NCoR nuclear receptor corepressor, UBC5 ubiquitin-conjugating enzyme 5

Of course, if liver X receptors will suppress inflammatory genes, appropriate agonists will have enormous potential. Cytokine release by monocytes in response to LPS has been investigated [104]. As yet, it is not clear how the process operates.

11.6 Cell Survival: Akt and Glycogen Synthase Kinase 3

Survival of a cell depends on a supply of growth factors from serum and on contact with neighbouring cells or adherence to extracellular matrix proteins [105]. The ERK pathway (Fig. 4.3) has the capacity for survival induction but is negatively regulated by phosphatases in conjunction with the proapoptotic p38/JNK paths. Hence, cancer cells should not emerge. Protein kinase B (Akt) confers the attributes of cell survival [106]. PI3 kinase (PI3K) produces PtdIns(3,4,5)P3 (Fig. 2.6), which binds to Akt so that it can translocate to the cell membrane, where it is phosphorylated by 3-phosphoinositide-dependent kinase 1 (PDK-1). When IL-2 acts on immune cells, Akt ensures their survival. Either a growth factor like PDGF (platelet-derived growth factor) or a cytokine like TNF α will activate Akt, which then associates with and phosphorylates IKK so that NF- κ B is activated. Concurrently, Akt suppresses proapoptotic influences [107]. Akt sequesters the proapoptotic protein Bad, it inhibits caspase 9, and it reduces transcription of Fas ligand (cf. Fig. 8.3). Conversely, since ceramide-activated PKC ζ and protein phosphatase 2A (PP2A) negatively regulate the Akt survival pathway [108], this explains how ceramide often mediates cell apoptosis. Nevertheless, the α 4-subunit of PP2A (and other phosphatases) works to keep apoptosis in check and to maintain cell survival [109].

It will help to be precise about what Akt can contribute. Akt can inactivate (a) death receptor-mediated apoptosis, (b) mitochondrial-dependent apoptosis and (c) p53-induced apoptosis. Akt via Cot serine-threonine kinase activates NF- κ B. NF- κ B supports antioxidant enzymes. Akt phosphorylates glycogen synthase kinase 3 β (GSK3 β), ensuring glucose uptake and metabolism. GSK3 is important in embryonic development. When Akt promotes the survival of adult cells, Akt phosphorylates and inhibits GSK3 [110] and thus enables supportive changes in glucose metabolism. GSK3 β is essential for cell survival, but too much or too little will cause apoptosis. On the one hand, components of the NF- κ B system are phosphorylated by GSK3 to promote NF- κ B activity. Conversely, if p53 binds nuclear GSK3, it is activated and promotes the transcriptional and apoptotic actions of p53 [111]. GSK3 supports formation of proinflammatory cytokines by inducing NF- κ B and CREB interaction with coactivator CBP. GSK3 inhibitors are thus anti-inflammatory [112].

Both growth factor and cytokine survival factors for cells, like insulin-like growth factor 1 (IGF-1) or IL-3 activate Akt, which phosphorylates Bad at three sites, and it then binds 14-3-3 protein, and it (Bad) is sequestered in the cytosol. So, apoptosis cannot occur. When macrophages meet endotoxin, they have to survive. When LPS interacts with TLR4 receptors (Figs. 12.2, 12.5), RIP links TLR4 to PI3K and so to Akt and cell survival [113]. Monocyte survival derives from surface CEACAM1 (CD66a), which connects with PI3K and hence to Akt activation.

Insulin growth factors I and II direct protein anabolism, and they are mitogens. Haemopoietic cells that are destined to die when IL-2 or IL-3 is withdrawn are protected by IGF-1. IGF-1 is antiapoptotic for many cells [114], and in many cancers an autophosphorylated IGF-1R (IGF-1 receptor) mediates antiapoptosis. The biochemical scenario is that when IGF-1 acts on its IGF-1R there is recruitment of a docking protein insulin receptor substrate 2 (IRS-2) (as for insulin) and that leads to activation of PI3K. PI3K activates a PDK-1, and in turn that activates Akt kinase. Naturally, the kinase-regulated signalling pathways are controlled by phosphatases. One is calcineurin, which will dephosphorylate Bad and so reverse the effect of Akt phosphorylation. The other is PTEN (phosphatase as in Fig. 2.6), the product of a tumour suppressor gene [115, 116]. The function of PTEN is to hydrolyze 3-phosphorylated inositol phospholipids [117]. Thus, PTEN acts in opposition to PI3K (Fig. 2.6). In malignancies, both copies of PTEN are deleted or mutated. Loss of PTEN in cells leads to Akt drive, and this is accompanied by resistance of cells to apoptotic stimuli [118]. Hence, cancer is likely to arise.

There are other survival kinases, suppressors of Myc-induced apoptosis, called the *Pim kinases* [119]. Myc induces proliferation but also increased apoptosis of nonmalignant cells. However, defects in control of apoptosis allow Myc to act as an oncogene. Akt or Pim control this. Both Akt and Pim inactivate Bad. Pim kinases are implicated in cancers and lymphomas. Pim kinase was discovered by its ability to stop Myc-induced apoptosis in a mouse model of lymphoma. Cytokines via Jak-Stat action (Sect. 12.9) induce Pim kinases (Pim1, -2, -3).

This section emphasizes the importance of NF- κ B for cell survival. NF- κ B suppresses apoptosis (Fig. 8.2) since NF- κ B supports bcl-2 and bcl-X1, it supports the inhibitor of apoptosis proteins cIAPc and xIAP, and c-FLIP works in its favour [120] it room. Yet, the JNK cascade can work in opposition to the (GADD45) and Xiap proteins, which promote antiapoptosis [120] (Table 11.3). Likewise, the p38 MAP kinase pathway is proapoptotic [121, 122]. Actually, induction of NF- κ B in some cells enables them to stay alive long enough to acquire genetic errors. Hence, NF- κ B activity can promote carcinogenesis [5] and help metastases [123]. Such carcinogenesis is oxygen radical mediated [124]. Yet generally, NF- κ B is acting to suppress tumour cell growth for it aids triggering of p53 and ARF (Fig. 17.5). However, if cells lose their tumour suppressor genes, NF- κ B then promotes oncogenesis [125].

11.7 Insulin Receptors

Insulin is a potent anabolic hormone. Whereas glucose elicits proinflammatory effects, insulin is anti-inflammatory for its action suppresses NF- κ B, AP-1 and early growth response 1 (EGR-1) TFs. Of course, insulin is a principal pancreatic autoantigen relevant to diabetes [126]. Yet, one could be interested in how signalling by the insulin receptors compares with other mechanisms discussed in this text.

Insulin receptor stimulation in insulin-responsive cells leads to translocation of GLUT4 and glucose transport. The insulin receptor is an $\alpha_2\beta_2$ heterotetrameric complex in which two α -subunits and two β -subunits are linked by disulphide bonds. Insulin binds to the extracellular α -subunits, and it thereby transmits a signal across the plasma membrane that activates the intracellular tyrosine kinase of the β -subunit. The one β -subunit phosphorylates its partner on specific tyrosine residues. Accordingly, the receptor tyrosine kinase induces phosphorylation of IRSs on multiple tyrosine residues. These IRS phosphotyrosine residues act as docking sites for -SH2 domain containing proteins, including the p85 regulatory unit of PI3K (Fig. 2.7). Hence, the p110 catalytic unit of PI3K, the lipid kinase unit, will lead to glucose transport [127, 128]. Other proteins that dock at the insulin receptor include adaptor proteins Shc and Grb2 (growth factor receptor-bound protein), leading to Ras-MAP kinase signalling for the mitogenic action of insulin, the SHP2 phosphatase [129] and Cbl (casitas B lineage lymphoma protein).

The PI3K (Fig. 11.5) p110 catalytic unit creates phosphoinositides PI(3,4,5)P3, PI(3,4)P2 and PI(3)P at the plasma membrane (Fig. 2.6). They increase GLUT4 translocation but do not actually increase glucose transport. That could be mediated by AS160 (Akt substrate of 160 kDa) [130]. The plasma membrane target for the GLUT4 vesicle is the t-SNARE called syntaxin 4. Inactivation of Glut4 in muscle leads to raised plasma glucose and glucose toxicity that results in insulin resistance [131]. Insulin action is facilitated by the formation of ROS at multiple signalling targets [132], but substantial ROS produced in response to TNF α or dexamethasone creates insulin resistance [133].

There is subtle control by suppressor of cytokine signalling (SOCS) proteins. First, SOCS3 reduces the phosphorylation of IRS-1 and its subsequent association with the p85 of PI3K. Second, the SOCS1 and -6 can inhibit insulin receptor tyrosine kinase activity, thereby reducing the phosphorylation of IRS-1/2 and the downstream events. IRS-1 and IRS-2 are targeted for proteasomal degradation. SOCS1 and -3 are elevated in rodent models of insulin resistance and diabetes.

In adipocytes of obese persons with type 2 diabetes, IRS-2 becomes the main docking protein for PI3K [134]. Local TNF α acting on the adipocytes is a cause of




Fig. 11.5 Scheme for insulin receptor activation. *PDK-1* 3-phosphoinositide-dependent kinase 1. Active JNK opposes IRS 1, 2 action. In this scheme, phosphatase Src homology inositol polyphosphate 5-phosphatase 2 (*SHIP2*) regulates the PI3 kinase-dependent signalling. Insulin-like growth factor 1 receptor (*IGF-IR*) also uses insulin receptor substrate 1 (*IRS-1*), which can become intranuclear and thereby activate cell cycle genes. This explains the relevance of IGF-1 to cancers

insulin resistance [135]. Ganglioside GM3 on fat cells corresponds to insulin resistance. TNF α -induced insulin resistance involves activation of inhibitory serine kinases or tyrosine phosphatases, which inhibit the insulin signalling path. It works through H₂O₂ generation. Also, there is serine phosphorylation of IRS-1, rather than IRS-1 tyrosine phosphorylation, that causes reduced insulin receptor kinase activity. In skeletal muscle, TNF α inhibits Akt phosphorylation, causing insulin resistance [136]. Ceramide, by blocking insulin stimulation of Akt, leads to insulin resistance [137]. TNF α can also produce insulin resistance in skeletal muscle by activation of I κ B kinase inhibitor of NF- κ B in a p38-dependent manner [138]. Most particularly there is upregulated JNK which impairs the action of IRS 1/2 [138b].

Crucially, we now realize that adiponectin receptors mediate antidiabetic metabolic effects. Adiponectin promotes glucose uptake, fatty acid oxidation and PPAR α ligand activities. Adiponectin is decreased in obesity, in type 2 diabetes and whenever there is insulin resistance [139]. There is more interest in adipokines from fat stores. The cytokine “resistin” inhibits insulin signalling in liver and muscle in mice [140, 141].

Insulin resistance is essentially due to decreased insulin-stimulated glucose uptake into skeletal muscle. The muscle PI3K (with its p110 catalytic unit and p85 regulatory unit) actually shows increased transcription of the p85 α and of splice variants p55 and p50, and the increase of these subunits must reduce PI3K activity by competing for phosphotyrosine targets [142]. Insulin resistance is accompanied by endothelial cell dysfunction, and there is enhanced aggregability of platelets. Insulin-resistant macrophages are integral to the problem of metabolic syndrome [143]. When they are controlled by knockout of JNK1, insulin resistance is ameliorated, and mice are resistant to high-fat diet-induced obesity [144].

Colca [145] rationalized the complex situation by proposing that insulin resistance is a physiological compensation to excessive oxidative metabolism. A new successful set of antidiabetic drugs, the thiazolidinediones (TZDs) [146], enhances PPAR γ -retinoid X receptor heterodimer formation (Fig. 4.14) and offsets insulin resistance.

11.8 Cross-Talk between Insulin Signalling and the Angiotensin II System

Insulin and angiotensin II (Ang II) play pivotal roles in the metabolic and circulatory systems. Failure in their proper actions leads to diabetes mellitus and hypertension, respectively [147]. There is cross-talk at multiple levels between these systems that affects the action of antihypertensive drugs used in the control of cardiac hypertrophy and energy acquisition by heart muscle. Thus, insulin actually promotes a significant increase of AT1 receptor protein expression in vascular smooth muscle cells (cf. Fig. 5.10).

The insulin signalling system described stems from a balance between tyrosine phosphorylation of IRS-1 and the negative effect of IRS-1 serine phosphorylation.

Too much of the latter occurs in insulin resistance (Fig. 11.5). In fact, PI3K, ERK and JNK as activated by AngII catalyse the serine phosphorylation of the insulin receptor and the IRS-1, and indeed of PI3K, impairing insulin-promoted activation toward Akt. Hence, in contrast to the effect of insulin, IRS-1/2-associated PI3K activity, and thereby endothelial nitric oxide synthase (eNOS) activity, is inhibited by Ang II in a dose-dependent manner [148]. So, insulin-induced vasodilatation is thwarted by Ang II. Furthermore, Ang II induces SOCS3 activity to impair signal transduction through the Jak2/Stat5 signal pathway [149].

11.9 Understanding mTOR

mTOR is mammalian target of rapamycin. It is an integrator of cellular energy status, nutrient and growth factor signals and the coordinator of cell growth and cell cycle progression [150]. TOR directs cell surface expression of various nutrient transporters. Amino acids also stimulate mTOR via the class 3 PI3Ks [151]. Since insulin and IGF-1 are anabolic, Fig. 11.6 shows how those receptors connect via IRS proteins to activation of PI3K and hence Akt. In its turn, Akt phosphorylates and inhibits substrates like the Foxo (Forkhead TFs), Bad, GSK3 and tuberin or hamartin. Tuberin, characteristic of benign tumours called hamartomas as seen in tuberose sclerosis [152, 153], inhibits TOR, but tuberin is inhibited by Akt. This means that Akt activity stimulates glucose uptake and cell growth and proliferation, and apoptosis is inhibited. Actually, mTOR exerts negative feedback on PDGF receptors [154]. The TOR pathway requires nutrients like the amino acid leucine, and then Tor complex 1 (mTORC1) activates S6 kinase and promotes the eukaryotic translation initiation factor 4E (eIF4E) [155]. Additionally, mTORC2 regulates polarization of the actin cytoskeleton. Furthermore, should cell ATP levels fall, the energy-sensing kinase AMPK inhibits hepatic gluconeogenesis (Fig. 1.1) because it promotes TORC2 phosphorylation, which blocks its nuclear accumulation [156].




Fig 11.6 Model of the PI3 kinase-Akt-tuberose sclerosis (*TSC*)-mammalian target of rapamycin (*mTOR*) pathway. *Tuberin (*TSC2*) or hamartin (*TSC1*) are tumour suppressors. **AMPK, which is activated by energy deprivation, stimulates *TSC2*/TOR complex 2 (*Torc2*). The eukaryotic translation initiation factors (*eIFs*) are versatile scaffolds for translation initiation complexes [139]

AMPK is a fuel sensor. In muscle AMPK is activated by exercise [157]. It is a key metabolic regulator in liver, heart and muscle. AMPK is activated by cell cAMP, and it will stimulate fatty acid β -oxidation. That means that if hepatic AMPK is inactivated, as occurs in diabetes, there is fatty acid synthesis and development of hyperlipidaemia. In T cells, AMPK controls metabolic steps required for cell proliferation [158].

The tuberose sclerosis 1 and 2 (TSC1-TSC2) protein complex heterodimers integrate cues from growth factors, the cell cycle and nutrients to regulate the activity of mTOR, p70 S6 kinase, 4E-BP1 \rightarrow eIF4E and ribosomal S6 protein [159]. As you will predict, hypertrophy of cells is achieved by augmentation of their RNA and protein content. Formation of mRNA is determined (a) by growth factor Ras-ERK-Mnk-1 \rightarrow eIF4E and (b) by the path mTOR \rightarrow p70 S6 kinase.

Note that these molecules in Fig. 11.6 do not form a linear signal pathway, but they act as agonists and antagonists of each other's activities. Stimulation of mTOR and S6 kinase by amino acids requires participation of a class 3 PI3K called hVps34 (human vacuolar protein sorting 34) [160]. PI3Ks promote glycolysis and thus production of ATP. As in other situations, PTEN controls PI3K activity. Loss of PTEN in mice leads to hyperactive signalling to TOR and disturbance of the control of haemopoietic stem cells and a liability to leukaemia [161]. Understandably, mTOR must be implicated in carcinogenesis [162].

GSK3 β inhibitors are anti-inflammatory, witness that inhibitors like thiazolidinediones (TZDS) attenuate carrageenan-induced lung injury [163].

In T cells, Akt is activated maximally when there is coligation of the TCR and CD28 (Fig. 13.5). Moreover, T cells have AMPK α 1, which is rapidly activated with TCR triggering via the adapter molecules LAT and SLP76 (Fig. 13.4) [164]. When amino acids are supplied, mTOR phosphorylates components of the translational apparatus like the eukaryotic initiation factor eIF4E and p70 S6 kinase. Cytokines also induce PIM1/PIM2 kinases, which help maintain a high rate of glycolysis and protein translation [165].

Rapamycin is a useful immunosuppressant for allograft transplants. Yet, rapamycin does not phenocopy the effects of nutrient or growth factor deprivation. Actually, it fails to prevent the early stages of T lymphocyte activation, including cell cycle entry and upregulation of cell surface activation markers. Outstandingly, rapamycin, by its inhibition of mTOR and thereby of the phosphorylation of S6 kinase and eIF, prevents cell hypertrophy of a remaining kidney after unilateral nephrectomy [166].

S6 kinase is integral to multiple pathways in which cells are rendered unresponsive to insulin, as during chronic insulin exposure, elevation of free fatty acids and perhaps TNF α exposure. A feedback leads to IRS1/2 serine phosphorylation and hence downregulation. Accordingly, there is insulin resistance and impaired cell survival.

Chronic localised activation of mTOR results from genetic loss of certain tumour suppressor genes like TSC1, PTEN in Cowden's disease and 1KB1 (Peutz-Jegher's syndrome) [167].

11.10 Alveolar Macrophage Survival

Armed as we are now with so much new knowledge, we can tackle the question of how alveolar macrophages (AMs) survive so well, even though they are exposed continually to chemical pollutants, ROS, inflammatory mediators and microbial invaders. In fact, AMs have two survival pathways: (a) the PI3K → Akt pathway and (b) ERK activity that leads to inhibition of proapoptotic proteins but good antiapoptosis. The Hunninghake group [168] investigated how it is that ERK activity leads to sustained protein translation initiation via eIF4E or hypophosphorylated (active) eIF2 α . Actually, ERK decreases proapoptotic JNK activity by stabilizing MPK7 dual phosphatase. When JNK activity is low, there is higher activity of phosphatase PP1 α . Accordingly, there is hypophosphorylation of translation initiation factor eIF2 α , so there is ongoing protein translation that means survival of the AMs.

References

1. Perkins MD. The Rel/NF κ B family: friend and foe. *Trends Biochem Sci* 2000;25:434–440
2. Fanton E, Traincard F, Levy SG et al. Nemo oligomerization in the dynamic assembly of the I κ B kinase core complex. *FASEB J* 2007;27:2540–2551
- 3a. Mukhopadhyay D, Riezman H. Proteasome independent functions of ubiquitin in endocytosis and signaling. *Science* 2007;315:201–205
- 3b. Wullaert A, Heyninck K, Janssens S, Beyaert R. Ubiquitin: tool and target for intracellular NF- κ B inhibitors. *Trends Immunol* 2006;27:533–540
- 4a. Sun S-C, Ley SC. New insights into NF- κ B regulation and function. *Trends Immunol* 2008; 29:469–479
- 4b. Pereisa SG, Oakley F. Nuclear Factor Kappa B: regulation and function. *Int J Biochem Cell Biol* 2008;40:1425–1430
5. Kumar A. Nuclear factor κ B: its role in health and disease. *J Mol Med* 2004;82:434–438
6. Barnes PJ. Nuclear factor kappa-B. *Int J Biochem* 1997;29:415–435
7. Bonizzi G, Karin M. The two NF- κ B activation pathways and their role in innate and adaptive immunity. *Trends Immunol* 2004;25:280–289
8. Mercurio F, Manning AM. NF κ B as a primary regulator of the stress response. *Oncogene* 1999;18:6163–6171
9. Bauerle PA, Henkel T. Function and activation of NF κ B in the immune system. *Annu Rev Immunol* 1994;12:141–179
10. Catley MC, Cambridge LM, Nasuhara Y et al. Inhibitors of PKC prevent activated transcription: downstream of NF κ B DNA binding. *J Biol Chem* 2004;279:18457–18466
11. Sec CK, Parker L. Antioxidant and redox regulation of gene transcription. *FASEB J* 1996;10:709–720
12. Kamata H, Hirata H. Redox regulation of cellular signalling. *Cell Signal* 1999;11:1–14
13. Marshall HE, Hess DT, Stamler JS. S-Nitrosylation: physiological regulation of NF κ B. *Proc Natl Acad Sci U S A* 2004;101:8841–8842
14. Friedland JS, Constantin D, Shaw TC, Stylanou E. Regulation of Il-8 gene expression after phagocytosis of zymosan by human monocytic cells. *J Leukoc Biol* 2001;70:600–605
15. Natarajan R, Gupta S, Fisher BJ et al. Nitric oxide suppresses Il-8 transcription by inhibiting c-Jun terminal kinase induced AP-1 activation. *Exp Cell Res* 2001;266:203–212

16. Sebban H, Yamaoka S, Gourtois G. Post-translational modifications of Nemo and its partners in NF- κ B signaling. *Trends Cell Biol* 2006;16(11):569–577
17. Woronicz JD, Gao X, Cao Z, Rothe M, Goeddel DV. NF κ B activation and complex formation with I κ B kinase and NIK. *Science* 1997;278:806–809
18. Israel A. The IKK complex: an integrator of all signals that activate NF- κ B. *Trends Cell Biol* 2000;10:129–132
19. Ling L, Cao Z, Goeddel DV. NF κ B inducing kinase activates IKK α by phosphorylation of Ser176. *Proc Natl Acad Sci U S A* 1998;95:2791–2797
20. Delhase M, Hayakawa M, Chen Y, Karin M. Positive and negative regulation of I κ B kinase activity through IKK β subunit phosphorylation. *Science* 1999;284:309–313
- 21a. Nelson DE, Ihekwa AEC, Ellitoo M et al. Oscillations in NF κ B signaling control the dynamics of gene expression. *Science* 2004;306:704–708
- 21b. Grey S T. The Ying and Yang of NF- κ B activation. *Immunol Cell Biol* 2008;86:299–300
- 22a. Wietek C, O’Neill LA. Diversity and regulation in the NF-kappaB system. *Trends Biochem Sci* 2007;32(7):311–319
- 22b. Enesa K, Ito K, Luong LA et al. Hydrogen peroxide Prolongs nucleus localization of NF- κ B in activated cells by suppressing negative regulatory mechanisms *J Biol Chem* 2008;283:18582–18590
- 22c. Salminen A, Paimela T, Surronen T, Kaarniranta K. Innate immunity meets with cellular stress at the IKK complex: regulation of the IKK complex by HSP10 and HSP 9d. *Immunol Cells* 2008;117:9–15
23. Olson CM, Hedrick MN, Izadi H et al. p38 MAP kinase controls NF-kappaB transcriptional activation and TNF α production. *Infect Immun* 2007;75:270–277
24. Liu B, Yang Y, Chermishof V et al. Proinflammatory IKK α phosphorylation of PIAS1 to restrict inflammation and immunity. *Cell* 2007;129:903–914
25. Gorska MM, Liang Q, Stafford SJ et al. MK2 controls the level of negative feedback in the NFkappaB pathway. *J Exp Med* 2007;204:1637–1652
26. Kolls JK. Caught between CYLD and Charybdis. *Immunity* 2007;27:187–189
27. Fessler MB, Arndt PG, Just I et al. Dual role for RhoA in suppression and induction of cytokines in the human neutrophil. *Blood* 2007;109:1248–1256
28. Gloire G, Dejardin E, Piette J. Extending the nuclear roles of I κ B kinase subunits. *Biochem Pharmacol* 2006;72:1081–1089
29. Hacker H, Karin M. Regulation and function of IKK and IKK-related kinases. *Sci STKE* 2006;Oct 17;2006(357):re13
30. Israel A. A regulator (IKK α) branches out. *Nature* 2003;423:596–597
31. Gloire G, Horion J, El Mjiyad N et al. Promoter dependent effect of IKK α on NF-kappaB/p65 DNA binding. *J Biol Chem* 2007;282:21308–21318
- 32a. Lawrence T, Bebie M. IKK α in regulation of inflammation and adaptive immunity. *Biochem Soc Trans* 2007;35(2):270–272
- 32b. Ghosh S, Hayden MS. New regulators of NF- κ B in inflammation. *Nat Revs Immunol* 2008;8:837–848
- 33a. Lawrence T, Bebie M, Liu GY, Nizet V, Karin M. IKK α limits macrophage NF κ B activation and contributes to resolution of inflammation. *Nature* 2005;434:1138–1143
- 33b. Rius J, Guma M, Schachtrup C et al. NF- κ B links immunity to the hypoxic response through transcriptional regulation of HIF-1 α . *Nature* 2008;453:807–811
34. Greten FR, Arkan MC, Bollrath J et al. NF- κ B is a negative regulator of Il-1beta secretion as revealed by genetic and pharmacological inhibition of IKK β . *Cell* 2007;130:918–931
- 34b. Fong C, Bebie M, Didierlaurent A et al. An anti-inflammatory role for IKK β through the inhibition of classical macrophage activation. *J Exp Med* 2008;205:1269–1276
- 34c. Timmer AM, Nizet V. IKK β /NF- κ B and the miscreant macrophage. *J Exp Med* 2008;205:1255–1259
35. Muller JR, Siebenlist U. Lymphotoxin receptor induces sequential activation of distinct NF κ B factors via separate signaling paths. *J Biol Chem* 2003;278:12006–12012

36. Cupedo T, Mebius RE. Cellular interactions in lymph node development. *J Immunol* 2005;174:21–25
37. Zhu M, Chin RK, Christiansen PA et al. NF- κ B for establishment of central tolerance through an Aire dependent pathway. *J Clin Invest* 2006;116:2964–2971
38. Novak DV, Yin L, Hagen-Stapleton A et al. The I κ B function of NF κ B2 p100 controls stimulated osteoclastogenesis. *J Exp Med* 2003;198:771–781
39. Aya K, Alhawaari M, Hagen-Stapleton A et al. NF- κ B inducing kinase controls lymphocyte and osteoclast activities in inflammatory arthritis. *J Clin Invest* 2005;115:17–22
40. Yamaguchi T, Sakaguchi S. Skin controls immune regulators. *Nat Med* 2006;12:1358–1359
41. Legarda-Addison D, Ting AT. Negative regulating of T cell receptor signaling by NF κ B2/p100. *J Immunol* 2007;178:7767–7778
42. Neumann M, Naumann M. Alternative regulation of NF κ B. *FASEB J* 2007;21:2642–2644
43. Saha R, Jana M, Pahan K. MAPkinase p38 regulates transcriptional activity of NF- κ B in primary human astrocytes via acetylation of p65. *J Immunol* 2007;179:7101–7109
44. Chen L-F, Greene WC. Regulation of NF κ B transcription factor complex by acetylation. *J Mol Med* 2003;81:549–557
45. Lu Y, Wahl LM. Oxidative stress augments production of MMP1, Cox2 and PGE2 through enhancement of NF κ B in LPS activated human monocytes. *J Immunol* 2005;175:5423–5429
46. Qu J, Zhang J, Pan J et al. Endotoxin tolerance inhibits lipopolysaccharide initiated acute pulmonary inflammation and lung injury in rats by the mechanism of NF κ B. *Scand J Immunol* 2003;58:613–619
47. Yoza BK, Hu JY-Q, Cousart SL et al. Induction of Rel B participates in endotoxin tolerance. *J Immunol* 2006;177:4080–4085
48. Cuschieri J, Billigren J, Maier RV. Endotoxin tolerance attenuates LPS induced TLR4 mobilization to lipid rafts. *J Leukoc Biol* 2006;80:1289–1297
49. Zheng S-J, Lamhamedi-Cherradi S-E, Wang P, Yu L, Chen YH. Tumor suppressor p53 inhibits autoimmune inflammation and macrophage function. *Diabetes* 2005;54:1423–1428
50. Totzke G, Essman F, Pohlmann S et al. I κ B- ζ as a regulator of NF- κ B activity. *J Biol Chem* 2006;281:12645–12654
51. Wessells J, Baer M, Young HA et al. Bcl-3 and NF κ B p50 attenuate lipopolysaccharide induced inflammatory responses in macrophages. *J Biol Chem* 2004;279:49995–50003
52. Carmody RJ. Negative regulation of TLR signaling by NF κ B p50 blockade ubiquitination. *Science* 2007;317:675–678
53. Toney LM, Cattoretti G, Graf JA et al. Bcl-6 regulates chemokine gene transcription in macrophages. *Nat Immunol* 2000;1:214–220
54. Dent AL, Shaffer AL, Yu X et al. Control of inflammation, cytokine expression and germinal centre formation by Bcl-6. *Science* 1997;276:589–592
55. Phan RT, Saito M, Basso K et al. Bcl-6 interacts with transcription factor Miz-1 to suppress the CDKI p21 and cell cycle arrest in germinal centre B cells. *Nat Immunol* 2005;6:1054–1058
56. Seto M. Genetic and epigenetic factors involved in B cell lymphomagenesis. *Cancer Sci* 2004;95:704–710
57. Pasqualucci L, Migliazza A, Basso K et al. Mutations of the Bcl-6 protooncogene disrupt its negative autoregulation in diffuse large B cell lymphoma. *Blood* 2003;101:2914–2919
58. Beg AA. Compartmentalising NF- κ B in the gut. *Nat Immunol* 2004;5:14–16
59. Schenk M, Bouchen A, Birrer S et al. Macrophages expressing triggering receptor expressed on myeloid cells-1 are underrepresented in the human intestine. *J Immunol* 2005;174:517–524
60. Visekruna A, Joeris T, Seidel D et al. Proteasome mediated degradation of I κ B α and processing of p105 in Crohns disease and ulcerative colitis. *J Clin Invest* 2006;116:3195–3203
61. Harris J, Olieri S, Sharma S et al. Nuclear accumulation of c-Rel following C-terminal phosphorylation by TBK1/IKK ϵ . *J Immunol* 2006;177:2527–2535

62. Kaisho T, Tanaka T. Turning NF- κ B and IRFs on and off in DC. *Trends Immunol* 2008; 29:329–337
63. Khachigian LM, Collins T, Fries JW. N-Acetylcysteine blocks mesangial VCAM-1 and NF κ B expression in vivo. *Am J Pathol* 1997;151:1225–1229
64. Loscher CE, Draper E, Leavy O et al. Conjugated linoleic acid suppresses NF κ B activation and IL-12 production in dendritic cells. *J Immunol* 2005;175:4990–4998
65. Nishiyama S, Manabe N, Kuhota Y et al. Cyclosporin inhibits the early phase of NF κ B/RelA activation induced by CD238 costimulation signaling to reduce IL-2 expression in human peripheral T cells. *Int Immunopharmacol* 2005;5:699–710
66. D'Acquisto F, Ianaro L. From willow bark to peptides: the ever widening spectrum of NF κ B inhibitors. *Curr Opin Pharmacol* 2006;6:387–392
67. Kim E-H, Singh Y-J. 15Deoxy-PGJ2 as potential endogenous regulator of redox-sensitive transcription factors. *Biochem Pharmacol* 2007;72:1516–1528
68. Li J, Liu Z, Jiang S et al. T suppressor lymphocytes inhibit NF κ B mediated transcription of CD86 gene in antigen presenting cells. *J Immunol* 1999;163:6386–6371
69. Fiorini E, Schmitz I, Morrison WE et al. Peptide induced negative selection of thymocytes activates transcription of an NF κ B inhibitor. *Mol Cell* 2002;9:637–648
70. Quaaaz F, Arron J, Zheng Y et al. Dendritic cells require distinct NF κ B subunits. *Immunity* 2002;16:257–270
71. Tato CM, Hunter CA. Subversion and utilization of the NF κ B pathway during infection. *Infect Immunol* 2002;70:3311–3317
72. Sacani S, Pontano S, Natoli G. p38 dependent marking of inflammatory genes for increased NF κ B recruitment. *Nat Immun* 2002;3:69–75
73. Adcock IM, Ito K, Barnes PJ. Glucocorticoids: effects on gene transcription. *Proc Am Thorac Soc* 2004;1:247–254
74. Gomez PF, Pillinger MH, Attur M et al. Resolution of inflammation: prostaglandin E2 dissociates nuclear trafficking of NF- κ B subunits. *J Immunol* 2005;175:6924–6930
75. Cohen-Lahav M, Douvdevani A, Chaimovitz C, Shany S. The anti-inflammatory activity of 1,25 dihydroxyvitamin D3 in macrophages. *J Steroid Biochem* 2007;103:558–562
76. Cuzzocrea S, Mazzon E, Di Paola R et al. The role of PPAR-alpha in the regulation of inflammation. *J Leukoc Biol* 2006;79:999–1010
77. Belaovaj A et al. Mice lacking neutrophil elastase reveal impaired host defence against gram-negative bacterial sepsis. *Nat Med* 1998;4:615–618
78. Reeves EP et al. Killing activity of neutrophils is mediated through activation of proteases by potassium flux. *Nature* 2002;416:291–297
79. Perkins ND. Oncogenes, tumour suppressors and p52NF κ B. *Oncogene* 2003;22: 7553–7556
80. Kataoka T et al. Caspase 8 inhibitor FLIP promotes activation of NF κ B and ERK signaling pathways. *Curr Biol* 2000;10:640–648
81. Iyer SS, Agrawal RS, Thompson CR et al. PLD-1 regulates phagocyte adhesion. *J Immunol* 2006;176:3686–3696
82. Yiu GK, Toker A. NFAT induces breast cancer cell invasion by promoting the induction of Cox2. *J Biol Chem* 2006;281:12210–12217
83. Pettus BJ, Bielawski S, Porcelli AM et al. The sphingosine kinase: sphingosine-1-phosphate pathway mediates Cox2 induction in response to TNF α . *FASEB J* 2003;17:1411–1421
84. Vega A, Chacon P, Alba G et al. Modulation of IgE dependent Cox2 gene expression by reactive oxygen species in human neutrophils. *J Leukoc Biol* 2006;80:152–163
85. Seo JH, Kim H, Kim KH. Cyclooxygenase 2 expression by transcription factors in *H. pylori* infected gastric epithelial cells. *Ann N Y Acad Sci* 2002;973:477–480
86. Ki SH, Choi MJ, Lee CH, Kim SG. G α 12 regulates Cox-2 induction by sphingosine-1-phosphate. *J Biol Chem* 2007;282:1938–1947
87. Dixon DA, Tolley ND, Bemis-Standoli K et al. Expression of Cox2 in platelet-monocyte interactions involving adhesion and cytokine signaling. *J Clin Invest* 2006; 116:2727–2732

88. Kititoshi S, Nishikowa T, Sonoda K et al. Reactive oxygen species from mitochondria induce Cox-2 gene expression in human mesangial cells. *Diabetes* 2003;52:2570–2576
89. Caivano M, Gorgoni B, Cohen P et al. The induction of Cox2 in macrophages is biphasic. *J Biol Chem* 2001;276:48693–48701
90. Kang Y-J, Wingerd BA, Arakawa T, Smith WL. Cox2 gene transcription in a macrophage model of inflammation. *J Immunol* 2006;177:8111–8122
91. Nie M, Pang L, Inoue H, Knox AJ. Transcriptional regulation of Cox-2 by bradykinin and interleukin 1 beta in human airway smooth muscle cells: involvement of different promoter elements, transcription factors and histone H4 acetylation. *Mol Cell Biol* 2003;13:9233–9244
92. Vila-del Sol V, Fresno M. Involvement of TNF α and NF κ B in the transcriptional control of Cox-2 expression by IFN γ in macrophages. *J Immunol* 2005;174:2825–2833
93. Rodriguez-Barbero A, Dorado F, Velasco S et al. TGF β 1 induces Cox2 expression and PGE2 synthesis in human mesangial cells. *Kidney Int* 2006;70:901–909
94. Steer SA, Moran JM, Christmann BS et al. MAPK in regulation of ds-RNA and virus induced Cox2 expression by macrophages. *J Immunol* 2006;177:3413–3420
95. Chen B-C, Liao C-C, Hsu M-J et al. Peptidoglycan induced Il-6 production in Raw 264.7 macrophages is mediated by Cox-2. *J Immunol* 2006;177:681–693
96. Han C, Demetris AJ, Stolz DB et al. Modulation of Stat3 activation by the cPLA2 and Cox2 controlled PGE2 signaling pathway. *J Biol Chem* 2006;281:24831–24846
97. Clark RB. The role of PPARs in inflammation and immunity. *J Leukoc Biol* 2002;71:388–400
98. Jo S-H, Yang C, Miao Q et al. PPAR γ promotes lymphocyte survival via actions on cell metabolic activities. *J Immunol* 2006;177:3737–3745
99. Faveeuw C, Fougeray S, Angeli V et al. PPAR γ activators inhibit interleukin 12 production in murine dendritic cells. *FEBS Lett* 2000;486:261–266
100. Fahmi H, Pelletier J-P, Martel-Pelletier J. PPAR γ ligands as modulators of inflammatory and catabolic responses in arthritis. *Arthritis Rheum* 2002;29:3–12
101. Yamasaki S, Nakashima T, Kawakami A et al. Functional changes in rheumatoid fibroblast synovial cells through activation of PPAR γ signaling pathway. *J Immunol* 2002;169:1145–1150
102. Glass CK, Ogawa S. Combinatorial roles of nuclear receptors in inflammation and immunity. *Nat Rev Immunol* 2006;6:44–55
103. Poynter ME, Davies RA. PPAR γ activation modulates cellular redox status, represses nuclear factor-B signaling, and reduces inflammatory cytokine production in ageing. *J Biol Chem* 1998;273:32833–32841
104. Myhre AE, Agren J et al. Liver X receptor regulator of cytokine release. *Shock* 2008;29:465–474
105. Yamada KM, Clar K. Survival in three dimensions. *Nature* 2002;419:790–791
106. Brazil DP, Hemmings BA. 10 years of protein kinase B signaling. *Trends Biochem Sci* 2001;26:657–664
107. Franke TF, Kaplan DR, Cantley LC. PI 3-kinase downstream AKTion blocks apoptosis. *Cell* 1997;88:435–437
108. van Blitterswijk WJ, van der Luit AH, Veldman RJ et al. Ceramide: second messenger or modulator of membrane structure. *Biochem J* 2003;369:199–211
109. Kong M, Fox CJ, Mu J et al. The PP2A associated protein α 4 is an essential inhibitor of apoptosis. *Science* 2004;306:695–698
110. Hoeflich KP, Luo J, Rubie EA et al. Requirement for glycogen synthase kinase 3 in cell survival and NF κ B activation. *Nature* 2000;406:86–90
111. Jope RS, Johnson GVW. The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem Sci* 2004;29:95–101
112. Martin M, Rehane K, Jore RS, Michalek SM. Toll-like receptor and cytokine production is differentially regulated by GSK3. *Nat Immun* 2005;6:777–784
113. Vivarelli MS, McDonald D, Miller M et al. RIP links TLR4 to Akt and is essential for cell survival in response to LPS stimulation. *J Exp Med* 2004;200:399–404

114. O'Connor R, Fennelly C, Krause D. Regulation of survival signals from the IGF-1 receptor. *Biochem Soc Trans* 2000;28:47–51
115. Wang X, Gjorloff-Wingren A, Saxen M et al. The tumor suppressor PTEN regulates T cell survival and antigen receptor signaling. *J Immunol* 2000;164:1939–1943
116. Choi Y, Zhang J, Murga C et al. PTEN suppresses the PI3K/Akt pathway and induces growth inhibition and apoptosis of myeloma cells. *Oncogene* 2002;21:5289–5300
117. Stambolic V, Mak T, Woodgett JR. Modulation of cellular apoptotic potential: contributions to oncogenesis. *Oncogene* 1999;18:6094–6103
118. Stambolic V. PTEN: a new twist on β -catenin. *TIPS* 2002;23:104–106
119. Cibull TL, Jones TD, Li L et al. Overexpression of Pim.1 during progression of prostate adenocarcinoma. *J Clin Pathol* 2006;59:285–288
120. Kyriakis JM. Life or death decisions. *Nature* 2001;414:265–266
121. Ono K, Han J. The p38 signal transduction pathways. *Cell Signal* 2000;12:1–13
122. Li S-P, Junttila MR, Han J et al. The p38 MAP kinase pathway suppresses cell survival by inducing dephosphorylation of MAP/ERK1,2. *Cancer Res* 2003;63:3473–3477
123. Mayo MW, Baldwin AS. The transcription factor NF κ B: control of oncogenes and cancer therapy resistance. *Biochim Biophys Acta* 2000;1470:M55–M62
124. Klaumig JE, Kamendulis LM. The role of oxidative stress in carcinogenesis. *Annu Rev Pharmacol Toxicol* 2004;44:239–267
125. Perkins ND. NF- κ B: tumour promoter/suppressor. *Trends Cell Biol* 2004;16:64–69
126. You S, Chatenoud L. Proinsulin: a unique autoantigen triggering auto-immune diabetes. *J Clin Invest* 2006;116:3108–3110
127. White MF. The IRS signaling system: a network of docking proteins that mediate insulin and cytokine action. *Recent Prog Horm Res* 1998;53:119–138
128. Pessin JE, Sattiel AR. Signaling pathways in insulin action: molecular targets of insulin resistance. *J Clin Invest* 2000;106:165–169
129. Goldstein BJ, Ahmad F, Ding W et al. Regulation of the insulin signaling pathway by protein tyrosine phosphatases. *Mol Cell Biochem* 1998;182:91–99
130. Watson RT, Pessin JE. Bridging the GAP between insulin signaling and GLUT4 translocation. *Trends Biochem Sci* 2006;31:215–222
131. Kim JK, Zisman A, Fillimore JJ et al. Glucose toxicity and development of diabetes in mice with muscle specific inactivation of Glut4. *J Clin Invest* 2001;108:153–160
132. Goldstein BJ, Kalankar M, Wu Y. Insulin antigen is facilitated by insulin stimulated ROS with multiple potential signaling targets. *Diabetes* 2005;54:311–321
133. Houston N, Rosen ED, Lander ES. ROS have a causal role in multiple forms of insulin resistance. *Nature* 2006;440:944–946
134. Rondinone CM, Wang M, Lonroth P et al. IRS-1 is reduced and IRS-2 is the docking protein for PI3kinase in adipocytes from subjects with NIDDM. *Proc Natl Acad Sci U S A* 1997;94:4171–4175
135. Brindley DN, Wong C-N, Mei J et al. TNF α and ceramide in insulin resistance. *Lipids* 1999;34(Suppl):S85–86
136. Plomgaard P, Bouzahri K, Krogh-Madsen R et al. TNF α induces skeletal muscle insulin resistance via inhibition of Akt phosphorylation. *Diabetes* 2005;54:2939–2945
137. Stratford S, Hoehn KL, Liu F, Sumners SA. Regulation of insulin action by ceramide. *J Biol Chem* 2004;279:36608–36615
- 138a. de Alvaro C, Tervel T, Hernandez R, Lorenzo M. TNF α produces insulin resistance in skeletal muscle by activation of inhibitor kappa-B kinase. *J Biol Chem* 2004;279:17070–17078
- 138b. Yang R, Trevillyan JM-Jun N-terminal kinase pathways is diabetes. *Int J Biochem Cell Biol* 2008;40:2702–2706
139. Yamaudi T, Kaman J, Ito Y et al. Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 2003;423:762–768
140. Yudkin JS, Eringa E, Stehouwer CDA. Vasocrine signalling from perivascular fat: mechanism linking insulin resistance to vascular disease. *Lancet* 2005;365:1817–1820
141. Arner P. Resistin. *Diabetologia* 2005;48:2203–2205

142. Bandyopadhyay GK, Yu JG, Ofrecio J, Olefsky JM. Increased p85/55/50 expression and decreased PI3kinase activity in insulin-resistant human skeletal muscle. *Diabetes* 2005;54:2351–2359
143. Liang C-P, Hans S, Senokuchi T, Tall AR. The macrophage at the crossroads of insulin resistance and atherosclerosis. *Circ Res* 2007;100:1546–1555
144. Solinas G, Viku C, Neels JG et al. JNK1 in hematopoietically derived cells contributes to diet induced inflammation and insulin resistance. *Cell Metab* 2007;6:386–397
145. Colca JR. Insulin sensitizers may prevent metabolic inflammation. *Biochem Pharmacol* 2006;72:125–131
146. Stumvoll M, Haring H-U. Glitazones: clinical effects and molecular mechanisms. *Ann Med* 2002;34:217–223
147. Velloso LA, Folli F, Perego L, Saad MJA. The multi-faceted cross-talk between the insulin and angiotensin II signal systems. *Diabetes Metab Res Rev* 2006;22:98–107
148. Velloso LA, Folli F, Sun XJ et al. Crosstalk between the insulin and angiotensin signaling systems. *Proc Natl Acad Sci U S A* 1996;93:12490–12495
149. Calegari VS, Alves M, Picardi PK et al. SOCS3 provides a novel interface in crosstalk between angiotensin II and insulin signaling systems. *Endocrinology* 2005;146:579–588
150. Fingar DC, Blenis J. Target of rapamycin (TOR). *Oncogene* 2004;23:3151–3171
151. Nobukuni T, Joaquin M, Rocco M et al. Amino acids mediate mTOR signaling through activation of class 3 PI3kinase. *Proc Natl Acad Sci U S A* 2005;102:14238–14243
152. Manning BD. Balancing Akt with S6K: implications for both metabolic diseases and tumorigenesis. *J Cell Biol* 2004;167:399–403
153. Crino B, Nathanson KL, Henske EP. The tuberose sclerosis complex. *N Engl J Med* 2006;355:1345–1356
154. Zhang H, Bajraszewski N, Wu E et al. PDGF-receptors are critical for PI3kinase/Akt activation: negative regulation by mTOR. *J Clin Invest* 2007;117:730–738
155. Martin DE, Hall MN. The expanding TOR network. *Curr Opin Cell Biol* 2005;17:158–166
156. Koo S-H, Flechner L, Qi L et al. The CREB coactivator TORC2 is a key regulator of fasting glucose metabolism. *Nature* 2005;437:1109–1113
157. Hardie DG, Sakamoto K. AMPK: a key sensor of fuel and energy status in skeletal muscle. *Physiology* 2006;21:48–60
158. Jones RG, Thompson CB. Signal transduction fuels T cell activation. *Immunity* 2007;27:173–178
159. Hinnebusch AG. eIF3: a versatile scaffold for translation initiation complexes. *Trends Biochem Sci* 2006;31(10):553–562
160. Nobukuni T, Kozman SC, Thomas G. Hvps34 enters a growing game: mTOR complex 1/ S6K1 signaling. *Curr Opin Cell Biol* 2007;19:135–141
161. Cheung AM, Mak TW. PTEN in the haematopoietic system and its therapeutic indications. *Trends Mol Med* 2006;12:503–505
162. Guertin DA, Sabatini DM. Defining the role of mTOR in cancer. *Cancer Cell* 2007;12:9–22
163. Cuzzocrea S, Crisafulli C, Mazzone E et al. Inhibition of GSK-3 β attenuates development of carrageenan-induced lung injury. *Br J Pharmacol* 2006;35(5):965–970
164. Tamas P, Hawley SA, Clarke RG et al. Regulation of the energy sensor AMP-activated protein kinase by antigen receptor and Ca²⁺ in T lymphocytes. *J Exp Med* 2006;203:1665–1672
165. Fox CJ, Hammerman PS, Thompson CB. The Pim-kinases control rapamycin resistant T cell survival and activation. *J Exp Med* 2005;201:259–266
166. Chen J-K, Chen J, Neilson EG, Harris RC. Role of mammalian target of rapamycin signaling in compensatory renal hypertrophy. *J Am Soc Nephrol* 2005;16:1384–1313
167. Petroulakis E, Mamone Y, Le Bacquer O et al. mTOR signaling: implications for cancer. *Br J Cancer* 2006;94:195–199
168. Monick MM, Powers LS, Gross TJ et al. Active ERK contributes to protein translation by preventing JNK-dependent of protein phosphatase-1. *J Immunol* 2006;177:1636–1645