

Differential Expression of Two Basement Membrane Collagen Genes, COL4A6 and COL4A5, Demonstrated by Immunofluorescence Staining Using Peptide-specific Monoclonal Antibodies

Yoshifumi Ninomiya,* Megumi Kagawa,[§] Ken-ichi Iyama,** Ichiro Naito,^{||} Yumiko Kishiro,[§] Jerome M. Seyer,^{‡‡} Manabu Sugimoto,^{**} Toshitaka Oohashi,* and Yoshikazu Sado[§]

Departments of *Molecular Biology and Biochemistry and [‡]Ophthalmology, Okayama University Medical School, Okayama 700 Japan; Divisions of [§]Immunology and ^{||}Ultrastructure Research, Shigei Medical Research Institute, Okayama 700 Japan; **Department of Developmental Neurobiology, Kumamoto University Medical School, Kumamoto 860 Japan; and ^{‡‡}Connective Tissue Section, University of Tennessee, Memphis, Tennessee 38104

Abstract. Genes for the human $\alpha 5(\text{IV})$ and $\alpha 6(\text{IV})$ collagen chains have a unique arrangement in that they are colocalized on chromosome Xq22 in a head-to-head fashion and appear to share a common bidirectional promoter. In addition we reported a novel observation that the COL4A6 gene is transcribed from two alternative promoters in a tissue-specific manner (Sugimoto, M., T. Oohashi, and Y. Ninomiya. 1994. *Proc. Natl. Acad. Sci. USA.* 91:11679–11683). To know whether the translation products of both genes are colocalized in various tissues, we raised $\alpha 5(\text{IV})$ and $\alpha 6(\text{IV})$ chain-specific rat monoclonal antibodies against synthetic peptides reflecting sequences near the carboxy terminus of each noncollagenous (NC)1 domain. By Western blotting $\alpha 6(\text{IV})$ chain-specific antibody recognized 27-kD monomers and associated dimers of the human type IV collagen NC1 domain, which is the first demonstration of the presence in tissues of the $\alpha 6(\text{IV})$ polypeptide as

predicted from its cDNA sequence. Immunofluorescence studies using anti- $\alpha 6(\text{IV})$ antibody demonstrated that in human adult kidney the $\alpha 6(\text{IV})$ chain was never detected in the glomerular basement membrane, whereas the basement membranes of the Bowman's capsules and distal tubules were positive. The staining pattern of the glomerular basement membrane was quite different from that obtained with the anti- $\alpha 5(\text{IV})$ peptide antibody. The $\alpha 5(\text{IV})$ and $\alpha 6(\text{IV})$ chains were colocalized in the basement membrane in the skin, smooth muscle cells, and adipocytes; however, little if any reaction was seen in basement membranes of cardiac muscles and hepatic sinusoidal endothelial cells. Thus, both genes are expressed in a tissue-specific manner, perhaps due to the unique function of the bidirectional promoter for both genes, which is presumably different from that for COL4A1 and COL4A2.

BASEMENT membranes are continuous sheets of specialized extracellular matrix composed of collagen IV, laminin, nidogen, heparan sulphate proteoglycan, and other glycoproteins that are found wherever cells meet extracellular matrix (Rohrbach and Timpl, 1993). This means that basement membranes are located outside of most of the cells: at the dermal-epithelial junction; at the base of all lining epithelia throughout the digestive, respiratory, reproductive, and urinary tracts; underlying endothelia of capillaries and venules; around Schwann cells, adipocytes, skeletal, cardiac, and smooth muscle cells; and at the base of parenchymatous exocrine and endocrine glands (Junqueira et al., 1992). Thus, they are the

natural substrates on which most of the cells except blood cells grow and develop. They closely adjoin and are products of the overlying cells, and serve to compartmentalize the extracellular matrix and provide a barrier between cells and matrix.

Collagen type IV is a major structural component of basement membrane. The major form of this protein is a heterotrimer containing $\alpha 1(\text{IV})$ and $\alpha 2(\text{IV})$ chains, and this form appears to be ubiquitous in all basement membranes (Rohrbach and Timpl, 1993). Microsequencing of peptides made it possible to identify the human $\alpha 3(\text{IV})$ and $\alpha 4(\text{IV})$ chains (Wieslander et al., 1985; Butkowski et al., 1990). The primary structure of the human $\alpha 3(\text{IV})$ chain has been reported just recently (Mariyama et al., 1994) and cDNAs encoding the entire human $\alpha 4(\text{IV})$ chain have been isolated and characterized as well (Sugimoto et al., 1994; Leinonen et al., 1994). Lately, the nature of the $\alpha 5(\text{IV})$ chain has been identified by cDNA isolation

Address all correspondence to Yoshifumi Ninomiya, Department of Molecular Biology and Biochemistry, Okayama University Medical School, 2-5-1, Shikata-cho, Okayama 700 Japan. Tel: 81 86 223 7151 (ext. 2390). Fax: 81 86 222 7768.

(Hostikka et al., 1990; Zhou et al., 1992), and linkage between the $\alpha 5(\text{IV})$ gene (COL4A5) and Alport's Syndrome has been clarified (Barker et al., 1990). Zhou et al. (1993) and Oohashi et al. (1994) cloned genomic DNA and cDNAs coding for yet another $\alpha(\text{IV})$ chain, $\alpha 6(\text{IV})$. These newly discovered α chains still contain the amino-terminal 7S domain, the central COL1 domain, and the carboxy-terminal noncollagenous (NC)1¹ domain (Oohashi et al., 1994) as found in the classical $\alpha 1(\text{IV})$ and $\alpha 2(\text{IV})$ chains; however we do not know the mechanism to explain which of the six α chains are selected, or how they come together, to form collagen type IV molecules.

Of interest is that the genes encoding the six individual α chains in humans are paired by two on three different chromosomes: the $\alpha 1(\text{IV})$ gene (COL4A1) and the $\alpha 2(\text{IV})$ gene (COL4A2) are located on chromosome 13 (Pihlajaniemi et al., 1985; Boyd et al., 1988; Griffin et al., 1987); COL4A3 and COL4A4 are on chromosome 2 (Mariyama et al., 1994; Kamagata et al., 1992); and COL4A5 and COL4A6 are on chromosome X (Hostikka et al., 1990; Zhou et al., 1993; Oohashi et al., 1994). Further, COL4A1 and COL4A2 are aligned in a head-to-head fashion sharing a common promoter between the two genes (Poschl et al., 1988; Soininen et al., 1988), whereas the precise upstream structure of COL4A3 and COL4A4 has not been reported yet. COL4A5 and COL4A6 have been found together on chromosome X at the segment q22 (Hostikka et al., 1990; Zhou et al., 1993; Oohashi et al., 1994) and have been reported to be also arranged in a head-to-head fashion and to share a bidirectional promoter (Zhou et al., 1993). Furthermore, we reported the novel observation that the COL4A6 gene is transcribed from two alternate promoters in a tissue-specific fashion (Sugimoto et al., 1994).

Presently we report for the first time the presence of the COL4A6 gene products as detected by Western blotting and immunofluorescence staining of various tissues by use of peptide-specific monoclonal antibodies. Our results demonstrate that the COL4A5 and COL4A6 genes are not necessarily coexpressed in several types of tissues.

Materials and Methods

Synthetic Peptides

Earlier we determined the complete primary structure of the human $\alpha 6(\text{IV})$ chain (Oohashi et al., 1994; Zhou et al., 1994). When the amino acid sequences of all six $\alpha(\text{IV})$ chains were compared, their noncollagenous (NC)1 domain portions showed the highest homology with each other. However, the amino acid sequences of certain subregions of NC1 were quite different from one another. We selected the peptide TTVEERQQFGELPVSSETLKAGQLHTRV from the $\alpha 6(\text{IV})$ chain (Oohashi et al., 1994) and ATVDVSDMFSKQPQSEILKAGDLRTRIS from the $\alpha 5(\text{IV})$ chain (Hostikka et al., 1990), both of which are close to the carboxyl ends, as shown in Fig. 1. We also synthesized a peptide, CPSTGELEFMGFPGK, derived from the third imperfection of the Gly-X-Y repeating sequence in the COL1 domain of the $\alpha 6(\text{IV})$ chain. A cysteinyl residue was added at the amino terminus to allow coupling of the peptide to the carrier protein. Oligopeptides were chemically synthesized by the solid-phase procedure of Merrifield (1963) with the aid of an automated peptide synthesizer (model 430; Applied Biosystems, Inc., Foster City, CA). The synthetic peptides were conjugated to a carrier protein, keyhole

1. Abbreviations used in this paper: COL, collagen; KLH, keyhole limpet hemocyanin; NC, noncollagenous; PVDF, polyvinylidene difluoride.




Figure 1. Schematic representation of the two genes COL4A5 and COL4A6 on chromosome Xq22. (*Top*) COL4A5 and COL4A6 are arranged head-to-head in close proximity on chromosome Xq22 (Sugimoto et al., 1994). Arrows indicate the direction of the transcripts for COL4A5 and COL4A6. There are two transcription start sites for COL4A6. One transcription start site (from exon 1') for COL4A6 is 442 bp away from the transcription start site of COL4A5, while an alternative transcription start site (from exon 1) is located 1050 bp apart from the first one and drives the expression of a second transcript that encodes an $\alpha 6(\text{IV})$ chain with a different signal peptide. (*Bottom*) Because of the direction of the two genes, the translation products, $\alpha 5(\text{IV})$ and $\alpha 6(\text{IV})$ chains, of the two genes are drawn accordingly. Chain-specific monoclonal antibodies were raised against synthetic peptides (*short bars*) of the NC1 domains close to the carboxyl termini of the two chains. The third peptide deduced from the third imperfection of the Gly-X-Y repeating sequence in the central COL1 domain was synthesized and used for monoclonal antibodies. The sequences of the three synthetic peptides are shown with cysteinyl residues at the amino termini. Using the Geysen's peptide scanning technique (Geysen et al., 1987) we determined the exact location of the epitopes for H61, H62, and H63. Relative location of the epitopes are shown by three thick bars under the $\alpha 6(\text{IV})$ peptide.

limpet hemocyanin, by the maleimide method (Ishikawa et al., 1983). In some experiments the peptides were directly used as antigen without conjugation to hemocyanin, and the antigenic efficiency was found not to differ much. Other peptides, ATIERSEMFKKPTSTLKAGELRTHVS from the $\alpha 1(\text{IV})$ chain (Pihlajaniemi et al., 1985), TTPEQSFQGSFADTLKAGLIRTHIS from the $\alpha 2(\text{IV})$ chain (Hostikka and Tryggvason, 1988), ASLNPERMFRKPIPSTVKAGELEKIIS from the $\alpha 3(\text{IV})$ chain (Morrison et al., 1991), and TVKADFEFSSAPADTLKESQAQRQKI from the $\alpha 4(\text{IV})$ chain (Sugimoto et al., 1993), were used for raising chain-specific antibodies against the other human α chains.

Peptide-specific Monoclonal Antibodies

We introduced new steps to our previous method and developed a new efficient procedure, the rat lymph node method, for raising monoclonal antibodies (Kishiro et al., 1995). Briefly, WKY/NCrj rats (Charles River Japan, Yokohama, Japan) were immunized in the hind footpads with 50 mg of hemocyanin-coupled synthetic peptide emulsified with Freund's complete adjuvant. Three or four weeks later the rats were killed, and lymphocytes obtained from the medial iliac lymph nodes of the rats were fused with mouse myeloma cells (SP2/0-Ag14). Supernatants from hybridoma cultures were screened by ELISA, using hemocyanin-free peptides from individual $\alpha(\text{IV})$ chains.

The initial screening was performed by ELISA with the synthetic peptides themselves and/or native NC1 fractions isolated from the human kidney and other peptides of the same region from all $\alpha(\text{IV})$ NC1 domains. 2,000 clones were screened and we picked 145 positive clones that reacted specifically with the $\alpha 6(\text{IV})$ peptide. The second screening was performed by indirect immunofluorescence using human tissue sections. Thus, three

clones, H61, H62, and H63, out of the 12 positive clones for both ELISA and immunofluorescence were established.

Preparation of the NC1-Domain Fraction from Renal Basement Membrane

Renal basement membrane was prepared by the same method as that used for bovine renal basement membrane (Sado et al., 1991). In order to recover the NC1 domains from the collagen IV molecules we used bacterial collagenase and removed the central collagenous (COL) domains from the molecules. Namely, lyophilized basement membrane was solubilized with collagenase (Seikagaku Kogyo, Tokyo, Japan) at 47°C for 20 h by a method similar to that used for bovine nephritogenic antigen. After solubilization, the sample was centrifuged at 27,000 g for 10 min and insoluble material was removed. The supernatant was dialyzed against distilled water for 8 h, lyophilized, dissolved in PBS, and dialyzed against PBS overnight. The material was then centrifuged to remove insoluble materials formed during the dialysis. The supernatant was applied to a gel filtration column of UltrogelAcA 34(26 × 950 mm) at a flow rate of 30 ml/h. The main peak fraction was collected and concentrated.

Western Blotting

The NC1 fractions of collagen type IV thus prepared were analyzed by SDS-polyacrylamide gel electrophoresis under nonreducing conditions (Laemmli, 1970). SDS-containing 5% stacking and 11.5% homogeneous running polyacrylamide slab gels were used. Proteins separated on the SDS gels were blotted onto polyvinylidene difluoride (PVDF) membranes. The membrane was incubated with the primary antibodies first and then with the peroxidase-conjugated secondary antibodies (Sado et al., 1991).

Immunohistochemical Staining

Indirect immunofluorescence staining was performed as described previously (Sado et al., 1991). In the preliminary experiments we have tested different conditions varying the concentration of the urea, pH, incubation time, and temperature using the representative tissues including kidney, muscles, and skin. We set up to the standard acid-urea denaturation condition: incubating ethanol-fixed sections with 6 M urea-HCl buffer (pH 3.5) for 1 h at room temperature. 6 M urea treatment is necessary for most of the antibodies probably due to the linearized synthetic peptide antigens. Control sections for immunofluorescence staining were reacted with nonimmune rat sera or with the secondary antibody alone. These controls were negative.

The staining method for the kidney and skin. Normal portions of human kidneys were obtained from two patients with renal tumors (both males, 56 and 85 yr old). Renal biopsies were performed from two Alport's syndrome patients (males, 14 and 9 yr old). Skin specimens were obtained from the patients with Alport's syndrome and from patients with non-renal diseases by surgical biopsy.

The tissues from the kidney and skin were snap-frozen in liquid nitrogen. 3-mm cryosections were fixed with acetone for 10 min. To reveal possible masked epitopes, we pretreated the sections with 6 M urea in 0.1 M glycine-HCl buffer (pH 3.5) for 10 min at room temperature. After having been washed with PBS, the sections were incubated with monoclonal antibodies for 1 h at room temperature. After another washing with PBS, they were incubated with FITC-labeled goat anti-rat IgG (Cappel Labs., Cochranville, PA) for 1 h, and then were examined under a fluorescence microscope (Axiophot, Zeiss, Germany). Control sections were treated by the same protocol except for the step of the incubation with the monoclonal antibody.

The staining for the extrarenal tissues. Various organs obtained from an autopsy case (64-yr-old female) were used as normal tissues. Acid-urea denaturation was performed by incubating acetone-fixed sections with 0.1 M glycine-HCl buffer (pH 3.5) containing 6 M urea for 2 to 30 min at room temperature. Positive control sections were stained with anti- α 1(IV) or α 2(IV) collagen antibody, as it is well known that both α 1(IV) and α 2(IV) chains are usually colocalized in basement membrane. Negative control sections were reacted with non-immune rat sera, with the secondary antibody alone, or with the primary antibody preabsorbed with the synthetic peptide.

Alport's Syndrome Patients

We examined two cases of Alport's syndrome diagnosed by clinical mani-

festations and histological examinations. One case, a 14-yr-old boy, had proteinuria, hearing loss, and eye lesions. Electron microscopic examination of the glomeruli showed basement membrane thickening with lamination and splitting. The second case, a 9-yr-old boy, had proteinuria (0.2–0.6 g/d) and hematuria since the age of three years. He displayed farsightedness of the eyes and hearing loss as well. Several relatives had renal diseases, and some of them had died. Histological and electron microscopic examination of the kidney demonstrated irregular thickening of the glomerular basement membrane, confirming the diagnosis.

Results


Production of α 6(IV) Chain-specific Antibodies

We selected a relatively nonconserved region, close to the carboxy termini of the NC1 domains, as a pattern to prepare synthetic peptides for use as immunogens to elicit α (IV) chain-specific monoclonal antibodies. Keyhole limpet hemocyanin (KLH)-conjugated peptide emulsified with Freund's complete adjuvant was injected into hind footpads, and medial iliac lymph nodes of the immunized rat were used for fusion with myeloma cells. For α 6(IV)-specific antibodies we repeated the cell fusion five times. In each experiment fused cells were plated on four 96-well plates. 145 positive wells from the five experiments were obtained by direct reaction with the synthetic peptides. We further screened by staining frozen sections of human kidney and selected 12 strongly positive clones, which were then tested with the synthetic peptides from the same region of the other α 1 through α 5 chains by ELISA. All of the 12 clones were specific for the α 6(IV) chain. Three antibody clones out of these 12 clones for the α 6(IV) peptide, H61, H62, and H63, were established after subcloning by limiting dilution. Monoclonal antibodies H11, H21, H31, H43, and H52 to the human α 1(IV), α 2(IV), α 3(IV), α 4(IV), and α 5(IV) chains, respectively, were also obtained in the study. Significant similarity was not found between the peptide sequences used for antibodies and the known amino acid sequence in proteins by DDBJ or Swiss Plot sequence analysis. The specificity of the antibodies against the synthetic peptides was tested by ELISA. A typical specific reaction of each antibody to individual antigen peptides is demonstrated in Fig. 2. Using the Geysen peptide scanning technique (Geysen et al., 1987), we narrowed down the epitopes for H61, H62, and H63 to the amino acid sequences of VSETLK, FGEL, and GELP, respectively, within the synthetic peptide sequence. Relative location of the epitopes for H61, H62, and H63 is drawn in Fig. 1.

Immunoblot Analysis

The structure of the entire human α 6(IV) collagen polypeptide was predicted by cDNA sequence analysis (Oohashi et al., 1994). The deduced collagen polypeptide contained 1690 amino acid residues, including a 21-residue signal peptide, a 24-residue amino-terminal NC domain, a central 1417-residue collagenous (COL1) domain, and a 228-residue carboxy-terminal NC1 domain. To detect the presumptive peptide at the protein level we utilized monoclonal antibody H63, specific for the cDNA-derived polypeptide that was mentioned above. Kidney was used as the starting material for Western blotting analysis. Since the predicted α 6(IV) collagen polypeptide structure was simi-

A



indicated by the shaded area. (B) The synthetic peptides used as immunogens for α chain-specific monoclonal antibodies were fixed onto the well bottoms of a 96-well plate. *Top row*: Control, only bovine serum albumin; *second row*: $\alpha 1(\text{IV})$ peptide; *third row*: $\alpha 2(\text{IV})$ peptide; *fourth row*: $\alpha 3(\text{IV})$ peptide; *fifth row*: $\alpha 4(\text{IV})$ peptide; *sixth row*: $\alpha 5(\text{IV})$ peptide; and *seventh row*: $\alpha 6(\text{IV})$ peptide. Nonimmune serum (7 wells in lane 1), monoclonal antibodies of H11 (lane 2), H21 (lane 3), H31 (lane 4), H43 (lane 5), H52 (lane 6), H63 (lane 7), and dH2O (lane 8) were added to the wells individually and incubated for 1 h. After the unbound antibodies had been washed out, the individual wells were blocked with bovine serum albumin. The peroxidase-conjugated secondary antibodies were added to all of the wells and reacted for 1 h. Note the specific reactions between the antigens and antibodies even though there is some similar sequence between peptide sequences used for raising the monoclonal antibodies.

B




Figure 2. Specificity of the $\alpha(\text{IV})$ chain-specific antibodies. (A) The carboxy-terminal ~ 40 amino acid residues of the $\alpha 1(\text{IV})$, $\alpha 2(\text{IV})$, $\alpha 3(\text{IV})$, $\alpha 4(\text{IV})$, $\alpha 5(\text{IV})$, and $\alpha 6(\text{IV})$ chains are aligned to have the maximum homology. Dashes indicate the amino acid residues that are the same as for the $\alpha 1(\text{IV})$ chain. Gaps (-) are introduced to maintain alignment. The amino acid sequences of peptides used as immunogens for α chain-specific monoclonal antibodies are

lar to that of the other known α chains of collagen IV, we digested away the central collagenous domain by bacterial collagenase; and the materials were then electrophoresed and transferred to PVDF membranes. The membrane strips containing the blotted protein were incubated with antibody H63 made against the $\alpha 6(\text{IV})$ -derived peptide (see Materials and Methods). As shown in Fig. 3, the antiserum stained a band of ~ 27 kDa in size, which is consistent with the NC1 size of the cDNA-derived $\alpha 6(\text{IV})$: 228 residues with a calculated molecular weight of 25,418 (Oohashi et al., 1994). This staining was blocked by preincubation of the antiserum with the antigen (data not shown). Preimmune serum was completely negative. When the other five antibodies against $\alpha 1(\text{IV})$ through $\alpha 5(\text{IV})$ peptides were reacted as controls with the same materials, several bands similar in size to those seen with $\alpha 6(\text{IV})$ sera and relatively distinct bands detected by $\alpha 1$ to $\alpha 5$ sera were obtained. The size of these bands was also reason-

able for each NC1 domain. Each antibody also detected some bands at ~ 50 kDa, which we consider to be the dimeric forms (Johansson et al., 1992). Thus, $\alpha 6(\text{IV})$ antibody reacted indeed with the NC1 domain of the $\alpha 6(\text{IV})$ polypeptide, which is the first demonstration of the presence of the polypeptide in human kidney.

Indirect Immunofluorescence Staining of Kidney Basement Membranes with Monoclonal Antibody against $\alpha 6(\text{IV})$ -derived Peptide

Immunofluorescence studies on frozen sections of human adult kidney with the $\alpha 6(\text{IV})$ peptide antiserum surprisingly gave a negative reaction in the glomerular basement membrane, whereas the basement membrane of the Bowman's capsules and of some tubules was positive, as shown in Fig. 4 F. The staining was completely blocked by preincubation of the antibody with the $\alpha 6(\text{IV})$ peptide used for raising antibody, H63 (Fig. 5 B). This staining pattern was quite in contrast with that obtained with the $\alpha 5(\text{IV})$ peptide antiserum (Fig. 4 E), in which basement membranes of glomerulus, Bowman's capsules, and some tubules were all positive. Thus, the two genes COL4A5 and COL4A6, controlled by a bidirectional promoter, are not always expressed at the same time; the cells that produce glomerular basement membrane express the $\alpha 5(\text{IV})$ gene but not the $\alpha 6(\text{IV})$ one at least in adult humans. Control experiments confirmed the specificity of these staining reactions. No reaction was obtained with the preimmune sera (data not shown). In contrast, antisera against $\alpha 1(\text{IV})$ and $\alpha 2(\text{IV})$ peptides gave a strong and similar staining pattern of all basement membranes of the glomeruli, tubules, and Bowman's capsules as well (Fig. 4, A and B). Intriguingly, the staining pattern for $\alpha 3(\text{IV})$ and $\alpha 4(\text{IV})$ antibodies was the same; both strongly stained the glomerular basement membrane and some distal tubules but mostly gave faintly positive reactions for basement membranes of Bowman's capsules (Fig. 4, C and D). The staining pattern for the kidney is summarized in Table I.




Figure 3. Western-blot analysis for the NC1-domain fraction from renal basement membrane. The NC1-domain fraction was prepared from renal basement membrane (see Materials and Methods), and SDS-PAGE was conducted on 11.5% polyacrylamide gels. After blotting of the proteins onto a PVDF membrane, the membrane was sliced into seven strips.

Lanes 1 and 8 were stained with Coomassie-Brilliant Blue, and the rest of them were incubated with the individual α chain-specific monoclonal antibodies. Lanes 2-7 were incubated with H11, H21, H31, H43, H52, and H63, respectively. Note the sizes of the reactive peptides are different from strip to strip. The sizes of the peptides for lane 6 (for $\alpha 5(\text{IV})$ NC1) and 7 (for $\alpha 6(\text{IV})$ NC1) are estimated as ~ 26 and 27 kDa, respectively. Several reactive peptides around the size of 50 kDa probably represent dimeric forms.




Figure 4. Differential staining of kidney basement membranes obtained with α (IV) chain-specific antibodies. α (IV) chain-specific antibodies were used to stain kidney basement membranes. Antibodies of H11, H21, H31, H43, H52, H63, and H64 were used for photos A–F, respectively. Note that α 1(IV) and α 2(IV) chains are coexpressed in all of the basement membranes in the kidney: glomeruli, Bowman’s capsules, distal tubules, proximal tubules, and some capillaries. Antibodies specific for α 3(IV) and α 4(IV) chains demonstrated rather restricted staining pattern, i.e., glomerular, some tubular basement membranes; however these two chains are expressed together. Interestingly, α 5(IV) and α 6(IV) chains are not coexpressed; no staining for α 6(IV) chain is seen in glomerular basement membrane (F), whereas α 5(IV) chains are clearly expressed in the glomerulus (E).

To confirm that the monoclonal antibodies are recognizing the α 6(IV) chain, we have done two experiments. One of them was to define the epitopes of the H61, H62, and H63 antibodies using Geysen’s technique (Geysen et al., 1987). As mentioned above, the epitope sequences of VSETLK, FGEL, and GELP for H61, H62, and H63 antibodies, respectively, were all within the original peptide sequence but different from each other. However, the im-

munostaining pattern using the antibodies for the kidney sections were the same (data not shown). This suggested that the different antibodies recognizing different epitopes were reacting with the same molecule. The second experiment was to raise monoclonal antibodies to the peptide sequence within the COL1 domain of the α 6(IV) chain. We selected CPSTGELEFMGFPPKG in the third imperfection of the Gly-X-Y repeating sequence (Oohashi et al., 1994)

Table I. Distribution of Type IV Collagen α Chains in Kidney Basement Membranes

	$\alpha 1(IV)$	$\alpha 2(IV)$	$\alpha 3(IV)$	$\alpha 4(IV)$	$\alpha 5(IV)$	$\alpha 6(IV)$
Glomerular BM	++	++	+++	+++	++	-
Mesangium	++	++	-	-	-	-
Bowman's BM	+++	+++	\pm	\pm	++	++
Tubular BM	++	++	+++*	+++*	+++*	+++*
Capillary BM	+	+	-	-	-	-

+, weakly positive; ++, moderately positive; +++, strongly positive; -, negative.
*Not all the tubular basement membranes were positive for $\alpha 3(IV)$, $\alpha 4(IV)$, $\alpha 5(IV)$, and $\alpha 6(IV)$ antibodies.

within the COL1 domain. Among 11 positive clones by ELISA, the two clones, H64 and H65, were subcloned and purified. The immunostaining pattern for the kidney section using these two clones was the same as that of H63 shown in Fig. 5. These results indicate that the different antibodies derived from the different sequences still recognize the same material, the $\alpha 6(IV)$ chain, in the tissues.

Differential Localization of the $\alpha 5(IV)$ and $\alpha 6(IV)$ Chains in Tissues Outside of the Kidney

To investigate whether the two genes are expressed in basement membranes in tissues other than the kidney, we stained several other organs and tissues, as shown in Fig. 6.

Skin. Epidermal basement membranes are thought to be produced by the keratinocytes, aligning at the base of the epithelial layers. Since the staining pattern for the $\alpha 1(IV)$ and $\alpha 2(IV)$ chains were almost the same, only the $\alpha 2$ staining pattern is shown in Fig. 6 A. As shown in Fig. 5, B and C, this basement membrane was stained by both antibodies against $\alpha 5(IV)$ and $\alpha 6(IV)$ chains, indicating that the keratinocytes express both genes together in human adult skin.

Muscle. Esophageal smooth muscle, cardiac muscle, and abdominal skeletal muscle were examined by using anti-

$\alpha 1(IV)$, $\alpha 2(IV)$, $\alpha 5(IV)$, and $\alpha 6(IV)$ antibodies. Both $\alpha 5(IV)$ and $\alpha 6(IV)$ chains colocalized in the basement membranes of the smooth muscle cells (Fig. 6, E and F). However, these two chains were never detected in cardiac muscle cells (Fig. 6, H and I), although both $\alpha 1(IV)$ and $\alpha 2(IV)$ chains were stained substantially (Fig. 6 G). Unexpectedly, in basement membranes surrounding the skeletal muscle cells, $\alpha 5(IV)$ was negative and $\alpha 6(IV)$ was weakly positive (Fig. 6 L) when the sections were stained with H-61 antibody. Further, this staining was blocked by incubating the antibody with the synthetic peptide of the $\alpha 6(IV)$ chain. Relative to the $\alpha 2(IV)$ staining, the $\alpha 6$ staining intensity was quite weak; therefore the level of the $\alpha 6(IV)$ chain is presumed to be fairly low.

Adipocytes. Adipocytes expressed both $\alpha 1(IV)$ and $\alpha 2(IV)$ chains strongly (Fig. 6 M). Since $\alpha 5(IV)$ and $\alpha 6(IV)$ staining was quite low (Fig. 6, N and O), we had to expose the film longer to get a clear staining pattern. The level of $\alpha 5(IV)$ and $\alpha 6(IV)$ expression around adipocytes also thus appears to be quite low.

Liver. $\alpha 2(IV)$ chain was present in the space of Disse (Fig. 6 P); however, neither $\alpha 5(IV)$ nor $\alpha 6(IV)$ chains were detected in this space underneath the sinusoidal endothelial cells (Fig. 5, Q and R).

Table II provides a summary of the above-mentioned staining patterns.

$\alpha 5(IV)$ and $\alpha 6(IV)$ Chains in Basement Membranes of Alport's Syndrome Patient

Immunohistochemical analysis was performed in kidney and skin from patients with X-linked Alport's syndrome. As shown in Fig. 7, antibodies against $\alpha 1(IV)$ (photo not shown) and $\alpha 2(IV)$ chains clearly stained basement membranes in glomeruli, Bowman's capsules, and tubules. However, neither of the antibodies against the $\alpha 5(IV)$ and $\alpha 6(IV)$ chains stained the basement membranes in the Alport kidney (Fig. 7, B and C). Antibodies against




Figure 5. Staining of kidney basement membranes by $\alpha 6(IV)$ antibodies. Serial sections of human kidney were stained by the two monoclonal antibodies to the different part of the $\alpha 6(IV)$ chain: one (H63) is to the synthetic peptide from the NC1 domain (A) and the other (H65) is to the COL domain (C). Note the staining pattern using the two monoclonal antibodies is the same. Positive fluorescence staining using H63 antibody was completely blocked by the original synthetic peptide, TTVEER-QQFGELPVSETLKAGQLH-TRV(B). D shows nonstained phase contrast microscopic image of the neighboring section.

Table II. Distribution of Type IV Collagen $\alpha 1(IV)$, $\alpha 2(IV)$, $\alpha 5(IV)$, $\alpha 6(IV)$ Chains in Basement Membranes of Extrarenal Tissues

	$\alpha 1(IV)$	$\alpha 2(IV)$	$\alpha 5(IV)$	$\alpha 6(IV)$
Skin	+++	+++	+++	+++
Muscle				
Skeletal muscle	++	++	-	+
Cardiac muscle	++	++	-	-
Smooth muscle	++	++	++	++
Adipocyte	++	++	+	+
Liver	+	+	-	-

+, weakly positive; ++, moderately positive; +++, strongly positive; -, negative.

$\alpha 3(IV)$ and $\alpha 4(IV)$ chains also did not give positive staining in basement membranes in the kidney of the two patients examined (data not shown).

In Fig. 8, distribution of $\alpha(IV)$ chains in skin specimens from a normal individual and an Alport patient is shown. Antibodies against $\alpha 1(IV)$ and $\alpha 2(IV)$ chains demonstrated a clear linearized staining pattern in the epidermal basement membrane and basement membranes around capillaries and sweat glands in normal skin. $\alpha 5(IV)$ and $\alpha 6(IV)$ chains were detected only in the basement membrane of the epidermis but not in basement membranes around capillaries and sweat glands. In contrast to the normal skin, Alport skin showed negative staining with $\alpha 5(IV)$ (Fig. 8 H) and $\alpha 6(IV)$ (Fig. 8 I) antibodies in epidermal basement membranes, whereas $\alpha 1(IV)$ (data not shown) and $\alpha 2(IV)$ (Fig. 8 G) antibodies clearly stained the basement membranes.

Discussion

Diversity has been revealed among the subunits that form collagen IV molecules with the discovery of the $\alpha 3$, $\alpha 4$, $\alpha 5$, and $\alpha 6$ chains (Hudson et al., 1993). But chain assemblies that make up type IV molecules have been difficult to demonstrate since the $\alpha 3$ - $\alpha 6(IV)$ chains were studied primarily in the context of the disease-related human tissues. The $\alpha 3(IV)$ chain was discovered as the Goodpasture antigen, which attacks primarily the alveolar basement membranes in the lung and the glomerular basement membranes in the kidney (Butkowski et al., 1987). Further, trials to purify the $\alpha 3(IV)$ chain resulted in the discovery of another α -chain, $\alpha 4(IV)$ (Gunwar et al., 1990). cDNA cloning of these two chains led to the identification of mutations in COL4A3 and COL4A4 in patients with autosomal recessive type Alport syndrome (Mochizuki et al., 1994; Lemmink et al., 1994). Most recently COL4A6 was


shown to be located next to COL4A5 (Zhou et al., 1993; Sugimoto et al., 1994) and both genes were mutated in several X-linked cases of Alport syndrome combined with leiomyomatosis (Zhou et al., 1993).

Immunohistochemical analyses have shown that the $\alpha 1(IV)$ and $\alpha 2(IV)$ chains are abundant in all tubular basement membranes and in the glomerular mesangial matrix (Kleppel et al., 1989). However the $\alpha 3$ - $\alpha 5(IV)$ collagen chains have a rather limited distribution in human tissues; highly enriched in glomerular basement membranes but found only in a subset of tubular basement membranes (Hostikka et al., 1990; Sanes et al., 1990; Hudson et al., 1992; Yoshioka et al., 1994). Just recently, Miners and Sanes (1994) reported the distribution of $\alpha 3$, $\alpha 4$, and $\alpha 5(IV)$ collagen chains in rodent basal laminae. They suggested from their immunohistochemical studies of various tissues that many combinations of $\alpha(IV)$ chains were possible, but $\alpha 3$ and $\alpha 4(IV)$ chains were always coexpressed together, and appeared only in basal laminae that were $\alpha 5(IV)$ chain positive. However, they did not examine the distribution of the newly discovered $\alpha 6(IV)$ chain with respect to the other $\alpha(IV)$ chains. In the light of gene regulation of the two neighboring genes, COL4A5 and COL4A6, we herein examined and focused on the expression of the two genes at the protein level.

We first raised monoclonal antibodies specific for the human $\alpha 6(IV)$ collagen chain and used them with antibodies produced by a previously isolated clone for $\alpha 5(IV)$ chain to identify location of $\alpha 5(IV)$ and $\alpha 6(IV)$ chains in basement membranes of kidney, skin, muscle, liver, and fat cells. In addition, we used the monoclonal antibodies to identify the NC1 domains of $\alpha 5(IV)$ and $\alpha 6(IV)$ by Western blotting.

Our immunohistochemical survey with monoclonal antibodies to $\alpha 5(IV)$ and $\alpha 6(IV)$ chains indicated a very restricted distribution of these proteins in basement membranes. The basement membrane in the glomerulus is physiologically and clinically a carefully studied matrix. This basement membrane contains specialized structures that face cell layers of epithelial cells and endothelia or other cells on both surfaces. As shown in Fig. 4, the glomerular basement membranes were strongly positive when stained with $\alpha 5(IV)$ chain-specific antibody, H51; however, antibody against the $\alpha 6(IV)$ peptide did not show a fluorescence staining within the glomerular basement membrane. In contrast, basement membranes in Bowman's capsule and distal tubules were positive with both antibodies. There are at least two and possibly more different molecular forms in the glomerulus: [$\alpha 1(IV)$]₂ $\alpha 2(IV)$ and one or more molecular forms of the combination of $\alpha 3$, $\alpha 4$, and $\alpha 5$ chains. In an Alport's nephritis case, neither of the

Figure 6. Staining of basement membranes in extrarenal tissues using $\alpha 1(IV)/\alpha 2(IV)$, $\alpha 5(IV)$, and $\alpha 6(IV)$ antibodies. Expression of $\alpha 5(IV)$, and $\alpha 6(IV)$ chains was examined in basement membranes in extrarenal tissues; skin (A-C), smooth muscle cells (D-F), cardiac muscle cells (G-I), abdominal skeletal muscle cells (J-L), adipocytes (M-O), and liver (P-R). Since antibodies for $\alpha 1(IV)$ and $\alpha 2(IV)$ chains gave quite similar patterns, only the $\alpha 2(IV)$ staining pattern using H21 is shown in A, D, G, J, M, and P. Expression of $\alpha 5(IV)$ (B, E, H, K, N, and Q, using H52 antibody) and $\alpha 6(IV)$ (C, F, I, L, O, and R, with H63 antibody) chains is compared in various basement membranes. Around the smooth muscle cells and at the epidermal/dermal junction, the $\alpha 5(IV)$ and $\alpha 6(IV)$ chains are coexpressed, but no expression of these genes was detected in cardiac muscle and liver. Intriguingly, only $\alpha 6(IV)$ is weakly stained in skeletal muscle, whereas no staining is detected for $\alpha 5(IV)$ chain.






Figure 7. Distribution of α (IV) chains in kidney from Alport's syndrome patient. Similar staining pattern was obtained with α 1(IV) (not shown) and α 2(IV) (A) antibodies, but no clear staining is shown in any basement membranes in the kidney sections stained with monoclonal antibodies for α 5(IV) (B, with H52) and α 6(IV) (C, with H63) chains.

three chains, α 3, α 4, and α 5, were found in the glomerulus (Fig. 6). We do not know the precise mutation of α (IV) gene in this case. However, since more than 50 different mutations have been identified in the COL4A5 gene but none in the COL4A6 gene, it is possible that a heterotrimeric molecule composed of the three chains, α 3, α 4, and α 5, cannot be formed due to abnormal α 5 chain synthesized from a mutated COL4A5 gene.

In addition to the human and murine α 1(IV) and α 2(IV) collagen genes (Killen et al., 1988; Soininen et al., 1988), there are several clustered genes in vertebrates that are organized in a head-to-head configuration and directed by bidirectional promoters. These include the histone H2a and H2b genes (Hentschel and Birnstiel, 1981), DHFR and mismatch repair protein 1 gene (Fujii et al., 1992), Wilms' tumor locus (Huang et al., 1990), a proliferation cell nuclear antigen gene (Rizzo et al., 1990), an SV40-like monkey genomic locus (Saffer et al., 1984), the GPAT/AIRC genes (Gavalas et al., 1993; Gavalas and Zalkin, 1995), and *Drosophila* ras2/rop genes (Lightfoot et al., 1994). Some of the promoters of these genes are bidirectional but the opposite genes are not identified. The promoter activity for one direction is quite different from that for the other direction, 10-fold different in some cases (Gavalas and Zalkin, 1995). Most of the paired genes are not homologous at all, however, the COL4A1/COL4A2 and COL4A5/COL4A6 genes are homologous, indicating that the clusters did evolve by gene duplication. Divergent transcription of the two genes from a bidirectional promoter can thus provide coexpression and coregulation of expression in some tissues. Proximal *cis*-acting elements as well as *trans*-acting factors such as CTC binding factor (Fischer et al., 1993) have the potential to modulate the expression of both genes simultaneously.

The genes for α 5(IV) and α 6(IV) chains are encoded on

opposite DNA strands and have a common promoter region (Sugimoto et al., 1994). A question that arose immediately when we noticed the specialized relation of these genes was whether these genes always produce the two chains at a steady rate as COL4A1 and COL4A2 genes do. As we demonstrated in Fig. 6, keratinocytes, esophageal smooth muscle cells, fat cells, striated muscle cells and placental epithelial cells express α 5(IV) and α 6(IV) genes, whereas hepatocytes and cardiac muscle cells do not express either gene. However, of interest was that skeletal muscle cells only synthesized α 6(IV) chain and that the glomerular basement membranes were positive with only α 5(IV) antibody in adult humans. These results are consistent with the analysis of transcripts found in a variety of tissues except for the heart (Zhou et al., 1993), although we have not analyzed tissues from human fetuses. The two genes are transcribed in the opposite direction in a tissue-specific manner, and regulation of expression for the two genes is controlled at the transcription level. One of the clues for the gene expression could be the presence of the alternative promoters on the COL4A6 gene (Sugimoto et al., 1994). The two different transcription forms for the α 6(IV) gene driven by the alternative promoters are found in a tissue-specific manner. The tissue-specific distribution of the newly discovered α 5(IV) and α 6(IV) chains are probably related to the specialized function of individual basement membranes in different tissues.

We are grateful to Dr. H. Yoshioka for his help and continued interest in the project.

This work was supported by a Grant-in-Aid for Scientific Research (B), 06454250, from the Ministry of Education, Science and Culture of Japan, the Nakatomi Foundation, and by National Institutes of Health grant EY07334 from the National Eye Institute.

Received for publication 19 November 1994 and in revised form 4 April 1995.




Figure 8. Distribution of α (IV) chains in dermal basement membrane from Alport syndrome patient. Since a similar staining pattern was obtained with α 1(IV) and α 2(IV) antibodies in basement membranes of dermal epidermal junction and capillaries, only the α 2(IV) staining pattern (A, with H21) is shown in normal skin. Both α 5(IV) (B, H52) and α 6(IV) (C, H63) antibodies stain only the dermal basement membrane but not that of the capillaries. In the Alport patient case, a similar staining pattern was obtained for α 1(IV) (not shown) and α 2(IV) chains as in normal skin (G); however no clear staining was shown by monoclonal antibodies for α 5(IV) (H) and α 6(IV) (I) chains at all (H and I). Phase-contrast images are shown for individual samples in normal skin (D–F) and Alport skin (J–L).

References

- Barker, D. F., S. L. Hostikka, J. Zhou, L. T. Chow, A. R. Oliphant, S. C. Gerken, M. C. Gregory, M. H. Skolnick, C. L. Atkin, and K. Tryggvason. 1990. Identification of mutations in the COL4A5 collagen gene in Alport syndrome. *Science (Wash. DC)*. 248:1224–1227.
- Boyd, C. D., K. Weliky, S. B. Deak, A. M. Christiano, J. M. Mackenzie, L. Sandell, K. Tryggvason, and E. Magenis. 1988. The single copy gene coding for human α 1(IV) procollagen is located at the terminal end of the long arm of chromosome 13. *Human Genet.* 74:121–125.
- Butkowski, R. J., J. P. M. Langeveld, J. Wieslander, J. Hamilton, and B. G. Hudson. 1987. Localization of the Goodpasture epitope to a novel chain of basement membrane collagen. *J. Biol. Chem.* 262:7874–7877.
- Butkowski, R. J., G.-Q. Shen, J. Wieslander, A. F. Michael, and A. J. Fish. 1990. Characterization of type IV collagen NC1 monomers and Goodpasture antigen in human renal basement membranes. *J. Lab. Clin. Med.* 115:365–373.
- Fischer, G., C. Schmidt, J. Opitz, Z. Cully, K. Kuhn, and E. Poschl. 1993. Identification of a novel sequence element in the common promoter region of human collagen type IV genes, involved in the regulation of divergent transcription. *Biochem. J.* 292:687–695.
- Fujii, H., E. Shinya, and T. Shimada. 1992. A GC box in the bidirectional promoter is essential for expression of the human dihydrofolate reductase and mismatch repair protein 1 genes. *FEBS (Fed. Eur. Biochem Soc.) Lett.* 314: 33–36.
- Gavalas, A., and H. Zalkin. 1995. Analysis of the chicken GPAT/AIRC bidirectional promoter for de novo purine nucleotide synthesis. *J. Biol. Chem.* 270:

- 2403–2410.
- Gavalas, A., J. E. Dixon, K. A. Brayton, and H. Zalkin. 1993. Coexpression of two closely linked avian genes for purine nucleotide synthesis from a bidirectional promoter. *Mol. Cell. Biol.* 13:4784–4792.
- Geysen, H. M., S. J. Rodda, T. J. Mason, G. Tribbick, and P. G. Schoofs. 1987. Strategies for epitope analysis using peptide synthesis. *J. Immunol. Methods.* 102:259–274.
- Griffin, C. A., B. S. Emmanuel, J. R. Hansen, W. K. Canevee, and J. C. Myers. 1987. Human collagen genes encoding basement membrane $\alpha 1(IV)$ and $\alpha 2(IV)$ chains map to the distal long arm of chromosome 13. *Proc. Natl. Acad. Sci. USA.* 84:512–516.
- Gunwar, S., J. Saus, M. E. Noelken, and B. G. Hudson. 1990. Glomerular basement membrane: identification of a fourth chain, $\alpha 4$, of type IV collagen. *J. Biol. Chem.* 265:5466–5469.
- Hentschel, C. C., and M. L. Birnstiel. 1981. The organization and expression of histone gene families. *Cell.* 25:301–313.
- Hostikka, S. L., and K. Tryggvason. 1988. The complete primary structure of the $\alpha 2$ chain of human type IV collagen and comparison with the $\alpha 1(IV)$ chain. *J. Biol. Chem.* 263:19488–19493.
- Hostikka, S. L., R. L. Eddy, Byers, M. Hoyhtya, T. B. Shows, and K. Tryggvason. 1990. Identification of a distinct type IV collagen chain with restricted kidney distribution and assignment of its gene to the locus of X chromosome-linked Alport syndrome. *Proc. Natl. Acad. Sci. USA.* 87:1606–1610.
- Huang, A., C. E. Campbell, L. Bonnetta, M. S. Hill, S. McNeill, M. J. Coppes, D. J. Law, A. P. Feinberg, H. Yeager, and B. R. G. Williams. 1990. Tissue, developmental, and tumor-specific expression of divergent transcripts in Wilms tumor. *Science (Wash. DC).* 250:991–994.
- Hudson, B. G., R. Kalluri, S. Gunwar, M. Weber, F. Ballester, J. K. Hudson, M. E. Noelken, M. Sarras, W. R. Richardson, J. Saus et al. 1992. The pathogenesis of Alport syndrome involves type IV collagen molecules containing the $\alpha 3(IV)$ chain: evidence from anti-GBM nephritis after renal transplantation. *Kidney Int.* 42:179–187.
- Hudson, B. G., S. T. Reeders, and K. Tryggvason. 1993. Type IV collagen: Structure, gene organization, and role in human diseases. *J. Biol. Chem.* 268:26033–26036.
- Ishikawa, E., M. Imagawa, S. Hashida, S. Yoshitake, Y. Hamaguchi, and T. Ueno. 1983. Enzyme-labeling of antibodies and their fragments for enzyme immunoassay and immunohistochemical staining. *J. Immunoassay.* 4:209–327.
- Johansson C., R. Butkowsky, and J. Wieslander. 1992. The structural organization of type IV collagen. Identification of three populations in the glomerular basement membrane. *J. Biol. Chem.* 267:24533–24537.
- Junqueira, L. C., J. Carneiro, and J. A. Long. 1992. Basic Histology. 7th ed. Lange Medical Publications, Los Altos. 66–131.
- Kamagata, Y., M.-G. Mattei, and Y. Ninomiya. 1992. Isolation and sequencing of cDNAs and genomic DNAs encoding the $\alpha 4$ chain of basement membrane collagen type IV and assignment of the gene to the distal long arm of human chromosome 2. *J. Biol. Chem.* 267:23753–23758.
- Killen, P. D., P. Burbelo, Y. Sakurai, and Y. Yamada. 1988. Structure of the amino-terminal portion of the murine $\alpha 1(IV)$ collagen chain and the corresponding region of the gene. *J. Biol. Chem.* 263:8706–8709.
- Kishiro, Y., M. Kagawa, I. Naito, and Y. Sado. 1995. A novel method of preparing rat monoclonal antibody-producing hybridomas by using rat medial iliac lymph node cells. *Cell Struct. Funct.* 20:151–156.
- Kleppel, M. M., C. Kashtan, P. A. Santi, J. Wieslander, and A. F. Michael. 1989. Distribution of familial nephritis antigen in normal tissue and renal basement membranes of patients with homozygous and heterozygous Alport familial nephritis. *Lab. Invest.* 61:278–289.
- Laemmli, U. K. 1970. Cleavage of structural protein during the assembly of the head of bacteriophage T4. *Nature (Lond.)*. 227:680–685.
- Leinonen, A., M. Mariyama, T. Mochizuki, K. Tryggvason, and S. T. Reeders. 1994. Complete primary structure of the human type IV collagen $\alpha 4(IV)$ chain. Comparison with structure and expression of the other $\alpha(IV)$ chains. *J. Biol. Chem.* 269:26172–26177.
- Lemmink, H. H., T. Mochizuki, L. P. W. J. van den Heuvel, C. H. Schroder, A. Barrientos, L. A. H. Monnens, B. A. van Oost, H. G. Brenner, S. T. Reeders, and H. J. M. Smeets. 1994. Mutations in the type IV collagen $\alpha 3$ gene (COL4A3) in autosomal recessive Alport syndrome. *Hum. Mol. Genet.* 3:1269–1273.
- Lightfoot, K., L. Maltby, R. Duarte, R. Veale, and O. Segev. 1994. Conserved cis-elements bind a protein complex that regulates Drosophila ras2/rop bidirectional expression. *Br. J. Cancer.* 69:264–273.
- Mariyama, M., A. Linonen, T. Mochizuki, K. Tryggvason, and S. T. Reeders. 1994. Complete primary structure of the human $\alpha 3(IV)$ collagen chain. Co-expression of the $\alpha 3(IV)$ and $\alpha 4(IV)$ collagen chains in human tissues. *J. Biol. Chem.* 269:23013–23017.
- Merrifield, R. B. 1963. Solid phase peptide synthesis. I. The synthesis of a tetrapeptide. *J. Am. Chem. Soc.* 85:2149–2154.
- Miners, J. H., and J. R. Sanes. 1994. Collagen IV $\alpha 3$, $\alpha 4$, and $\alpha 5$ chains in rodent basal laminae: sequence, distribution, association with laminins, and developmental switches. *J. Cell Biol.* 127:879–891.
- Mochizuki, T., H. H. Lemmink, M. Mariyama, C. Antignac, M.-C. Gubler, Y. Pirson, C. Verellen-Dumoulin, B. Chan, C. H. Schroder, H. J. H. J. Smeets, and S. T. Reeders. 1994. Identification of mutations in the $\alpha 3(IV)$ and $\alpha 4(IV)$ collagen genes in autosomal recessive Alport syndrome. *Nature Genet.* 8:77–82.
- Morrison, K. E., M. Mariyama, T. L. Yang-Feng, and S. T. Reeders. 1991. Sequence and localization of a partial cDNA encoding the human $\alpha 3$ chain of type IV collagen. *Am. J. Hum. Genet.* 49:545–554.
- Oohashi, T., M. Sugimoto, M.-G. Mattei, and Y. Ninomiya. 1994. Identification of a new collagen IV chain, $\alpha 6(IV)$, by cDNA isolation and assignment of the gene to chromosome Xq22, which is the same locus for COL4A5. *J. Biol. Chem.* 269:7520–7526.
- Pihlajaniemi, T., K. Tryggvason, J. Myers, M. Kurkinen, R. Lebo, M. Cheung, D. J. Prockop, and C. D. Boyd. 1985. cDNA clones coding for the pro- $\alpha 1(IV)$ chain of human type IV procollagen reveal an unusual homology of amino acid sequences in two halves of the carboxyl-terminal domain. *J. Biol. Chem.* 260:7681–7687.
- Poschl, E., R. Pollner, and K. Kuhn. 1988. The gene for the $\alpha 1(IV)$ and $\alpha 2(IV)$ chains of human basement membrane collagen type IV are arranged head-to-head and separated by a bidirectional promoter of unique structure. *EMBO J.* 7:2687–2695.
- Rizzo, M. G., L. Offavio, S. Travali, C. Chang, B. Kaminska, and R. Baserga. 1990. The promoter of the human proliferating cell nuclear antigen(PCNA) is bidirectional. *Exp. Cell Res.* 188:286–293.
- Rohrbach, D. H., and R. Timpl. 1993. Molecular and Cellular Aspects of Basement Membranes. Academic Press, Inc., San Diego. 1–437.
- Sado, Y., M. Kagawa, I. Naito, and T. Okigaki. 1991. Properties of bovine nephritogenic antigen that induces anti-GBM nephritis in rats and its similarity to the Goodpasture antigen. *Virchows Arch. B Cell Pathol.* 60:345–351.
- Saffer, J. D., and M. F. Singer. 1984. Transcription from SV40-like monkey DNA sequences. *Nucleic Acids Res.* 12:4769–4788.
- Sanes, J. R., E. Engvall, R. Butkowsky, and D. D. Hunter. 1990. Molecular heterogeneity of basal laminae: isoform of laminin and collagen IV at the neuromuscular junction and elsewhere. *J. Cell Biol.* 111:1685–1699.
- Soininen, R., M. Houtari, S. L. Hostikka, D. J. Prockop, and K. Tryggvason. 1988. The structural genes for $\alpha 1$ and $\alpha 2$ chains of human type IV collagen are divergently encoded on opposite DNA strands and have an overlapping promoter region. *J. Biol. Chem.* 263:17217–17220.
- Sugimoto, M., T. Oohashi, H. Yoshioka, N. Matsuo, and Y. Ninomiya. 1993. cDNA isolation and partial gene structure of the human $\alpha 4(IV)$ collagen chain. *FEBS Lett.* 330:122–128.
- Sugimoto, M., T. Oohashi, and Y. Ninomiya. 1994. The genes COL4A5 and COL4A6, coding for basement membrane collagen chains $\alpha 5(IV)$ and $\alpha 6(IV)$, are located head-to-head in close proximity on human chromosome Xq22 and COL4A6 is transcribed from two alternative promoters. *Proc. Natl. Acad. Sci. USA.* 91:11679–11683.
- Wieslander, J., J. Langeveld, R. Butkowsky, M. Jodkowski, M. Noelken, and B. G. Hudson. 1985. Physical and immunohistochemical studies of the globular domain of the type IV collagen. *J. Biol. Chem.* 260:8564–8570.
- Yang-Feng, T. L., and S. T. Reeders. 1992. Colocalization of the genes for the $\alpha 3(IV)$ and $\alpha 4(IV)$ chains of type IV collagen to chromosome 2 bands q35–q37. *Genomics.* 13:809–813.
- Yoshioka, K., S. Hino, T. Takemura, S. Maki, J. Wieslander, Y. Takekoshi, H. Makino, M. Kagawa, Y. Sado, and C. E. Kashtan. 1994. Type IV collagen $\alpha 5$ chain: Normal distribution and abnormalities in X-linked Alport syndrome revealed by monoclonal antibody. *Am. J. Pathol.* 144:986–996.
- Zhou, J., J. M. Hertz, A. Leinonen, and K. Tryggvason. 1992. Complete amino acid sequence of the human $\alpha 5(IV)$ collagen chain and identification of a single-base mutation in exon 23 converting glycine521 in the collagenous domain to cysteine in an Alport syndrome patient. *J. Biol. Chem.* 267:12475–12481.
- Zhou, J., T. Mochizuki, H. Smeets, C. Antignac, P. Laurila, A. Paeppe, K. Tryggvason, and S. T. Reeders. 1993. Deletion of the paired $\alpha 5(IV)$ and $\alpha 6(IV)$ collagen genes in inherited smooth muscle tumors. *Science (Wash. DC).* 261:1167–1169.
- Zhou, J., M. Ding, and S. Reeders. 1994. Complete primary structure of the sixth chain of human basement membrane collagen, $\alpha 6(IV)$. *J. Biol. Chem.* 269:13193–13199.