

POSTER PRESENTATION

Open Access

Activity of the dietary flavonoid, apigenin, against multidrug-resistant tumor cells as determined by pharmacogenomics and molecular docking

Mohamed Saeed^{1*}, Thomas Efferth¹, Onat Kadioglu¹, Hassan Khalid², Yoshikazu Sugimoto³

From Breast Cancer Immunotherapy Symposium (BRECIS), part of the Sidra Symposia Series, held in partnership with the Society for Immunotherapy of Cancer Doha, Qatar. 13-14 April 2015

Natural products have been extensively studied and involved in cancer therapy field [1], apigenin has considerable cytotoxic activity in vitro and in vivo. Despite many mechanistic studies, less is known about resistance factors hampering apigenin's activity. We investigated the ATP-binding cassette (ABC) transporters BCRP/ABCG2, P-glycoprotein/ABCB1 and its close relative ABCB5. Apigenin inhibited not only P-glycoprotein, but also BCRP by increasing cellular uptake of doxorubicin and showed synergistic inhibitory effect in combination with doxorubicin or docetaxel against multidrug-resistant cells. To perform in silico studies, we first generated homology models for human P-glycoprotein and ABCB5 based on the crystal structure of murine P-glycoprotein. Their nucleotide binding domains (NBDs) revealed the highest degrees of sequence homologies (89%-100%), indicating that ATP binding and cleavage is of crucial importance for ABC transporters. In silico studies showed a pigenin bound to the NBDs of P-glycoprotein and ABCB5. Hence, apigenin may compete with ATP for NBD-binding leading to energy depletion to fuel the transport of ABC transporter substrates. Furthermore, we performed COMPARE and hierarchical cluster analyses of transcriptome-wide mRNA expression profiles of the National Cancer Institute tumor cell line panel. Microarray-based mRNA expressions of genes of diverse biological functions significantly predicted responsiveness of tumor cells to apigenin [2].

Authors' details

¹Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry - Johannes Gutenberg-University, Mainz, Germany. ²Department of Pharmacognosy, University of Khartoum, Khartoum, Sudan. ³Division of Chemotherapy, Faculty of Pharmacy - Keio University, Tokyo, Japan.

Published: 14 August 2015

References

1. Newman DJ, Cragg GM: Natural products as sources of new drugs over the 30 years from 1981 to 2010. *J Nat Prod.* 2012, **75**(3):311-35.
2. Saeed M, Kadioglu O, Khalid H, Sugimoto Y, Efferth T: Activity of the dietary flavonoid, apigenin, against multidrug-resistant tumor cells as determined by pharmacogenomics and molecular docking. *J Nutr Biochem* 2015, **26**(1):44-56.

doi:10.1186/2051-1426-3-S1-P10

Cite this article as: Saeed et al.: Activity of the dietary flavonoid, apigenin, against multidrug-resistant tumor cells as determined by pharmacogenomics and molecular docking. *Journal for ImmunoTherapy of Cancer* 2015 **3**(Suppl 1):P10.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry - Johannes Gutenberg-University, Mainz, Germany
Full list of author information is available at the end of the article