

Factors to Consider When Determining Whether Lipotoxicity Solely Causes B-Cell Decompensation in KPT2D [Letter]

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dan Geltser 

Imperial College London, School of
Medicine, London, UK

Dear editor

I read with great interest the paper by Ye et al,¹ which identified the significance of raised triglycerides in the treatment of ketosis-prone type 2 diabetes (KPT2D). The study concluded the need to implement lifestyle changes and administration of lipid-lowering medication for paediatric, hyperglycaemic KPT2D patients. As a 5th year medical student, who completed an intercalated BSc focusing on lipidology in diabetes mellitus, I would like to discuss further factors which should be considered when evaluating the conclusion.

I commend the paper on importantly highlighting the significant difference in triglyceride levels and obesity between KPT2D and type 2 diabetes mellitus (T2DM) patients. The paper interestingly identified that both glucotoxicity and lipotoxicity lead to β -cell decompensation. However, the study chose patients in tertiary care who were highly hyperglycaemic; this would make it difficult to determine whether it is exclusively the poor glycaemic control or the hypertriglyceridemia contributing to pancreatic cell destruction and therefore it is unknown whether lipid-lowering medications would in fact help.

Furthermore, the medication regimen of patients was not accounted for in this paper. Lipid-lowering medications such as statins both have affect on triglyceride levels as well as glycaemic control. Triglyceride levels may have been underestimated for patients currently on statins. A meta-analysis found that statins generally cause a higher HbA1c, particularly atorvastatin, where as pitavastatin improves it; this could have affected their conclusion.² Moreover, certain T2DM medications, such as sulfonylureas, can cause a rise in a patients lipid profile, with raised total cholesterol, LDL and triglycerides.³

The study did not stratify patients based on gender or ethnicity which could be confounding factors. Research has shown that patients of black ethnicity have a less atherogenic lipid profile than those of white ethnicity. Patients of Asian ethnicity have higher triglyceride levels overall.⁴ Moreover, the exclusion criteria stated that all the patients in the study were autoantibody negative; however, there is an increasing understanding in the pathology of T2D and the autoimmune component of the disease.⁵

Lipid profiles change over time and oscillate day to day, therefore using a one-off measurement of triglyceride levels is not representative of the patients general

Correspondence: Dan Geltser
Imperial College London, School of
Medicine, London, UK
Email dg1116@ic.ac.uk

lipid profile. It would be useful to do a longitudinal study and identify the average triglyceride levels over a longer period of time for each patient.

In conclusion, the research findings are useful in identifying KPT2D patients who may require treatment earlier on to avoid atherogenic lipid profiles and develop better glycaemic control. However, this needs to be done in all sectors of care and in an ethnically diverse community in order to make the data more representative and useful to extrapolate. It is also imperative to understand the exact mechanisms behind how triglycerides and other lipoproteins lead to β -cell decompensation and whether results differ in patients with well-controlled glucose levels.

Disclosure

The author reports no conflicts of interest in this communication.

References

1. Ye S, Ran H, Zhang H, et al. Elevated serum triglycerides are associated with ketosis-prone type 2 diabetes in young individuals. *Diabetes Metab Syndr Obes.* 2021;14:497–504. doi:10.2147/DMSO.S296085
2. Cui JY, Zhou RR, Han S, Wang TS, Wang LQ, Xie XH. Statin therapy on glycemic control in type 2 diabetic patients: a network meta-analysis. *J Clin Pharm Ther.* 2018;43(4):556–570. doi:10.1111/jcpt.12690
3. Yue-hong C. Sulfonylureas may affect lipids in type 2 diabetes. *J Evid Based Med.* 2015. doi:10.1111/jebm.12157.11
4. Frank A, Zhao B, Jose P, Azar K, Fortmann S, Palaniappan L. Racial/ethnic differences in dyslipidemia patterns. *Circulation.* 2014;129(5):570–579. doi:10.1161/CIRCULATIONAHA.113.005757
5. Itariu BK, Stulnig TM. Autoimmune aspects of type 2 diabetes mellitus - a mini-review. *Gerontology.* 2014;60(3):189–196. doi:10.1159/000356747

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>