

Article

Mutagenicity of Ochratoxin A and Its Hydroquinone Metabolite in the *Sup*F Gene of the Mutation Reporter Plasmid Ps189

Steven A. Akman^{1,*}, Marissa Adams¹, Doug Case², Gyungse Park³ and Richard A. Manderville^{4,*}

- ¹ Department of Cancer Biology, Wake Forest University Health Sciences, Winston-Salem, North Carolina, NC, USA; E-Mail: madams@wakehealth.edu
- ² Department of Public Health Sciences, Wake Forest University Health Sciences, Winston-Salem, North Carolina, NC, USA; E-Mail: dcase@wakehealth.edu
- ³ Department of Chemistry, College of Science and Technology, Kunsan National University, Miryong-Dong, Kusan, Korea; E-Mail: parkg@kunsan.ac.kr
- ⁴ Departments of Chemistry and Toxicology, University of Guelph, Guelph, Ontario, Canada; E-Mail: rmanderv@uoguelph.ca
- * Authors to whom correspondence should be addressed;
 E-Mails: sakman@wakehealth.edu (S.A.A.); rmanderv@uoguelph.ca (R.A.M.);
 Tel.: +1-336-716-4464 (S.A.A.); +1-519-824-4120 ext. 53963 (R.A.M.);
 Fax: +1-336-716-5687 (S.A.A.); +1-519-766-1499 (R.A.M.).

Received: 9 February 2012; in revised form: 29 March 2012 / Accepted: 6 April 2012 / Published: 16 April 2012

Abstract: Ochratoxin A (OTA) is a mycotoxin that enhances renal tumor formation in the outer medulla of male rat kidney. Direct DNA damage and subsequent mutagenicity may contribute to these processes. In this study we have determined whether OTA in the absence or presence of activated rat liver microsomes (RLM) or redox-active transition metals (Fe(III) or Cu(II)) causes promutagenic DNA damage in the supF gene of the mutation reporter plasmid pS189 replicating in human Ad293 cells. In addition, we have assessed the mutagenicity of the hydroquinone metabolite (OTHQ) of OTA in the absence or presence of cysteine without added cofactors. Our results show that oxidation of OTA, either by RLM or by transition metal ions, activates OTA to a directly genotoxic mutagen(s). The Fe(III)/OTA system was the most potent mutagen in our experimental system, causing a 32-fold increase in mutant fraction (MF) above the spontaneous control MF. The Cu(II)/OTA system caused a 9-fold increase in MF, while a 6–10-fold increase in MF was observed for OTA in the presence of RLM. The OTHQ metabolite is also

mutagenic, especially in the presence of cysteine, in which a 6-fold increase in MF was observed. Our data provide further insight into OTA bioactivation that may account for its *in vivo* mutagenicity in male rat kidney.

Keywords: ochratoxin A; mutagenicity; DNA adduct; genotoxicity; carcinogenesis

1. Introduction

Ochratoxin A (OTA, Figure 1) is a human toxin produced by Penicillium verrucosum and several Aspergillus species [1,2]. It is found in a variety of human foods, including cereal grains, e.g., wheat and rye, as well as coffee, beer, and wine [3,4]. OTA has been implicated as a cause of nephropathies and urothelial tract tumors in the Balkans [5–7] and in North African countries [8]. The International Agency for Research on Cancer (IARC) has classified OTA as a possible human carcinogen (group 2B) [9], and OTA exposure has been associated with cancer of the urothelial tract in rats [10, and reviewed in 5] and chicks [11]. OTA has also been proposed as a cause of testicular cancer in young men [12,13].

OTA is a substrate for photochemical [14–16], electrochemical [17], and transition metal ion-mediated [18,19] oxidation. OTA is also a substrate for enzymatic oxidation by microsomal mixed function oxidases [20–22] and enzymes with peroxidase activity [23,24]. Exposure to OTA is cytotoxic to cultured cells [25–28] as well as *in vivo* in rodent models [29–31]. The cytotoxicity of OTA shows a close correlation with the onset of oxidative DNA damage mediated by the toxin through production of reactive oxygen species (ROS) [25,27,28].

Figure 1. Structure of OTA, OTHQ and the OTB-dG adduct.



Studies of OTA-mediated mutagenicity have produced inconsistent results. The original assessment in S. typhimurium in the presence of rat liver post-mitochondrial supernatants was negative [32]. However, De Groene et al. observed OTA-mediated mutagenesis in NIH 3T3 cells stably expressing certain P450s [33,34]. Additional studies by the Obrecht-Plumio et al. in the presence of high dose OTA (403–1210 µg/plate), kidney microsomes, NADPH, and arachidonate caused the formation of mutagens detectable in S. typhimurium [35]. However, Zepnick and coworkers found negative evidence for OTA-mediated mutagenicity [36]. Palma et al. found that OTA-mediated mutagenicity is consistent with oxidative DNA damage and that bioactivation of OTA is not a requirement for the observed mutagenicity [37]. Several other studies have concluded that enzymatic oxidation of OTA does not cause covalent adduction of OTA to DNA [38-40] and that mutagenicity stems from by-products of cellular cytotoxicity [41]. However, other recent studies have produced conflicting results that indicate that oxidation of OTA in the presence of nucleosides or DNA yields covalent OTA-DNA nucleoside adducts [42-44]. Included in these studies is liquid chromatography-mass spectrometry (LC-MS) results that support the presence of the nonchlorinated OTB-2'-deoxyguanosine (dG) adduct (Figure 1) in male rat kidney [44]. Hibi and coworkers have also demonstrated the mutagenicity of OTA in the target tissue (outer medulla) of male rats [45]. In these studies the gpt delta transgenic rat model was employed and the reporter gene mutation assay showed significantly higher levels of deletion mutations compared to controls using DNA extracted from the outer medulla of male rat. The extracted DNA was also examined for 8-oxo-dG formation derived from ROS generation and no evidence for 8-oxo-dG formation was found and OTA exposure failed to increase the frequencies of GC: TA transversion mutations, which are characteristic of 8-oxo-dG-mediated mutagenicity. The in vivo mutagenicity assays reported by Hibi et al. suggest that oxidative DNA damage does not contribute to OTA-mediated mutagenicity and favor a direct genotoxic mechanism [45].

The positive results on OTA-mediated *in vivo* mutagenicity [45] combined with the finding that OTA generates the OTB-dG adduct in male rat kidney [44] demonstrates that DNA adduction and mutagenicity remains a viable mechanism of action for OTA-mediated renal carcinogenesis [46]. These results prompted us to report the current study, in which we address the mutagenicity of OTA in cell culture, using the human mutation reporter plasmid pSP189 developed by Seidman [47]. The data presented herein indicate that oxidation of OTA, either by microsomal enzymes or by transition metal ions, activates OTA to a directly genotoxic mutagen(s). Synthetic ochratoxin hydroquinone (OTHQ, Figure 1), an OTA metabolite that forms covalent DNA adducts [43], is also mutagenic.

2. Experimental Section

2.1. Reagents

OTA (\geq 98%), β NADP, glucose-6-phosphate and glucose-6-phosphate dehydrogenase were purchased from Sigma Chemical Co. (St. Louis, MO, USA). Arochlor[®]-activated rat liver microsomes were purchased from In Vitro Technologies, Inc. (Baltimore, MD, USA). Plasmid pSP189 was received as a generous gift from Dr. Michael Seidman. The construction and properties of pSP189, including its "signature sequence", have been reported [48]. OTHQ was chemically synthesized as a mixture of diastereomers (3(R/S), Figure 1) using the synthetic protocol previously reported [22] and

was \geq 96% pure based on reverse-phase HPLC analysis. Stock solutions of OTA and OTHQ (13.7 mM) were prepared in dioxane. Stock solutions containing 10 mM OTA and 5 mM cupric acetate or ferric ammonium citrate (1:2 metal ion:OTA molar ratio) were made in 10 mM MOPS buffer, initially at pH 4, then adjusted to 7.4 with NaOH. Coordination of copper ions by OTA was verified by the appearance of an absorbance peak at 365 nm [18,49].

2.2. Treatment of Plasmid PSP189 with OTA

In general reactions were carried out in 50 mM potassium phosphate buffer, pH 7.4, except those involving Cu(II)- or Fe(III)-OTA complexes, in which case the buffer was 50 mM HEPES, pH 7.4. For reactions utilizing rat liver microsomes (RLM), a 25 mg/mL microsome suspension and a nucleotide regenerating system consisting of 0.7 mM β NADP, 7.7 mM glucose-6-phosphate, plus 1.5 units/mL glucose-6-phosphate dehydrogenase were pre-warmed to 37 °C for 5 min. All reactions were carried out in 500 μ L volumes at 37 °C for 60 min and included 25 μ g pSP189, OTA or OTHQ, 625 μ g activated microsomes and 125 μ L nucleotide regenerating system, where appropriate. Reactions were quenched by cooling on ice, followed by phenol:chloroform:isoamyl alcohol (24:21) extraction, and ethanol precipitation. Precipitated plasmid was washed twice with 70% ethanol, dried, and dissolved in 10 mM Tris-HCl, pH 8 for analysis.

2.3. Transfection of Target Cells

Human Ad293 cells, which are immortal, but not malignant, human cells derived from embryonic kidney, were grown as a monolayer in Dulbecco's modified Eagle's medium (Gibco, Grand Island, NY, USA) containing 5% heat-inactivated fetal bovine serum in a humidified 5% CO₂ atmosphere. Target Ad293 cells, grown to ca. 50% confluence in 100 mm³ dishes, were transfected with 10 μ g pSP189 DNA by the diethylaminoethyl dextran technique [50].

2.4. Mutation Analysis

Low molecular weight DNA was isolated from target Ad293 cells 48 h after transfection by the procedure of Hirt [51]. Plasmid DNA was purified and analyzed for the mutant fraction (MF) as previously described [52]. Briefly, purified plasmid DNA was treated with 0.1 units/ μ L Dpn I to remove unreplicated plasmid DNA, then used to electrotransform Escherichia coli strain MBM7070. MBMB7070 cells contain an amber mutation in lacZ that is suppressed by wild type supF tRNA, producing functional β -galactosidase. Transformants were plated on Luria broth plates containing 50 μ g/mL ampicillin plus isopropyl- β -D-thiogalactoside (IPTG) and 5-bromo-4-chloro-3-indoyl- β -D-galactoside (X-gal, the chromogen). Total (blue + white) and mutant supF-containing (white) colonies are enumerated. Mutant (M) white colonies were confirmed by secondary streaking.

2.5. Statistical Analysis

Poisson regression was used to assess the effect of treatment on the mean number of M colonies. The log of the number of colonies was used as an offset in the models, and pair-wise treatment differences were tested using linear contrasts. The Deviance statistic was used to assess goodness of

271

fit, and a likelihood ratio test comparing the log likelihoods from Poisson and Negative Binomial models was used to test for overdispersion. Weighted means and standard errors (SE) are presented in the text and tables.

3. Results

3.1. Activation with Rat Liver Microsomes

Exposure to OTA did not significantly enhance the mutant fraction (MF) of plasmid pSP189 replicated in human Ad293 cells above the background value of 3.5×10^{-6} (Table 1). Exposure to 0.5–5 mM OTA in the presence of activated rat liver microsomes (RLM) enhanced the MF 6–10 fold to $3.5 \pm 0.93 \times 10^{-5}$ (Table 1, p < 0.02 for each comparison). However, no dependence of the MF on OTA concentration in the range 0.5–5 mM was discernible. Replacing activated RLM with boiled microsomes totally abrogated the OTA/microsome associated enhancement of mutagenicity.

Expt.	Treatment	Colonies	M Colonies ^a	MF (×10 ⁴) ^b	Mean ± SE
1	Plasmid alone	211267	2	0.09	0.035 ± 0.025
2		81533	0	0.00	
3		274100	0	0.00	
1	1 mM OTA	145467	1	0.07	0.061 ± 0.035
2		48367	1	0.21	
3		295233	1	0.04	
1	RLM + 0.5 mM OTA	165233	3	0.18	0.31 ± 0.087
2		81767	2	0.24	
3		169300	8	0.47	
1	RLM + 1 mM OTA	104167	2	0.19	0.23 ± 0.072
2		57767	2	0.35	
3		275867	6	0.21	
1	RLM + 5 mM OTA	88900	2	0.22	0.35 ± 0.093
2		31333	2	0.64	
3		281700	10	0.36	
1	Boiled RLM + 1 mM OTA	56067	0	0.00	0.032 ± 0.032
2		51567	1	0.19	
3		204900	0	0.00	

Table 1. Mutagenicity of OTA in the presence of rat liver microsomes.

^a The number of mutant (M) supF-containing (white) colonies; ^b Mutant Fraction (MF) = number of M colonies (white)/total (blue + white) colonies.

3.2. Mutagenicity of OTHQ

The OTHQ metabolite of OTA can undergo an autoxidative process to generate the quinone electrophile OTQ [22], that reacts with DNA to generate adduct spots, as evidenced by ³²P-postlabeling [43]. Thus, the OTHQ metabolite of OTA was tested for mutagenicity in the absence of metabolism. Its mutagenicity was also tested in the presence of 1 equiv. cysteine. These experiments were prompted by our earlier findings that the OTQ electrophile reacts covalently with cysteine to

form a conjugate that is unstable and undergoes further reactions to afford electrophilic species that may react with DNA [16]. A one hour exposure to 1 mM OTHQ senhanced the MF of plasmid pSP189 replicated in human Ad293 cells 3.6 fold above background to $1.4 \pm 0.51 \times 10^{-4}$ (p = 0.049, Table 2). Addition of 1 mM cysteine to 1 mM OTHQ further increased the MF to 6.3 fold above background to $2.5 \pm 0.68 \times 10^{-4}$ (p = 0.003 vs. background, Table 2).

Expt.	Treatment	Colonies	M Colonies	$MF (\times 10^4)$	Mean ± SE
1	Plasmid alone	40150	2	0.50	0.39 ± 0.22
2		18050	1	0.55	
3		19600	0	0.00	
1	1 mM cysteine	30500	3	0.98	0.73 ± 0.28
2		32700	3	0.92	
3		32700	1	0.32	
1	1 mM OTHQ	12400	4	3.2	1.41 ± 0.45
2		29200	3	1.0	
3		29350	3	1.0	
1	1 mM OTHQ + 1 mM cysteine	11200	6	5.4	2.45 ± 0.60
2		29050	7	2.4	
3		29050	4	1.4	

Table 2. Mutagenicity of OTHQ in the absence or presence of cysteine.

3.3. Mutagenicity of OTA-Transition Metal Ion Complexes

The mutagenicity of OTA in the presence of Fe(III) and Cu(II) was also determined, given that OTA is oxidized by Fe(III) into the quinone electrophile OTQ [22] that provides a rationale for the mutagenicity of OTHQ (Table 2). OTA also reacts with dG in the presence of Fe to generate the OTB-dG adduct shown in Figure 1 [53], and forms an OTA-Fe complex that has been implicated in lipid peroxidation mediated by the toxin [54]. OTA also forms a complex with Cu(II) [18,49] that can facilitate oxidative DNA strand scission [18,19]. Thus, we were interested to determine whether OTA would show mutagenicity in the presence of these redox-active transition metal ions. For these experiments, OTA was kept in a two-fold excess over the transition metal in the non-metal-coordinating HEPES buffer pH 7.4. Both Fe(III) [54] and Cu(II) [18] form 1:1 complexes with OTA and at pH 6.0 the equilibrium binding constant ($K_{1:1}$) for Cu(II) is ~2.5 × 10⁶ M⁻¹ [18], while a $K_{1,1}$ value $\sim 2 \times 10^8$ M⁻¹ has been determined for Fe(III) binding [54], suggesting complete metal ion coordination by OTA under our experimental conditions. Free Fe chelated to diethylenetriamine-pentacetic acid does produce a small dose dependent increase in MF in the mutation reporter plasmid assay [55]. In these experiments, 5 mM Fe showed a 3-fold increase in MF compared to control, while 0.1 mM Fe failed to enhance the MF [55]. Free Cu(II) has also been shown to act as a mutagen [56]. In our experiments, exposure to 1 mM Fe(III):2 mM OTA enhanced the MF of plasmid pSP189 replicated in human Ad293 cells 32-fold above background to $2.5 \pm 0.52 \times 10^{-4}$ (p = 0.001, Table 3). Exposure to the 1 mM Cu(II):2 mM OTA complex enhanced mutagenicity to a lesser extent, *ca.* 9-fold above background to $7.1 \pm 2.7 \times 10^{-5}$ (*p* = 0.041, Table 3).

Expt.	Treatment	Colonies	M Colonies	MF (× 10 ⁴)	Mean ± SE
1	Plasmid alone	12100	0	0.00	0.080 ± 0.080
2		37200	0	0.00	
3		76050	1	0.13	
1	1 mM Cu(II)/2 mM OTA	10067	1	0.99	0.71 ± 0.27
2		50700	3	0.59	
3		38350	3	0.78	
1	1 mM Fe(III)/2 mM OTA	24800	12	4.8	2.52 ± 0.52
2		35600	4	1.1	
3		34650	8	2.3	

Table 3. Mutagenicity of transition metal ion/OTA complexes.

4. Discussion

There has been considerable debate whether OTA is directly genotoxic [38,39,44,46,57,58]. OTA can promote oxidative DNA damage through ROS generation that causes cytotoxicity [25,27,28] and oxidative DNA damage has been proposed to play an important role in carcinogenicity [41]. However, other lines of evidence favour DNA adduction by OTA [23,42–44,46,57]. The *in vivo* mutagenicity findings for OTA do not support oxidative base damage as the initiation event and favour direct genotoxicity in OTA-induced renal carcinogenicity [44]. However, the link between DNA adduction and OTA-mediated mutagenicity has not been firmly established [46].

The data presented here demonstrate that OTA is activated to a species that is a directly genotoxic mutagen. The free toxin lacked mutagenicity in the absence of external cofactors, such as RLM (Table 1) or redox-active transition metal ions (Table 3). These findings differ from the mutagenicity data presented by Palma *et al.*, which suggested that bioactivation in not required for mutagenicity [37], and are more in line with the data from Tozlovanu *et al.*, 2006 showing that DNA adduction by OTA is not observed directly with OTA in the absence of oxidative metabolism [43].

That RLM could be used for the conversion of OTA to a genotoxic mutagen in our experimental system (Table 1), suggested that the genotoxic metabolite of OTA is an oxidation product. In this regard, we previously demonstrated that the oxidation of OTA (100 μ M) by RLM (1 mg/mL) generated a GSH-conjugate that suggested the intermediacy of the quinone electrophile (OTQ) in the oxidation of OTA [59]. Pfohl-Leszkowicz and coworkers have also outlined a role for GSH in conjugation of OTA-derived electrophiles following bioactivation of the parent toxin [60]. The oxidation of OTA by activated RLM in the presence of a reducing agent (ascorbate) also generates the hydroquinone metabolite OTHQ [59]. OTHQ autoxidizes to OTQ that reacts covalently with DNA to generate DNA adducts [43]; although the structure(s) of such adducts are unknown. This process occurs spontaneously in aqueous media in the presence of molecular O₂, and no other cofactors are required for conversion of OTHQ into OTQ [22]. While this process has the potential to generate genotoxic ROS, we have recently demonstrated that OTHQ lacks cytotoxicity in mammalian kidney cells, suggesting that the metabolite is ineffective at stimulating ROS production [61]. Thus, DNA damage mediated by OTHQ is mainly due to covalent DNA adduction stimulated by OTQ formation [43]. This background information implies that OTQ formation is likely responsible for mutagenicity stimulated by the RLM/OTA system (Table 1).

Authentic OTHQ was also mutagen in our experimental system, albeit a weak one (Table 2). The weakness of OTHQ mediated mutagenicity may be a reflection of its slow rate of autoxidation [22]. Prolonged incubation of pSP189 with OTHQ may have enhanced its mutagenicity. However, interestingly, addition of cysteine enhanced the mutagenicity of OTHQ. A possible mechanism for enhancement of OTHQ-mediated mutagenicity by cysteine is suggested by our previous work [16] and outlined by Monks and Lau for the nephrotoxicity of polyphenolic-gluathione conjugates [62]. Thus, as outlined in Figure 2, autoxidation of OTHQ generates OTQ that reacts with cysteine to afford the conjugate OTHQ-Cys [16]. This conjugate is not stable [16] and can also undergo an autoxidative process to reform the quinone structure that reacts covalently with the amine group of the attached cysteine to afford a quinonimine that upon tautomerization will afford the imine electrophile in the new six-membered ring system. The imine functional group forms the basis for DNA adduction by the pyrrolobenzodiazepines, such as anthramycin, that have potential use in anticancer therapies [63]. This proposed pathway may play a role for the heightened mutagenicity of OTHQ in the presence of cysteine (Table 2).





In our experimental system, Fe(III)/OTA was the most potent mutagen showing a 32-fold MF above background levels (Table 3). This observation may correlate with earlier studies on the reactivity of OTA in the presence of dG to yield OTB-dG [53]. These studies were prompted by reports that OTA forms guanine-specific DNA adducts [64,65] and demonstrated that OTB-dG formation resulted from reaction of OTA/dG in the presence of Fe, Cu, or peroxidase enzymes [53]. Interestingly, Fe generated OTB-dG in the highest yield, ~5 orders of magnitude greater than Cu and peroxidase. The greater reactivity of OTA toward guanine in the presence of Fe, as opposed to Cu, may provide a rationale for the enhanced MF of the Fe/OTA system (Table 3). However, Fe(III) also oxidizes OTA to the quinone electrophile OTQ that reacts with DNA [43] and provides a rationale for the mutagenicity of OTHQ (Table 2). Thus, there are at least two possible mechanisms for Fe(III)/OTA mediated mutagenesis: (1) OTB-dG formation, and (2) OTQ formation with subsequent

adduct formation. At present, the DNA adduct(s) formed by OTQ have not been structurally characterized and it is unknown whether the OTB-dG adduct is mutagenic. Our goal is to first characterize the DNA adducts formed by OTQ, and incorporate site-specifically the OTQ adducts and OTB-dG into oligonucleotide substrates so that their biological consequences can be assessed.

The observation of OTA-mediated mutations in the supF gene of plasmid DNA replicating in human Ad293 cells seemingly is not concordant with the work of Hibi et al. [45], who did not observe enhanced mutagenesis in the gpt gene in kidneys of OTA-exposed mice. The explanation for the disparate results is not known; however, methodologic differences may be contributory gpt mutations are detected phenotypically by resistance to 6-thioguanine due to loss of hypoxanthine-guanine phosphoribosyl transferase (HGPRT) catalytic activity. Phenotypically silent gpt mutations that do not affect HGPRT catalytic activity and are therefore not detected in a 6-thioguanine resistance assays may have contributed to underestimation of OTA-mediated mutations in mouse kidney DNA. In contrast, there are no phenotypically silent mutations in the supF gene in the reporter assay used in this study. Hibi et al. [45] did observe enhance OTA-mediated production of large (ca. 10 kb) deletions in mouse kidney; such mutations are not observable in the plasmid pSP189-based assay. The occurrence of large deletions in the mouse kidney vs. base substitutions and frameshift mutations in the in vitro plasmid based assay may reflect differences in the OTA activation pathways to a proximate mutagen. Base substitution and/or frameshift mutations mediated by OTA in vitro are enhanced by conversion to the hydroquinone in the presence of cysteine or to the phenoxyl radical in the presence of Fe(III), both of which produce DNA base-reactive OTA intermediates. It is possible that OTA activation pathways in the mouse kidney generate phosphodiester backbone reactive intermediates, producing double strand breaks that lead to large deletions, rather than DNA base reactive intermediates. In this regard, differences in OTA activation pathways in the kidney as compared to those utilized in the in vitro mutation reporting system may be attributable to the fact that liver, as opposed to kidney, microsomes were used for *in vitro* activation. Future experiments using the *in vitro* mutation reporting system described herein will focus on the impact of OTA bioactivation by kidney microsomes, enabling comparison with the mutagenicity of liver microsome-activated OTA in vitro and the in vivo OTA-mediated genetic alterations described by Hibi et al.

Acknowledgments

Support from the National Cancer Institute (R01-CA080787) is acknowledged.

References

- 1. van der Merwe, K.J.; Steyn, P.S.; Fourie, L.; Scott, D.B.; Theron, J.J. Ochratoxin A, a toxic metabolite produced by *Aspergillus ochraceus* Wilh. *Nature* **1965**, *205*, 1112–1113.
- 2. Pohland, A.E.; Nesheim, S.; Friedman, L. Ochratoxin A, a review. *Pure Appl. Chem.* **1992**, *64*, 1029–1046.
- 3. Jørgensen, K. Survey of pork, poultry, coffee, beer and pulses for ochratoxin A. *Food Addit*. *Contam.* **1998**, *15*, 550–554.
- 4. Trucksess, M.; Giler, J.; Young, K.; White, K.D.; Page, S.W. Determination and survey of ochratoxin A in wheat, barley, and coffee—1997. *J. AOAC Int.* **1999**, *82*, 85–89.

- 5. Pfohl-Leszkowicz, A.; Manderville, R.A. Ochratoxin A: An overview on toxicity and carcinogenicity in animals and humans. *Mol. Nutr. Food Res.* **2007**, *51*, 61–99.
- 6. Stoev, S.D. The role of ochratoxin A as a possible cause of Balkan endemic nephropathy and its risk evaluation. *Vet. Human Toxicol.* **1998**, *40*, 352–360.
- 7. Pfohl-Leszkowicz, A.; Petkova-Bocharova, T.; Chernozemsky, I.N.; Castegnaro, M. Balkan endemic nephropathy and the associated urinary tract tumours: Review on etiological causes, potential role of mycotoxins. *Food Add. Contam.* **2002**, *19*, 282–302.
- 8. Grosso, F.; Saïd, S.; Mabrouk, I.; Fremy, J.M.; Castegnaro, M.; Jemmali, M.; Dragacci, S. New data on the occurrence of ochratoxin A in human sera from patients affected or not by renal diseases in Tunisia. *Food Chem. Toxicol.* **2003**, *41*, 1133–1140.
- IARC. Monographs on the evaluation of carcinogenic risks to humans. In Some Naturally Occurring Substances: Food Items and Constituents, Heterocyclic Aromatic Amines and Mycotoxins, No. 56, Ochratoxin A; International Agency for Research on Cancer: Lyon, France, 1993; pp. 489–521.
- Boorman, G.A. Toxicology and Carcinogenesis Studies of Ochratoxin A (CAS No 303-47-9) in F344/N Rats; Technical Report for National Toxicology Program, NIH Publication No. 89-2813; US Department of Health and Human Services, National Institutes of Health: Research Triangle Park, NC, USA, 1989.
- 11. Stoev, S.D. Studies on carcinogenic and toxic effects of ochratoxin A in chicks. *Toxins* **2010**, *2*, 649–664.
- 12. Schwartz, G.G. Hypothesis: Does ochratoxin A cause testicular cancer? *Cancer Causes Control* **2002**, *13*, 91–100.
- Jennings-Gee, J.E.; Tozlovanu, M.; Manderville, R.; Miller, M.S.; Pfohl-Leszkowicz, A.; Schwartz, G.G. Ochratoxin A: In utero exposure in mice induces adducts in testicular DNA. *Toxins* 2010, *2*, 1428–1444.
- 14. Gillman, I.G.; Yezek, J.M.; Manderville, R.A. Ochratoxin A acts as a photoactivatable DNA cleaving agent. *Chem. Commun.* **1998**, 647–648.
- 15. Il'ichev, Y.V.; Perry, J.L.; Manderville, R.A.; Chignell, C.F.; Simon, J.D. The pH-dependent primary photoreactions of ochratoxin A. *J. Phys. Chem. B* **2001**, *105*, 11369–11376.
- Brow, M.E.; Dai, J.; Park, G.; Wright, M.W.; Gillman, I.G.; Manderville, R.A. Photochemically catalyzed reaction of ochratoxin A with D- and L-cysteine. *Photochem. Photobiol.* 2002, *76*, 649–656.
- 17. Calcutt, M.W.; Gillman, I.G.; Noftle, R.E.; Manderville, R.A. Electrochemical oxidation of ochratoxin A: Correlation with 4-chlorophenol. *Chem. Res. Toxicol.* **2001**, *14*, 1266–1272.
- 18. Ardus, J.A.; Gillman, I.G.; Manderville, R.A. On the role of copper and iron in DNA cleavage by ochratoxin A. Structure-activity relationships in metal binding and copper-mediated DNA cleavage. *Can J. Chem.* **1998**, *76*, 907–918.
- Manderville, R.A.; Calcutt, M.W.; Dai, J.; Park, G.; Gillman, I.G.; Noftle, R.E.; Mohammed, A.K.; Dizdaroglu, M.; Rodriguez, H.; Akman, S.A. Stoichiometric preference in copper-promoted oxidative DNA damage by ochratoxin A. J. Inorg. Biochem. 2003, 95, 87–96.

- Stormer, F.C.; Storen, O.; Hansen, C.E.; Pedersen, J.I.; Aasen, A.J. Formation of (4*R*)- and (4*S*)-4-hydroxyochratoxin A and 10-hydroxyochratoxin A from ochratoxin A by rabbit liver microsomes. *Appl. Environ. Microbiol.* **1983**, *45*, 1183–1191.
- 21. Omar, R.F.; Hasinoff, B.B.; Mejilla, F.; Rahimtula, A.D. Effect of cytochrome P450 induction on the metabolism and toxicity of ochratoxin A. *Biochem. Pharmac.* **1996**, *40*, 1183–1191.
- 22. Gillman, I.G.; Clark, T.N.; Manderville, R.A. Oxidation of ochratoxin A by an Fe-porphyrin system: Model for enzymatic activation and DNA cleavage. *Chem. Res. Toxicol.* **1999**, *12*, 1066–1076.
- 23. Pfohl-Leszkowicz, A.; Grosse, Y.; Kane, A.; Creppy, E.E.; Dirheimer, G. Differential DNA adduct formation and disappearance in three mouse tissues after treatment with the mycotoxin ochratoxin A. *Mutat. Res.* **1993**, *289*, 265–273.
- 24. Pinelli, E.; Adlouni, C.E.; Pipy, B.; Quartulli, F.; Pfohl-Leszkowicz, A. Roles of cyclooxygenase and lipoxygenases in ochratoxin A genotoxicity in human epithelial lung cells. *Environ. Toxicol. Pharmacol.* **1999**, *7*, 95–107.
- Kamp, H.G.; Eisenbrand, G.; Schlatter, J.; Würth, K.; Janzowski, C. Ochratoxin A: Induction of (oxidative) DNA damage, cytotoxicity and apoptosis in mammalian cell lines and primary cells. *Toxicology* 2005, 206, 413–425.
- Faucet-Marquis, V.; Pont, F.; Størmer, F.; Rizk, T.; Castegnaro, M.; Pfohl-Leszkowicz, A. Evidence of a new dechlorinated OTA derivative formed in opossum kidney cell cultures after pre-treatment by modulators of glutathione pathways. Correlation with DNA adducts formation. *Mol. Nutr. Food Res.* 2006, *50*, 531–542.
- Arbillaga, L.; Azqueta, A.; Ezpeleta, O.; Lopez de Cerain, A. Oxidative DNA damage induced by ochratoxin A in the HK-2 human kidney cell line: evidence of the relationship with cytotoxicity. *Mutagenesis* 2007, 22, 35–42.
- Ali, R.; Mittelstaedt, R.A.; Shaddock, J.G.; Ding, W.; Bhalli, J.A.; Khan, Q.M.; Heflich, R.H. Comparative analysis of micronuclei and DNA damage induced by ochratoxin A in two mammalian cell lines. *Mutat. Res.* 2011, 723, 58–64.
- 29. Baudrimont, I.; Betbeder, A.M.; Gharbi, A.; Pfohl-Leszkowicz, A.; Dirheimer, G.; Creppy, E.E. Effect of superoxide dismutase and catalase on the nephrotoxicity of subchronical administration of ochratoxin A in rats. *Toxicology* **1994**, *89*, 101–111.
- Gautier, J.C.; Holzhaeuser, D.; Markovic, J.; Gremaud, E.; Schilter, B.; Turesky, R.J. Oxidative damage and stress response from ochratoxin A exposure in rats. *Free Rad. Biol. Med.* 2001, *30*, 1089–1098.
- Kamp, H.G.; Eisenbrand, G.; Janzowski, C.; Kiossev, J.; Latendresse, J.R.; Schlatter, J.; Turesky, R.J. Ochratoxin A induces oxidative DNA damage in liver and kidney after oral dosing to rats. *Mol. Nutr. Food Res.* 2005, 49, 1160–1167.
- 32. Wurgler, F.E.; Friederich, U.; Schlatter, J. Lack of mutagenicity of ochratoxin A and B, citrinin, patulin and cnestine in *Salmonella typhimurium* TA102. *Mutat. Res.* **1991**, *261*, 209–216.
- De Groene, E.M.; Hassing, I.G.A.M.; Blom, M.J.; Seinen, W.; Fink-Gremmels, J.; Horbach, G.J. Development of human cytochrome P450-expressing cell lines: Application in mutagenicity testing of ochratoxin A. *Cancer Res.* 1996, 56, 299–304.

- 34. De Groene, E.M.; Jahn, A.; Horbach, G.J.; Fink-Gremmels, J. Mutagenicity and genotoxicity of the mycotoxin ochratoxin A. *Environ. Toxicol. Pharmacol.* **1996**, *1*, 21–26.
- Obrecht-Plumio, S.; Chassat, T.; Dirheimer, G.; Marzin, D. Genotoxicity of ochratoxin A by Salmonella mutagenicity test after bioactivation by mouse kidney microsomes. *Mutat. Res.* 1999, 446, 95–102.
- 36. Zepnik, H.; Pähler, A.; Schauer, U.; Dekant, W. Ochratoxin A-induced tumor formation: Is there a role of reactive ochratoxin A metabolites? *Toxicol. Sci.* **2001**, *59*, 59–67.
- Palma, N.; Cinelli, S.; Sapora, O.; Wilson, S.H.; Dogliotti, E. Ochratoxin A-induced mutagenesis in mammalian cells is consistent with the production of oxidative stress. *Chem. Res. Toxicol.* 2007, 20, 1031–1037.
- Gross-Steinmayer, K.; Weymann, J.; Hege, H.C.; Metzler, M. Metabolism and lack of DNA reactivity of the mycotoxin ochratoxin A in cultured rat and human primary hepatocytes. *J. Agric. Food Chem.* 2002, *50*, 938–945.
- 39. Mally, A.; Zepnik, H.; Wanek, P.; Eder, E.; Dingley, K.; Ihmels, H.; Völkel, W.; Dekant, W. Ochratoxin A: Lack of formation of covalent DNA adducts. *Chem. Res. Toxicol.* **2004**, *17*, 234–242.
- Mally, A.; Pepe, G.; Ravoori, S.; Fiore, M.; Gupta, R.C.; Dekant, W.; Mosesso, P. Ochratoxin A causes DNA damage and cytogenetic effects but no DNA adducts in rats. *Chem. Res. Toxicol.* 2005, 18, 1253–1261.
- 41. Marin-Kuan, M.; Ehrlich, V.; Delatour, T.; Cavin, C.; Schilter, B. Evidence for a role of oxidative stress in the carcinogenicity of ochratoxin A. *J. Toxicol.* **2011**, 1–15.
- 42. Faucet, V.; Pfohl-Leszkowicz, A.; Dai, J.; Castegnaro, M.; Manderville, R.A. Evidence for covalent DNA adduction by ochratoxin A following chronic exposure to rat and subacute exposure to pig. *Chem. Res. Toxicol.* **2004**, *17*, 1289–1296.
- 43. Tozlovanu, M.; Faucet-Marquis, V.; Pfohl-Leszkowicz, A.; Manderville, R.A. Genotoxicity of the hydroquinone metabolite of ochratoxin A: Structure-activity relationships for covalent DNA adduction. *Chem. Res. Toxicol.* **2006**, *19*, 1241–1247.
- 44. Mantle, P.G.; Faucet-Marquis, V.; Manderville, R.A.; Squillaci, B.; Pfohl-Leszkowicz, A. Structures of covalent adducts between DNA and ochratoxin A: A new factor in debate about genotoxicity and human risk assessment. *Chem. Res. Toxicol.* **2010**, *23*, 89–98.
- 45. Hibi, D.; Suzuki, Y.; Ishii, Y.; Jin, M.; Watanabe, M.; Sugita-Konishi, Y.; Yanai, T.; Nohmi, T.; Nishikawa, A.; Umemura, T. Site-specific *in vivo* mutagenicity in the kidney of gpt delta rats given a carcinogenic dose of ochratoxin A. *Toxicol. Sci.* 2011, *122*, 406–414.
- 46. Pfohl-Leszkowicz, A.; Manderville, R.A. An update on direct genotoxicity as a molecular mechanism of ochratoxin A carcinogenicity. *Chem. Res. Toxicol.* **2012**, *25*, 252–262.
- 47. Seidman, M.M.; Dixon, K.; Razzaque, A.; Zagursky, R.J.; Berman, M.L. A shuttle vector plasmid for studying carcinogen-induced point mutations in mammalian cells. *Gene* **1985**, *38*, 233–237.
- 48. Parris, C.N.; Seidman, M.M. A signature element distinguishes sibling and independent mutations in a shuttle vector plasmid. *Gene* **1992**, *117*, 1–5.
- 49. Gillman, I.G.; Day, C.S.; Manderville, R.A. Stepwise formation of a nonsymmetric dinuclear copper complex of ochratoxin A. *Inorg. Chem.* **1998**, *37*, 6385–6388.

- 50. McCuthen, J.A.; Pagano, J.S. Enhancement of the infectivity of SV40 deoxyribonucleic acid with diethylaminoethyl dextran. *J. Natl. Cancer Inst.* **1968**, *41*, 351–357.
- 51. Hirt, B. Selective extraction of polyoma DNA from infected mouse cell cultures. *J. Mol. Biol.* **1967**, *20*, 365–369.
- 52. Waters, S.B.; Akman, S.A. A new assay to quantify *in vivo* repair of G:T mispairs by base excision repair. *Mutat. Res.* **2001**, *487*, 109–119.
- 53. Dai, J.; Wright, M.W.; Manderville, R.A. Ochratoxin A forms a carbon-bonded C8-deoxyguanosine nucleoside adduct: Implication for C8-reactivity by a phenolic radical. *J. Am. Chem. Soc.* **2003**, *125*, 3716–3717.
- 54. Omar, R.F.; Hasinoff, B.B.; Mejilla, F.; Rahimtula, A.D. Mechanism of ochratoxin A stimulated lipid peroxidation. *Biochem. Pharmacol.* **1990**, *40*, 1183–1191.
- 55. Akman, S.A.; Forrest, G.P.; Doroshow, J.H.; Dizdaroglu, M. Mutation of potassium permanganate-and hydrogen peroxide-treated plasmid pZ189 replicating in CV-1 monkey kidney cells. *Mutat. Res.* **1991**, *261*, 123–130.
- 56. Akman, S.A.; Doroshow, J.H.; Kensler, T.W. Copper-dependent site-specific mutagenesis by benzoyl peroxide in the supF gene of the mutation reporter plasmid pS189. *Carcinogenesis* **1992**, *13*, 1783–1787.
- 57. Manderville, R.A. A case for the genotoxicity of ochratoxin A by bioactivation and covalent DNA adduction. *Chem. Res. Toxicol.* **2005**, *18*, 1091–1097.
- 58. Turesky, R.J. Perspective: Ochratoxin A is not a genotoxic carcinogen. *Chem Res. Toxicol.* 2005, *18*, 1082–1090.
- 59. Dai, J.; Park, G.; Wright, M.W.; Adams, M.; Akman, S.A.; Manderville, R.A. Detection and characterization of a glutathione conjugate of ochratoxin A. *Chem. Res. Toxicol.* **2002**, *15*, 1581–1588.
- Pfohl-Leszkowicz, A.; Grosse, Y.; Kane, A.; Gharbi, A.; Baudrimont, I.; Obrecht, S.; Creppy, E.E.; Dirheimer, G. Is the oxidative pathway implicated in the genotoxicity of ochratoxin A? In *Human Ochratoxicosis and Its Pathologies*; Creppy, E.E., Castegnaro, M., Dirheimer, G., Eds.; John Libbey Euro-Text: Montrouge, France, 1993; Volume 231, pp. 177–187.
- Hadjeba-Medjdoub, K.; Tozlovanu, M.; Pfohl-Leszkowicz, A.; Frenette, C.; Paugh, R.J.; Manderville, R.A. Structure-activity relationships imply different mechanisms of action for ochratoxin A-mediated cytotoxicity and genotoxicity. *Chem. Res. Toxicol.* 2012, 25, 181–190.
- 62. Monks, T.J.; Lau, S.S. Biological reactivity of polyphenolic-glutathione conjugates. *Chem. Res. Toxicol.* **1997**, *10*, 1296–1313.
- 63. Rahman, K.M.; James, C.H.; Thurston, D.E. Observation of the reversibility of a covalent pyrrolobenzodiazepine (PDB) DNA adduct by HPLC/MS and CD spectroscopy. *Org. Biomol. Chem.* **2011**, *9*, 1632–1641.
- 64. Obrecht-Pflumio, S.; Dirheimer, G. *In vitro* DNA and dGMP adducts formation caused by ochratoxin A. *Chem. Biol. Interact.* **2000**, *127*, 29–44.

65. Obrecht-Pflumio, S.; Dirheimer, G. Horseradish peroxidase mediates DNA and deoxyguanosine 3'-monophosphate adduct formation in the presence of ochratoxin A. *Arch. Toxicol.* **2001**, *75*, 583–590.

 \bigcirc 2012 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (http://creativecommons.org/licenses/by/3.0/).