



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review article

The nervous system—A new territory being explored of SARS-CoV-2

Meng Zhang^{a,1}, Lingyan Zhou^{a,1}, Jing Wang^b, Kun Wang^a, Yuan Wang^a, Xudong Pan^a, Aijun Ma^{a,*}^a Department of Neurology, The Affiliated Hospital of Qingdao University, Qingdao 266000, China^b Department of Rheumatology, The Affiliated Hospital of Qingdao University, Qingdao 266000, China

ARTICLE INFO

Article history:

Received 22 July 2020

Accepted 21 October 2020

Keywords:

Nervous system

COVID-19

SARS-CoV-2

Neuroinvasion

Epidemic

ABSTRACT

In December 2019, COVID-19 outbreaked in Wuhan, then sweeping the mainland of China and the whole world rapidly. On March 4, Beijing Ditan Hospital confirmed the existence of SARS-CoV-2 in the cerebrospinal fluid by gene sequencing, indicating the neurotropic involvement of SARS-CoV-2. Meanwhile, neurological manifestations in the central nervous system, peripheral nervous system and skeletal muscular were also observed, indicating the potential neuroinvasion of SARS-CoV-2. In particular, we focused on its neurological manifestations and specific pathogenesis, as well as its comparison with other viral respiratory infections. Finally, we further summarized the significance of the neuroinvasion and the follow-up issues that need to be paid attention to by scientists, so as to help neurologists understand the influence of SARS-CoV-2 on nervous system better and promote the accurate diagnosis and efficient treatment of COVID-19.

© 2020 Elsevier Ltd. All rights reserved.

1. Introduction

In December 2019, a novel pneumonia of coronavirus, Corona Virus Disease 2019 (COVID-19) outbreaked in Wuhan, China, then sweeping the mainland of China and the whole world rapidly, seriously threatening individuals' life and health [1–4]. Due to the high homology with the severe acute respiratory syndrome coronavirus (SARS-CoV), the pathogen was named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) [5].

The main clinical symptoms involved by SARS-CoV-2 is pulmonary manifestations. However, there is growing evidence that SARS-CoV-2 can result in a broad spectrum of neurologic diseases [6–9], which is not surprising, as neurological manifestations have been reported in other respiratory viral infections, including coronavirus, but the nervous system manifestations of COVID-19 are more common and disabling, raising the worldwide concerns about its potential long-term complications to humans [10,11].

In particular, we focused on its neurological manifestations and specific pathogenesis, as well as its comparison with other viral respiratory infections. Finally, we further summarized the significance

of the neuroinvasion and the follow-up issues that need to be paid attention to by scientists, so as to help neurologists understand the influence of SARS-CoV-2 on nervous system better and promote the accurate diagnosis and efficient treatment of COVID-19.

2. SARS-CoV-2

SARS-CoV-2 is a non-segmented, large enveloped, positive single-stranded RNA virus, which is the member of Betacoronavirus genus, with an overall genomic sequence consistency of 96.2% and 79.5% sequence homology with the SARS coronavirus [12]. It has been discovered that SARS-CoV-2 combined with a metalloproteinase, angiotensin-converting enzyme 2 (ACE2) receptors to infect the target cells on their surfaces in line with SARS-CoV, which requires the participation of spike protein(S) and transmembrane protein serine protease 2 (TMPRSS2) [12,13]. Therefore, neurons, endothelial cells, glial cells and arterial smooth muscle cells in the brain, which expressing ACE2 receptors become vulnerable to it [14,15]. Besides, recent researches found that SARS-CoV-2 has a higher affinity for human ACE2 than SARS-CoV [16], which further elucidate the reason for the strong pathogenicity and transmissibility of it.

People are generally susceptible to SARS-CoV-2 [17]. It has shown the aged has higher susceptibility to develop severe sickness, especially with specific comorbidities. Additionally, about

* Corresponding author.

E-mail address: maaijun541@163.com (A. Ma).¹ These authors contributed to this work equally and should be regarded as co-first authors.

half of the patients infected with SARS-CoV2 had chronic diseases, including cardiocerebrovascular disease as well as diabetes [3].

3. The potential neuroinvasion of SARS-CoV-2

3.1. Initial evidence

A retrospective study [18] showed that among 214 patients diagnosed with COVID-19, more than 30% of them had neurological manifestations. On March 4, Beijing Ditan Hospital confirmed the existence of SARS-CoV-2 in the cerebrospinal fluid by gene sequencing, indicating SARS-CoV-2 could invade the nervous system directly. Moreover, it became the first evidence that SARS-CoV-2 may have the potential neuroinvasion. In fact, the earliest cases of encephalitis/meningitis occurred in Japan [19]. A man in twenties was confirmed as a victim of SARS-CoV-2, who started with fever and fatigue at first, and he was found unconscious at home. During emergency treatment, he suffered from epileptic seizures several times. It was also found SARS-CoV-2 in his cerebrospinal fluid in 8 March, which is the further evidence that the virus is attacking the nervous system.

3.2. Speculated pathogenesis

The central nervous system(CNS) is essential to regulate the whole human body, but it has no immune function against most infections, especially the virus infection. The blood–brain barrier (BBB) is a natural cover for defence, preventing from the intracranial disease of the foreign pathogen. CoVs can enter the CNS from the respiratory tract via hematogenous and transneuronal routes. Increasing evidence shows that a variety of CoVs can not only infect the respiratory system but also have the neurotropism feature [19–24].

The existing researches mainly focus on SARS-CoV. In 2004, the virus RNA was detected in the cerebrospinal fluid of a SARS patient, which led to the scientists' suspicion that SARS-CoV was neuroinvasive [25]. One year later, Xu et al. isolated the virus from the brain tissue of a patient, and the neuropathologic manifestation was nerve cell damage. Meanwhile, the patient showed neurological symptoms, making the hypothesis confirmed [26]. Furthermore, extensive neuronal infections can be observed in animal models.

Furthermore, SARS-CoV entered human cells through a receptor named as angiotensin-converting enzyme 2 (ACE2), which is consistent with SARS-CoV-2 [27]. Therefore, SARS-CoV is neuroinvasive and neurotropic in both animals and humans, which may be related to the development of neurological diseases. Given SARS-CoV2 has similar gene sequencing and pathologic anatomy with SARS-CoV, it could be assumed that SARS-CoV-2 has the same neuroinvasion and pathogenesis.

SARS-CoV-2 might cause a series of neurological damage by two mechanisms; one is the hypoxic brain injury. It is well known that hypoxia, hypercapnia, endocrine and metabolic disorders, and the accumulation of toxic substances caused by respiratory failure in severe COVID-19 patients can cause neuronal swelling and brain edema, leading to brain damage [28]. The other is the parainfectious mechanisms, that is, immune-mediated brain damage. It is mainly referred to cytokine storm [29]. Cytokine storm syndrome is a highly inflammatory state characterized by sharply elevated cytokine levels, overshooting immune responses and explosive multi-organ failure, leading to vasodilatory dysfunction, membrane leakage, coagulation dysfunction, multiple organ failure, and severe vasospasm shocks. Besides, if ECMO is used, the contact between fluids in vivo and

circuits in vitro might activate clotting and inflammatory pathways, leading to disseminated intravascular coagulation in severe cases [30,31].

3.3. Significance of neuroinvasion

Firstly, given that SARS-CoV-2 has the underlying neuroinvasion, carrying out the therapy to resist it as soon as possible can block the entry of the virus to the CNS. Moreover, the ability of antiviral agents to cross the blood–brain barrier is also an issue to consider in the future. Although the progression of new antiviral drugs requires a lot of workforce and material resources, it can not benefit the general public a lot in the current epidemic. However, as the outbreak fades, the virus may mutate into a coronavirus, which is similar to the influenza virus, spreading from person to person over the long term. Therefore, it is necessary to accelerate the research and development of specific antiviral drugs and vaccines steadily.

Secondly, considering that SARS-CoV2 might be latent in some neurons to get away with immune surveillance, we can not guarantee that the virus has been eliminated. Actually, SARS-CoV-2 is still can be detected; even the COVID-19 patients are under the convalescent period [32]. If existing in the nervous system for the long term, it is likely to serve as a trigger for some human neurological diseases in those genetically predisposed individuals. Thus the long-term follow-up of patients is an essential task in the future.

Thirdly, the blood-brain barrier degrades gradually with age, which indicates that the nerve invasion ability of the virus to the elderly increases. Once the virus invades the nervous system, demyelination, cell apoptosis and neurodegeneration will occur, then further aggravating the brain ageing and neurodegenerative diseases [33]. Neurologists will face more age-related neurodegenerative diseases after we control the outbreak. Therefore, more long-term neurological follow-up of the elderly is needed after SARS-COV-2 infection.

The final concern is the immuno-therapies for COVID-19 with autoimmune diseases of the nervous system. Once diagnosed, neurologists must assess the risks and benefits of immunotherapy in light of the different conditions of the disease. For example, high doses or long-term use of steroid hormones may put patients at further risk for COVID-19 infection. Protective immunoglobulin is recommended if possible. Additionally, given that the crucial step to control the pandemic is the application of SARS-COV-2 vaccination, it is indispensable to weigh the safety and effectiveness of vaccination during immunotherapy.

4. Neurological manifestations of COVID-19

An increasing number of evidence indicated that SARS-CoV-2 might have potential neuroinvasion, hence we summarized the existing related researches and data of COVID-19, the neurological signs of it could be divided into three categories: symptoms involving the central nervous system, peripheral nervous system, and skeletal, muscular symptoms.

4.1. Central nervous system (CNS) diseases or symptoms

4.1.1. Acute cerebrovascular diseases

The virus infections can cause vascular endothelial injury and vascular system damage, then leading to ischemic and hemorrhagic infarcts through overactive inflammation response, thrombosis and vasculitis [34,35].

A retrospective case series study in Wuhan, China [18], showed that the severe patients of COVID-19 commonly had acute cere-

brovascular diseases, and the pathophysiological changes during the infection may render the victims prone to it.

Firstly, in the latest study, 52% of patients of COVID-19 had elevated IL-6 levels, and 86% of them had elevated CRP(3), suggesting a significant inflammatory response in their bodies. Meanwhile, inflammation is also essential in the occurrence, development as well as prognosis of cerebrovascular diseases and may trigger cerebrovascular events [36]. Secondly, according to the laboratory examination of the severe patients who had coagulation dysfunction and elevated D-dimer [37], those people may have more possibility of venous thrombosis and hemorrhagic stroke [38]. A 75-year-old woman infected by SARS-CoV-2 without any predisposing factors suffered from severe bilateral pneumonia and acute pulmonary embolism, suggesting the severe infections is a precipitant factor of acute venous embolism and stroke [40,41].

Thirdly, SARS-CoV-2 readily attacks the lung, then causes dyspnea and decreased blood oxygen saturation [3]. Hypoxemia attributes the alteration of consciousness, confusion or delirium, causing an acute or subacute stroke or intracerebral hemorrhage [41,42]. Additionally, overexpression of ACE2 reduces the risk of ischemic stroke, which explains the reason why elderly COVID-19 patients are more likely to have a stroke [43,44].

4.1.2. Viral encephalitis and meningitis

Although the virus seems to have a hard time penetrating the central nervous system, pathogens can be abnormally active in spreading and replicating, then may induce overreacting immune response and lead to fatal meningitis and encephalitis. The discovery of the positive RT-PCR of CSF [19] suggested that meningitis and encephalitis may associated with viral invasion of the CNS. Elevated temperature, headache, vomiting, and consciousness disorders are common signs, which are similar to acute encephalitis [45]. Autopsy reports also showed that there was the edema of brain tissue and degeneration in the neurons of those victims [20]. Presumably, SARS-CoV-2 may enter the CNS from the respiratory tract via hematogenous and transneuronal routes like other CoVs [46].

4.1.3. Acute hemorrhagic necrotizing Encephalopathy/Acute disseminated encephalomyelitis

Acute necrotizing encephalopathy is a rare acute and severe explosive encephalopathy, which usually occurs after a viral infection. The most common hypothesis for its pathogenesis is the destruction of the blood–brain barrier caused by cytokine storm [27], which is also the possible pathogenesis of COVID-19 [47]. Neo Poyiadji et al. reported the first case of acute necrotizing hemorrhagic encephalopathy associated with COVID-19 from a woman in her fifties. For the first three days, her symptoms were fever, cough and altered mental state. Laboratory tests confirmed the SARS-CoV-2 infection and neuroimaging showed characteristic changes: symmetrical multifocal changes and thalamic invasion [48]. Reichard et al also found pathological findings which are similar to acute disseminated encephalomyelitis in a patient who died from complications of COVID-19 [49]. However, whether the neuropathological lesions resulted from primary vascular disease secondary to white matter damage or demyelination disease caused via a parainfectious mechanism.

4.1.4. Leukoencephalopathy

Radmanesh et al. observed two kinds of imaging characteristics by cerebral MRI in 11 patients with severe COVID-19: diffuse leukoencephalopathy and microhemorrhages, speculating that they were late complications in severe COVID-19 patients [50]. Given that similar involvement patterns have been described in patients after anoxic injury, it was considered delayed post hypoxic leukoencephalopathy (DPHL) which was associated with oligoden-

drocyte damage and demyelination [51,52]. Besides, Sachs et al also reported a COVID-19 patient who suffered from leukoencephalopathy with microhemorrhage. It is a 59-year-old man who was hospitalized for upper respiratory symptoms and fever. Brain MRI mainly showed large posterior fused white matter lesions, significant microhemorrhage in the corpus callosum and high posterior circulation perfusion [53], which could also be seen in acute disseminated encephalomyelitis, acute hemorrhagic encephalomyelitis, vasculitides and posterior reversible encephalopathy syndrome [54,55]. Recently, Franceschi et al describe 2 COVID-19 patients whose brain imaging suggested hemorrhagic posterior reversible encephalopathy syndrome [56]. The authors summarized three mechanisms of PRES: cytokine release syndrome which breaks down the BBB; hypoxia caused by inflammation; unstable blood pressure [57,58].

4.1.5. Acute myelitis

A 66-year-old male in Wuhan was diagnosed as COVID-19 with acute myelitis [59], presenting with fever, pain, and acute delayed paralysis of the lower extremities along with urinary and fecal incontinence. The patients responded well to immunoglobulin and hormone. Meanwhile, the high levels of C-reactive protein, inflammatory cytokines, and serum ferritin suggested that his myelitis was a result of a cytokine storm.

4.1.6. Others

There are also other central nervous system manifestations reported [18] such as headache, dizziness, ataxia, impaired consciousness, and epilepsy. Dizziness and headache are the most common complaints. A headache expert pointed out that there are two stages of COVID-19-related headache. The first stage is an acute headache caused by a viral infection, primary cough, tension-type and heterophoria. The headache in the second stage (bounded by 7–10 days) could attribute to hypoxia, which could be used as a predictor of cytokine storm onset [60]. In the patients who had severe virus infection also showed agitation, corticospinal tract signs as well as a dysexecutive syndrome [61]. A COVID-19 patient with meningitis/encephalitis also developed epilepsy [19], and another present as focal status epilepticus as the first symptom [62]. Therefore, it is reasonable to speculate that COVID-19 patients may develop epilepsy under conditions of hypoxia, endocrine and metabolic disorders, and multiple organ damage.

In addition, anosmia may occur before the symptoms of viral infection. It has been reported that CoVs can enter the CNS in the early stages of infection through the olfactory nerve [63], which indicates the same transneuronal routes of SARS-CoV-2. A multicenter study in European [64] found olfactory disorder in patients who did not need intensive cares. And the impaired functions do not always recover as the disappearance of the typical infection symptoms. Meanwhile, compared to the work of MAO et al. in Wuhan, China, the prevalence of the dysfunctions was significantly higher in European. Considering the different affinity of SARS-CoV-2 to different populations and the diversity of ACE2 expression [65], studies should focus more on the polymorphisms of ACE2 in different races.

4.2. Peripheral nervous system (PNS) symptoms

4.2.1. Guillain-Barré syndrome

Guillain-Barrésyndrome is a kind of neuroimmune disease secondary to respiratory or gastrointestinal infections. It is known that there is a molecular imitation mechanism. The invading virus has similar epitopes with some components of the nervous system, stimulating the immune system to produce antibodies which can not only bind to the virus, but also cross-react with components

of the nervous system, then causing a series of neurological dysfunction [66].

The first Guillain-Barré syndrome associated with COVID-19 case was found in Wuhan, China [68]. A 61-year-old woman suddenly developed progressive lower limb weakness without fever or cough. Three days later, her condition worsened, and the weakness spread to the upper limbs. Typical blood and cerebrospinal fluid test and nerve conduction have confirmed the presence of Guillain-Barré syndrome. On the eighth day, she started to have the characteristic fever, cough, and chest CT findings of COVID-19. Laboratory tests confirmed the SARS-CoV-2 infection. Then the northern Italy hospitals reported similar cases, involving motor and sensory disorders of the limbs as well as ataxia [67]. Jessica et al reported a COVID-19 patient admitted to hospital with an abnormal gait. Although the neurologist did not observe ophthalmoplegia, in consideration of her excellent response to immunoglobulin, apparent ataxia and hypoxia of the lower extremities, she was eventually diagnosed as Miller-Fisher-like syndrome overlapping with COVID-19 infection [69]. Consider that these are all observational studies, more epidemiological evidence is needed to support the potential relationship between Guillain-Barré syndrome and COVID-19 in the future.

4.2.2. Others

Hypoxemia is also a common symptom in COVID-19 patients. Acute respiratory symptoms associated with coronavirus are characterized by hypoxia, one of the ways that coronavirus associated with the nervous system, causing endocrine and metabolic disorders, edema in the brain and subsequent neurological manifestations. Besides, taste, vision impairment, and facial pain have also been reported [44].

4.3. Skeletal muscular symptoms

Skeletal muscular symptoms were also reported in COVID-19 patients [18]. Given the high levels of lactate dehydrogenase and creatine kinase, it is speculated that the symptoms resulted from skeletal muscle injury which might be associated with ACE2 existing in it [44]. Nevertheless, whether SARS-CoV-2 damages skeletal muscle through the receptor on the cells still needs further researches. Moreover, the elevated pro-inflammatory cytokines, which indicated abnormal immune response mediated by infection, might be another reason for it.

5. Comparison with other viral respiratory infections

A great many viral infections could cause varying degrees of damage to the nervous system, including its the structure and function. Such as acute disseminated encephalomyelitis, toxic encephalopathy and severe acute demyelinating diseases in connection with the post-infectious inflammatory response. The presence of the virus in cerebrospinal fluid also confirms the nerve invasiveness of the virus [70,71]. As SARS-COV-2 has stronger binding ability to ACE2, it is more pathogenic and transmissible as well as debilitating than SARS-CoV and MERS-CoV [72]. At the same time, COVID-19 and acute respiratory distress syndrome patients had twice the rate of thrombosis complications compared to the ARDS from other causes, which explains why the rates of COVID-19 associated d-dimer elevation and concurrent stroke appear to be much higher than other viruses. Therefore, timely anticoagulant therapy is essential for the prognosis of patients [73].

6. Conclusion

In conclusion, SARS-CoV-2 has potential neuroinvasion and may cause a series of nervous system diseases or symptoms. Once involving the nervous system, it usually predicts poor prognosis. Therefore, patient with COVID-19 requires more physicians'attention to evaluate their neurologic manifestations as early as possible. Paying attention to the patients who have neurological symptoms as the first manifestation other than the typical cough, vomiting, diarrhea, and fever is imperative for the control of the rapidly evolving epidemic. Gene sequencing of cerebrospinal fluid and other neuroimaging examinations should also be taken in necessity.

7. Fundings

No funding.

8. Ethics approval

This study was based on publicly available data and did not require ethical approval.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Chan J-W, Yuan S, Kok K-H, To K-W, Chu H, Yang J, Xing F, Liu J, Yip C-Y, Poon R-S, Tsoi H-W, Lo S-F, Chan K-H, Poon V-M, Chan W-M, Ip JD, Cai J-P, Cheng V-C, Chen H, Hui C-M, Yuen K-Y. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;395(10223):514–23. [https://doi.org/10.1016/S0140-6736\(20\)30154-9](https://doi.org/10.1016/S0140-6736(20)30154-9).
- [2] Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KSM, Lau EHY, Wong JY, Xing X, Xiang N, Wu Y, Li C, Chen Qi, Li D, Liu T, Zhao J, Liu M, Tu W, Chen C, Jin L, Yang R, Wang Qi, Zhou S, Wang R, Liu H, Luo Y, Liu Y, Shao Ge, Li H, Tao Z, Yang Y, Deng Z, Liu B, Ma Z, Zhang Y, Shi G, Lam TTY, Wu JT, Gao GF, Cowling BJ, Yang Bo, Leung GM, Feng Z. Early transmission dynamics in wuhan, china, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020;382(13):1199–207. <https://doi.org/10.1056/NEJMoa2001316>.
- [3] Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang Li. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395(10223):507–13. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7).
- [4] Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet* 2020;395(10223):470–3. [https://doi.org/10.1016/S0140-6736\(20\)30185-9](https://doi.org/10.1016/S0140-6736(20)30185-9).
- [5] Lu R, Zhao X, Li J, Niu P, Yang Bo, Wu H, Wang W, Song H, Huang B, Zhu Na, Bi Y, Ma X, Zhan F, Wang L, Hu T, Zhou H, Hu Z, Zhou W, Zhao Li, Chen J, Meng Y, Wang Ji, Lin Y, Yuan J, Xie Z, Ma J, Liu WJ, Wang D, Xu W, Holmes EC, Gao GF, Wu G, Chen W, Shi W, Tan W. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020;395(10224):565–74. [https://doi.org/10.1016/S0140-6736\(20\)30251-8](https://doi.org/10.1016/S0140-6736(20)30251-8).
- [6] De Felice FG, Tovar-Moll F, Moll J, Munoz DP, Ferreira ST. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the central nervous system. *Trends Neurosci* 2020;43(6):355–7. <https://doi.org/10.1016/j.tins.2020.04.004>. Epub 2020 Apr 21. PMID: 32359765; PMCID: PMC7172664.
- [7] Abboud H, Abboud FZ, Kharbouch H, Arkha Y, El Abbadi N, El Ouahabi A. COVID-19 and SARS-Cov-2 infection: pathophysiology and clinical effects on the nervous system. *World Neurosurg* 2020;140:49–53. <https://doi.org/10.1016/j.wneu.2020.05.193>. Epub 2020 May 28. PMID: 32474093; PMCID: PMC7255736.
- [8] Pennisi M, Lanza G, Falzone L, Fiscaro F, Ferri R, Bella R. SARS-CoV-2 and the nervous system: from clinical features to molecular mechanisms. *Int J Mol Sci* 2020;21(15):5475. <https://doi.org/10.3390/ijms21155475>. PMID: 32751841; PMCID: PMC7432482.
- [9] Li H, Xue Q, Xu X. Involvement of the nervous system in SARS-CoV-2 infection. *Neurotox Res* 2020;38(1):1–7. <https://doi.org/10.1007/s12640-020-00219-8>. Epub 2020 May 13. PMID: 32399719; PMCID: PMC7220627.

- [10] Pleasure SJ, Green AJ, Josephson SA. The spectrum of neurologic disease in the severe acute respiratory syndrome coronavirus 2 pandemic infection: neurologists move to the frontlines. *JAMA Neurol* 2020;77(6):679–80. <https://doi.org/10.1001/jamaneurol.2020.1065>. PMID: 32275291.
- [11] Marshall M. How COVID-19 can damage the brain. *Nature* 2020;585(7825):342–3. <https://doi.org/10.1038/d41586-020-02599-5>. PMID: 32934351.
- [12] Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A, Müller MA, Drosten C, Pöhlmann S. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181(2):271–280.e8. <https://doi.org/10.1016/j.cell.2020.02.052>.
- [13] Zou X, Chen Ke, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* 2020;14(2):185–92. <https://doi.org/10.1007/s11684-020-0754-0>.
- [14] Alenina N, Bader M. ACE2 in brain physiology and pathophysiology: evidence from transgenic animal models. *Neurochem Res* 2019;44(6):1323–9. <https://doi.org/10.1007/s11064-018-2679-4>.
- [15] Baig AM, Khaleeq A, Ali U, Syeda H. Evidence of the COVID-19 virus targeting the CNS: tissue distribution, host–virus interaction, and proposed neurotropic mechanisms. *ACS Chem Neurosci* 2020;11(7):995–8. <https://doi.org/10.1021/acscchemneuro.0c00122>.
- [16] Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, Graham BS, McLellan JS. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367(6483):1260–3. <https://doi.org/10.1126/science.abb2507>.
- [17] Wang D, Hu Bo, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, Zhao Y, Li Y, Wang X, Peng Z. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061. <https://doi.org/10.1001/jama.2020.1585>.
- [18] Mao L, Jin H, Wang M, Hu Yu, Chen S, He Q, Chang J, Hong C, Zhou Y, Wang D, Miao X, Li Y, Hu Bo. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol* 2020;77(6):683. <https://doi.org/10.1001/jamaneurol.2020.1127>.
- [19] Moriguchi T, Harii N, Goto J, Harada D, Sugawara H, Takamino J, Ueno M, Sakata H, Kondo K, Myose N, Nakao A, Takeda M, Haro H, Inoue O, Suzuki-Inoue K, Kubokawa K, Ogihara S, Sasaki T, Kinouchi H, Kojin H, Ito M, Onishi H, Shimizu T, Sasaki Yu, Enomoto N, Ishihara H, Furuya S, Yamamoto T, Shimada S. A first case of meningitis/encephalitis associated with SARS-Coronavirus-2. *Int J Infect Dis* 2020;94:55–8. <https://doi.org/10.1016/j.ijid.2020.03.062>.
- [20] Li Y-C, Bai W-Z, Hirano N, Hayashida T, Hashikawa T. Coronavirus infection of rat dorsal root ganglia: ultrastructural characterization of viral replication, transfer, and the early response of satellite cells. *Virus Res* 2012;163(2):628–35. <https://doi.org/10.1016/j.virusres.2011.12.021>.
- [21] M. Dubé A. Le Coupancec A.H.M. Wong J.M. Rini M. Desforges P.J. Talbot M.S. Diamond Axonal Transport Enables Neuron-to-Neuron Propagation of Human Coronavirus OC43. *J Virol* 92 17 10.1128/JVI.00404-18 <https://jvi.asm.org/content/92/17/e00404-18>.
- [22] Talbot PJ, Ekandé S, Cashman NR, Mounir S, Stewart JN. Neurotropism of human coronavirus 229E. *Adv Exp Med Biol* 1993;342:339–46. https://doi.org/10.1007/978-1-4615-2996-5_52.
- [23] Lau KK, Yu WC, Chu CM, Lau ST, Sheng B, Yuen KY. Possible central nervous system infection by SARS coronavirus. *Emerg Infect Dis* 2004;10(2):342–4. <https://doi.org/10.3201/eid1002.030638>.
- [24] Xu J, Zhong S, Liu J, Li L, Li Y, Wu X, Li Z, Deng P, Zhang J, Zhong N, Ding Y, Jiang Y. Detection of severe acute respiratory syndrome coronavirus in the brain: potential role of the chemokine mig in pathogenesis. *Clin Infect Dis* 2005;41(8):1089–96.
- [25] Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. *J Virol* 2008;82(15):7264–75.
- [26] Tu H, Tu S, Gao S, Shao A, Sheng J. The epidemiological and clinical features of COVID-19 and lessons from this global infectious public health event. *J Infect* 2020. <https://doi.org/10.1016/j.ijinf.2020.04.011>. pii: S0163-4453(20)30222-X.
- [27] Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395(10229):1033–4.
- [28] Tveito K. Cytokinstormer ved covid-19? *Tidsskriftet* 2020. <https://doi.org/10.4045/tidsskr.20.0239>.
- [29] Chousterman BG, Swirski FK, Weber GF. Cytokine storm and sepsis disease pathogenesis. *Semin Immunopathol* 2017;39(5):517–28. <https://doi.org/10.1007/s00281-017-0639-8>.
- [30] von Bahr V, Millar JE, Malferrtheiner MV, Ki KK, Passmore MR, Bartnikowski N, Redd MA, Cavaye M, Suen JY, McAuley DF, Fraser JF. Mesenchymal stem cells may ameliorate inflammation in an ex vivo model of extracorporeal membrane oxygenation. *Perfusion* 2019;34(1_suppl):15–21. <https://doi.org/10.1177/0267659119830857>.
- [31] Rothe C, Schunk M, Sothmann P, Bretzel G, Froeschl G, Wallrauch C, Zimmer T, Thiel V, Janke C, Guggemos W, Seilmaier M, Drosten C, Vollmar P, Zwirgmaier K, Zange S, Wölfel R, Hoelscher M. Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *N Engl J Med* 2020;382(10):970–1. <https://doi.org/10.1056/NEJMc2001468>.
- [32] Hascup ER, Hascup KN. Does SARS-CoV-2 infection cause chronic neurological complications?. *GeroScience* 2020;42(4):1083–7. <https://doi.org/10.1007/s11357-020-00207-y>.
- [33] Montagne A, Barnes S, Sweeney M, Halliday M, Sagare A, Zhao Z, Toga A, Jacobs R, Liu C, Amezcua L, Harrington M, Chui H, Law M, Zlokovic B. Blood-brain barrier breakdown in the aging human hippocampus. *Neuron* 2015;85(2):296–302. <https://doi.org/10.1016/j.neuron.2014.12.032>.
- [34] Manousakis G, Jensen MB, Chacon MR, Sattin JA, Levine RL. The interface between stroke and infectious disease: infectious diseases leading to stroke and infections complicating stroke. *Curr Neurol Neurosci Rep* 2009;9(1):28–34. <https://doi.org/10.1007/s11910-009-0005-x>.
- [35] Naveed M, Zhou Q-G, Han F. Cerebrovascular inflammation: a critical trigger for neurovascular injury?. *Neurochem Int* 2019;126:165–77.
- [36] Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *J Med Virol* 2020;92(6):568–76. <https://doi.org/10.1002/jmv.25748>.
- [37] Prandoni P, Vedovetto V, Ciammaichella M, Bucherini E, Corradini S, Enea I, Cosmi B, Mumoli N, Visonà A, Barillari G, Bova C, Quintavalla R, Zanatta N, Pedrini S, Villalta S, Camporese G, Testa S, Parisi R, Becattini C, Cuppini S, Pengo V, Palareti G. Residual vein thrombosis and serial D-dimer for the long-term management of patients with deep venous thrombosis. *Thromb Res* 2017;154:35–41. <https://doi.org/10.1016/j.thromres.2017.04.002>.
- [38] Zheng J, Cheng C, Zhou C, Chen H, Guo Z, Sun X. The Clinical Value of d-Dimer Level in Patients with Nonaneurysmal Subarachnoid Hemorrhage. *World Neurosurgery* 2018;114:e1161–7. <https://doi.org/10.1016/j.wneu.2018.03.165>.
- [39] G.B. Danzi M. Loffi G. Galeazzi E. Gherbesi Acute pulmonary embolism and COVID-19 pneumonia: a random association? 41 19 2020 2020 1858 1858 <https://academic.oup.com/eurheartj/article/41/19/1858/5813284>.
- [40] Rowat AM, Dennis MS, Wardlaw JM. Hypoxaemia in acute stroke is frequent and worsens outcome. *Cerebrovasc Dis* 2006;21(3):166–72. <https://doi.org/10.1159/000090528>.
- [41] Tang Y, Lu A, Aronow BJ, Wagner KR, Sharp FR. Genomic responses of the brain to ischemic stroke, intracerebral haemorrhage, kainate seizures, hypoglycemia, and hypoxia: Genomic responses of the brain to injury. *Eur J Neurosci* 2002;15(12):1937–52. <https://doi.org/10.1046/j.1460-9568.2002.02030.x>.
- [42] Chen Ji, Xiao X, Chen S, Zhang C, Chen J, Yi D, Shenoy V, Raizada MK, Zhao B, Chen Y. Angiotensin-converting enzyme 2 priming enhances the function of endothelial progenitor cells and their therapeutic efficacy. *Hypertension* 2013;61(3):681–9. <https://doi.org/10.1161/HYPERTENSIONAHA.111.00202>.
- [43] Chen Ji, Zhao Y, Chen S, Wang J, Xiao X, Ma X, Penchikala M, Xia H, Lazartigues E, Zhao B, Chen Y. Neuronal over-expression of ACE2 protects brain from ischemia-induced damage. *Neuropharmacology* 2014;79:550–8. <https://doi.org/10.1016/j.neuropharm.2014.01.004>.
- [44] Cabello-Verrugio C, Morales MG, Rivera JC, Cabrera D, Simon F. Renin-Angiotensin System: An Old Player with Novel Functions in Skeletal Muscle: ras regulates the physiopathology of skeletal muscle. *Med Res Rev* 2015;35(3):437–63. <https://doi.org/10.1002/med.21343>.
- [45] Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu Li, Tai Y, Cai C, Gao T, Song J, Xia P, Dong J, Zhao J, Wang F-S. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respirat Med* 2020;8(4):420–2.
- [46] Mizuguchi M, Yamanouchi H, Ichiyama T, Shiomi M. Acute encephalopathy associated with influenza and other viral infections. *Acta Neurol Scand* 2007;115(s186):45–56.
- [47] Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B. COVID-19-associated Acute Hemorrhagic Necrotizing Encephalopathy: CT and MRI Features [published online ahead of print, 2020 March 31]. *Radiology* 2020;201187. <https://doi.org/10.1148/radiol.2020201187>.
- [48] Reichard RR, Kashani KB, Boire NA, Constantopoulos E, Guo Y, Lucchinetti CF. Neuropathology of COVID-19: a spectrum of vascular and acute disseminated encephalomyelitis (ADEM)-like pathology. *Acta Neuropathol* 2020;140(1):1–6. <https://doi.org/10.1007/s00401-020-02166-2>.
- [49] Radmanesh A, Derman A, Lui YW, Raz E, Loh JP, Hagiwara M, Borja MJ, Zan E, Fatterpekar GM. COVID-19-associated Diffuse Leukoencephalopathy and Microhemorrhages. *Radiology* 2020;297(1):E223–7.
- [50] Breit H, Jhaveri M, John S. Concomitant delayed posthypoxic leukoencephalopathy and critical illness microbleeds. *Neurol Clin Pract* 2018;8(5):e31–3.
- [51] Ginsberg MD, Hedley-Whyte ET, Richardson Jr EP. Hypoxic-ischemic leukoencephalopathy in man. *Trans Am Neurol Assoc* 1974;99:213–6.
- [52] Sachs JR, Gibbs KW, Swor DE, Sweeney AP, Williams DW, Burdette JH, West TG, Geer CP. COVID-19-associated leukoencephalopathy. *Radiology* 2020;296(3):E184–5.
- [53] Banerjee A, Suthar R, Vyas S, Singh MP. Acute disseminated encephalomyelitis: complication of a vaccine preventable disease. *BMJ Case Rep* 2018;11(1):e225710. <https://doi.org/10.1136/bcr-2018-225710>.
- [54] Pinto PS, Taipa R, Moreira B, Correia C, Melo-Pires M. Acute hemorrhagic leukoencephalitis with severe brainstem and spinal cord involvement: MRI features with neuropathological confirmation. *J Magn Reson Imaging* 2011;33(4):957–61.
- [55] Abdel Razek AK, Alvarez H, Bagg S, Refaat S, Castillo M. Imaging Spectrum of CNS Vasculitis. *RadioGraphics* 2014;34(4):873–94.

- [56] Franceschi AM, Ahmed O, Giliberto L, Castillo M. Hemorrhagic Posterior Reversible Encephalopathy Syndrome as a Manifestation of COVID-19 Infection. *AJNR Am J Neuroradiol* 2020;41(7):1173–6.
- [57] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Yi, Zhang Li, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo Li, Xie J, Wang G, Jiang R, Gao Z, Jin Qi, Wang J, Cao B. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020;395(10223):497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- [58] Bartels K, Grenz A, Eltzschig HK. Hypoxia and inflammation are two sides of the same coin. *Proc Natl Acad Sci* 2013;110(46):18351–2. <https://doi.org/10.1073/pnas.1318345110>.
- [59] Munz M, Wessendorf S, Koretsis G, Tewald F, Baegi R, Krämer S, Geissler M, Reinhard M. Acute transverse myelitis after COVID-19 pneumonia. *J Neurol* 2020;267(8):2196–7. <https://doi.org/10.1007/s00415-020-09934-w>.
- [60] Belvis R. Headaches During COVID-19: my clinical case and review of the literature. *Headache J Head Face Pain* 2020;60(7):1422–6. <https://doi.org/10.1111/head.13841>.
- [61] Helms J, Kremer S, Merdji H, Clere-Jehl R, Schenck M, Kummerlen C, Collange O, Boulay C, Fafi-Kremer S, Ohana M, Anheim M, Meziani F. Neurologic Features in Severe SARS-CoV-2 Infection. *N Engl J Med* 2020;382(23):2268–70.
- [62] Vollono C, Rollo E, Romozzi M, Frisullo G, Servidei S, Borghetti A, Calabresi P. Focal status epilepticus as unique clinical feature of COVID-19: A case report. *Seizure* 2020;78:109–12. <https://doi.org/10.1016/j.seizure.2020.04.009>.
- [63] Giacomelli A, Pezzati L, Conti F, et al. Self-reported olfactory and taste disorders in SARS-CoV-2 patients: a cross-sectional study [published online ahead of print, 2020 March 26] *ciaa330. Clin Infect Dis* 2020. <https://doi.org/10.1093/cid/ciaa330>.
- [64] Lechien JR, Chiesa-Estomba CM, De Siati DR, Horoi M, Le Bon SD, Rodriguez A, Dequanter D, Blecic S, El Afia F, Distinguin L, Chekkoury-Idrissi Y, Hans S, Delgado IL, Calvo-Henriquez C, Lavigne P, Falanga C, Barillari MR, Cammaroto G, Khalife M, Leich P, Souchay C, Rossi C, Journe F, Hsieh J, Edjlali M, Carlier R, Ris L, Lovato A, De Filippis C, Coppee F, Fakhry N, Ayad T, Saussez S. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol* 2020;277(8):2251–61. <https://doi.org/10.1007/s00405-020-05965-1>.
- [65] Cao Y, Li L, Feng Z, Wan S, Huang P, Sun X, Wen F, Huang X, Ning G, Wang W. Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* 2020;6(1). <https://doi.org/10.1038/s41421-020-0147-1>.
- [66] Goodfellow JA, Willison HJ. Guillain-Barré syndrome: a century of progress. *Nat Rev Neurol* 2016;12(12):723–31. <https://doi.org/10.1038/nrneurol.2016.172>.
- [67] Sedaghat Z, Karimi N, Guillain Barre syndrome associated with COVID-19 infection: a case report. *J Clin Neurosci* 2020;76:233–5. <https://doi.org/10.1016/j.jocn.2020.04.062>.
- [68] Zhao H, Shen D, Zhou H, Liu J, Chen S. Guillain-Barré syndrome associated with SARS-CoV-2 infection: causality or coincidence?. *Lancet Neurol* 2020;19(5):383–4. [https://doi.org/10.1016/S1474-4422\(20\)30109-5](https://doi.org/10.1016/S1474-4422(20)30109-5).
- [69] Fernández-Domínguez J, Ameijide-Sanluis E, García-Cabo C, García-Rodríguez R, Mateos V. Miller-Fisher-like syndrome related to SARS-CoV-2 infection (COVID 19). *J Neurol* 2020;267(9):2495–6. <https://doi.org/10.1007/s00415-020-09912-2>.
- [70] Gerges Harb J, Noureldine HA, Chedid G, Eldine MN, Abdallah DA, Chedid NF, Nour-Eldine W. SARS, MERS and COVID-19: clinical manifestations and organ-system complications: a mini review. *Pathog Dis*. 2020 Jun 1;78(4):ftaa033. doi: 10.1093/femspd/ftaa033. PMID: 32633327; PMCID: PMC7454523.
- [71] Hochberg FH, Nelson K, Janzen W. Influenza type B-related encephalopathy. The 1971 outbreak of Reye syndrome in Chicago. *JAMA*. 1975 Feb 24;231(8):817–21. doi: 10.1001/jama.231.8.817. PMID: 1172750.
- [72] Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, Yu J, Kang M, Song Y, Xia J, Guo Q, Song T, He J, Yen H-L, Peiris M, Wu J. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382(12):1177–9.
- [73] Merkler AE, Parikh NS, Mir S, Gupta A, Kamel H, Lin E, Lantos J, Schenck EJ, Goyal P, Bruce SS, Kahan J, Lansdale KN, LeMoss NM, Murthy SB, Stieg PE, Fink ME, Iadecola C, Segal AZ, Cusick M, Campion TR Jr, Diaz I, Zhang C, Navi BB. Risk of Ischemic Stroke in Patients With Coronavirus Disease 2019 (COVID-19) vs Patients With Influenza. *JAMA Neurol*. 2020 Jul 2:e202730. doi: 10.1001/jamaneurol.2020.2730. Epub ahead of print. PMID: 32614385; PMCID: PMC7333175.