



Review Article

Research Progress on Chemical Constituents of *Zingiber officinale* Roscoe

Yan Liu ¹, Jincheng Liu,² and Yongqing Zhang ¹

¹School of Pharmacy, Shandong University of Traditional Chinese Medicine, Jinan 250355, China

²School of Chemical and Chemical Engineering, Shandong University of Technology, Zibo 255049, China

Correspondence should be addressed to Yongqing Zhang; zyq622003@126.com

Received 19 August 2019; Revised 4 November 2019; Accepted 26 November 2019

Academic Editor: Guiming Liu

Copyright © 2019 Yan Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Zingiber officinale Roscoe is commonly used in food and pharmaceutical products but can also be used in cosmetics and daily necessities. In recent years, many scholars have studied the chemical composition of *Zingiber officinale* Roscoe; therefore, it is necessary to comprehensively summarize the chemical composition of *Zingiber officinale* Roscoe in one article. The purpose of this paper is to provide a comprehensive review of the chemical constituents of *Zingiber officinale* Roscoe. The results show that *Zingiber officinale* Roscoe contains 194 types of volatile oils, 85 types of gingerol, and 28 types of diarylheptanoid compounds, which can lay a foundation for further applications of *Zingiber officinale* Roscoe.

1. Introduction

Zingiber officinale Roscoe (ZOR, also Shengjiang in Chinese) is a perennial herb from the Zingiberaceae family, native to the Pacific Islands. It can be found in the Chinese provinces of Shandong, Henan, Hubei, Yunnan, Guangdong, Sichuan, and Jiangsu. ZOR is the fresh root of ginger, which is not only an important condiment but also one of the most commonly used Chinese medicines in clinical practice. Traditional Chinese medicine believes that ZOR has effects of releasing exterior and dissipating cold, arresting vomiting, resolving phlegm, and relieving coughs and can be used to treat fish and crab poison, stomach colds and vomiting, and cold sputum cough [1]. Modern pharmacological studies have shown that ZOR can promote digestion, improve blood circulation, lower blood lipids, lower blood sugar, relieve vestibular stimulation, and provide anti-inflammatory, antitumor, antimicrobial, and antioxidant effects [2–5]. Due to its rich active constituents, ZOR has been used in cosmetics [6], toothpaste [7], and health foods [8–10].

All development and utilization of ZOR are based on its material composition. The chemical composition of ZOR is complex, includes more than 300 types of species, and can be broadly divided into three categories: volatile oils, gingerol, and

diarylheptanoids [11–13]. In this paper, the existing research literature of ZOR is systematically summarized, and each chemical composition and its chemical structure are listed in detail, with a view to providing references for quality control, cultivation production, and further development of ZOR.

2. Constituents

2.1. Volatile Oils. Volatile oils, also known as ginger essential oils, are generally composed of terpenoids [14]. Ginger essential oils give ZOR a unique aromatic smell [11]. The volatile oil composition varies based on where the ZOR is harvested. Currently, the ingredients identified in the volatile oils of ZOR and their chemical structures are shown in Table 1.

2.2. Gingerol. Gingerol is the spicy component of ZOR. It is a mixture of various substances, all of which contain the 3-methoxy-4-hydroxyphenyl functional group. Gingerols can be divided into gingerols, shogaols, paradols, zingerones, gingerdiones, and gingerdiols, according to the different fatty chains connected by this functional group [28, 29]. The structural formulas are given in Table 2.

TABLE 1: Volatile oils in ZOR.















No.	Type	Name	Structure	Reference
1	Terpene	α -Terpinene		[15]
2	Terpene	α -Terpineol		[15]
3	Terpene	4-Terpineol		[15]
4	Terpene	Terpinolene		[15]
5	Terpene	γ -Terpinolene		[15]
6	Alcohol	Cineole		[15]
7	Alcohol	β -Eudesmol		[15]
8	Alcohol	Nerol		[15]
9	Alcohol	<i>trans</i> -Nerolidol		[15]
10	Alcohol	4-Isopropylbenzyl alcohol		[15]
11	Alcohol	3,7-Dimethylocta-1,6-dien-3-ol		[15]
12	Alcohol	3,7-Dimethyloct-6-en-1-yn-3-ol		[15]
13	Alcohol	3-Methylhexan-2-ol		[15]
14	Alcohol	<i>cis</i> -Piperitol		[15]

TABLE 1: Continued.
















No.	Type	Name	Structure	Reference
15	Alcohol	Borneol		[15]
16	Alcohol	Elemol		[15]
17	Alcohol	<i>tau</i> -Muurolol		[15]
18	Alcohol	2-Methoxy-1,7,7-trimethylbicyclo[2.2.1]heptane		[15]
19	Alcohol	1-Isopropyl-4-methylcyclohex-3-enol		[15]
20	Alcohol	2-Tetradecanol		[15]
21	Alcohol	Myrtenol		[15]
22	Alcohol	Citronellol		[15]
23	Alcohol	Geraniol		[15]
24	Alcohol	<i>cis</i> -Linalool oxide		[15]
25	Alcohol	4-Ethoxybutan-1-ol		[15]
26	Alcohol	α -Eudesmol		[15]
27	Alcohol	Nerolidol		[15]
28	Alcohol	Farnesol		[15]
29	Alcohol	<i>trans</i> -4-Isopropyl-1-methyl-2-cyclohexen-1-ol		[15]

TABLE 1: Continued.















No.	Type	Name	Structure	Reference
30	Alcohol	<i>cis</i> -4-Isopropyl-1-methyl-2-cyclohexen-1-ol		[15]
31	Alcohol	2-Heptanol		[16]
32	Alcohol	1-Methoxy-2-methyl		[16]
33	Alcohol	<i>cis</i> -Sesquisabinene hydrate		[17]
34	Alcohol	<i>cis</i> -2- <i>p</i> -Menthen-1-ol		[17]
35	Alcohol	endo-Borneol		[17]
36	Alcohol	<i>trans</i> -Sabinene hydrate		[17]
37	Alcohol	2-Nonanol		[18]
38	Alcohol	Propanol		[18]
39	Alcohol	<i>cis</i> - β -Sesquiphellandrol		[18]
40	Alcohol	<i>trans</i> - β -Sesquiphellandrol		[18]
41	Alcohol	β -Santalol		[19]
42	Alcohol	Zingiberol		[19]
43	Alcohol	<i>tau</i> -Cadinol		[20]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
44	Alcohol	Zingiberenol		[21]
45	Alcohol	2-Pinen-5-ol		[21]
46	Alcohol	Bornyl methyl ether		[21]
47	Alcohol	Isoborneol		[22]
48	Alcohol	2-Decanol		[22]
49	Alcohol	Fenchol		[22]
50	Alcohol	Linalool		[23]
51	Alcohol	Plinol		[23]
52	Alcohol	Camphenol		[23]
53	Alcohol	<i>trans</i> -2-Decen-1-ol		[23]
54	Alcohol	Hentriacontanol		[24]
55	Alcohol	10-O- β -D-Glucopyranosyl-hydroxyl cineole		[25]
56	Aldoketone	Butanal		[15]
57	Aldoketone	Germacrone		[15]
58	Aldoketone	2,6-Dimethylhept-5-enal		[15]
59	Aldoketone	2-Heptanone		[15]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
60	Aldoketone	(<i>E</i>)-Citral		[15]
61	Aldoketone	(<i>Z</i>)-Citral		[15]
62	Aldoketone	2-Nonanone		[15]
63	Aldoketone	3-((3 <i>E</i> ,5 <i>E</i>)-Deca-3,5-dienyl)cyclopentanone		[15]
64	Aldoketone	β -Cyclocitral		[15]
65	Aldoketone	2-Undecanone		[15]
66	Aldoketone	1,7,7-Trimethylbicyclo[2.2.1]heptan-2-one		[15]
67	Aldoketone	(1 <i>R</i>)-(-)-Myrtenal		[15]
68	Aldoketone	β -Citronellal		[15]
69	Aldoketone	Crypton		[15]
70	Aldoketone	4-Isopropylcyclohex-2-enone		[15]
71	Aldoketone	Camphor		[15]
72	Aldoketone	6-Methyl-5-hepten-2-one		[15]
73	Aldoketone	<i>trans,trans</i> -Farnesal		[15]
74	Aldoketone	Hexanal		[16]
75	Aldoketone	Neral		[17]

TABLE 1: Continued.





















No.	Type	Name	Structure	Reference
76	Aldoketone	Geranial		[17]
77	Aldoketone	Octanal		[17]
78	Aldoketone	Methyl heptenone		[18]
79	Aldoketone	Nonyl aldehyde		[18]
80	Aldoketone	Acetaldehyde		[18]
81	Aldoketone	Propionaldehyde		[18]
82	Aldoketone	Valeraldehyde		[18]
83	Aldoketone	Perillal		[19]
84	Aldoketone	(E)-Dodec-2-enal		[21]
85	Aldoketone	(Z)-3,7-Dimethylocta-3,6-dienal		[21]
86	Aldoketone	(E)-3,7-Dimethylocta-3,6-dienal		[21]
87	Aldoketone	(E)-Dec-2-enal		[21]
88	Aldoketone	Decanal		[23]
89	Aldoketone	Citronella		[23]
90	Aldoketone	2-Octenal		[21]
91	Aldoketone	Octanal		[26]
92	Aldoketone	Acetone		[26]
93	Acid	L-Bornyl acetate		[15]
94	Acid	Geranic acid		[15]
95	Acid	Undecanoic acid		[16]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
96	Ester	Neryl acetate		[15]
97	Ester	Methyl 11-(cyclopent-2-enyl)undecanoate		[15]
98	Ester	Geranyl propionate		[15]
99	Ester	endo-Bornyl acetate		[15]
100	Ester	sec-Butyl acetate		[15]
101	Ester	3,7-Dimethyl-2,6-octadienyl acetate		[15]
102	Ester	Neryl propionate		[15]
103	Ester	Geraniol formate		[15]
104	Ester	Myrtenyl acetate		[15]
105	Ester	Geranyl acetate		[15]
106	Ester	Formic acid ethyl ester		[16]
107	Ester	Ethyl butanoate		[17]
108	Ester	Citronellyl acetate		[17]
109	Ester	Heptyl acetate		[17]
110	Ester	Methyl acetate		[18]
111	Ester	Ethyl acetate		[18]

TABLE 1: Continued.
















No.	Type	Name	Structure	Reference
112	Ester	Butyl acetate		[21]
113	Ester	2-Octyl acetate		[21]
114	Fat hydrocarbon	allo-Aromadendrene		[15]
115	Fat hydrocarbon	β -Sesquiphellandrene		[15]
116	Fat hydrocarbon	α -Cedrene		[15]
117	Fat hydrocarbon	β -Thujene		[15]
118	Fat hydrocarbon	Cadina-5,8-diene		[15]
119	Fat hydrocarbon	Bicyclo[2.2.1]heptane		[15]
120	Fat hydrocarbon	(<i>E</i>)-2,7-Dimethyloct-3-en-5-yne		[15]
121	Fat hydrocarbon	(<i>Z</i>)-2,6-Dimethylocta-2,6-diene		[15]
122	Fat hydrocarbon	(<i>E</i>)-3,7-Dimethylocta-1,3,6-triene		[15]
123	Fat hydrocarbon	β -Phellandrene		[15]
124	Fat hydrocarbon	α -Bergamotene		[15]
125	Fat hydrocarbon	α -Gurjunene		[15]
126	Fat hydrocarbon	Sabinene		[15]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
127	Fat hydrocarbon	(+)-Cyclosativene		[15]
128	Fat hydrocarbon	(Z)- β -Farnesene		[15]
129	Fat hydrocarbon	(E)- β -Farnesene		[15]
130	Fat hydrocarbon	(Z,Z)- α -Farnesene		[15]
131	Fat hydrocarbon	Zingiberene		[15]
132	Fat hydrocarbon	α -Farnesene		[15]
133	Fat hydrocarbon	(E)-5-Methylocta-1,6-diene		[15]
134	Fat hydrocarbon	5-Methyloct-3-yne		[15]
135	Fat hydrocarbon	7-Methylocta-3,4-diene		[15]
136	Fat hydrocarbon	γ -Elemene		[15]
137	Fat hydrocarbon	γ -Humulene		[15]
138	Fat hydrocarbon	Thujopsene		[15]
139	Fat hydrocarbon	β -Elemene		[15]
140	Fat hydrocarbon	β -Bisabolene		[15]
141	Fat hydrocarbon	α -Pinene		[15]
142	Fat hydrocarbon	β -Pinene		[15]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
143	Fat hydrocarbon	Caryophyllene		[15]
144	Fat hydrocarbon	β -Caryophyllene		[15]
145	Fat hydrocarbon	Tricyclene		[15]
146	Fat hydrocarbon	Moslene		[15]
147	Fat hydrocarbon	Cedrene		[15]
148	Fat hydrocarbon	(-)-allo-Aromadendrene		[15]
149	Fat hydrocarbon	Neoclovene		[15]
150	Fat hydrocarbon	3-Octyne		[15]
151	Fat hydrocarbon	1-Octene		[15]
152	Fat hydrocarbon	β -Myrcene		[15]
153	Fat hydrocarbon	β -Eudesmene		[15]
154	Fat hydrocarbon	Eudesma-3,7(11)-diene		[15]
155	Fat hydrocarbon	Caryophyllene		[15]
156	Fat hydrocarbon	Bicyclo[3.1.1]heptane		[15]
157	Fat hydrocarbon	1-Cyclopropylpentane		[15]
158	Fat hydrocarbon	3-Carene		[15]

TABLE 1: Continued.

















No.	Type	Name	Structure	Reference
159	Fat hydrocarbon	2-Carene		[15]
160	Fat hydrocarbon	(+)-Aromadendrene		[15]
161	Fat hydrocarbon	Fenchene		[16]
162	Fat hydrocarbon	δ -Elemene		[17]
163	Fat hydrocarbon	D-Limonene		[18, 22]
164	Fat hydrocarbon	β -Phellandrene		[18, 26]
165	Fat hydrocarbon	10-Epizonarene		[18]
166	Fat hydrocarbon	Octane		[18]
167	Fat hydrocarbon	Nonane		[18]
168	Fat hydrocarbon	α -Bergamotene		[19]
169	Fat hydrocarbon	β -Bisabolene		[20]
170	Fat hydrocarbon	τ -Epi- α -selinene		[20]
171	Fat hydrocarbon	4-Carene		[22]
172	Fat hydrocarbon	Camphene		[23]
173	Fat hydrocarbon	α -Phellandrene		[23]
174	Fat hydrocarbon	(Z)-3,7-Dimethylocta-1,3,6-triene		[27]

TABLE 1: Continued.





















No.	Type	Name	Structure	Reference
175	Fat hydrocarbon	Germacrene		[27]
176	Fat hydrocarbon	δ -Cadinene		[26]
177	Fat hydrocarbon	α -Cubebene		[26]
178	Fat hydrocarbon	α -Copaene		[26]
179	Arene	α -Curcumene		[15]
180	Arene	2-Isopropyltoluene		[15]
181	Arene	<i>o</i> -Cymene		[15]
182	Arene	Styrene		[17]
183	Arene	Methylbenzene		[17]
184	Arene	Cumene		[18]
185	Arene	<i>p</i> -Cymene		[19]
186	Others	<i>p</i> -Cymen-8-ol		[15]
187	Others	2-Acetoxy-1,8-cineole		[17]
188	Others	Diethyl sulphide		[18]
189	Others	Ethyl isopropyl sulphide		[18]
190	Others	Methyl allyl sulphide		[18]

TABLE 1: Continued.

No.	Type	Name	Structure	Reference
191	Others	Dibutyl phthalate		[20]
192	Others	2-(3'-Methyl-2'-butenyl)-3-methylfuran		[21]
193	Others	Isoeugenol		[21]
194	Others	2-(2',3'-Epoxy-3'-methylbutyl)-3-methylfuran		[21]

2.3. Diarylheptanoids. Diarylheptanoid is a group of compounds with 1,7-disubstituted phenyl groups and heptane skeletons in its parent structure. Currently, it can be divided into linear diphenyl heptane and cyclic diphenyl heptane compounds with antioxidant activity [53]. The structural formulas are shown in Table 3.

2.4. Others

2.4.1. Proteins and Amino Acids. ZOR contains a variety of amino acids, including glutamate, aspartic acid, serine, glycine, threonine, alanine, cystine, valine, methionine, isoleucine, leucine, tyrosine, phenylalanine, lysine, histidine, arginine, proline [22, 60], and tryptophan [51].

2.4.2. Sugars. ZOR also contains polysaccharides [44], cellulose, and soluble sugar.

2.4.3. Organic Acids. ZOR contains oxalic acid, tartaric acid, lactic acid, acetic acid, citric acid, succinic acid, formic acid, and malonic acid [61].

2.4.4. Inorganic Elements. ZOR has been shown to contain more than 20 inorganic elements such as K, Mg, Ga, Mn, P, Al, Zn, Fe, and Ba [44].

3. Discussion

Various gingers have different regions and chemical compositions. Jolad [30] conducted quantitative analysis on the extracts of dichloromethane from Chinese white ginger and Japanese turmeric and found that the highest content of 6-gingerol was 28% and 34%, respectively. The next highest concentrations were 8-gingerol and 10-gingerol, and the lowest content of 6-shogaol was only 0.35%. Onyenekwe

[62] determined that the main components of the volatile oils of Nigerian ginger were terpenoids such as zingiberene (29.5%) and β -sesquiphellandrene (18.4%), which were quite different from those of ginger grown in other regions. Another study showed the volatile oil content of ginger grown in five different areas of China (Shandong Laiwu, Anhui Tongling, Shandong Anqiu, Guangdong Guangzhou, and Hunan Rucheng) was 0.13%, 0.23%, 0.30%, 0.14%, and 0.17% [63], respectively. 6-Gingerol is often the quality standard for ginger, where the ginger found in Qianwei, Sichuan Province, shows a higher effective content of 6-gingerol than that of the pharmacopoeia standard of the People's Republic of China [64, 65]. The concentrations of 6-gingerol and 6-zingiberol of ginger grown in different regions of China vary greatly, which may be related to the growth environment [66]. Mature and fresh ginger extracts contain the same chemical components, but the difference is in the relative content of each component. Ginger oleoresin in mature ginger is significantly higher than that in fresh ginger. In aromatic terpenoids, the contents of 2-acetoxy-1,8-cineole, β -citronellal, citral, geraniol, geranyl acetate, and zingiberene in mature ginger are lower than those in fresh ginger. The relative content of α -curcumene in mature ginger was higher than that in fresh ginger. In spicy gingerol compounds, the relative content of gingerol in mature ginger is higher than that in fresh ginger, which may be the result of further synthesis and accumulation of gingerol components in the process of continued growth of mature ginger in the second year [48]. The varieties of ginger with the highest oil content are Laiwu ginger, Japanese ginger, Shannong 1 ginger, Shannong 2 ginger, and Anqiu big ginger, with concentrations of 4.56%, 4.42%, 4.52%, 4.50%, and 4.35%, respectively. Average oil contents of 3.45% and 3.16% were found in Jinchang ginger and Chinger, respectively. The lowest oil extraction rates were found in Anqiu small ginger, Fangzhou ginger, and Jinshi ginger, which were 2.95%, 2.60%, and 1.55%, respectively [48].






TABLE 2: Gingerols in ZOR.

No.	Name	Structure	R	R ₁	R ₂	n	Reference
195	3-Gingerol		H	H	—	1	[30]
196	4-Gingerol		H	H	—	2	[30]
197	5-Gingerol		H	H	—	3	[30]
198	6-Gingerol		H	H	—	4	[31, 32]
199	8-Gingerol		H	H	—	6	[31, 33]
200	10-Gingerol		H	H	—	8	[33]
201	12-Gingerol		H	H	—	10	[31, 32]
202	5-Methoxy-6-gingerol		H	CH ₃	—	4	[34]
203	Acetoxy-4-gingerol		H	COCH ₃	—	2	[34]
204	Acetoxy-6-gingerol		H	COCH ₃	—	4	[35]
205	Acetoxy-8-gingerol		H	COCH ₃	—	6	[34]
206	Acetoxy-10-gingerol		H	COCH ₃	—	8	[34]
207	4-Gingeryl methyl ether		CH ₃	H	—	2	[34]
208	6-Gingeryl methyl ether		CH ₃	H	—	4	[34]
209	6-Gingeryl methyl ether acetate		CH ₃	COCH ₃	—	4	[34]
210	6-Gingeryl diacetate		COCH ₃	COCH ₃	—	4	[36]
211	8-Gingeryl diacetate		COCH ₃	COCH ₃	—	6	[36]
212	10-Gingeryl diacetate		COCH ₃	COCH ₃	—	8	[36]
213	Zingerone		H	—	—	0	[36]
214	1-Paradol		H	—	—	1	[20]
215	2-Paradol		H	—	—	2	[20]
216	3-Paradol		H	—	—	3	[20]
217	4-Paradol		H	—	—	4	[20]
218	6-Paradol		H	—	—	6	[36, 37]
219	7-Paradol		H	—	—	7	[20]
220	8-Paradol		H	—	—	8	[36]
221	9-Paradol		H	—	—	9	[20]
222	10-Paradol		H	—	—	10	[36]
223	11-Paradol		H	—	—	11	[34]
224	13-Paradol		H	—	—	13	[34]
225	Methyl-6-paradol		CH ₃	—	—	6	[34, 38]
226	Methyl-8-paradol		CH ₃	—	—	8	[39]
227	Zingerone acetate		COCH ₃	—	—	0	[36]
228	6-Paradyl monoacetate		COCH ₃	—	—	6	[34]
229	8-Paradyl monoacetate		COCH ₃	—	—	8	[36]
230	6-Paradyl benzoate		COPh	—	—	6	[36]
231	1-Dehydro-3-gingerdione		—	—	—	1	[34]
232	1-Dehydro-6-gingerdione		—	—	—	4	[35, 40]
233	1-Dehydro-8-gingerdione		—	—	—	6	[35, 41]
234	1-Dehydro-10-gingerdione		—	—	—	8	[37, 42]
235	12-Dehydrogingerdione		—	—	—	10	[43]
236	6-Gingerdione		—	—	—	4	[44]
237	10-Gingerdione		—	—	—	8	[45]
238	4-Shogaol		H	—	—	2	[30, 44]
239	5-Shogaol		H	—	—	3	[39]
240	6-Shogaol		H	—	—	4	[35, 37]
241	8-Shogaol		H	—	—	6	[36, 37]
242	10-Shogaol		H	—	—	8	[36, 37]
243	12-Shogaol		H	—	—	10	[30, 44]
244	Methyl-4-shogaol		CH ₃	—	—	2	[39]
245	Methyl-6-shogaol		CH ₃	—	—	4	[34]
246	Methyl-8-shogaol		CH ₃	—	—	6	[34]

TABLE 2: Continued.

No.	Name	Structure	R	R ₁	R ₂	n	Reference
247	4-Gingerdiol		H	H	H	2	[30]
248	6-Gingerdiol		H	H	H	4	[44]
249	8-Gingerdiol		H	H	H	6	[44]
250	10-Gingerdiol		H	H	H	8	[44]
251	5-Acetoxy-4-gingerdiol		H	COCH ₃	H	2	[34]
252	5-Acetoxy-6-gingerdiol		H	COCH ₃	H	4	[46]
253	5-Acetoxy-7-gingerdiol		H	COCH ₃	H	5	[34]
254	Diacetoxy-4-gingerdiol		H	COCH ₃	COCH ₃	2	[46, 47]
255	Diacetoxy-6-gingerdiol		H	COCH ₃	COCH ₃	4	[46, 47]
256	Methyl-5-acetoxy-4-gingerdiol		CH ₃	COCH ₃	H	2	[34]
257	Methyl-5-acetoxy-6-gingerdiol		CH ₃	COCH ₃	H	4	[34]
258	Methyl diacetoxy-4-gingerdiol		CH ₃	COCH ₃	COCH ₃	2	[34]
259	Methyl diacetoxy-6-gingerdiol		CH ₃	COCH ₃	COCH ₃	4	[34]
260	Methyl diacetoxy-10-gingerdiol		CH ₃	COCH ₃	COCH ₃	8	[34]
261	6-Dihydroparadol		H	—	—	6	[34]
262	Acetoxy-6-dihydroparadol		Ac	—	—	6	[34]
263	1-(4'-Hydroxy-3'-methoxyphenyl)-7-octen-3-one		H	—	—	—	[34]
264	1-(4'-Hydroxy-3'-methoxyphenyl)-7-decen-3-one		CH ₂ CH ₃	—	—	—	[34]
265	1-(4'-Hydroxy-3'-methoxyphenyl)-7-dodecen-3-one		(CH ₂) ₃ CH ₃	—	—	—	[34]
266	4-Isogingerol		H	—	—	2	[34]
267	6-Isogingerol		H	—	—	4	[48]
268	Methyl-6-isogingerol		CH ₃	—	—	4	[34]
269	6-Zingerine		—	—	—	4	[49]
270	8-Zingerine		—	—	—	6	[49]
271	10-Zingerine		—	—	—	8	[49]
272	3-Dihydro-6-demethoxy shogaol		—	—	—	—	[34]
273	6-Isoshogaol		—	—	—	—	[44]
274	Dehydrozingerone		—	—	—	—	[36]

TABLE 2: Continued.

No.	Name	Structure	R	R ₁	R ₂	n	Reference
275	1-Dehydro-3-dihydro-10-gingerdione		—	—	—	—	[34, 44]
276	(Z)-10-Isoshogaol		—	—	—	—	[48]
277	(E)-10-Isoshogaol		—	—	—	—	[48]
278	β -Sitosterol		—	—	—	—	[50, 51]
279	Tetracosanoic acid		—	—	—	—	[50, 52]

Ginger, as a kind of food and medicine, has many functions, such as antioxidant, anti-inflammatory, antimicrobial, anticancer, antiobesity, antidiabetic, anti-nausea, antiemetic, antiallergic, neuroprotective, hepatoprotective, cardiovascular protective, and respiratory protective activities [67]. Currently, most studies of the bioactive components of ginger focus on ginger volatile oil, gingerol, shogaol, and zingerone compounds. Ginger essential oil can effectively improve the antioxidant capacity of the liver, reduce inflammatory response, and protect against fatty liver [68]. The antioxidant compounds in ginger are primarily gingerol and diarylheptanoid. Substituents on alkyl chains contribute to free radical scavenging and oxidation inhibition of lipids [69]. Antioxidant activity is typically derived from gingerols, shogaols, and some related phenolic ketone derivatives [70]. Gingerols are spicy ingredients in which 6-gingerol shows the highest biological activity, so 6-gingerol is often used as an indicator of ginger quality [71]. 6-Gingerol has been used to inhibit angiogenesis in vivo and in vitro [72]. It has been shown to have anticancer and antagastric ulcer properties while suppressing central nervous stimulation and various pharmacological activities [73, 74]. 6-Gingerol has been used to treat tumors by regulating the apoptosis gene by reversing the abnormal expression of tumor cell genes. It can also affect the apoptosis signal transduction pathway and induce apoptosis [75]. 8-Gingerol and 10-gingerol have good inhibitory effects on the activity of various tumor cells, where the inhibitory effects are somewhat different. The two may affect the phosphorylation level of the MAPK pathway proteins ERK and P38, leading to G1 phase arrest of breast

cancer cells, thus applying inhibitory effects on the proliferation of tumor cells [76]. The main components of strong heart are gingerol and 6-shogaol [77]. The effects of 6-gingerol and 6-shogaol on blood pressure have been shown to induce a hypotensive effect at low doses, while high doses have shown a three-phase reaction. Initially, blood pressure drops rapidly, then rises, and then provides a hypotensive effect at later stages [78]. Ginger polysaccharide has biological activities such as antitumor, hypoglycemic, lipid-lowering, immune regulation, antiviral, and antifungal [79].




4. Conclusion

ZOR is a widely used drug and food in clinical and daily life and has been used in the prevention and treatment of the digestive, circulatory, respiratory, and central nervous system diseases and other diseases. In this paper, the chemical constituents found in ZOR in recent years are summarized, and the results show that more than 300 chemical constituents are identified from the extracts of ZOR, including 194 types of volatile oil, 85 types of gingerol, and 28 types of diarylheptanoids compounds. From this, it can be clearly observed that ZOR has a complex chemical composition. The interactions between the components provide the clinical effects; therefore, it is necessary to further study the chemical composition and pharmacological action of ginger, for further applications. Exploring the mechanism by which different components perform the same effects is a new way to develop drugs in the future; for example, 4-terpineol and beta-sitosterol can act on the two targets of the 5-

TABLE 3: Diarylheptanoids in ZOR.

No.	Name	Structure	R ₁	R ₂	R ₃	Reference
280	5-Hydroxy-1-(4'-hydroxy-3'-methoxyphenyl)-7-(4''-hydroxyphenyl)heptan-3-one		H	H	H	[32]
281	5-Hydroxy-1,7-bis(4'-hydroxy-3'-methoxyphenyl)heptan-3-one		H	OCH ₃	H	[32]
282	7-(3',4'-Dihydroxy-5''-methoxyphenyl)-5-hydroxy-1-(4''-hydroxy-3''-methoxyphenyl)heptan-3-one		H	OCH ₃	OH	[54]
283	5-Hydroxy-7-(4'-hydroxy-3',5'-dimethoxyphenyl)-1-(4''-hydroxy-3''-methoxyphenyl)heptan-3-one		H	OCH ₃	OCH ₃	[32]
284	5-Hydroxy-1-(4'-hydroxy-3',5'-dimethoxyphenyl)-7-(4''-hydroxy-3''-methoxyphenyl)heptan-3-one		OCH ₃	OCH ₃	H	[55]
285	5-Hydroxy-1,7-bis(4'-hydroxy-3',5'-dimethoxyphenyl)heptan-3-one		OCH ₃	OCH ₃	OCH ₃	[32]
286	(E)-7-(3',4'-Dihydroxyphenyl)-1-(4''-hydroxy-3''-methoxyphenyl)hept-4-en-3-one		H	—	V	[55]
287	1,7-bis(4'-Hydroxy-3'-methoxyphenyl)-4-heptene-3-one		CH ₃	—	—	[55, 56]
288	3,5-Dihydroxy-1,7-bis(4'-hydroxy-3'-methoxyphenyl)heptane		H	H	CH ₃	[54–56]
289	1,7-bis(3',4'-Dihydroxyphenyl)-3,5-diacetate heptane		COCH ₃	H	H	[55]
290	1,7-bis(4'-Hydroxy-3'-methoxyphenyl)-3,5-diacetate heptane		COCH ₃	H	CH ₃	[55–57]
291	1,7-bis(4'-Methoxy-3'-acetatephenyl)-3,5-diacetate heptane		COCH ₃	CH ₃	COCH ₃	[55]
292	1,7-bis(3',4'-Diacetatephenyl)-3,5-diacetate heptane		COCH ₃	COCH ₃	COCH ₃	[55]
293	5-(6-(4-Hydroxy-3-methoxyphenethyl)-4-hydroxy-tetrahydro-2H-pyran-2-yl)-3-methoxybenzene-1,2-diol		H	—	—	[54, 58]
294	2-(4'-Hydroxy-3'-methoxyphenethyl)-6-(3'',4''-dihydroxy-5''-methoxyphenyl)-tetrahydro-2H-pyran-4-yl acetate		COCH ₃	—	—	[58]
295	7-(3',4'-Dihydroxyphenyl)-1-(4''-hydroxy-3''-methoxyphenyl)-3,5-diacetate heptane		H	H	—	[55]
296	7-(4'-Hydroxy-3'-methoxyphenyl)-1-(4'',5''-dihydroxy-3''-methoxyphenyl)-3,5-diacetate heptane		OH	CH ₃	—	[54]
297	7-(4'-Hydroxy-3'-methoxyphenyl)-1-(4''-hydroxy-5''-methyl-3''-methoxyphenyl)-3,5-diacetate heptane		CH ₃	CH ₃	—	[55]
298	7-(4'-Hydroxy-3'-methoxyphenyl)-1-(4''-hydroxy-3'',5''-dimethoxyphenyl)-3,5-diacetate heptane		OCH ₃	CH ₃	—	[55, 56]
299	5-(6-(4-Hydroxyphenethyl)-4-hydroxy-tetrahydro-2H-pyran-2-yl)-3-methoxybenzene-1,2-diol		H	H	H	[25]
300	5-(6-(4-Hydroxy-3-methoxyphenethyl)-4-hydroxy-tetrahydro-2H-pyran-2-yl)-3-methoxybenzene-1,2-diol		H	H	OCH ₃	[54, 58]
301	5-(6-(4-Hydroxy-3-methoxyphenethyl)-4-hydroxy-tetrahydro-2H-pyran-2-yl)-2-hydroxy-3-methoxyphenyl acetate		H	COCH ₃	OCH ₃	[58]
302	2-(4'-Hydroxy-3'-methoxyphenethyl)-6-(3'',4''-dihydroxy-5''-methoxyphenyl)-tetrahydro-2H-pyran-4-yl acetate		COCH ₃	H	OCH ₃	[58]
303	1,7-bis(4'-Hydroxy-3'-methoxyphenyl)-5-oxoheptan-3-yl acetate		H	—	—	[57]
304	1,7-bis(3'-Methoxy-4'-acetatephenyl)-5-oxoheptan-3-yl acetate		COCH ₃	—	—	[57]

TABLE 3: Continued.

No.	Name	Structure	R ₁	R ₂	R ₃	Reference
305	1,7-bis(4'-Hydroxy-3'-methoxyphenyl)-3,5-heptadione		—	—	—	[34]
306	(1E,6E)-1,7-bis(4-Hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione		—	—	—	[56]
307	2,4-bis(3,4-Dihydroxyphenethyl)pentanedioic acid		—	—	—	[59]

hydroxytryptamine receptor 3A and the mu-type opioid receptor, respectively, and provide corresponding therapeutic effects on diarrhea and dysentery. This can provide ideas for the research and development of new drugs and lay a foundation for further applications of ZOR.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was financially supported by the Key Research and Development Technology of Shandong Province (Industry Key Technology) (2016CYJS08A01).

References

- [1] National Pharmacopoeia Commission, *Pharmacopoeia of the People's Republic of China: 2015 Edition*, China Pharmaceutical Science and Technology Press, Beijing, China, 2015.
- [2] X. F. Wang, G. T. Wu, T. H. Niu et al., "Chemistry, pharmacology and application of ginger," *Chinese Fruit and Vegetable*, vol. 36, no. 6, pp. 23–29, 2016.
- [3] W. Y. Hu, R. P. Zhang, L. P. Tang et al., "Advances in chemical and pharmacological research on ginger," *Chinese Folk Medicine*, vol. 9, pp. 10–14, 2008.
- [4] M. M. Zhu, D. Liu, K. X. Lu et al., "Research progress on pharmacological action of ginger horseradish," *Journal of Hubei Institute of Medicine*, vol. 37, no. 4, pp. 390–394, 2018.
- [5] Y. J. Sun, "Research progress on pharmacological action of Ginger," *Modern Journal of Integrated Chinese and Western Medicine*, vol. 16, no. 4, pp. 561–564, 2007.
- [6] Y. Chen, Y. Y. Ni, and T. Y. Cai, "Study on comprehensive utilization and deep processing of ginger extract," *Food Industry Technology*, vol. 21, no. 4, pp. 76–78, 2000.
- [7] J. P. Li, "Application of ginger extract in toothpaste," *Toothpaste Industry*, no. 2, pp. 10–11, 2008.
- [8] Z. Y. Peng, W. Huang, Y. F. Guan et al., "Development of ginger health jelly," *Food and Fermentation Technology*, vol. 45, no. 4, pp. 74–76, 2009.
- [9] W. M. Dong, Q. Y. Tang, and W. Yuan, "Study on processing technology of ginger brown sugar Beverage," *Modern Food Technology*, vol. 23, no. 2, pp. 57–58, 2007.
- [10] B. Zhan and F. Z. Wang, "Processing technology of Ginger Beverage series products," *Agro-Processing*, vol. 1, p. 77, 2013.
- [11] H. Wohlmuth, M. K. Smith, L. O. Brooks, S. P. Myers, and D. N. Leach, "Essential oil composition of diploid and tetraploid clones of ginger (*Zingiber officinale Roscoe*) grown in Australia," *Journal of Agricultural and Food Chemistry*, vol. 54, no. 4, pp. 1414–1419, 2006.
- [12] W. S. He, Y. X. Yan, and B. J. Guo, "General situation of study on chemical constituents and bioactivity of ginger," *Chinese Herbal Medicine*, vol. 24, no. 5, pp. 376–379, 2001.
- [13] S. D. Jolad, R. C. Lantz, G. J. Chen, R. B. Bates, and B. N. Timmermann, "Commercially processed dry ginger (*Zingiber officinale*): composition and effects on LPS-stimulated PGE₂ production," *Phytochemistry*, vol. 66, no. 13, pp. 1614–1635, 2005.
- [14] E. Langner, S. Greifenberg, and J. Gruenwald, "Ginger: history and use," *Advances in Therapy*, vol. 15, pp. 25–44, 1998.
- [15] J. N. Tan, "GC-MS Study on chemical constituents of volatile oils from ginger in different producing areas," *Asia-Pacific Traditional Medicine*, vol. 7, no. 4, pp. 23–25, 2011.
- [16] H. Zhang, J. C. Yan, H. G. Zhanget et al., "Extraction of volatile oil from ginger and analysis of its chemical constituents by gas chromatography-mass spectrometry," *Analytical Chemistry*, vol. 24, no. 3, pp. 348–352, 1996.
- [17] X. Pang, J. Cao, D. Wang, J. Qiu, and F. Kong, "Identification of ginger (*Zingiber officinale Roscoe*) volatiles and localization of aroma-active constituents by GC-olfactometry," *Journal of Agricultural and Food Chemistry*, vol. 65, no. 20, pp. 4140–4145, 2017.
- [18] V. S. Govindarajan and D. W. Connell, "Ginger—chemistry, technology, and quality evaluation: Part 1," *CRC Critical Reviews in Food Science and Nutrition*, vol. 17, no. 1, pp. 1–96, 1983.
- [19] Z. K. Lin and Y. F. Hua, "Study on chemical constituents of ginger essential oil in Sichuan," *Organic Chemistry*, vol. 6, pp. 444–448, 1987.
- [20] J. P. Li, Y. S. Wang, H. Ma et al., "Comparative study on the main chemical components of dried ginger and ginger," *Chinese Journal of Chinese Medicine*, vol. 26, no. 11, pp. 748–751, 2001.

- [21] O. Nishimura, "Identification of the characteristic odorants in fresh Rhizomes of ginger (*Zingiber officinale Roscoe*) using aroma extract dilution analysis and modified multidimensional gas chromatography-mass spectroscopy," *Journal of Agricultural and Food Chemistry*, vol. 43, no. 11, pp. 2941–2945, 1995.
- [22] T. Jia, R. L. Xiao, and H. Jiao, "Analysis of volatile components in rhizome zingibers, *Zingiber officinale Roscoe* and ginger pee by GC-MS and chemometric resolution," *JCMRD*, vol. 1, no. 2, pp. 47–53, 2012.
- [23] J. F. Wu, C. Zhang, X. M. Zhang et al., "Extraction and composition analysis of ginger flavor substances," *Deep Processing of Fruits and Vegetables*, vol. 22, no. 3, pp. 94–99, 2006.
- [24] W. D. Xuan, J. Bian, B. Yuan et al., "Study on chemical constituents of ginger," *Chinese Herbal Medicine*, vol. 39, no. 11, pp. 1616–1619, 2008.
- [25] Y. Zhao, Q. F. Tao, R. P. Zhang et al., "Two new compounds from *Zingiber officinale*," *Chinese Chemical Letters*, vol. 18, no. 10, pp. 1247–1249, 2007.
- [26] M. K. Kim, M. S. Na, J. Hong, and S. T. Jung, "Volatile flavor components of Korean ginger (*Zingiber officinale Roscoe*) extracted with liquid carbon," *Hanguk Nonghwa Hakhorchi*, vol. 35, no. 1, pp. 59–63, 1992.
- [27] Q. X. Cui and Y. Dong, "GC-MS analysis of chemical constituents of ginger volatile oil," *Journal of Liaocheng University (Natural Science Edition)*, vol. 19, no. 2, pp. 43–45, 2006.
- [28] C. J. Lu, M. Ou, and N. S. Wang, "Summary of chemical composition analysis of ginger," *New Drugs and Clinical Pharmacology of Traditional Chinese Medicine*, vol. 14, no. 3, pp. 215–217, 2003.
- [29] H. L. Luo, "Extraction, separation and component study of the efficacy components of ginger," M.S. dissertation, Anhui Agricultural University, Anhui, China, 2010.
- [30] H. Xiong, "Comparative study of components in ginger extracts by different extraction methods," M.S. dissertation, Xihua University, Sichuan, China, 2006.
- [31] G. Vernin and C. Parkanyl, "Chemistry of ginger," in *Ginger: The Genus Zingiber*, P. N. Ravindran and K. Nirmal Babu, Eds., CRC Press, New York, NY, USA, pp. 87–180, 2005.
- [32] X. S. Huang, R. A. Yan, and J. Z. Wu, "A review of the bioactivity of ginger phenol," *Journal of Jinan University*, vol. 26, no. 3, pp. 434–435, 2005.
- [33] E. Katsuya, K. Emi, and O. Yoshiteru, "Structures of antifungal diarylheptenones, gingerenones A, B, C and isogingerenone B, isolated from the rhizomes of *Zingiber officinale*," *Phytochemistry*, vol. 29, no. 3, pp. 797–799, 1990.
- [34] S. D. Jolad, R. C. Lantz, A. M. Solyom, G. J. Chen, R. B. Bates, and B. N. Timmermann, "Fresh organically grown ginger (*Zingiber officinale*): composition and effects on LPS-induced PGE₂ production," *Phytochemistry*, vol. 65, no. 13, pp. 1937–1954, 2004.
- [35] L. Wang, L. Fang, H. Q. Zhao et al., "High performance liquid chromatography-electrospray four-stage mass spectrometry analysis of ginger horseradish compounds in ginger," *Chinese Journal of Traditional Chinese Medicine*, vol. 36, no. 24, pp. 3467–3471, 2011.
- [36] D. W. Connell and R. McLachlan, "Natural pungent compounds: IV. Examination of the gingerols, shogaols, paradols and related compounds by thin-layer and gas chromatography," *Journal of Chromatography A*, vol. 67, no. 1, pp. 29–35, 1972.
- [37] J. Zhang, Y. S. Chang, C. Zeng et al., "Study on chemical constituents in ginger extract," *Anhui Agricultural Science*, vol. 43, no. 25, pp. 287–290, 2015.
- [38] A. M. Galal, "Antimicrobial activity of 6-paradol and related compounds," *International Journal of Pharmacognosy*, vol. 34, no. 1, pp. 64–69, 1996.
- [39] S. B. Tan, "Study on water-soluble chemical constituents of ginger and shear," M.S. dissertation, Lanzhou University of Technology, Gansu, China, 2017.
- [40] R. Charles, S. N. Garg, and S. Kumar, "New gingerdione from the rhizomes of *Zingiber officinale*," *Fitoterapia*, vol. 71, no. 6, pp. 716–718, 2000.
- [41] K. Masanori, U. U. Kaoru, H. Tatsuo et al., "Cell differentiation inducers and antiandrogenic active compounds from *Zingiberis* rhizome," *Natural Medicines*, vol. 56, no. 2, pp. 47–50, 2002.
- [42] X.-G. He, M. W. Bernart, L.-Z. Lian, and L.-Z. Lin, "High-performance liquid chromatography-electrospray mass spectrometric analysis of pungent constituents of ginger," *Journal of Chromatography A*, vol. 796, no. 2, pp. 327–334, 1998.
- [43] Y. A. Han, C. W. Song, W. S. Koh et al., "Anti-inflammatory effects of the *Zingiber officinale Roscoe* constituent 12-dehydrogingerdione in lipopolysaccharide-stimulated raw 264.7 cells," *Phytotherapy Research*, vol. 27, no. 8, pp. 1200–1205, 2013.
- [44] W. Z. Zhao, R. X. Zhang, Z. P. Yu et al., "Research progress on chemical constituents and bioactivity of Ginger," *Food Industry Technology*, vol. 37, no. 11, pp. 383–387, 2016.
- [45] E. Koh, H. Kim, S. Kim et al., "Modulation of macrophage functions by compounds isolated from *Zingiber officinale*," *Planta Medica*, vol. 75, no. 02, pp. 148–151, 2009.
- [46] H. Kikuzaki, S.-M. Tsai, and N. Nakatani, "Gingerdiol related compounds from the rhizomes of *Zingiber officinale*," *Phytochemistry*, vol. 31, no. 5, pp. 1783–1786, 1992.
- [47] Y. Masada, T. Inoue, K. Hashimoto, M. Fujioka, and C. Uchino, "Studies on the constituents of ginger (*Zingiber officinale Roscoe*) by GC-MS," *Yakugaku Zasshi*, vol. 94, no. 6, pp. 735–738, 1974.
- [48] K. Y. Zhan, "Chemical composition analysis of ginger oil and study on isolation and purification of ginger horseradish," M.S. dissertation, Shandong Agricultural University, Shandong, China, 2009.
- [49] J. Juan, H. P. Zhang, T. E. Prisinzano, L. A. Mitscher, and B. N. Timmermann, "Identification of unprecedented purine-containing compounds, the zingerines, from ginger rhizomes (*Zingiber officinale Roscoe*) using a phase-trafficking approach," *Phytochemistry*, vol. 72, no. 9, pp. 935–941, 2011.
- [50] J. W. Purselglove, E. G. Brown, C. L. Green, and S. R. J. Robbins, "Spices," M.S. dissertation, Longman, New York, NY, USA, 1981.
- [51] J. Li, "Processing principle of ginger," M.S. dissertation, Henan College of Traditional Chinese Medicine, Henan, China, 2007.
- [52] D. Liu, C. H. Zhang, R. H. An et al., "Research progress on extraction and application of main bioactive components of ginger," *Food Industry Science and Technology*, vol. 37, no. 20, pp. 391–393, 2016.
- [53] X. Wang, "Research progress on active sites and components of ginger," *Research on Traditional Chinese Medicine*, vol. 22, no. 12, pp. 53–55, 2009.
- [54] L. X. Yang, C. X. Zhou, K. X. Huang et al., "Study on anti-oxidant and cytotoxic activity of diphenyl heptane compounds in Ginger," *Chinese Journal of Traditional Chinese Medicine*, vol. 34, no. 3, pp. 319–322, 2009.

- [55] H. Kikuzaki, M. Kobayashi, and N. Nakatani, "Diarylheptanoids from rhizomes of *Zingiber officinale*," *Phytochemistry*, vol. 30, no. 11, pp. 3647–3651, 1991.
- [56] H. Kikuzaki and N. Nakatani, "Antioxidant effects of some ginger constituents," *Journal of Food Science*, vol. 58, no. 6, pp. 1407–1410, 1993.
- [57] J. P. Ma, X. L. Jin, L. Yang et al., "Two new diarylheptanoids from the rhizomes of *zingiber officinale*," *Chinese Chemical Letters*, vol. 15, no. 11, pp. 1306–1308, 2004.
- [58] H. Kikuzaki and N. Nakatani, "Cyclic diarylheptanoids from rhizomes of *Zingiber officinale*," *Phytochemistry*, vol. 43, no. 1, pp. 273–277, 1996.
- [59] J. C. Luo, Y. F. Feng, and X. Ji, "Research progress of natural linear diphenyl heptane compounds," *Chinese Herbal Medicine*, vol. 39, no. 12, pp. 1913–1916, 2012.
- [60] Y. F. Zhang and Z. C. Ma, "Ingredients and applications of ginger," *Chemistry Teaching*, vol. 8, pp. 73–80, 2012.
- [61] S. H. Li, F. Z. Chen, Z. Liu et al., "Determination of Vc, nitrate, nitrite, total sugar, organic acid content in greenhouse ginger," *Anhui Agricultural Science*, vol. 34, no. 14, pp. 3346–3347, 2006.
- [62] P. C. Onyenekwe and S. Hashimoto, "The composition of the essential oil of dried Nigerian ginger (*Zingiber officinale Roscoe*)," *European Food Research and Technology*, vol. 209, no. 6, pp. 407–410, 1999.
- [63] W. M. Zhang, H. F. Jiang, and J. Zhang, "Study on gas chromatographic fingerprints of different aromatic parts of ginger," *China Wild Plant Resources*, vol. 22, no. 5, pp. 53–55, 2003.
- [64] National Pharmacopoeia Commission, *Pharmacopoeia of the People's Republic of China (Part I)*, China Medical Science and Technology Press, Beijing, China, 2015.
- [65] X. F. Liao, Y. Y. Ying, W. Y. Guo et al., "Analysis of the characteristics of qianwei ginger and determination of its main components," *Chinese Journal of Traditional Chinese Medicine*, vol. 34, no. 6, pp. 2680–2683, 2019.
- [66] W. D. Xuan, J. Bian, C. W. Wang et al., "Comparison of extraction methods and quality control of 6-gingerol in ginger," *Chinese Journal of Pharmacy of the People's Liberation Army*, vol. 24, no. 4, pp. 329–331, 2008.
- [67] Q.-Q. Mao, X.-Y. Xu, S.-Y. Cao et al., "Bioactive compounds and bioactivities of ginger (*Zingiber officinale Roscoe*)," *Foods*, vol. 8, no. 6, p. 185, 2019.
- [68] Y.-S. Lai, W.-C. Lee, Y.-E. Lin et al., "Ginger essential oil ameliorates hepatic injury and lipid accumulation in high fat diet-induced nonalcoholic fatty liver disease," *Journal of Agricultural and Food Chemistry*, vol. 64, no. 10, pp. 2062–2071, 2016.
- [69] Y. Masuda, H. Ikuzaki, M. Hisamoto, and N. Nakatani, "Antioxidant properties of gingerol related compounds from ginger," *Biofactors*, vol. 21, no. 1–4, pp. 293–296, 2004.
- [70] N. Liu, G. C. Huo, and L. Zhang, "Effects of ginger on lipid peroxidation in hyperlipidemia rats," *Health Research*, vol. 32, no. 1, pp. 22–23, 2003.
- [71] X. H. Zhang and H. X. Liu, "Research progress of gingerol," *Journal of Guangxi Normal University*, vol. 26, no. 1, pp. 110–113, 2009.
- [72] E.-C. Kim, J.-K. Min, T.-Y. Kim et al., "[6]-Gingerol, a pungent ingredient of ginger, inhibits angiogenesis in vitro and in vivo," *Biochemical and Biophysical Research Communications*, vol. 335, no. 2, pp. 300–308, 2005.
- [73] R. C. Lantz, G. J. Chen, M. Sarihan, A. M. Solyom, S. D. Jolad, and B. N. Timmermann, "The effect of extracts from ginger rhizome on vegetation," *Phytomedicine*, vol. 14, no. 2-3, pp. 123–128, 2007.
- [74] H. M. Qian, M. Wang, and Z. X. Su, "Preliminary experimental study on anti-tumor effects of ginger extract," *Jiangsu Pharmaceutical & Clinical Research*, vol. 7, no. 3, pp. 14–16, 1999.
- [75] J. Xu, "Research progress on anti-tumor effects of ginger," *Journal of Youjiang Medical College for Nationalities*, vol. 37, no. 3, pp. 496–497, 2015.
- [76] X. Liu, H. W. Zhang, R. Q. Fu et al., "Anti-tumor effects and mechanism of active components of gingerol in ginger," *Journal of Third Military Medical University*, vol. 39, no. 9, pp. 884–890, 2017.
- [77] C. J. Lu, M. Ou, and N. S. Wang, "Pharmacological effects of ginger on cardiovascular and cerebrovascular system," *Chinese Medicine New Drug and Clinical Pharmacology*, vol. 14, no. 5, pp. 356–357, 2003.
- [78] M. Suekawa, A. Ishige, K. Yuasa, K. Sudo, M. Aburada, and E. Hosoya, "Pharmacological studies on Ginger. I. Pharmacological actions of pungent constituents, (6)-gingerol and (6)-shogaol," *Journal of Pharmacobio-Dynamics*, vol. 7, no. 11, pp. 836–848, 1984.
- [79] J. A. Wu, X. Y. Wang, K. L. Wang et al., "Research progress on nutritional value and pharmacological action of ginger," *Food Industry*, vol. 40, no. 2, pp. 237–239, 2019.