

POSTER PRESENTATION

Open Access

# Unifocal and multifocal chronic non-bacterial osteomyelitis (CNO) in children

G Zanon<sup>1\*</sup>, A Insalaco<sup>2</sup>, G Martini<sup>1</sup>, S Corradin<sup>3</sup>, P Zucchetta<sup>4</sup>, L Tanturri<sup>2</sup>, M Garganese<sup>2</sup>, F De Benedetti<sup>2</sup>, F Zulian<sup>1</sup>

From 18th Pediatric Rheumatology European Society (PReS) Congress  
Bruges, Belgium. 14-18 September 2011

## Background

Chronic non bacterial osteomyelitis (CNO) is a rare condition in children and little is known about its clinical assessment and course.

## Aim

To describe the clinical characteristics and long-term outcome of pediatric patients with CNO.

## Methods

We retrospectively evaluated patients with sterile bone inflammation, lasting longer than 6 month, referred to two tertiary care pediatric rheumatology units. Information on family history, clinical features at disease onset and course, laboratory and outcome were collected. A comparison between unifocal (UF-CNO) and multifocal course (MF-CNO), based on bone scintiscan result, was performed.

## Results

29 CNO patients entered the study. Ten had UF-CNO, 19 had MF-CNO, mean age at disease onset 9,2 years (range 0,8-17), males (52%). Disease duration at diagnosis was longer in pts with UF-CNO (11,1 vs 4,6 months). Localized bone pain was the leading symptom at onset in all patients; systemic symptoms, such as fever and fatigue, were more frequent in MF-CNO. 30% presented associated skin disease and positive family history for autoimmune disease. At onset WBC normal, CRP was elevated in 38%, ESR in 72%, especially in MF-CNO. Scintiscan allowed us to identify multiple lesions in five patients with one-site symptoms. MF-CNO involve more frequently the lower limbs than the UF-CNO (18/19 vs 4/10, p 0,001). After three years follow up, 70% of

patients had no symptoms and 90% were off-therapy. UF-CNO developed complications, such as hyperostosis, vertebral collapse or limb dysmetry, significantly more often than MF-CNO (p 0,002).

## Conclusion

UF-CNO and MF-CNO belong to the same spectrum but present different clinical features and outcome.

### Author details

<sup>1</sup>Dept. of Pediatrics, University of Padua, Italy. <sup>2</sup>IRCCS Ospedale Pediatrico Bambino Gesù, Rome, Italy. <sup>3</sup>Dept. of Radiology, University of Padua, Italy.

<sup>4</sup>Dept. of Nuclear Medicine, University of Padua, Italy.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P32

Cite this article as: Zanon et al.: Unifocal and multifocal chronic non-bacterial osteomyelitis (CNO) in children. *Pediatric Rheumatology* 2011 **9** (Suppl 1):P32.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: giulia\_zeta@libero.it

<sup>1</sup>Dept. of Pediatrics, University of Padua, Italy

Full list of author information is available at the end of the article