

POSTER PRESENTATION

Open Access

Saccharomyces boulardii ingestion increases the humoral response of a DNA vaccine against leptospirosis in mice

Marcelle Silveira¹, Rafael Rosa^{1*}, Marcelo Mendonça¹, Daiane Hartwig², Alexandre Bilhalva¹, Gustavo Moreira¹, Patrícia Diaz³, Luisa Scapin¹, Fabrício Conceição¹, Ângela Moreira⁴

From 5th Congress of the Brazilian Biotechnology Society (SBBIOTEC)
Florianópolis, Brazil. 10-14 November 2013

Background

DNA vaccines are a good option to generate a desired antigen using the cellular machinery of the vaccinated subject. This kind of vaccine can induce both of the humoral and cellular immune response, and show high stability and easy working, as well as offer a low-cost production. and safety for immune compromised patients [1]. However, some disadvantages, such as the low transfection rate and low immunogenicity, makes necessary the use of multiple doses [2,3]. Therefore, several studies have been conducted trying to increase the immune response generated by the DNA vaccines. *Saccharomyces boulardii*, a probiotic yeast that is capable to increase the host immune response [4], was not previously evaluated as an adjuvant for DNA vaccines. Thereby, in the present study we evaluated the *S. boulardii* capacity to increase the humoral response of mice using DNA vaccines against a fragment of the leptospiral antigen LigA cloned in the mammalian expression plasmid pTARGET [4,5].

Material and methods

Four to six month-old Female BALB/c mice were separated into 2 groups of 12 animals each. Group 1 (G1, control) were fed with antimicrobial-free ration; group 2 (G2) with the same ration containing 10^8 cfu.g⁻¹ of *S. boulardii*, 14 days before the first immunization, for adaptation, and during all the experiment. The immunization protocol consisted in 3 doses (days 0, 14 and 21) of 100 µg of the pTARGET/ligAni plasmid injected intramuscularly. Blood samples were collected from the retro eye venous plexus of the animals on days 0, 13, 20 and 27

for the humoral immune response evaluation of each immunization through indirect ELISA, using recombinant LigAni (5 µg.mL⁻¹) as antigen and serum dilutions 1:30 in PBS-T. ELISA units were calculated by dividing the mean absorbance of each group by the mean absorbance of the day 0. The student's test-t was performed to determine significant differences ($p < 0.05$) between the groups.

Results and conclusions

Sera from G2 has shown ELISA units statistically higher than the sera from G1 after the second ($p < 0.0001$) and third immunization ($p = 0.0264$). In addition, there was no significant difference between the second and third immunizations of G2. This way, the results indicate that using *S. boulardii* as adjuvant, only two immunizations are sufficient to obtain an adequate immune response. Therefore, the *S. boulardii* probiotic was able to enhance the humoral immune response to a DNA vaccine against leptospirosis in mice.

Acknowledgements

We would like to thank CAPES, CNPq and FAPERGS for the financial support and scholarships.

Authors' details

¹Centro de Desenvolvimento Tecnológico, Núcleo de Biotecnologia, Universidade Federal de Pelotas, Pelotas, RS, Brazil. ²Instituto de Biologia, Departamento de microbiologia e parasitologia, Universidade Federal de Pelotas, Pelotas, RS, Brazil. ³Engenharia de materiais, CDTec, Universidade Federal de Pelotas, Pelotas, RS, Brazil. ⁴Faculdade de Nutrição, Universidade Federal de Pelotas, Pelotas, RS, Brazil.

¹Centro de Desenvolvimento Tecnológico, Núcleo de Biotecnologia, Universidade Federal de Pelotas, Pelotas, RS, Brazil

Full list of author information is available at the end of the article

Published: 1 October 2014

References

1. Mazumder S, Maji M, Das A, Ali N: Potency, efficacy and durability of DNA/DNA, DNA/protein and protein/protein based vaccination using gp63 against *Leishmania donovani* in BALB/c mice. *Plos One* 2011, **6**(2):e14644; doi:10.1371/journal.pone.0014644.
2. Zhang X, Divangahi M, Ngai P, Santosuosso M, Millar J, Zganiacz A, Wang J, Bramson J, Xing Z: Intramuscular immunization with a monogenic plasmid DNA tuberculosis vaccine: Enhanced immunogenicity by electroporation and coexpression of GM-CSF transgene. *Vaccine* 2005, **25**(7):1342-52; doi:10.1016/j.vaccine.2006.09.089.
3. Donnelly JJ, Wahren B, Liu MA: DNA vaccines: progress and challenges. *J Immunol* 2005, **175**(2):633-9; doi: 10.1007/s10620-005-9016-x.
4. Buts JP, De Keyser N: Effects of *Saccharomyces boulardii* on intestinal mucosa. *Digestive Disease and Sciences* 2006, **51**(8):1485-1492; doi: 10.1007/s10620-005-9016-x.
5. Thomas S, Przesdzing I, Metzke D, Schmitz J, Radbruch A, Baumgart DC: *Saccharomyces boulardii* inhibits lipopolysaccharide induced activation of human dendritic cells and T cell proliferation. *Clin Exp Immunol* 2009, **156**(1):78-87; doi: 10.1111/j.1365-2249.2009.03878.

doi:10.1186/1753-6561-8-S4-P160

Cite this article as: Silveira et al.: *Saccharomyces boulardii* ingestion increases the humoral response of a DNA vaccine against leptospirosis in mice. *BMC Proceedings* 2014 **8**(Suppl 4):P160.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

